


SciELO





SciELO



SciELO

MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO

(FORMERLY DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

80

FAMILY DIASTATIDAE

ANGELO PIRES DO PRADO

*Departamento de Parasitologia,
Universidade Estadual de Campinas, São Paulo*

This small family is very poorly known. Only three species of the genus *Diastata* Meigen are recorded from the Neotropical Region.

Nothing is known about their habits or immature stages.

Genus DIASTATA Meigen

DIASTATA Meigen, 1830:94. Type-species, *Geomyza obscurella* Fallén (Westwood, 1840: 152; misidentification) = *vagans* Loew.

ARGENTINA (Malloch), 1934: 439, pl. 8, fig. 4 (*Calopterella*). Type-locality: Argentina, Buenos Aires. Distr. - Argentina.

COSTARICENSIS Hennig, 1956:150, fig. 4. Type-locality: Costa Rica, Farm La Caja, 8 km west San José. Distr. - Costa Rica.

DIMIDIATA (Hendel), 1913:617, fig. 1 (*Trixoscelis*). Type-locality: Peru, Cusco, 3.700 m; Bolivia, Sorata, 2.300 m.

LITERATURE CITED

HENDEL, F.

1913. Neue amerikanische Dipteren. 1. *Deutsch Ent. Zeitschr.* 1913: 617-636, 5 figs.

HENNIG, W.

1956. Neue neotropische Acalyptrata aus dem Deutschen Entomologischen Institut. (Diptera: Acalyptrata). *Beitr. Ent.* 6 (1/2): 146-154, 8 figs.

MALLOCH, J.R.

1934. *Acalyptrata*, pp. 393-489, figs. 69-84, pl. 8. In: British Museum (Natural History), *Diptera of Patagonia and South Chile*, Pt. 6, Fasc. 5, 499 pp., 84 figs., 8 pls., London.

MEIGEN, J.W.

1830. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten*. Vol. 6, iv-401 pp., pls. 56-66, Hamm.

WESTWOOD, J.O.

1840. Order XIII. Diptera Aristotle (Anfliata Fabricius, Halteriptera Clairv.). pp. 145-158 (= signature L). In his *An Introduction to the modern classification of insects. Synopsis of the genera of British Insects*, 158 pp., London.



595.77
S239c
v.3

DEDALUS - Acervo - MZ
Catalogue of the Diptera of the Americas South of the United States.



12400006019



MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO
(FORMERLY DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA
AGRICULTURA, -SÃO PAULO)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS SOUTH OF THE UNITED STATES

81

FAMILY CHLOROPIDAE

CURTIS W. SABROSKY

*Systematic Entomology Laboratory, Agricultural Research Service
U. S. Department of Agriculture, c/o U. S. National Museum of Natural History
Washington, D. C. 20560*

CELUTA H. PAGANELLI*

*Museu de Zoologia
Universidade de São Paulo*

Chloropids, commonly called "grass flies", are small (1,5-5,0 mm) and often found in grass, sedges, other low vegetation, and sometimes flowers. A few species, such as the eye gnats or Hippelates flies, are persistent annoyances of man and other animals, and some of them feed at wounds or body openings.

The food habits of the neotropical species are virtually unknown, and a great deal of biological work remains to be done. Data from other regions indicate that the larvae of many species of Chloropidae are saprophagous, often scavenging from damage by other insects or feeding on decaying leaf sheaths or decaying vegetable matter in the soil. Some well-known species are phytophagous, including such pests of cereals as the frit fly of Europe and North America, *Oscinella frit* (Linnaeus), and the stem maggots, *Meromyza* spp. A few species are gall formers, and a few are reared from fungi. Some larvae are predaceous: e. g., *Thaumatomyia* spp. prey on root aphids, and *Pseudogaurax* spp. prey on the egg masses of spiders, mantids, and some Lepidoptera.

For the Neotropical Region, the family was monographed by Theodor Becker in 1912 and again by O. Duda in 1930. Although Duda described 133 new species and 26 new genera, he missed — or at least did not mention — 75 species and 12 genera already described. Even so, many of his new species in the rich neotropical fauna were undoubtedly valid, but the ranges and possible synonymy of other species remain to be worked out.

* Bolsista da Fundação de Amparo à Pesquisa do Estado de São Paulo.



The growth in recognized neotropical genera and species has been as follows:

Becker, 1912	28 genera, 152 species
Duda, 1930	42 genera, 288 species
Present catalogue	57 genera, 430 species

Unfortunately, Duda (1930) published only the keys of the large monograph, and the provenance of many new species was not stated. It is desirable here to remedy this lack of type localities as far as possible. Duda regularly labeled each specimen of his new taxa as new, but he did not label holotypes and considered every specimen in his type series to be a "typus". In collections, Duda material has thus been labeled "typus" or "cotypus" (i. e., syntype) by various curators. Many of these 'types' of Duda's species were seen by the authors in various European museums (Sabrosky in 1937 and 1938, and Paganelli in 1982) or borrowed for study. The lists of type material of Chloropidae in the Zoologisches Museum in Berlin by Wendt (1975, 1978) added useful information, because Duda's private collection came to that museum after Sabrosky's visits.

If only one type specimen is known, we here name it as the "presumed holotype". If two or more 'type' specimens are known, including those no longer in existence, we record them as "presumed syntypes". However, we regard as without standing any specimens labeled "typus" or "cotypus" that disagree with Duda's published statements of provenance. Apparently a sizeable part of Duda's work was based on material borrowed from the Museum at Hamburg, mainly extensive series collected by Heinrich Schmidt at Farm La Caja, 8 km west of San José, Costa Rica. This collection was destroyed during World War II, but detailed notes were taken by Sabrosky in 1937. Some of that material also went to other museums, through exchange or perhaps from Duda himself, and such surviving syntypes are also recorded here.

Lectotypes in Oscinellinae have been designated by Sabrosky and in Chloropinae by Paganelli, and they are to be credited accordingly. This is a final step in the clarification of those species described by Duda.

The location of Duda's material has also been stated, except for the few cases, from his papers subsequent to 1930, in which provenance and type location were recorded. For brevity, the location has been given by city:

Berlin: Zoologisches Museum der Humboldt-Universität.

Budapest: Magyar Nemzeti Múzeum.

Cambridge, Mass.: Museum of Comparative Zoology, Harvard University.

Dresden: Staatliches Museum für Tierkunde.


Eberswalde: Institut für Pflanzenschutzforschung, Abteilung Taxonomie der Insekten, formerly the Deutsches Entomologisches Institut.

Hamburg: Zoologisches Museum.

Helsinki: private collection of Richard Frey, now in the Museum of Zoology, University of Helsinki.

London: British Museum (Natural History).

Milan: Museo Civico di Storia Naturale, Milano.



São Paulo: Museu de Zoologia, Universidade de São Paulo.

Stuttgart: Staatliches Museum für Naturkunde in Stuttgart, Zweigstelle Ludwigsburg.

Vienna: Naturhistorisches Museum.

Warsaw: Polska Akademia Nauk, Instytut Zoologii.

Numerous new generic combinations are proposed in this catalogue. Three new names are necessary: *Hippelates catharinae*, *Olcella paraguayensis*, and *Chlorops yungas*.

Keys to species by Duda (1930), the most recent monograph, could be cited for most of the genera; however, overlooked species, synonymy, recombinations, and known misidentifications make the keys of limited value. A few that are inclusive or especially useful are cited. Keys in Enderlein (1911a), Curran (1928), and Malloch (1934) may be useful for limited areas.

Refs. — Enderlein, 1911a (revision of Oscinellinae; almost all neotropical material from Santa Catarina, Brazil); Becker, 1912a (monograph, Nearctic and Neotropical Regions); Curran, 1928 (keys, Puerto Rico); Duda, 1930 and supplements (monograph, Neotropical Region); Malloch, 1934 (keys, Patagonia and South Chile); Sabrosky, 1941b (type species).

Subfamily SIPHONELLOPSINAE

Genus APOTROPINA Hendel

Ectropa Schiner, 1868: 242 (preocc. Wallengren, 1863). Type species, *viduata* Schiner, 1868 (mon.).

APOTROPINA Hendel, 1907: 98 (nom. nov. for *Ectropa* Schiner). Type species, *Ectropa viduata* Schiner, 1868 (aut.).

Lasiopleura Becker, 1910: 130. Type species, *Oscinis longepilosa* Strobl, 1893 (mon.).

Pseudohippelates Malloch, 1913c: 261. Type species, *Hippelates capax* Coquillett, 1898 (orig. des.).

Omochaeta Duda, 1930: 58. Type species, *Oscinosoma specularifrons* Enderlein, 1911 (Sabrosky, 1941b: 759).

Ref. — Duda, 1930 (key, Neotropical spp.), Malloch, 1934 (partial key); Sabrosky, 1951 (key, Nearctic spp.), 1980a (name of genus).

CAPAX (Coquillett), 1898: 48 (*Hippelates*). Type locality: U.S.A., Illinois. Distr. — U.S.A. (eastern), Mexico (Tabasco). N. COMB.

CIRCUMDATA (Duda, 1930: 60 (*Omochaeta*)). Type locality: "Paraguay" [San Bernardino, from presumed holotype in Vienna].

FORTIS (Becker), 1916: 451 (*Oscinella*). Type locality: Peru, Callanga. N. COMB.

GRACILIS (Malloch), 1913c: 262 (*Pseudohippelates*). Type locality: Panama, Canal Zone. Distr. — Panama, Peru. N. COMB.

- GRISEA (Malloch), 1934: 417 (*Lasiopleura*). Type locality: U.S.A., Arizona, Bill Williams Fork (lectotype, Sabrosky, 1951: 338). Distr. — U.S.A. (western), Mexico (Puebla, Sinaloa, Sonora, Veracruz), Guatemala. N. COMB.
- HIRTIVENTRIS (Malloch), 1934: 418 (*Lasiopleura*). Type locality: "Panama" [Trinidad Rio, from lectotype by Sabrosky, 1951: 343]. N. COMB.
- INFUMATA (Becker), 1916: 452 (*Oscinella*). Type locality: Colombia, Aracataca. N. COMB.
- LONGIPENNIS (Becker), 1912a: 211 (*Oscinella*). Type locality: Bolivia, Sorata. N. COMB.
- LONGULA (Becker), 1912a: 89 (*Hippelates*). Type locality: Grenada (as "Canada", error, see Aldrich, 1931: 71). Distr. — Honduras, Grenada, Guyana, Brazil (Pará).
- NIGRICORNIS (Duda), 1930: 60 (*Omochaeta*). Type locality: "Peru". N. COMB.
- PALLIATA (Curran), 1926: 8 (*Botanobia*). Type locality: Puerto Rico, Adjuntas. N. COMB.
- PANAMENSIS (Malloch), 1934: 417 (*Lasiopleura*). Type locality: "Panama" [Alhajuelo, from lectotype by Sabrosky, 1951: 343]. N. COMB.
- RUFITHORAX (Duda), 1930: 59 (*Omochaeta*). Type locality: "Brazil" [Rio de Janeiro, Petrópolis, from holotype in São Paulo]. N. COMB.
- SENILIS (Duda), 1930: 60 (*Omochaeta*). Type locality: "Paraguay" [San Bernardino, from presumed syntypes in Vienna]. Distr. — Paraguay, Argentina (Malloch, 1934). N. COMB.
- SPECULARIFRONS (Enderlein), 1911a: 220 (*Oscinosoma*). Type locality: Brazil, Santa Catarina. Distr. — Paraguay, Brazil. N. COMB.
- pubescens* Becker, 1916: 452 (*Oscinella*). Type locality: Paraguay, San Bernardino. N. COMB.

Subfamily OSCINELLINAE

Genus ACANTHOPELTASTES Enderlein

- ACANTHOPELTASTES Enderlein, 1911a: 194. Type species, *spineus* Enderlein, 1911 (orig. des.).
- Dactylothyrea* Meijere of Becker (1912a) and Duda (1930), misident. Ref. — Duda, 1930 (key).
- CIRCULARIS (Becker), 1912a: 175, 176 (*Dactylothyrea*). Type locality: Peru, Callanga. N. COMB.
- CURVINERVIS (Duda), 1930: 75 (*Dactylothyrea*). Type locality: "Costa Rica" [Suiza de Turrialba, from syntypes in Budapest (Duda, 1933a: 43)]. N. COMB.



- HASTATUS (Duda), 1930: 74 (*Dactylothyrea*). Type locality: "Costa Rica" [Suiza de Turrialba, from holotype in Budapest (Duda, 1933a: 42)]. N. COMB.
- HORVATHI (Becker), 1912a: 175 (*Dactylothyrea*). Type locality: Peru, Callanga. N. COMB.
- HYALINIPENNIS Enderlein, 1911a: 195. Type locality: Brazil, Santa Catarina. Distr. — Costa Rica, Brazil.
hyalipennis Becker, 1912a: 122, 177 (*Dactylothyrea*), error or emend.
incolorata Becker, 1912a: 177 (*Dactylothyrea*) (Unnecessary replacement new name; *D. hyalipennis* Meijere does not preoccupy *hyalinipennis*). N. COMB.
- SCHNUSEI (Becker), 1912a: 175 (*Dactylothyrea*). Type locality: Peru, Meshagua, Rio Urubamba. N. COMB.
- SPINEUS Enderlein, 1911a: 194. Type locality: Brazil, Santa Catarina. Distr. — Costa Rica, Brazil (Rio de Janeiro, Santa Catarina).
- STRANDI (Duda), 1930: 75 (*Dactylothyrea*). Type locality: "Costa Rica". [Suiza de Turrialba, from holotype in Budapest (Duda, 1933a: 44)]. N. COMB.

Genus APALLATES Sabrosky

- APALLATES Sabrosky, 1980b: 417. Type species, *Oscinis dissidens* Tucker, 1908 (orig. des.).

Hippelates, part, and *Oscinella*, part, of authors.

Hippelates, *dissidens* group of Sabrosky, 1941a: 25.

There is no comprehensive key, but portions will be covered by *Hippelates* keys of Becker (1912a), Malloch (1913c), Duda (1930), and Sabrosky (1941a). Considerable synonymy may be found when the genus has been studied comprehensively.

- AENEUS (Duda), 1931: 162 (*Oscinella coxendix* var.). Type locality: Not stated [Provenance unknown; presumed syntypes in Vienna are labeled "Otto. 1895 Tambillo"]. N. COMB.

aenea Duda, 1930: 106 (*Oscinella coxendix* var.). Unavailable name (no separate diagnosis).

- ANTIGUANUS (Duda), 1933b: 196 (*Liohippelates convexus* var.). Type locality: Guatemala, Antigua and El Salto Antigua. N. COMB.

- COLUMBIANUS (Enderlein), 1911a: 217 (*Oscinosoma*). Type locality: "Colombia". Distr. — Mexico (Veracruz), Costa Rica, Panama, Colombia. N. COMB.

- CONVEXUS (Loew), 1866a: 184 (Cent. 6, no. 94) (*Hippelates*). Type locality: "Cuba". Distr. — Mexico (Sonora, Tamaulipas, Veracruz, Yucatán), Guatemala, Nicaragua, Panama, Bermuda, Bahamas, Cuba, Jamaica, Haiti, Dominican Republic, Puerto Rico, Virgin Islands, St. Vincent,



- Trinidad, Colombia, Venezuela, Ecuador, Bolivia, Brazil. (Goiás, Rio de Janeiro, Pernambuco), and Argentina (Corrientes, Misiones, Salta).
- convexiformis* Duda, 1936: 108 (*Liohippелates*). Type locality: Brazil, Pernambuco, Recife. N. SYN., N. COMB.
- simulator* Séguy, 1940: 353 (*Hippelates*). Type locality: Cuba, Havana. N. SYN., N. COMB.
- COSTARICANUS (Duda), 1930: 106 (*Oscinella coxendix* var.). Type locality: "Costa Rica" (inferred from the name) [Farm La Caja, near San José, from presumed syntypes in Hamburg, now destroyed, and in Berlin]. Distr. — Guatemala, Honduras, Costa Rica. N. COMB.
- COXENDIX (Fitch), 1856: 533 (1856: 301) (*Oscinis*). Type locality: U.S.A., New York. Distr. — Canada, U.S.A., Mexico (Aguascalientes, Distrito Federal, Durango, Hidalgo, México, Michoacán, Morelos, Nayarit, Zatecas).
- DISSIDENS (Tucker), 1908: 274 (*Oscinis*). Type locality: Texas, Collin County, Plano. Distr. — U.S.A. (eastern), Mexico (Morelos, Puebla).
- texana* Malloch, 1913c: 251 (*Hippelates*). Type locality: U.S.A., Texas, Brownsville.
- DIVERSIPES (Curran), 1926: 7 (*Botanobia*). Type locality: Puerto Rico, Arecibo. N. COMB.
- FORBESI (Curran), 1931: 12 (*Oscinella*). Type locality: Puerto Rico, Vieques Id. Distr. — Cuba, Puerto Rico, Virgin Islands. N. COMB.
- coxendix* (Fitch) of Curran, 1928: 58 (*Botanobia*), misident.
- FRONTALIS (Becker), 1912a: 167 (*Hippelates*). Type locality: Bolivia, Mapiri, Sarampiuni. N. COMB.
- FUR (Williston), 1896: 425 (*Oscinis*). Type locality: St. Vincent. Distr. — Mexico (Colima, Guanajuato, Jalisco, San Luis Potosí, Sinaloa, Veracruz), Guatemala, El Salvador, Nicaragua, Costa Rica, Colombia, St. Vincent. N. COMB.
- strandii* Duda, 1930: 61, 64, 105; 1933a: 46 (*Oscinella*). Type locality: Costa Rica [Farm La Caja, near San José (Duda, 1933a: 47)]. N. COMB.
- HERMSI (Sabrosky), 1941a: 27 (*Hippelates*). Type locality: California, Coachella. Distr. — U.S.A. (California to Texas), Mexico (Jalisco, Nayarit, Sinaloa, Sonora).
- LUTEIPES (Becker), 1912a: 210 (*Oscinella*). Type locality: Peru, mouth of Rio Pachitea. Distr. — Panama, Peru. N. COMB.
- LUTZI (Curran), 1926: 6 (*Botanobia*). Type locality: Puerto Rico, Arecibo. Distr. — Dominican Republic, Puerto Rico. N. COMB.
- MACULICORNIS (Duda), 1936: 111 (*Oscinella*). Type locality: Brazil, Pernambuco, Recife. N. COMB.
- MAGNICORNIS (Duda), 1930: 109 (*Lioscinella*). Type locality: Not stated [Costa Rica, Farm La Caja, near San José, from presumed syntypes in Hamburg, now destroyed, and in Berlin and Eberswalde]. Distr. —



- Mexico (Veracruz), Guatemala, El Salvador, Costa Rica, Colombia, Ecuador. N. COMB.
- MAGNIPALPOIDES (Curran), 1926: 11 (*Botanobia*). Type locality: Puerto Rico, Arecibo and San Juan. N. COMB.
- MICCOTROGUS (Séguy), 1940: 342 (in key), 348 (as *miccotrogus*) (*Hippelates*). Type locality: Costa Rica, La Caja. N. COMB.
- miccotrogus* Séguy, 1940: 348, rejected alternate original spelling.
- MICROCENSTRUS (Coquillett), 1904a: 28 (*Hippelates*). Type locality: California, Stanford University. Distr. — U.S.A. (western), Mexico (Baja California, Jalisco).
- NEOCOXENDIX (Sabrosky), 1940: 215 (*Oscinella*). Type locality: U.S.A., Georgia, Okefenokee Swamp, Billy's Island. Distr. — U.S.A. (eastern), Canada (eastern), Mexico (Durango, Hidalgo, Michoacán, Morelos, Puebla, San Luis Potosí).
- NIGRICOXA (Malloch), 1913c: 250 (*Hippelates*). Type locality: West Indies, Montserrat. N. COMB.
- NIGRIPES (Duda), 1930: 68 (*Liohippelates*). Type locality: Chile, Corral. Distr. — Argentina (Mendoza, Rio Negro), Chile. N. COMB.
- buccalis* Duda, 1930: 109, part (*Lioscinella*). Type locality: Not stated [Chile, Talcahuano, Coronel, and Los Andes, from presumed syntypes in Berlin, Dresden, and Eberswalde; Peru presumed syntypes, Mamara and Cuzco, and probably Callao, Paso de Lares (as "Laristhal"), and Puno, are a different species]. N. COMB.
- particeps* Becker, 1912a: 208, in part (*Oscinella*).
- nigricoxa* of Malloch, 1934: 412, misident. (*Cadrema*).
- NOCENS (Becker), 1912a: 208 (*Oscinella*). Type locality: Bolivia, Mapiri, Sarapiquí, and Brazil, Santa Catarina. N. COMB.
- PAESSLERI (Duda), 1930: 65 (*Liohippelates*). Type locality: "Chile, [error?], Nicaragua" [Corinto, from presumed syntypes in Hamburg, now destroyed, and Berlin]. Distr. — Guatemala, Nicaragua.
- PARTICEPS (Becker), 1912a: 115 (*Oscinella*). Type locality: U.S.A., Pennsylvania. (lectotype, Sabrosky, 1936: 246). Distr. — U.S.A. (widespread), Mexico (Durango, Guerrero, Morelos, Oaxaca, Puebla, Veracruz).
- PARVULUS (Malloch), 1934: 436 (*Oscinosoma*). Type locality: Argentina, Buenos Aires. N. COMB.
- PRUINOSUS (Duda), 1930: 65 (*Liohippelates*). Type locality: "Brazil" [Rio de Janeiro, Petrópolis, lectotype ♂, here designated, in São Paulo]. Distr. — Ecuador, Brazil (Pará, Rio de Janeiro, Santa Catarina), Paraguay, Argentina (Misiones, Salta). N. COMB.
- punctulatus* (Duda), 1934: 104 (*Oscinella coxendix* var. *costaricana* subvar.). Type locality: Panama, Sabanas [published in error as "Subana"]. Unavailable name (4th term of quadrimomial).
- SICATRIX (Curran), 1926: 8 (*Botanobia*). Type locality: Puerto Rico, Mona Island. Distr. — Puerto Rico, Virgin Islands. N. COMB.

SUBTOMENTOSUS (Enderlein), 1911a: 228 (*Tropidoscinis*). Type locality: Brazil, Santa Catarina. Distr. — Colombia, Ecuador, Peru, Brazil (Rio de Janeiro, Santa Catarina). N. COMB.

TENER (Coquillett), 1900: 265 (*Hippelates*). Type locality: Puerto Rico, Utuado and Fajardo. Distr. — U.S.A. (Texas and Florida), Mexico (Tamaulipas, Veracruz, Yucatán), Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela.

TOMENTOSUS (Enderlein), 1911a: 228 (*Tropidoscinis*). Type locality: Brazil, Santa Catarina. N. COMB.

TRIANGULINUS (Duda), 1930: 109 (*Lioscinella*). Type locality: Not stated [Costa Rica, Suiza de Turrialba, from presumed holotype in Budapest]. N. COMB.

VARIPALPUS (Curran), 1926: 12 (*Botanobia*). Type locality: Puerto Rico, Mona Id. N. COMB.

VIRIDINIGER (Enderlein), 1911a: 192 (*Hippelates*). Type locality: Brazil, Santa Catarina. Distr. — El Salvador, Costa Rica, Colombia, Ecuador, Peru, Bolivia, Brazil (Rio de Janeiro, Santa Catarina). N. COMB.

Genus BIORBITELLA Sabrosky

BIORBITELLA Sabrosky, 1980b: 420. Type species, *Oscinella hesperia* Sabrosky, 1940 (orig. des.).

CONFUSA (Malloch), 1913a: 61 (*Botanobia*). Type locality: Puerto Rico, Bayamon. N. COMB.

virgata Coquillett of Coquillett, 1900: 266 (*Oscinis*).

COSTARICANA (Duda), 1930: 101 (*Conioscinella rubicunda* var.). Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed type series in Hamburg, now destroyed]. Distr. — Mexico (San Luis Potosí, Veracruz), Guatemala, El Salvador, Costa Rica, Panama. N. COMB.

FRONTOORBITALIS (Sabrosky), 1940: 221 (*Oscinella*). Type locality: U.S.A., Utah, St. George. Distr. — U.S.A. (southwestern), Mexico (Sonora).

HESPERIA (Sabrosky), 1940: 219 (*Oscinella*). Type locality: U.S.A., Utah, Cedar City. Distr. — U.S.A. (western), Mexico (Aguascalientes, Baja California, Distrito Federal, México, Morelos, San Luis Potosí, Veracruz).

NIGRONOTATA (Duda), 1930: 111 (*Lioscinella*). Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed syntypes in Hamburg, now destroyed, and syntypes in Berlin and a syntype in Eberswalde]. Distr. — Guatemala, Costa Rica, Colombia, Peru. N. COMB.

ORBITALIS (Duda), 1930: 98 (*Neoolcanabates*). Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed type series in Hamburg, now destroyed, and syntypes in Berlin and Eberswalde]. Distr. — Mexico (Hidalgo), Guatemala, Costa Rica, Colombia.

RUBICUNDA (Becker), 1912a: 213 (*Oscinella*). Type locality: Bolivia, Lorenzo-pata and Sarampiuní. Distr. — Colombia, Venezuela, Ecuador, Peru, Bolivia, Brazil (Rio de Janeiro). N. COMB.

Genus CADREMA Walker

- CADREMA Walker, 1859: 117. Type species, *lonchopteroides* Walker, 1859 (mon.).
Prohippелates Malloch, 1913c: 260. Type species, *Hippelates pallidus* Loew, 1866 (orig. des.).
Palaeogaurax Duda, 1930: 57. Type species, *Hippelates pallidus* Loew, 1866 (Sabrosky, 1941b: 761).
PALLIDA (Loew), 1866a: 184 (Cent. 6, no. 93) (*Hippelates*). Type locality: "Cuba". Distr. — U.S.A. (s. Florida), Bermuda, Bahamas, Cuba, Jamaica, Puerto Rico, Antigua, St. Vincent, Barbados, Panama, Colombia, Clipperton Island.

Genus CHAETHIPPUS Duda

- CHAETHIPPUS Duda, 1930: 58. Type species, *Hippelates mellinus* Becker, 1912 (sub. des. and sub. mon., Duda, 1931: 165).
MELLINUS (Becker), 1912a: 168 (*Hippelates*). Type locality: Peru, Meshagua and Umahuanquiali, both on Rio Urubamba.

Genus CHAETOCHLOROPS Malloch

- CHAETOCHLOROPS Malloch, 1914: 120. Type species, *Siphonella inquilina* Coquillett (orig. des.).
Ref. — Sabrosky, 1950c (key, biol.).
INQUILINUS (Coquillett), 1898: 48 (*Siphonella*). Type locality: U.S.A., Missouri and Virginia. Distr. — U.S.A. (eastern), Mexico (Michoacán).
SCUTELLARIS (Becker), 1916: 449 (*Siphonella*). Type locality: Costa Rica, Higuito, San Mateo. Distr. — Guatemala, El Salvador, Costa Rica, Panama, Colombia, Venezuela.

Genus CONIOSCINELLA Duda

- Oscinella*, subg. CONIOSCINELLA Duda, 1929: 166. Type species, *Oscinella soluta* Becker, 1912 (Sabrosky, 1941b: 751).
Ref.: Duda, 1930 (key).
AEQUISECTA (Enderlein), 1911a: 210 (*Tricimba*). Type locality: Brazil, Santa Catarina.
beckeri Enderlein, 1913: 364 (*Coryphisoptron*). Type locality: Brazil, Santa Catarina. N. SYN., N. COMB.
ANGULICORNIS Duda, 1936: 109. Type locality: Brazil, Pernambuco, Recife.
ANNULIFEMUR Duda, 1930: 102. Type locality: "Brazil" [Rio de Janeiro, Petrópolis, from presumed holotype in São Paulo]. Distr. — Brazil (Rio de Janeiro, Santa Catarina).
arcuata Duda, 1934: 59. Nomen nudum. Mentioned in a comparative note under *C. cricopus* (Enderlein), but never published, or published under another name but cited here as *arcuata*, through oversight.

- ARGENTINICA (Duda), 1929: 167 (*Oscinella*, subg. *Conioscinella*, as *quadrivittata* var.). Type locality: Argentina (northern), San José.
- CHILENICA Duda, 1930: 104 (as *perlineata* var.). Type locality: "Chile" [Callas, from presumed holotype in Berlin].
- CONFLUENS (Becker), 1919: 200 (*Oscinella*). Type locality: Ecuador, Alausi. N. COMB.
- CRICOPUS (Enderlein), 1911a: 210 (*Tricimba*). Type locality: Brazil, Santa Catarina.
- DIMIDIATA (Becker), 1912a: 206 (*Oscinella*). Type locality: Bolivia, Mapiri, Bellavista.
- FLAVISCUTELLATA (Enderlein), 1911a: 226 (*Gaurax*). Type locality: Brazil, Santa Catarina.
- FLAVOTRIGONA Duda, 1934: 60. Type locality: Panama, Corazal (Canal Zone) and Trinidad Rio; Costa Rica, Higuito, San Mateo.
- formosa* (Becker). Not Neotropical.
- frontella* (Fallén). Not Neotropical.
- GALAPAGENSIS (Curran), 1932: 357 (*Oscinella*). Type locality: Ecuador, Galapagos Is., Floreana, Post Office Bay. N. COMB.
- IMPROVISA (Becker), 1912a: 207 (*Oscinella*). Type locality: Bolivia, Mapiri, Sarampiuni.
- INCIPIENS (Williston), 1896: 424 (*Oscinis*). Type locality: West Indies, St. Vincent.
- INCONSPICUA (Becker), 1916: 452 (*Oscinella*). Type locality: Colombia, Aracataca.
- INCONSTANS (Becker), 1912a: 215 (*Oscinella*). Type locality: Peru, Cuzco; Ecuador, La Rinconada. Distr. — Ecuador, Peru, Chile, Argentina (Jujuy, La Rioja, Mendoza, Rio Negro, Tucumán).
- IOTA (Malloch), 1934: 435 (*Oscinosoma*). Type locality: Argentina, Rio Negro, Lake Gutiérrez near Bariloche. N. COMB.
- LACTEIHALTERATA (Enderlein), 1911a: 221 (*Oscinosoma*). Type locality: Brazil, Santa Catarina.
- MARS (Curran), 1926: 10 (*Botanobia*). Type locality: Puerto Rico, Naguabo. Distr. — Puerto Rico, Virgin Is., Guyana.
- NIGROHALTERATA (Duda), 1930: 97, 108 (*Lioscinella*). Type locality: Not stated [Costa Rica, Farm La Caja, near San José, from presumed holotype in Hamburg, now destroyed]. N. COMB.
- NUDA (Adams), 1905a: 110 (*Oscinis*). Type locality: U.S.A., Georgia, Tifton. Distr. — Arizona to southeastern U.S.A., Mexico (Santa Engracia River).
- PERLINEATA Duda, 1930: 103. Type locality: "Brazil" [Rio de Janeiro, Petrópolis, from presumed holotype in São Paulo]. Distr. — Brazil (Rio de



Janeiro, Santa Catarina), Argentina (Chaco, Corrientes, Entre Ríos, Jujuy, Misiones, Salta, Tucumán).

PSEUDOGAURACICA Duda, 1930: 94. Type locality: Not stated [Costa Rica, Suiza de Turrialba, from presumed holotype in Budapest].

PULVERULENTA (Duda), 1930: 98 (*Neoolcanabates*). Type locality: "Costa Rica" [Suiza de Turrialba, from presumed holotype in Budapest]. N. comb.

QUADRIVITTATA (Duda), 1929: 167 (*Oscinella*, subg. *Conioscinella*). Type locality: Not stated [Paraguay, from presumed holotype in Budapest, recorded as *quadrilineata* by Becker, 1912a].

quadrilineata Williston of Becker, 1912a: 212 (*Oscinella*), misident.

RUFITHORAX Duda, 1930: 100. Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed syntypes in Hamburg, now destroyed, and one in Berlin]. Distr. — Costa Rica, Panama.

SOLUTA (Becker), 1912a: 207 (*Oscinella*). Type locality: Bolivia, Mapiri, Sarampiuni. The status of Duda's varieties is not clear, and they are merely recorded here.

var. NIGRIPES Duda, 1930: 99. Type locality: Not stated [Costa Rica, Farm La Caja, near San José, from presumed syntypes in Hamburg, now destroyed, and Dresden]. Distr. — Costa Rica, Panama.

var. RUFOSCUTELLATA Duda, 1930: 95 (as *rufoscutulata*), 102. Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed syntypes in Hamburg, now destroyed, and Dresden].

var. SEMICIRCULARIS Duda, 1930: 94. Type locality: "Brazil". Distr. — Brazil (Pernambuco).

var. UNISERIATA (Duda), 1929: 167 (*Oscinella*, subg. *Conioscinella*). Type locality: Bolivia, Samuhuate.

SUBLINEATA Duda, 1930: 103. Type locality: "Brazil" [Rio de Janeiro, Petrópolis, from presumed holotype in São Paulo]. Distr. — Brazil (Rio de Janeiro, Santa Catarina).

TRIPUNCTATA (Curran), 1926: 10 (*Botanobia*). Type locality: Puerto Rico, Mona Id. Distr. — Jamaica, Puerto Rico.

TURBIDA (Becker), 1912a: 206 (*Oscinella*). Type locality: Peru, Cuzco; Bolivia, Camino de Yungas; Ecuador, Troya.

Genus CORYPHISOPTON Enderlein

CORYPHISOPTON Enderlein, 1911a: 204. Type species, *flavipes* Enderlein, 1911 (orig. des.).

ALBOHIRTUM Enderlein, 1913: 363. Type locality: Brazil, Santa Catarina.

COSTARICANUM (Duda), 1930: 88 (*Discogastrella*). Type locality: "Costa Rica" (inferred from specific name) [Suiza de Turrialba, from presumed holotype in Budapest]. N. comb.

DISCIVENTRE (Enderlein), 1911a: 202 (*Discogastrella*). Type locality: Brazil, Santa Catarina.

- DUBIUM (Duda), 1930: 89 (*Discogastrella*). Type locality: Not stated [Peru, Rio Urubamba, from presumed holotype in Dresden]. N. COMB.
- FLAVICEPS (Enderlein), 1911a: 201 (*Discogastrella*). Type locality: Brazil, Santa Catarina.
- flavifrons* (Duda), 1930: 88 (*Discogastrella*). Type locality: Not stated [Costa Rica, Suiza de Turrialba, from presumed holotype in Budapest]. N. COMB. Unavailable name (no separate diagnosis).
- FLAVIPES Enderlein, 1911a: 205. Type locality: Brazil, Santa Catarina.
- FLAVITARSE (Enderlein), 1911a: 203 (*Discogastrella*). Type locality: Brazil, Santa Catarina.
- FLAVITIBIA (Duda), 1930: 76 (*Gauracisoma*). Type locality: "Peru" [Rio Urubamba, from presumed holotype in Dresden]. N. COMB.
- FLAVIVENTRE (Malloch), 1934: 434 (*Oscinosoma*). Type locality: Argentina, Buenos Aires. N. COMB.
- RETRACTUM (Duda), 1930: 89 (*Discogastrella*). Type locality: Not stated [Paraguay, San Bernardino, from presumed syntypes in Vienna]. N. COMB.
- RUFOSCUTELLATUM (Duda), 1934: 58 (*Discogastrella retracta* var.). Type locality: Costa Rica, Higuato, San Mateo. N. COMB.
- SALEBROSUM (Enderlein), 1911a: 202 (*Discogastrella*). Type locality: Brazil, Santa Catarina.
- TRIGONALIFRONS (Enderlein), 1911a: 204 (*Discogastrella*). Type locality: Brazil, Santa Catarina.

Genus DISCOGASTRELLA Enderlein

- DISCOGASTRELLA Enderlein, 1911a: 200. Type species, *gruenbergi* Enderlein, 1911 (orig. des.).
- GRUENBERGI Enderlein, 1911a: 201. Type locality: Brazil, Santa Catarina. Distr. — Guatemala, Costa Rica, Panama, Trinidad, Colombia, Ecuador, Peru, Bolivia, Brazil (Distrito Federal, Minas Gerais, Paraná, Rio de Janeiro, Santa Catarina, São Paulo), Argentina (Chaco, Misiones, Salta, Tucumán).
- numerata* Becker, 1912a: 215, part (*Oscinella*). Type locality: "Brazil"; Argentina, Tucumán; Bolivia, Mapi, Sorata and Lorenzopata; also Ecuador, Pinu, but that syntype is a *Fiebrigella*.

Genus ELACHIPTERA Macquart

- ELACHIPTERA Macquart, 1835: 621. Type species, *Chorops brevipennis* Meigen, 1830 (orig. des.).
- Melanochaeta*, authors, in part.
- Gampsocera* Schiner of Becker (1912a, 1916) and Enderlein (1911).
Ref. — Sabrosky, 1938 (key, red spp.).

ATTENUATA (Adams), 1908: 152 (*Crassiseta*). Type locality: Costa Rica, San José. Distr. — Guatemala, Nicaragua, Costa Rica, Colombia, Cuba, Dominican Republic, Easter Id.

pilosa Duda, 1930: 81. Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed syntypes in Hamburg, now destroyed, and Berlin, and one in Eberswalde].

CONIOTRIGONA Duda, 1933b: 204. Type locality: Guatemala, El Salto, Antigua.

COSTATA (Loew), 1863: 33 (Cent. 3, no. 62) (*Crassiseta*). Type locality: U.S.A., District of Columbia. Distr. — Canada and U.S.A. (widespread), Mexico (Aguascalientes, Distrito Federal).

DUBIOSA (Becker), 1916: 448 (*Gampsocera*). Type locality: Colombia, Aracataca. Distr. — Colombia, Brazil (Rio de Janeiro).

FLAVIDA Williston, 1896: 417. Type locality: St. Vincent. Distr. — Colombia, St. Vincent.

mitis Williston, 1896: 424 (*Oscinis*). Type locality: St. Vincent, Windward side.

MELAENA (Becker), 1912a: 181 (*Melanochaeta*). Type locality: Peru, Cuzco and Oroya; Bolivia, Sorata.

POLLINOSA Sabrosky, 1938: 421. Type locality: Paraguay, Villarica. Distr. — Guatemala, Paraguay.

RUBIDA Becker, 1912a: 179. Type locality: Peru (as "Chile"), Tacna.

RUBROLIMBATA Duda, 1930: 82. Type locality: Not stated [Paraguay, San Bernardino, from presumed holotype in Vienna].

RUFICOLLIS (Frey), 1918: 23 (*Melanochaeta*). Type locality: Brazil, Rio de Janeiro.

SACCULICORNIS (Enderlein), 1911a: 237 (*Gampsocera*). Type locality: Brazil, Santa Catarina. Distr. — Peru, Bolivia, Brazil (Distrito Federal, Espírito Santo, Minas Gerais, Paraná, Rio de Janeiro, Santa Catarina, São Paulo), Paraguay, Uruguay, Argentina (Buenos Aires, Chaco, Córdoba, Corrientes, Formosa, Jujuy, La Rioja, Mendoza, Misiones, Salta, Tucumán), Chile; also St. Helena.


festina Becker, 1912a: 182 (*Melanochaeta*). Type locality: Bolivia, La Paz and Sorata; Chile, Quillota; Paraguay, Puerto Max; Argentina, Tucumán; Brazil.

SUBLINEATA (Becker), 1912a: 181 (*Melanochaeta*). Type locality: Paraguay, San Bernardino. Distr. — Paraguay, Brazil (Rio de Janeiro).

Genus ENDERLEINIELLA Becker

ENDERLEINIELLA Becker, 1912a: 192. Type species, *Tricimba longiventris* Enderlein, 1911 (orig. des.).

Anoscinella Becker, 1916: 448. Type species, *Anoscinella tripunctata* Becker, 1916 (orig. des.).



LONGIVENTRIS (Enderlein), 1911a: 207 (*Tricimba*). Type locality: "Costa Rica."
Distr. — Mexico (Colima, San Luis Potosí), Costa Rica.

TRIPUNCTATA (Becker), 1916: 448 (*Anoscinella*). Type locality: Costa Rica, Higuito, San Mateo. Distr. — Mexico (Jalisco, Veracruz), Costa Rica, Panama, Ecuador, Peru.

Genus ERIBOLUS Becker

ERIBOLUS Becker, 1910: 127. Type species, *sudeticus* Becker, 1910 (Enderlein, 1911: 206).

CALIFORNICUS Sabrosky, 1950b: 91. Type locality: U.S.A., California, San Diego.
Distr. — U.S.A. (California), Mexico (Estado México).

Genus EUGAURAX Malloch

EUGAURAX Malloch, 1913d: 46. Type species, *floridensis* Malloch, 1913 (orig. des.). Ref. — Sabrosky, 1950c (key).

BICOLOR Malloch, 1934: 424. Type locality: Argentina, Buenos Aires.

BIPUNCTATUS Malloch, 1934: 423. Type locality: Uruguay, Montevideo.

FASCIVENTRIS (Duda), 1930: 102 (*Conioscinella formosa* var.). Type locality: "Peru" [Callao, from presumed holotype in Hamburg (Sabrosky, 1950c: 188), now destroyed].

HYSTRIX (Duda), 1930: 99 (*Conioscinella*). Type locality: "Brazil" [Rio de Janeiro, Petrópolis, from presumed holotype (Sabrosky, 1950c: 188), in São Paulo].

INSULARIS Malloch, 1913a: 64. Type locality: Puerto Rico, Fajardo.

quadrilineata Williston of Coquillett, 1900: 266 (*Oscinis*).

INTERMEDIUS Sabrosky, 1950c: 187. Type locality: Costa Rica, Higuito, San Mateo.

PLEURALIS Malloch, 1934: 422. Type locality: Chile, Isla Chiloé, Castro.

QUADRILINEATUS (Williston), 1896: 422 (*Oscinis*). Type locality: St. Vincent.
Distr. — St. Vincent, Trinidad.

SETIGENA Sabrosky, 1974: 347. Type locality: Guyana, Georgetown. Distr. — Colombia, Guyana.

Genus FIEBRIGELLA Duda

FIEBRIGELLA Duda, 1921: 123, 143. Type species, *verrucosa* Duda, 1921 (orig. des., as "n. gen. n. sp.").

Goniopsis Duda, 1929: 166 (Preoccupied Haan, 1835, and Melichar, 1899).
Type species, *Goniopsis verrucosa* Duda, 1929 (mon.) = *boliviensis* Sabrosky.

Goniopsita Duda, 1930: 69, 72. Type species, *Goniopsis verrucosa* Duda, 1929
Sabrosky, 1941b: 753) = *boliviensis* Sabrosky.

- Lacometopa* Duda, 1930: 69, 72. Type species, *Siphonella breviventris* Becker, 1916 (mon.).
- Mimoscinis* Malloch, 1934: 432 (as *Oscinosoma* subgenus). Type species, *Oscinosoma maculiventris* Malloch (mon.). N. SYN.
- Teleocoma* Aldrich, 1924: 1. Type species, *crassipes* Aldrich, 1924 (orig. des.). N. SYN.
Ref. — Sabrosky, 1970: 182 (synonymy and revisionary note).
- ALBIFACIES (Duda), 1930: 73 (*Goniopsita*). Type locality: "Paraguay" [San Bernardino, from presumed holotype in Vienna].
- BOLIVIENSIS Sabrosky, 1970: 182 (nom. nov. for *Goniopsis verrucosa* Duda, 1929). Type locality: Bolivia, San José de Chiquitos.
verrucosa Duda, 1929: 166 (*Goniopsis*); preocc. in *Fiebrigella* by Duda, 1921). Type locality: Bolivia, San José de Chiquitos.
- BREVIVENTRIS (Becker), 1916: 449 (*Siphonella*). Type locality: Colombia, Baranquilla. N. COMB.
- DELICATA (Becker), 1912a: 209 (*Oscinella*). Type locality: Paraguay. N. COMB.
- MACULIVENTRIS (Malloch), 1934: 432 (*Oscinosoma*). Type locality: Argentina, Bariloche. N. COMB.
- VERRUCOSA Duda, 1921: 143. Type locality: Paraguay, San Bernardino. Distr. — Ecuador, Paraguay, Brazil.
crassipes Aldrich, 1924: 1 (*Teleocoma*). Type locality: Brazil, Rio de Janeiro. N. SYN., N. COMB.
- VIRIDIFRONS (Duda), 1930: 73 (*Goniopsita*). Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed syntype in Hamburg, now destroyed, and one in Berlin].

Genus GAURAX Loew

- GAURAX Loew, 1863: 35 (Cent. 3, no. 66). Type species, *festivus* Loew, 1863 (mon.).
- Botanobia* Lioy, 1864: 1125. Type species, *Oscinis dubia* Macquart, 1835 (mon.).
- Pseudochlorops* Duda, 1930: 106. Type species, *Pseudochlorops costaricana* Duda, 1930 (mon.). Preoccupied Malloch, 1914. N. SYN.
- Pseudoscinnella* Duda, 1931: 160 (nom. nov. for *Pseudochlorops* Duda, 1930). Type species, *Pseudochlorops costaricana* Duda (aut.). N. SYN.
- COSTARICANUS (Duda), 1930: 106 (*Pseudochlorops*). Type locality: "Costa Rica" (inferred from specific name) [Farm La Caja, near San José, from presumed syntypes in Hamburg, now destroyed, and Suiza de Turrialba, from presumed syntypes in Budapest]. Distr. — Guatemala, Costa Rica. N. COMB.
- FLAVITHORAX (Duda), 1930: 101, 1933a: 47 (description) (*Oscinella*). Type locality: "Costa Rica" [Farm La Caja, near San José, from holotype (Duda, 1933a: 47) in Hamburg, now destroyed] ?*Gaurax*, as n. comb.



PALLIDICORNIS (Duda), 1930: 112 (*Lioscinella*). Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed holotype in Hamburg, now destroyed]. N. COMB.

TINCTIPES (Becker), 1912a: 214 (*Oscinella*). Type locality: "Paraguay". N. COMB.

Genus GONIASPIS Duda

GONIASPIS Duda, 1930: 59. Type species, *Cadrema rubra* Becker, 1916 (Duda, 1931: 165).

Palaeoenderleiniella Duda, 1930: 57. Type species, *Cadrema rubra* Becker, 1916 (Sabrosky, 1941b: 761).

EQUALIS (Williston), 1896: 419 (*Hippelates*). Type locality: St. Vincent [Windward side, from holotype in London].

OBSCURATA Duda, 1930: 63. Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed syntypes in Hamburg, now destroyed, and in Berlin].

PERUANA Duda, 1930: 64. Type locality: Peru [Callao, from presumed holotype in Hamburg, now destroyed].

RUBRA (Becker), 1916: 451 (*Cadrema*). Type locality: Colombia, Sierra San Lorenzo. Distr. — Colombia, Ecuador.

SCUTELLARIS (Williston), 1896: 420 (*Hippelates*). Type locality: St. Vincent. Distr. — Puerto Rico, Peru.

SUBEQUALIS (Malloch), 1913c: 257 (*Hippelates*). Type locality: West Indies, Montserrat. Distr. — Honduras, Costa Rica, Panama, Colombia, Venezuela, Montserrat, Trinidad, Brazil (Rio de Janeiro).

TRUNCATA (Malloch), 1913c: 247 (*Hippelates*). Type locality: Guatemala, Cacao, Trece Aguas, Alto V. Paz. Distr. — Mexico (Durango, Veracruz), Guatemala, El Salvador, Honduras, Costa Rica.

Genus HIPPELATES Loew

HIPPELATES Loew, 1863: 35, 36 (Cent. 3, no. 67, 68). Type species, *plebejus* Loew, 1863 (Coquillett, 1910: 552).

Hippelates, subgenus *Siphomyia* Williston, 1896: 418. Type species, *proboscideus* Williston, 1896 (Coquillett, 1910: 606).


Palaeoconioscinella Duda, 1930: 57. Type species, *Hippelates proboscideus* Williston, 1896 (Sabrosky, 1941b: 24).

Hippelates, *plebejus* group of Sabrosky, 1941a: 24.
Refs. — Duda, 1930 (key, as *Siphomyia*); Sabrosky, 1941a (key, Nearctic species).

ANGULICORNIS (Malloch), 1934: 414 (*Cadrema*). Type locality: Argentina, Mendoza. Distr. — Uruguay, Argentina (Formosa, Jujuy, Mendoza, Salta, Tucumán).



- ANNOTATUS Becker, 1912a: 169. Type locality: Peru, mouth of Rio Pachitea.
- ANNULATUS Enderlein, 1911a: 192. Type locality: Brazil, Santa Catarina. Distr. — Guatemala, Costa Rica, Panama, Colombia, Ecuador, Brazil (Rio de Janeiro, Santa Catarina), Uruguay, Argentina (Buenos Aires, Corrientes, Jujuy), Chile.
- flavus* Séguy, 1940: 346 (preocc. Thomson, 1869). Type locality: Ecuador, Quito. N. SYN.
- BRACATUS (Duda), 1930: 62 (*Siphomyia*). Type locality: "Costa Rica" [Finca La Caja, near San José, from presumed syntypes in Hamburg, now destroyed, and in Berlin]. Distr. — Mexico (Veracruz), Guatemala, Costa Rica, Ecuador.
- benoisti* Séguy, 1940: 343 (*Hippelates*). Type locality: Ecuador, Santo Domingo de los Colorados. N. SYN.
- BRUMPTI Séguy, 1940: 344. Type locality: Venezuela, Valera. Distr. — Cuba, Dominican Republic, Puerto Rico, Virgin Islands, Bonaire, Venezuela.
- CATHARINAE Sabrosky & Paganelli, NOM. NOV.
- brasiliensis* Duda, 1933b: 195 (as *Hippelates braccatus* var.; primary homonym of *H. brasiliensis* Aldrich, 1931). Type locality: Brazil, Santa Catarina.
- DORSALIS Loew, 1869a: 42 (Cent. 8, no. 75). Type locality: "Cuba". Distr. — U.S.A. (California to Texas, Florida), Mexico (Baja California, Morelos, Nuevo León, Quintana Roo, Sonora, Tamaulipas, Veracruz), Bermuda, Bahamas, Cuba, Jamaica, Puerto Rico, Virgin Islands, Barbados, St. Vincent.
- dorsatus* Williston, 1896: 419. Type locality: St. Vincent.
- GENALIS Thomson, 1869: 608. Type locality: "California" [Possibly error for Ecuador].
- GRANDIUSCULUS Becker, 1912a: 165. Type locality: Peru, Unini, Rio Ucayali. Distr. — Ecuador (Becker, 1919), Peru, Chile (Malloch, 1934).
- ignobilis* Becker, 1912a: 166. Unrecognized, if available at all. Possibly lapsus for *nobilis* Loew.
- IMPRESSUS Becker, 1912a: 92. Type locality: U.S.A., Texas, Austin and Sta. Gertrude. Distr. — U.S.A. (southwestern), Mexico (Aguascalientes, Baja California, Chihuahua, Durango, Jalisco, México, Nuevo León, Sonora).
- INCIPIENS Curran, 1926: 5. Type locality: Puerto Rico, Naguabo and Coamo Springs. ? = *dorsalis* Loew.
- LINEARIS (Malloch), 1934: 415 (*Cadrema*). Type locality: Argentina, Mendoza. N. COMB.
- NABIS Séguy, 1940: 349. Type locality: Cuba, Havana. ? = *dorsalis* Loew.
- NIGRICOLLIS Becker, 1912a: 167. Type locality: Paraguay, Asunción.



- PARVISETA (Malloch), 1934: 413 (*Cadrema*). Type locality: Brazil, "Southeast" [Rio de Janeiro, Alto do Itatiaia, Serra de Itatiaia, from paratype in London].
- PEHLKEI Enderlein, 1911a: 191. Type locality: "Colombia" [Departamento Caldas, Hacienda Pehlke or Hacienda Hamburgo, near Victoria].
- PLEBEJUS Loew, 1863: 36 (Cent. 3, no. 68). Type locality: USA., District of Columbia. Distr. — Canada (southeast), U.S.A. (eastern), Mexico (Nuevo León).
- PLEURITICUS (Wiedemann), 1830: 596 (*Chlorops*). Type locality: "South America".
- PROBOSCIDEUS Williston, 1896: 418. Type locality: St. Vincent. Distr. — U.S.A. (Texas, Georgia, Florida), Mexico (Tamaulipas), Guatemala, Costa Rica, Panama, Bahamas, Cuba, Jamaica, Puerto Rico, St. Vincent, Trinidad, Guyana, Brazil (Bahia, Ceará).
- SAUNDERSI Kumm, 1936: 326. Type locality: Jamaica, St. Ann Parish, Alva. Distr. — U.S.A. (southeast), Jamaica.
- SELECTUS Becker, 1912a: 165. Type locality: Peru, Cuzco, and Callanga. Distr. — Colombia, Ecuador, Peru.
- exlineatus* Becker, 1912a: 164. Type locality: Peru, Cuzco. N. SYN.
- punctatus* Becker, 1912a: 171. Type locality: Ecuador, Tulcán. N. SYN.
- STIGMATICUS (Duda), 1930: 61 (*Siphomyia*). Type locality: "Chile" [Rancagua, Contulmo, and Talcahuano, from presumed syntypes in Berlin and Dresden]. Secondary homonym of *Hippelates stigmaticus* Lamb, 1912, but Lamb's species is not a *Hippelates* and Duda's has not been renamed, hence *stigmaticus* (Duda) can be maintained (International Code of Zoological Nomenclature, Article 59b(ii)). Distr. — Bolivia, Chile, Argentina (Buenos Aires, Córdoba, La Rioja, Mendoza, Rio Negro, Santa Fe).
- subannulatus* Malloch, 1934: 405 (*Chaethippus*). Type locality: Argentina, Rio Negro, eastern end of Lake Nahuel Huapi.
- TRIMACULATUS (Duda), 1930: 96 (*Conioscinella*). Type locality: Not stated [Bolivia, La Paz; Peru, Juliaca; Chile, Antofagasta, all from presumed syntypes in Dresden, Vienna, London, Berlin, and Eberswalde].
- TURNERI Kumm, 1936: 325. Type locality: Jamaica, Trelawny Parish, Good Hope. Distr. — Cuba, Jamaica.

Genus INCERTELLA Sabrosky

- INCERTELLA Sabrosky, 1980b: 420. Type species, *Oscinella incerta* Becker, 1912 (orig. des.).
- dorsata* (Loew). Not Neotropical.
- MINOR (Adams), 1905a: 110 (*Oscinis*). Type locality: U.S.A., Louisiana, Opelousas. Distr. — U.S.A. and Canada (widespread, east of Rocky Mts.), Mexico (Nuevo León).

MONA (Curran), 1926: 9 (*Botanobia*). Type locality: Puerto Rico, Mona Id. N. COMB.

RUFIPLEURA (Duda), 1934: 63 (*Conioscinella*). Type locality: Costa Rica, Higuito, San Mateo. Distr. — Costa Rica, Guyana. N. COMB.

Genus LIOHIPPELATES Duda

Hippelates, subg. LIOHIPPELATES Duda, 1929: 165, 169. Type species, *Hippelates pusio* Loew, 1872 (Sabrosky, 1941b: 755).

Stenoprosopon Duda, 1930: 59, 68. Type species, *Oscinella satanas* Becker, 1912 (mon.).

Stratiomicroneurum Duda, 1933b: 198, 199. Type species, *Stratiomicroneurum microneurinum* Duda, 1933 (orig. des.).

Neohippelates Roberts, 1934: 252, *nomen nudum*.

Hippelates, *pusio* group of Sabrosky, 1941a: 23.

Ref. — Duda, 1930 (key); Sabrosky, 1941a (key, nearctic spp.).

AENEIVENTRIS (Becker), 1912a: 185 (*Siphonella*). Type locality: Ecuador, Riobamba. N. COMB.

AEQUATORIALIS (Becker), 1912a: 171 (*Hippelates*). Type locality: Ecuador, El Angel, and Peru, Callanga. Distr. — Colombia, Ecuador, Peru, Brazil (Santa Catarina), Argentina (Jujuy, La Pampa, La Rioja, Rio Negro, San Luis, Tucumán). N. COMB.

APICATUS (Malloch), 1913c: 248 (*Hippelates*). Type locality: Puerto Rico, Mayaguez. Distr. — U.S.A. (Florida), Bahamas, Cuba, Jamaica, Puerto Rico, Virgin Is., Mexico (Guerrero, Tamaulipas), El Salvador, Nicaragua, Costa Rica, Panama, Venezuela, Trinidad.

illicis Curran, 1926: 4 (*Hippelates*). Type locality: Puerto Rico, Arecibo.

AUSTRALIS (Sabrosky), 1955: 45 (*Hippelates*). Type locality: Argentina, Rio Negro, Bariloche. Distr. — Colombia, Ecuador, Peru, Argentina (Chubut, Mendoza, Neuquén, Rio Negro, Tucumán), Chile, Islas Juan Fernandez.

partitus of Becker, 1912a: 169, in part.

pallipes of Malloch, 1934: 410 (*Cadrema*).

metallica of Enderlein, 1938: 658 (*Cadrema*).

BICOLOR (Coquillett), 1898: 48 (*Hippelates*). Type locality: U.S.A., Florida, Lake Worth. Distr. — U.S.A. (east coast), Bahamas, Puerto Rico, Virgin Islands.

BISHOPPI (Sabrosky), 1941a: 26 (*Hippelates*). Type locality: U.S.A., Florida, Zellwood. Distr. — U.S.A. (eastern), Mexico (Tamaulipas, Veracruz).

BREVISTYLUS (Malloch), 1934: 411 (*Cadrema*). Type locality: Uruguay, Montevideo. N. COMB.

- COLLUSOR** (Townsend), 1895: 619 (*Oscinis*). Type locality: Mexico, Baja California, San José del Cabo. Distr. — U.S.A. (southwestern), Mexico (Aguascalientes, Baja California, Chihuahua, Durango, Jalisco, Morelos, Nayarit, Nuevo León, Sonora, Tamaulipas).
- CURRANI** (Aldrich), 1931: 70 (*Hippelates*; nom. nov. for *collusor* Curran, 1926). Distr. — Mexico (Tamaulipas), Guatemala, Honduras, Costa Rica, Panama, Cuba, Jamaica, Dominican Republic, Puerto Rico, Virgin Islands, St. Vincent, Trinidad, Tobago, Colombia, Venezuela, Ecuador, Brazil (Espírito Santo, Mato Grosso, Pará, Paraíba, Rio de Janeiro).
- collusor* Curran, 1926: 4 (*Hippelates*; preocc. in *Hippelates* by Townsend, 1895). Type locality: Virgin Islands, St. Thomas, Charlotte Amalie.
- ESCOMELI** (Séguy), 1940: 346 (*Hippelates*). Type locality: Peru, Arequipa. N. COMB.
- FEMORALIS** (Duda), 1929: 165 (as *Hippelates pusio* var.). Type locality: Bolivia, Buena Vista, 80 km nw. Santa Cruz de la Sierra. Distr. — Colombia, Peru, Bolivia, Brazil (Rio de Janeiro), Argentina (Jujuy, Salta, Tucumán).
- femorata* Duda, 1929: 169, rejected alternate original spelling.
- FLAVIBUCCA** Duda, 1930: 67 (as *Liohippelates pusio* var.). Type locality: "Chile" [apparently error for Peru, from presumed holotype in Berlin, mislabeled "Chile/Callao"].
- FLAVICEPS** (Loew), 1863: 38 (Cent. 3, no. 71) (*Oscinis*). Type locality: "Cuba". Distr. — U.S.A. (Texas, Georgia, Florida), Mexico (Baja California, Tabasco, Veracruz), Guatemala, El Salvador, Nicaragua, Costa Rica, Bahamas, Cuba, Jamaica, Puerto Rico, Dominica, Martinique, St. Vincent, Venezuela, Ecuador, Peru, Brazil (Ceará, Rio de Janeiro, Santa Catarina, São Paulo), Paraguay, Argentina (Tucumán), Chile.
- metallicus* Becker, 1912a: 170 (*Hippelates*). Type locality: Peru, Unini (Rio Ucayali) and Mollendo; Paraguay, Asunción.
- nudifrons* Malloch, 1913c: 242 (*Hippelates*). Type locality: Mexico, Tabasco. N. COMB.
- FLAVIPES** (Loew), 1866a: 184 (Cent. 6, no. 95) (*Hippelates*). Type locality: "Cuba". Distr. — Mexico (Nayarit, Oaxaca, Veracruz), all Central America, Bahamas, Cuba, Jamaica, Haiti, Dominican Republic, Puerto Rico, Virgin Islands, Antigua, Guadeloupe, Dominica, St. Lucia, Barbados, St. Vincent, Grenada, Trinidad, Colombia, Venezuela, Ecuador, Peru, Bolivia, Brazil (Bahia, Mato Grosso, Pará, Paraná, Rio de Janeiro, Santa Catarina, São Paulo), Argentina (Misiones).
- angustibuccus* Duda, 1930: 66. Type locality: Not stated [Brazil, Pará; Peru, Meshagua on Rio Urubamba; Costa Rica, Suiza de Turrialba; from presumed syntypes in, respectively, Berlin, Dresden, and Budapest].
- brasilensis* Aldrich, 1931: 71 (*Hippelates*). Type locality: Brazil, Paraíba. N. COMB.
- pallipes* of some authors (e.g., Kumm, 1936), Neotropical records.
- papillipes*, error.



- GALAPAGENSIS (Curran), 1932: 356 (*Hippelates*). Type locality: Galapagos Islands, Floreana Id., Post Office Bay. N. COMB.
- INCOMPLETUS (Becker), 1912a: 170 (*Hippelates*). Type locality: Peru (as "Chile"), Tacna, lectotype in Dresden, here designated. Distr. — Peru, Chile.
- LUTZI (Curran), 1926: 5 (*Hippelates*). Type locality: Puerto Rico, Mona Island, and Virgin Islands, St. Croix. Distr. — Bahamas, Cuba, Puerto Rico, Virgin Islands, Panama.
- MICRONEURINUS (Duda), 1933b: 198 (*Stratiomicroneurum*). Type locality: Bolivia, Tumupasa. N. COMB.
- NIGRIBUCCA Duda, 1930: 67 (as *Liohippelates pusio* var.). Type locality: Not stated [Peru, Callao, from presumed syntypes in Hamburg, now destroyed, and Berlin, the latter mislabeled "Chile/Callao"].
- NIGRIFRONS Duda, 1930: 68. Type locality: Not stated [Paraguay, Hohenau, from presumed syntypes in Dresden, and Brazil, Rio de Janeiro, Petrópolis, from presumed syntype in São Paulo].
- NIGRIVERTEX Duda, 1936: 107. Type locality: Brazil, Pernambuco, Recife.
- PALLIPES (Loew), 1863: 37 (Cent. 3, no. 69) (*Oscinis*). Type locality: "Cuba". Distr. — U.S.A. (eastern), Cuba.
- partitus* Becker, 1912a: 89 (*Hippelates*). Type locality: U.S.A., Alabama.
- PERUANUS (Becker), 1912a: 170 (*Hippelates*). Type locality: Peru, Vilcanota; Argentina, San Pablo; Paraguay, Asunción. Distr. — U.S.A. (Florida), Mexico (Morelos, Tamaulipas, Veracruz), Guatemala, Nicaragua, Costa Rica, Bahamas, Cuba, Jamaica, Haiti, Dominican Republic, Puerto Rico, Virgin Islands, Guadeloupe, Trinidad, Tobago, Colombia, Venezuela, Ecuador, Peru, Bolivia, Brazil (Ceará, Pará, Paraná, Rio de Janeiro, Santa Catarina, São Paulo), Paraguay, Argentina (Chaco, Corrientes, Formosa, Jujuy, Misiones, Salta, Tucumán).
- circumdata* Becker, 1912a: 185 (*Siphonella*). Type locality: Haiti. N. SYN., N. COMB.
- PUSIO (Loew), 1872: 103 (Cent. 10, no. 87) (*Hippelates*). Type locality: U.S.A., Texas. Distr. — U.S.A. (widespread), Mexico (Coahuila, Durango, Hidalgo, Jalisco, Nuevo León, Oaxaca, San Luis Potosí, Tamaulipas, Veracruz), Guatemala, El Salvador, Honduras, Nicaragua, Costa Rica, Panama, Bermuda, Bahamas, Cuba, Jamaica, Haiti, Dominican Republic, Puerto Rico, Montserrat, Trinidad.
- ROBERTSONI (Sabrosky), 1941a: 26 (*Hippelates*). Type locality: U.S.A., California, San Bernardino National Forest. Distr. — U.S.A. (western), Mexico (Aguascalientes, Baja California, Chiapas, Chihuahua, Durango, Guanajuato, Hidalgo, Jalisco, México, Michoacán, Morelos, Nayarit, Nuevo León, Oaxaca, Puebla, San Luis Potosí, Tamaulipas, Veracruz, Zacatecas), Guatemala, El Salvador, Panama.
- SATANAS (Becker), 1912a: 210 (*Oscinella*). Type locality: Peru, Cuzco. N. COMB.

- STRANDI (Duda), 1930: 68, 74, 110; 1933a: 41, full description (*Goniopsita*). Type locality: Not stated in 1930; in 1933a, Peru, Cuzco and Urubamba, in Dresden; Costa Rica, Farm La Caja, near San José, in Hamburg and now destroyed, and in Berlin. N. COMB.
- TENACISSIMUS (Iches), 1906: 269 (*Chlorops*). Type locality: Argentina, Chaco. Unrecognized; type cannot be located. N. COMB. (from description).
- TIBIALIS Duda, 1930: 67. Type locality: Not stated [Peru, Chanchamayo and Meshagua (Rio Urubamba), from presumed syntypes in Dresden, and Brazil, Rio de Janeiro, Petrópolis, from presumed syntype in São Paulo]. Distr. — Guatemala, Panama, Jamaica, Puerto Rico, Colombia, Ecuador, Peru, Brazil (Ceará, Pará, Rio de Janeiro, São Paulo), Paraguay, Argentina (Misiones).

Genus LIOSCINELLA Duda

Oscinella subg. LIOSCINELLA Duda, 1929: 168, 169. Type species, *Oscinosoma sulfurihalterata* Enderlein, 1911 (Sabrosky, 1941b: 755).

This group is close to *Liohippелates*, and perhaps it should constitute a subgenus or a species group of that genus. If the two are combined, *Liohippелates* would obviously be the name of choice because of the association with *Hippelates* flies.

- BOLIVIENSIS Duda, 1930: 111. Type locality: "Bolivia" (inferred from name) [La Paz, from presumed syntypes in Dresden].
- DENTITIBIA (Enderlein), 1913: 367 (*Siphonella*). Type locality: "Costa Rica" [probably Farm La Caja, near San José, because the holotype was collected by H. Schmidt].
- IGNORATA (Duda), 1929: 168 (*Oscinella*, subg. *Lioscinella*, *sulfurihalterata* var.). Type locality: Argentina, Tapikiolé [lectotype ♂, here designated, in Stuttgart] and Bolivia, San José de Chiquitos [Bolivian syntypes appear to be a distinct but similar species]. Distr. — Argentina (Chaco, Jujuy, Tucumán).
- NIGRIROSTRA (Adams), 1904b: 103 (*Siphonella*). Type locality: West Indies, Grenada. N. COMB. [probable position; name made available in a key, and the type has not been found].
- PROSTHIOMELAS (Enderlein), 1913: 366 (*Siphonella*, as *speculigera* var.). Type locality: Colombia, Hacienda Pehlke [Caldas, near Victoria (Papavero, 1973: 299)]. Distr. — Costa Rica, Colombia.
- SPECULIGER (Enderlein), 1911a: 232, part (*Siphonella*). Type locality: Brazil, Santa Catarina [Two syntypes from Costa Rica were later (Enderlein, 1913) described as *Siphonella dentitibia* and *Oscinosoma longipalpus*]. Distr. — Brazil (Rio de Janeiro, Santa Catarina), Paraguay.
- SULFURIHALTERATA (Enderlein), 1911a: 222 (*Oscinosoma*). Type locality: "Costa Rica" [probably Farm La Caja, near San José, because the holotype was collected by H. Schmidt].



Genus LOXOBATHMIS Enderlein

LOXOBATHMIS Enderlein, 1911a: 205. Type species, *obliquans* Enderlein, 1911 (orig. des.).

OBLIQUANS Enderlein, 1911a: 205. Type locality: Brazil, Santa Catarina. Distr. — Costa Rica, Brazil.

Genus MALLOEWIA Sabrosky

MALLOEWIA Sabrosky, 1980b: 422. Type species, *Siphonella diabolus* Becker, 1912 (orig. des.).

COERULEIFRONS (Duda), 1930: 104 (*Oscinella*). Type locality: Not stated, but presumed syntypes listed by Duda, 1933a: 45: Bolivia, La Paz, syntypes in Dresden and Vienna; Paraguay, Asunción, Villa Morra, syntype in Budapest; Chile, Quillota, syntype in Eberswalde. Distr. — Bolivia, Paraguay, Argentina (Buenos Aires, Córdoba, Jujuy, Salta, Santa Fe), Chile. N. COMB.

INCOLUMIS (Becker), 1912a: 185 (*Siphonella*). Type locality: Bolivia, Sorata [Lectotype should be designated from Bolivia; a syntype from Argentina, Salta, Metán, was referred by Duda to his *Oscinella coeruleifrons*]. Distr. — Bolivia, Argentina (Córdoba, Corrientes, Entre Ríos, Jujuy, Salta, Tucumán). N. COMB.

NEGLECTA (Becker), 1912a: 100 (*Siphonella*). Type locality: U.S.A., California, Berkeley. Distr. — U.S.A. and Canada (both western), Mexico (Baja California, Chihuahua, Jalisco, México, Puebla, San Luis Potosí).

NITIDA Duda, 1930: 110 (*Lioscinella vulgaris* var.). Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed syntypes in Hamburg, now destroyed, and Berlin, Dresden]. Distr. — Guatemala, El Salvador, Costa Rica, Panama, N. STATUS, N. COMB.

VULGARIS (Duda), 1930: 110 (*Lioscinella*). Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed syntypes in Hamburg, now destroyed, Eberswalde (lectotype ♂, here designated, Farm La Caja, Sept. — Nov. 1924), and Berlin]. N. COMB.

Genus Melanochaeta Bezzi

eunota (Loew). Not Neotropical (Eastern U.S.A. and Canada). Neotropical records in error: Becker, 1912a: 121 (type locality is Texas, not Cuba), 181 (Havana, Illinois, not Cuba).

Genus METASIPHONELLA Duda

METASIPHONELLA Duda, 1930: 107. Type species, *magnifica* Duda, 1930 (mon.).

MAGNIFICA Duda, 1930: 107. Type locality: Not stated [Costa Rica, Farm La Caja, near San José, from presumed syntypes at Hamburg, now destroyed, and presumed syntype from Turrialba, at Budapest]. Distr. — Costa Rica, Panama, Venezuela, Ecuador.

Genus MONOCHAETOSCINELLA Duda

MONOCHAETOSCINELLA Duda, 1930: 107. Type species, *Oscinis anonyma* Williston, 1896 (Duda, 1931: 166).

ANONYMA (Williston), 1896: 423 (*Oscinis*). Type locality: St. Vincent. Distr. — U.S.A. (Arizona, Texas, Florida), Mexico (Aguascalientes, Baja California, Hidalgo, Jalisco, Michoacán, Morelos, San Luis Potosí, Sonora, Tamaulipas, Veracruz, Yucatán), Guatemala, El Salvador, Honduras, Nicaragua, Costa Rica, Panama, Bermuda, Bahamas, Cuba, Jamaica, Haiti, Dominican Republic, Puerto Rico, Virgin Islands, Dominica, St. Vincent, Trinidad, Colombia, Venezuela, Guyana, Surinam, Peru, Bolivia, Brazil (Pará, Rio de Janeiro, Santa Catarina, São Paulo), Argentina (Corrientes).

pura Curran, 1926: 7 (*Botanobia anonyma* var.). Type locality: Puerto Rico, San Juan (Curran, 1928: 54). N. SYN., N. COMB.

attingens Duda, 1930: 110 (*Lioscinella*). Type locality: "Surinam" [Paramaribo, from presumed syntypes in Hamburg, now destroyed, and in Eberswalde (lectotype ♀, here designated), and Berlin]. N. SYN., N. COMB.

NIGRIPES Duda, 1930: 104, 107. Type locality: Not stated [Argentina, Tucumán, Lules, from presumed holotype at Budapest]. Distr. — Brazil (Pará, Santa Catarina), Paraguay, Argentina (Corrientes, Jujuy, Tucumán).

ZERNYI Duda, 1936: 112. Type locality: Brazil, Pará, Belém. Distr. — Ecuador, Brazil (Rio de Janeiro, Santa Catarina).

Genus NOTAULACELLA Enderlein

NOTAULACELLA Enderlein, 1911a: 212. Type species, *armillata* Enderlein, 1911 (orig. des.).

Baseoneura Duda, 1930: 69, 71. Type species, *Notaulacella armillata* Enderlein, 1911 (Duda, 1931: 165).
Ref. — Duda, 1930: 71 (key).

ANGUSTATA (Enderlein), 1911a: 223 (*Oscinosoma*). Type locality: Brazil, Santa Catarina [lectotype ♂, here designated, labeled "type" by Enderlein, originally in Stettin, now in Warsaw].

ARMILLATA Enderlein, 1911a: 212. Type locality: Brazil, Santa Catarina.

COMPTA (Duda), 1930: 71 (*Baseoneura*). Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed syntype in Hamburg, now destroyed, and syntype in Berlin]. Distr. — Costa Rica.

FIEBRIGI (Duda), 1930: 72 (*Baseoneura*). Type locality: "Paraguay" [San Bernardino, from presumed syntypes in Vienna].

SCHILDI Duda, 1933b: 200 (as *N. trapezisoptron* var.). Type locality: Costa Rica, Higuito, San Mateo.

TRAPEZISOPTRON (Enderlein), 1911a: 224 (*Oscinosoma*). Type locality: Brazil, Santa Catarina.




Genus OLCANABATES Enderlein

- OLCANABATES Enderlein, 1911a: 189. Type species, *strigatus* Enderlein, 1911 (orig. des.).
- Hippelatinus* Enderlein, 1911a: 193. Type species, *H. calcaratus* Enderlein, 1911 (orig. des.).
- CALCARATUS (Enderlein), 1911a: 193 (*Hippelatinus*). Type locality: Brazil, Santa Catarina.
- sulcatus* Enderlein, 1911a: 190 (*Olcababates*). Type locality: Brazil, Santa Catarina.
- FLAVIDULUS (Malloch), 1913c: 256 (*Hippelates*). Type locality: West Indies, Montserrat. Distr. — El Salvador, Panama, Colombia, Guyana, Montserrat. N. COMB.
- RUFESCENS (Duda), 1934: 65 (*Tropidoscinis nitens* var.). Type locality: Costa Rica, Higuito, San Mateo. N. COMB.
- STRIGATUS Enderlein, 1911a: 189. Type locality: Brazil, Santa Catarina. Distr. — Brazil (Rio de Janeiro, Santa Catarina), Peru.

Genus OLCELLA Enderlein

- OLCELLA Enderlein, 1911a: 231. Type species, *mendozana* Enderlein, 1911 (orig. des.).
- Siphonella*, part, and *Madiza*, part, of authors.
- ANGUSTIBUCCA (Duda), 1930: 96 (*Conioscinella*). Type locality: Not stated [Paraguay, San Bernardino, from presumed holotype in Berlin]. N. COMB.
- AUSTRALIS (Malloch), 1934: 429 (*Madiza*). Type locality: Uruguay, Montevideo. N. COMB.
- CINEREA (Loew), 1863: 43 (Cent. 3, no. 81) (*Siphonella*). Type locality: U.S.A., Florida. Distr. — U.S.A. (eastern), Mexico (Jalisco, Nayarit, Tamaulipas).
- CONVEXA (Malloch), 1934: 429 (*Madiza mendozana* var.). Type locality: Uruguay, Montevideo. N. COMB.
- FASCIATA (Duda), 1930: 101 (*Conioscinella*). Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed holotype in Hamburg, now destroyed]. N. COMB.
- FLAVISETA (Malloch), 1934: 430 (*Madiza*). Type locality: Argentina, Lake Nahuel Huapi. N. COMB.
- MENDOZANA Enderlein, 1911a: 231. Type locality: Argentina, Mendoza.
- var. GRISEOSCUTELLATA (Duda), 1930: 92, 93 (*Conioscinella mendozana* var.). Type locality: "Argentina" [Mendoza] and "Paraguay" [San Bernardino], localities from presumed syntypes in Vienna. N. COMB.
- var. OBSCURIPILA (Duda), 1929: 166 (*Oscinella mendozana* var.). Type locality: Argentina, Aguaray. N. COMB.



- var. SCUTELLARIS (Malloch), 1934: 429 (*Madiza mendozana* var.). Type locality: Chile, Santiago. N. COMB.
- NEOTROPICA (Becker), 1912a: 183, 184 (as *nectropica*) (*Siphonella*). Type locality: "Argentina" [Tucumán, from holotype in Budapest]. N. COMB.
- NIGRISETA (Malloch), 1934: 431 (*Madiza*). Type locality: Chile, Santiago. N. COMB.
- NIGRITHORAX (Duda), 1930: 98 (*Neoolcanabates*). Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed holotype in Hamburg, now destroyed]. N. COMB.
- OPPOSITA (Becker), 1912a: 184 (*Siphonella*). Type locality: Chile, Palca, and Peru, Cuzco. N. COMB.
- PARAGUAYENSIS Sabrosky & Paganelli, NOM. NOV.
pleuralis Duda, 1930: 92, 100 (*Conioscinella*; preocc. in *Olcella* by *pleuralis* Becker, 1912). Type locality: Paraguay [San Bernardino, from presumed holotype in Vienna]. N. COMB.
- PARVA (Adams), 1904b: 104 (*Siphonella*). Type locality: U.S.A., Colorado. Distr. — Canada and U.S.A. (widespread), Mexico (Aguascalientes, Durango, Hidalgo, Jalisco, México, Michoacán, Morelos, Nayarit, Nuevo León, Puebla, San Luis Potosí, Tamaulipas, Zacatecas).
- PLEURALIS (Becker), 1912b: 645 (*Oscinella*; replacement name for *Oscinis nana* Williston). Distr. — Guatemala, Costa Rica, Panama, Cuba, Jamaica, Puerto Rico, St. Vincent. N. COMB.
- nana* Williston, 1896: 423 (*Oscinis*; preocc. Zetterstedt, 1838). Type locality: St. Vincent. N. COMB.
- griseolineata* Duda, 1930: 91 (*Conioscinella*). Type locality: Not stated [Costa Rica, presumed holotype (Sabrosky, 1939: 179), Farm La Caja near San José, in Hamburg, now destroyed; subsequently a presumed syntype, same locality, found in Berlin, and a presumed syntype (or syntypes), found in Budapest, from Costa Rica, Suiza de Turrialba]. N. COMB.
- PROJECTA (Malloch), 1913b: 283 (*Madiza*). Type locality: U.S.A., New Mexico, Las Cruces. Distr. — U.S.A. (western), Mexico (Aguascalientes, Jalisco, Nuevo León, Tamaulipas).
- PUNCTIFRONS (Becker), 1912a: 101 (*Siphonella*). Type locality: U.S.A., Utah, Cache County. Distr. — U.S.A. (western), Mexico (Durango).
- quinquelineata* (Adams). Not Neotropical. Puerto Rico records are probably either of *trilineata* or *pleuralis* (Becker).
- TRIGRAMMA (Loew), 1863: 42 (Cent. 3, no. 80) (*Oscinis*). Type locality: U.S.A., District of Columbia. Distr. — U.S.A. (eastern), Mexico (Guerrero, Tamaulipas).
- TRILINEATA (Duda), 1930: 94 (*Conioscinella*). Type locality: "Brazil" [Estado do Rio de Janeiro, Petrópolis, from presumed holotype in São Paulo]. Distr. — Guatemala, Costa Rica, Jamaica, Dominican Republic, Puerto Rico, Brazil (Rio de Janeiro). N. COMB.



TURBATA (Duda), 1930: 99 (*Neoolcanabates*). Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed holotype in Hamburg, now destroyed]. N. COMB.

Genus ONYCHASPIDIUM Enderlein

ONYCHASPIDIUM Enderlein, 1911a: 196. Type species, *sexdentatum* Enderlein, 1911 (orig. des.).

Leptopeltastes Enderlein, 1911a: 229. Type species, *L. longiscutellata* Enderlein, 1911 (orig. des.).

Meroscinis Meijere of Becker, 1912a, and Duda, 1930, misident. Ref. — Duda, 1930: 82 (key).

APICALE (Williston), 1896: 422 (*Oscinis*). Type locality: St. Vincent, west slope of Soufrière, 1800 ft. Distr. — Mexico (Hidalgo, México, Morelos, San Luis Potosí, Veracruz), Costa Rica, St. Vincent, Brazil (Rio de Janeiro).

quadridentata Duda, 1930: 83 (*Meroscinis*). Type locality: "Brazil" [Rio de Janeiro, Petrópolis, from lectotype (Sabrosky, 1947: 177) in São Paulo].

FLAVIPES (Duda), 1930: 83 (*Meroscinis*). Type locality: "Bolivia" [Mapiri, from presumed holotype in Dresden]. N. COMB.

LONGISCUTELLATUM (Enderlein), 1911a: 229 (*Leptopeltastes*). Type locality: Brazil, Santa Catarina.

NIGRICEPS (Duda), 1930: 83 (*Meroscinis*). Type locality: "Costa Rica" [Suiza de Turrialba, from presumed holotype in Budapest]. N. COMB.

SEXDENTATUM Enderlein, 1911a: 196. Type locality: Brazil, Santa Catarina. Distr. — Peru, Brazil (Santa Catarina).

Genus OPETIOPHORA Loew

OPETIOPHORA Loew, 1872: 105 (Cent. 10, no. 90). Type species, *straminea* Loew, 1872 (mon.).

STRAMINEA Loew, 1872: 105 (Cent. 10, no. 90). Type locality: U.S.A., Texas. Distr. — U.S.A. (west central), Mexico (Puebla).

Genus OSCINELLA Becker

OSCINELLA Becker, 1909: 120. Type species, *Musca frit* Linnaeus, designated under the plenary power (I. C. Z. N., 1978: 203).

CARBONARIA (Loew), 1869a: 42 (Cent. 8, no. 76) (*Oscinis*). Type locality: U.S.A., District of Columbia. Distr. — widespread, southern Canada and the U.S.A., Mexico (Tamaulipas).

DAMPFI Aldrich, 1929: 90. Type locality: Mexico, Distrito Federal, San Jacinto. Distr. — Mexico (Distrito Federal, México, Hidalgo).



FRIT (Linnaeus), 1758: 598 (*Musca*). Type locality: Sweden. Distr. — Hol-arctic, Canada and U.S.A., Mexico (Distrito Federal, México, Hidalgo, Nayarit).

OBSCURA (Coquillett), 1900: 266 (*Oscinis*). Type locality: Puerto Rico, Mayagüez [specimen labeled type by Coquillett].

PLESIA (Curran), 1926: 11 (*Botanobia*). Type locality: Puerto Rico, Arecibo.

UJHELYI Duda, 1930: 105. Type locality: Colombia [Sierra San Lorenzo, from presumed holotype in Budapest, now headless].

umbrosa (Loew). Not Neotropical.

Oscinella, blanda group

This small group of tiny species with linear genae and very large, polished black, convex-sided frontal triangle may deserve generic distinction but there is no available name and for the present it is left in *Oscinella* until adequate material is available for study. It seems likely that some of the species listed will prove to be synonyms.

BLANDA Becker, 1912a: 208. Type locality: Bolivia, Mapiri, Bellavista; Ecuador, Riobamba area.

BRACATA (Duda), 1930: 73 (*Goniopsita blanda* var.). Type locality: "Costa Rica" [Farm La Caja near San José, from presumed holotype in Hamburg, now destroyed]. Distr. — Mexico (Veracruz), Costa Rica. N. COMB.

CONCINNA (Williston), 1896: 424 (*Oscinis*). Type locality: St. Vincent.

FLAVITIBIA (Duda), 1933b: 200 (*Goniopsita blanda* var.). Type locality: Panama, Canal Zone, Monte Lirio. N. COMB.

MATTEA (Curran), 1926: 5 (*Madiza*). Type locality: Puerto Rico, Adjuntas. N. COMB.

VULGARIMONENS Duda, 1934: 110. Type locality: Costa Rica, Higuito, San Mateo. Distr. — Costa Rica, Panama (Canal Zone).

Genus OSCINOIDES Malloch


OSCINOIDES Malloch, 1916: 86. Type species, *arpidia* Malloch, 1916 (orig. des.).

ANNULICRUS (Duda), 1933b: 205 (*Gaurax*). Type locality: Costa Rica, Higuito, San Mateo. N. COMB.

Genus PARASIPHONELLA Enderlein

PARASIPHONELLA Enderlein, 1913: 367. Type species, *Scoliophthalmus dentigena* Enderlein, 1911 (orig. des.).

DENTIGENA (Enderlein), 1911a: 233 (*Scoliophthalmus*). Type locality: Brazil, Santa Catarina. Type lost; close to *Loxobathmis obliquans*.



PSEUDOGAURAX Malloch, 1915: 159. Type species, *Gaurax anchora* Loew, 1866 (orig. des.).

Pseudogaurax Duda, 1930: 86. Type species, *Gaurax interruptus* Becker, 1912 (mon.). Homonym and synonym of *Pseudogaurax* Malloch.

Mimogaurax Hall, 1937: 257. New name for *Pseudogaurax* Duda.

Gaurax Loew of Becker (1912a) and Duda (1930).

The species of *Pseudogaurax* are often reared from spider egg sacs and mantid oothecae. Ref. — Sabrosky, 1966 (key and revision, Western Hemisphere).

CINGULATUS Sabrosky, 1966: 124. Type locality: Brazil, Rio Grande do Sul, São Leopoldo. Distr. — Brazil (Rio Grande do Sul, São Paulo).

FLORIDENSIS Sabrosky, 1950a: 33. Type locality: U.S.A., Florida, Middle Cape of Cape Sable. Distr. — U.S.A. (Florida), Jamaica.

IMMACULATUS (Becker), 1916: 450 (*Gaurax*). Type locality: Colombia, Sierra San Lorenzo.

INTERRUPTUS (Becker), 1912a: 197 (*Gaurax*). Type locality: "Brazil".

LANCIFER (Coquillett), 1900: 265 (*Gaurax*). Type locality: Puerto Rico, San Juan, and Montserrat. Distr. — Haiti, Puerto Rico, Virgin Islands, Montserrat.

LATIMACULATUS Hall, 1937: 260. Type locality: Panama, Alhajuelo.

LONGILINEATUS Sabrosky, 1949: 33. Type locality: Brazil, Rio de Janeiro, Nova Friburgo.

LUCISCENS (Enderlein), 1911a: 219 (*Oscinosoma*). Type locality: Brazil, Santa Catarina.

luciscens, error, Duda, 1930: 86.

MISCEOMACULATUS Hall, 1937: 259. Type locality: Puerto Rico, Bayamón.

NIGROLINEATUS (Enderlein), 1911a: 218 (*Oscinosoma*). Type locality: Brazil, Santa Catarina. Distr. — Brazil (Rio de Janeiro, Santa Catarina).

nigrilineatus, error, Becker, 1912a: 198, and Duda, 1930: 86 (both *Gaurax*).

NIGROMACULATUS (Duda), 1933b: 206 (*Gaurax*). Type locality: Costa Rica, Higuito, San Mateo.

OCULATUS (Becker), 1912a: 214 (*Oscinella*). Type locality: Paraguay, San Bernardino. Distr. — Paraguay, Argentina (Tucumán).

OECETIPHAGUS (Blanchard), 1941: 20 (*Gaurax*). Type locality: Argentina, Buenos Aires, Vicente López.

oeceticola, error, Griot, et al., 1946: 320, 324 (*Gaurax*).

- PALLIPES (Duda), 1933b: 207 (*Gaurax tectus* var.). Type locality: Costa Rica, Higuito, San Mateo.
- PARALLELINERVIS (Duda), 1933b: 208 (*Botanobia*). Type locality: Costa Rica, Higuito, San Mateo.
- PLAUMANNI Sabrosky, 1966: 122. Type locality: Brazil, Santa Catarina, Nova Teutônia.
- RUFUS (Duda), 1930: 87 (*Gaurax*). Type locality: "Brazil" [presumed holotype in Berlin, no specific locality].
- SIGNATUS (Loew), 1876: 338 (*Gaurax*). Type locality: U.S.A., Texas. Distr. — U.S.A. (southern), Mexico (Morelos), Cuba.
- TECTUS (Becker), 1916: 450 (*Gaurax*). Type locality: Peru, Callanga.
- TESTACEUS (Enderlein), 1911a: 220 (*Oscinosoma*). Type locality: Brazil, Santa Catarina.
- TIBIALIS Malloch, 1934: 421. Type locality: Argentina, Bompland (= Bonpland).
- TRABEATUS (Enderlein), 1911a: 218 (*Oscinosoma*). Type locality: Brazil, Santa Catarina.
- TRIANGULARIS (Williston), 1896: 421 (*Oscinis*). Type locality: St. Vincent, Leeward side.
- TRIDENS Sabrosky, 1966: 121. Type locality: Brazil, Santa Catarina.
- TRIFIDUS (Duda), 1930: 87 (*Gaurax*). Type locality: "Costa Rica" [Suiza de Turrialba, from presumed holotype in Budapest].
- lyricus* Hall, 1937: 260. Type locality: Costa Rica, Higuito, San Mateo. N. SYN.
- TRILINEATUS (Duda), 1930: 86 (*Gaurax*). Type locality: "Brazil" (error? The only specimens found with Duda's label as new species are from Paraguay, San Bernardino, in Vienna). Distr. — Nicaragua, Paraguay.
- UNILINEATUS Hall, 1937: 259. Type locality: Panama (Canal Zone), Ancon.

Genus PTEROGAURAX Duda

- PTEROGAURAX Duda, 1930: 84. Type species, *Gaurax pennatus* Becker, 1912 (Duda, 1931: 165).
- PENNATUS (Becker), 1912a: 196 (*Gaurax*). Type locality: Bolivia, Mapiri, Sarampiuni. [The syntype from Peru, Marcapata, was described by Duda, 1930, as *peruanus*].
- PERUANUS Duda, 1930: 84. Type locality: "Peru" [Marcapata, from presumed holotype in Budapest, syntype of *pennatus*].

Genus RHODESIELLA Adams

- RHODESIELLA Adams, 1905b: 197. Type species, *tarsalis* Adams, 1905 (orig. des.).



Unidentified spp. — El Salvador, Costa Rica, Panama, Jamaica, Dominican Republic, Puerto Rico, Brazil (Rio de Janeiro, Santa Catarina, São Paulo).

Genus RHOPALOPTERUM Duda

Oscinella, subg. RHOPALOPTERUM Duda, 1929: 167. Type species, *Oscinella limitata* Becker, 1912 (mon.).

Rhopalopternum Duda, 1930: 107. Type species, *Oscinella limitata* Becker (Duda, 1931: 166). Probably error or emendation for *Rhopalopternum* Duda, 1929.

Ref. — Sabrosky, 1961 (key).

FLAVICORNE Duda, 1934: 68 (*Rhopalopternum*). Type locality: Costa Rica, Higuito, San Mateo.

GLABRUM (Duda), 1929: 167 (*Oscinella limitata* var.). Type locality: Brazil, Rio de Janeiro.

LIMITATUM (Becker), 1912a: 211 (*Oscinella*). Type locality: Haiti. Distr. — U.S.A. (Florida), Mexico (San Luis Potosí, Veracruz), Guatemala, Costa Rica, Panama, Bermuda, Cuba, Jamaica, Haiti, Dominican Republic, Puerto Rico, Trinidad, Colombia, Venezuela.

umbrosa Loew of Coquillett, 1900: 267 (*Oscinis*), misident.

Genus SIPHONELLOMYIA Séguy

SIPHONELLOMYIA Séguy, 1934: 13. Type species, *hemiptera* Séguy, 1934 (orig. des.).

HEMIPTERA Séguy, 1934: 13. Type locality: Argentina, La Plata.

Genus SIPHUNCULINA Rondani

SIPHUNCULINA Rondani, 1856: 128. Type species, *brevinervis* Rondani, 1856 (orig. des.).

STRIOLATA (Wiedemann), 1830: 597 (*Chlorops*). Type locality: China. Distr. — U.S.A. (Florida), Mexico (Baja California), Panama, Cuba, Jamaica, Haiti, Puerto Rico, Virgin Islands, St. Lucia, Trinidad, Venezuela, Guyana, Ecuador, Peru, Brazil (Ceará, Pará, Rio de Janeiro), Paraguay. Widespread, tropicopolitan species.

plumbella Wiedemann, 1830: 574 (*Homalura*). Type locality: West Indies. ?Synonym; type lost; description fits *striolata*.

signata Wollaston, 1858: 117 (*Oscinis*). Type locality: Madeira.

reticulata Loew, 1869a: 43 (Cent. 8, no. 78) (*Siphonella*). Type locality: Cuba.

Genus STENOSCINIS Malloch

STENOSCINIS Malloch, 1918: 21. Type species, *Oscinis longipes* Loew, 1863 (orig. des.).

Ref. — Sabrosky, 1961 (key).

ANTIGUENSIS (Duda), 1934: 67 (*Rhopalopternum*). Type locality: Guatemala. El Salto, Antigua.

MAJOR (Duda), 1934: 101 (*Rhopalopternum*, as *majus*). Type locality: Costa Rica, San José. Distr. — Mexico (Distrito Federal, Jalisco), Guatemala, El Salvador, Honduras.

Genus STRANDIMYIA Duda

STRANDIMYIA Duda, 1930: 59, 70. Type species, *flava* Duda, 1930 (mon.).

FLAVA Duda, 1930: 59, 100; 1931: 167 (description). Type locality: Brazil, Rio de Janeiro, Petrópolis (Duda, 1931).

Genus TRICIMBA Lioy

TRICIMBA Lioy, 1864: 1125. Type species, *Oscinis lineella* Fallén, 1820 (Enderlein, 1911a: 207).

Microchaetaspis Duda, 1930: 77. Type species, *crassiseta* Duda, 1930 (Duda, 1931: 165). N. SYN.

Hammaspis Duda, 1930: 76. Type species, *Tricimba spinigera* Malloch, 1913 (mon.).

Subgenus TRICIMBA Lioy

BREVIPIILA Duda, 1930: 78. Type locality: Not stated [Costa Rica, Farm La Caja, near San José, from presumed holotype in Hamburg, now destroyed]. Distr. — Honduras, Costa Rica, Colombia.

CRASSISETA (Duda), 1930: 78 (*Microchaetaspis*). Type locality: Chile [Valparaíso, from presumed holotype in Dresden]. N. COMB.

ENDERLEINI Duda, 1930: 77. Type locality: Not stated [Brazil, Santa Catarina, from presumed holotype, originally in Stettin, now in Warsaw]. Distr. — Peru, Brazil (Santa Catarina), Argentina (Salta, Tucumán).

cricopus Enderlein, in part.

FLAVIPES Duda, 1933b: 203 (as *enderleini* var.). Type locality: Costa Rica, Higuito, San Mateo.

FLAVOLINEATA (Duda), 1930: 78 (*Microchaetaspis*). Type locality: "Costa Rica", [Suiza de Turrialba, from presumed holotype in Budapest]. N. COMB.

MELANCHOLICA (Becker), 1912a: 109 (*Oscinella*). Type locality: U.S.A., Michigan, Battle Creek. Distr. — Widespread U.S.A. and Canada, Mexico (Durango, Hidalgo, Michoacán, Puebla, Tamaulipas).

NIGRA Malloch, 1934: 427 (as *niger*). Type locality: Argentina, Rio Negro, Lake Correntoso.

Subgenus PENTANOTAULAX Enderlein

PENTANOTAULAX Enderlein, 1911a: 197 (as genus). Type species, *virgulata* Enderlein, 1911 (orig. des.).

Gauracisoma Duda, 1930: 76 (as genus). Type species, *Pentanotaulax caviventris* Enderlein, 1911 (Duda, 1931: 165). N. SYN.

This should perhaps be considered a distinct genus, but most of the species were described in or referred to *Tricimba*, and for the present at least it is useful to retain them in *Tricimba*.

BREVIVENTRIS Malloch, 1934: 426. Type locality: Argentina, Buenos Aires.

CAVIVENTRIS (Enderlein), 1911a: 199 (*Pentanotaulax*). Type locality: Brazil, Santa Catarina. N. COMB.

NIGROFEMORATA Duda, 1930: 79. Type locality: Not stated [Costa Rica, Farm La Caja, near San José, from presumed syntypes in Hamburg, now destroyed, and Berlin, and from Suiza de Turrialba, in Budapest].

GCHRIPES Enderlein, 1911a: 209. Type locality: Brazil, Santa Catarina.

PALPALIS Becker, 1912a: 191. Type locality: "Paraguay".

PUBISETA (Becker), 1912a: 191 (*Pentanotaulax*). Type locality: "Paraguay". Distr. — Brazil (Rio de Janeiro), Paraguay.

PYGMAEA Duda, 1930: 80 (as *Tricimba virgulata* var.). Type locality: Not stated [Costa Rica, Suiza de Turrialba, from presumed syntypes in Budapest].

RIMATA Enderlein, 1911a: 211. Type locality: Brazil, Santa Catarina.

SIMILIS Enderlein, 1911a: 210. Type locality: Brazil, Santa Catarina.

SULCATA Duda, 1930: 80. Type locality: "Costa Rica" [Suiza de Turrialba, from presumed holotype in Budapest].

TECTA (Becker), 1912a: 190 (*Pentanotaulax*). Type locality: Bolivia, Mapiri, San Ernesto.

TOFINISTRIGATA (Enderlein), 1911a: 198 (*Pentanotaulax*). Type locality: Brazil, Santa Catarina.

tofinigastra, error.

VIRGULATA (Enderlein), 1911a: 197 (*Pentanotaulax*). Type locality: Brazil, Santa Catarina. Distr. — Mexico (Vera Cruz), Guatemala, Honduras, Costa Rica, Ecuador, Peru, Brazil (Rio de Janeiro, Santa Catarina).

Genus TROPIDOSCINIS Enderlein

TROPIDOSCINIS Enderlein, 1911a: 226. Type species, *luederwaldti* Enderlein, 1911 (orig. des.).

Neoolcanabates Duda, 1930: 91. Type species, *Tropidoscinis luederwaldti* Enderlein (Sabrosky, 1941b: 759).

Ref. — Sabrosky, 1980b (lectotype designation from mixed series).

LUEDERWALDTI Enderlein, 1911a: 227. Type locality: Brazil, Santa Catarina [lectotype designated Sabrosky, 1980b: 417].

Species *incertae sedis*, OSCINELLINAE

- CATHARINENSIS Enderlein, 1911a: 221 (*Oscinosoma*). Type locality: Brazil, Santa Catarina. Becker (1912a) considered that it was a synonym of *Discogastrella trigonalifrons* Enderlein (see *Coryphisopteron* in this catalogue). The holotype was lost during its return to Enderlein, but he continued to regard *catharinensis* as distinct from *trigonalifrons*, but in *Oscinosoma*.
- CURVINERVIS Duda, 1930: 91 (*Conioscinella*). Type locality: "Costa Rica" [Suiza de Turrialba, from presumed syntypes in Budapest, of which a male, V-2-1921, bearing Duda's identification label as "n.sp." is here designated lectotype, and in Berlin]. Not *Conioscinella*, but generic position uncertain.
- LEPRAE Linnaeus, 1758: 598 (*Musca*). Type locality: "Americae". Unrecognized. Wiedemann (1830: 598), as *Chlorops leprae*, questionably identified the species from a specimen that Becker (1912a: 172) placed in *Hippelates* (sens. lat.), but there is no assurance that the Linnaean species was even a chloropid.
- LINDNERI Duda, 1929: 168 (*Oscinella*, subg. *Lioscinella*). Type locality: Bolivia, Samuhuate [presumed syntypes in Stuttgart]. Not *Lioscinella* but generic position uncertain.
- LONGIPALPUS Enderlein, 1913: 360 (*Oscinosoma*). Type locality: (Brazil, Santa Catarina, or Costa Rica?) Originally included in the type series of *Siphonella speculiger*. On p. 361 he stated the holotype (only specimen) to be from Brazil, but on p. 366, in a note under *speculiger*, he stated that it was a specimen from Costa Rica. Holotype not found.
- NITENS Duda, 1930: 97 (*Neoolcanabates*). Type locality: "Costa Rica". Type(s) not found.

Subfamily CHLOROPINAE

Genus ANTHRACOPHAGA Loew

- ANTHRACOPHAGA Loew, 1866b: 15. Type species, *Musca strigula* Fabricius, 1794 (Coquillett, 1910: 508).

This group has been placed as a subgenus of *Chlorops* by Andersson (1977). Pending studies on the proper position of the species, the group has been left here as distinct.

Ref. — Duda, 1930: 113 (key).

- CUNEATA (Becker), 1912a: 156 (*Leptotrigonum*). Type locality: Argentina, Salta, Metan. N. COMB.

INSIGNIS Becker, 1912a: 144. Type locality: Peru, Chanchamayo.

LONGICEPS Duda (as "Becker"), 1930: 119. Type locality: Not stated [Paraguay, Chaco, at Río Negro, from lectotype ♀, here designated, in Vienna].



Genus CHLOROPS Meigen

CHLOROPS Meigen, 1803: 278. Type species, *Musca pumilionis* Bjerkander, 1778 (I. C. Z. N., 1955: 423).

Chlorops yungas is not a true *Chlorops*, but in the absence of information it is left as described.

FERRUGINEUS Duda, 1930: 122. Type locality: "Cordilleren" [Colombia, from presumed holotype in Berlin].

OBSCURICORNIS Loew, 1863: 49 (Cent. 3, no. 90). Type locality: U.S.A., D.C., Washington. Distr. — Nearctic (widespread), Mexico (Veracruz).

UNICORNIS Giglio-Tos, 1893: 14, 1895: 65. Type locality: "Mexico".

YUNGAS Sabrosky & Paganelli, NOM. NOV.

maculata Duda, 1930: 124 (preocc. Walker, 1849). Type locality: Not stated [Bolivia, Yungas de Coroico, from presumed holotype in Dresden].

maculosa Duda, 1931: 171 (*maculata*, 1930, said to be an oversight, but *maculosa* is also preoccupied, by Loew, 1872).

Genus DIPLOTOXA Loew

DIPLOTOXA Loew, 1863: 54 (Cent. 3, no. 97). Type species, *Chlorops versicolor* Loew, 1863 (mon.).

GLABRICOLLIS (Thomson), 1869: 604 (*Chlorops*). Type locality: Argentina, Buenos Aires. Distr. — Ecuador, Peru, Bolivia, Chile, Argentina, Uruguay.

RUFOMARGINATA Duda, 1930: 120. Type locality: Not stated [Peru, Puno on Lake Titicaca, from lectotype ♂, here designated, in Dresden, and Cuzco, Juliaca, Sicuani, Lake Titicaca; Bolivia, Guaqui on Lake Titicaca, from type series in Berlin, Dresden, Eberswalde, London and Vienna]. Distr. — Peru, Bolivia.

UNICOLOR Becker, 1912a: 41. Type locality: U.S.A., California, Three Rivers. Distr. — western U.S.A., Mexico (México, Oaxaca, Zacatecas).

versicolor (Loew). Not Neotropical. The Havana, Cuba record of Becker, 1912a: 137, is an error for Havana, Illinois.

Genus ECTECEPHALA Macquart

ECTECEPHALA Macquart, 1851: 253 (1851: 280). Type species, *albistylum* Macquart, 1851 (orig. des.).

Leptotrigonum Becker, 1912a: 155. Type species, *Platycephala obscura* Schiner, 1868 (Sabrosky, 1941b: 743).

Most of the species fall into one of two groups, an *albistylum* group (*Ectecephala* s. str.) with narrow frontal triangle and elongated antennae, and a *capillata* group with broad frontal triangle and shorter antennae, but there are intermediate species.



ALBISTYLUM Macquart, 1851: 253 (1851: 280). Type locality: "North America".
Distr. — Widespread, U.S.A., Mexico (Veracruz, Nayarit). Specimens
from Bolivia and Argentina recorded as *albistylum* and reidentified by
Duda as *transatlantica* are neither species.

albistyla of Duda, 1930: 115.

ARGENTINA (Malloch), 1934: 400 (*Chlorops*). Type locality: Argentina, Misiones,
"Bompland" (Bonpland). N. COMB.

BOLIVIENSIS (Becker), 1912a: 149 (*Chlorops*). Type locality: "Bolivia".
N. COMB.

BORGMEIERI Duda, 1930: 115. Type locality: "Brazil" [Rio de Janeiro, Petró-
polis, from presumed holotype in São Paulo].

BRASILIENSIS Becker, 1912a: 154. Type locality: Brazil, Santa Catarina, lectotype
♂, here designated, in Warsaw. Distr. — Brazil, Paraguay.

CAPILLATA (Coquillett), 1904b: 98 (*Chlorops*). Type locality: Nicaragua, Gra-
nada. Distr. — Mexico and Central America.

COERULEIFRONS (Frey), 1918: 22 (*Leptotrigonum*). Type locality: Brazil, Rio
de Janeiro. N. COMB.

CONFLUENS Becker, 1916: 447. Type locality: Bolivia, Coroico.

FLAVESCENS (Duda), 1930: 114 (*Leptotrigonum*). Type locality: Not stated
[Argentina, Salta, from presumed holotype in Berlin]. N. COMB.

HIRTA (Duda), 1930: 114 (*Leptotrigonum*). Type locality: Not stated [Peru,
Rio Ucayali, from presumed holotype in Dresden]. N. COMB.

INTERMEDIA Duda, 1930: 116. Type locality: Not stated [Belize, Manatee, from
presumed holotype in Berlin].

LAEVIFRONS Becker, 1912a: 153. Type locality: Paraguay, San Bernardino,
lectotype ♀ (here designated) in Budapest. The species was originally
described from a mixture of Nearctic (Kansas, Texas) and Neotropical
specimens (Paraguay, Argentina).

NITIDIFRONS Duda, 1930: 116. Type locality: Not stated [Brazil, Rio de Janeiro,
Petrópolis, from lectotype ♀, here designated, in São Paulo].


OBSCURA (Schiner), 1868: 245 (*Platycephala*). Type locality: "South America"
[Venezuela, from presumed holotype in Vienna]. Distr. — Mexico
(Oaxaca, Puebla), Guatemala, Costa Rica, Panama, Peru, Brazil (Santa
Catarina), Paraguay, Argentina.

PICTA Duda, 1930: 115. Type locality: Not stated [Argentina, Tucumán, from
presumed holotype in Budapest].

similis Becker. Nearctic.

TENUISETA Duda, 1930: 115. Type locality: Not stated [Bolivia, Mapiri, Sa-
rampiuní, 700 m., from presumed holotype in Dresden].

TINCTA Becker, 1912a: 153. Type locality: Brazil, São Paulo.



TRANSATLANTICA (Schiner), 1868: 245 (*Platycephala*). Type locality: "South America". Distr. — Bolivia.

TRIPUNCTATA Aldrich, 1923: 22. Type locality: Costa Rica, Talamanca.

Genus ELLIPONEURA Loew

ELLIPONEURA Loew, 1869a: 44 (Cent. 8, no. 79). Type species, *debilis* Loew, 1869 (mon.).

Unidentified spp. Belize, Costa Rica, Panama, Galapagos Is., Jamaica, Ecuador, Brazil.

Genus HOMALUROIDES Sabrosky

HOMALUROIDES Sabrosky, 1980b: 426. Type species, *Chlorops gramineus* Coquillett, 1898 (orig. des.).

Homalura of Becker, 1912a, Frey, 1918, and Duda, 1930, for Neotropical species.

CHAMAELEON (Becker), 1912a: 141 (*Homalura*). Type locality: "Brazil" [Salto Grande, now in Uruguay, Prov. de Salto, from holotype in Berlin]. N. COMB.

disciventris Enderlein, 1911: 124 (*Homalura*). Type locality: Madagascar (listed in error as Brazil by Becker, 1912a: 122).

DISTINCTUS (Becker), 1912a: 141 (*Homalura*). Type locality: Argentina, Tucumán. N. COMB.

GIGANTEUS (Duda), 1930: 117 (*Homalura*). Type locality: "Brazil" [Paraná, Ti-bagi, from presumed holotype in Berlin]. N. COMB.

INCOMPLETUS (Becker), 1912a: 140 (*Homalura*). Type locality: "Paraguay", lectotype ♀, here designated, in Budapest. N. COMB.

MENDOZAE (Malloch), 1934: 400 (*Chlorops*). Type locality: Argentina, Mendoza. Distr. — Argentina (Córdoba, Mendoza). N. COMB.

MEXICANUS (Duda), 1930: 117 (*Homalura*). Type locality: Not stated, but inferred as "Mexico" [Orizaba, from presumed holotype in Vienna]. N. COMB.

PUNCTATUS (Duda), 1930: 117 (*Siphilus*). Type locality: Not stated [Uruguay, from lectotype ♀, here designated, in Vienna]. N. COMB.

QUINQUEPUNCTATUS (Loew), 1863: 51 (Cent. 3, no. 94) (*Chlorops*). Type locality: U.S.A., Nebraska. Distr. — eastern U.S., Mexico.

RUFIPECTUS (Becker), 1912a: 147 (*Chlorops*). Type locality: "Paraguay". N. COMB.

laevis Becker, 1912a: 142 (*Siphilus*). Type locality: Paraguay, Asunción. N. SYN., N. COMB.

STENROTHI (Frey), 1918: 20 (*Homalura*). Type locality: Argentina, Misiones, near Bonpland. Distr. — Argentina (Corrientes, Misiones). N. COMB.



Genus MEROMYZA Meigen

- MEROMYZA Meigen, 1830: 163. Type species, *Musca saltatrix* Linnaeus, 1761, (Macquart, 1835: 589).
- COMMUNIS Fedoseeva, 1971: 913. Type locality: U.S.A., North Dakota, White Lake. Distr. — Western U.S.A., Mexico (Sonora).
- OPACULA Fedoseeva, 1978: 912. Type locality: Mexico, Chiapas, 3 miles sw Lomatan.
- PILOSA Fedoseeva, 1971: 919. Type locality: U.S.A., California, Woodfords. Distr. — Western U.S.A., Mexico (Sonora).

Genus NEODIPLOTOXA Malloch


- NEODIPLOTOXA Malloch, 1914: 116. Type species, *Chlorops nigricans* Loew, 1872 (orig. des.).
- Merectecephala* Duda, 1930: 113, 117. Type species, *Meromyza albiseta* Becker, 1912 (Sabrosky, 1941b: 743). Ref. — Duda, 1930 (key).
- ALBISETA (Becker), 1912a: 136 (*Meromyza*). Type locality: Chile, Arica, lectotype ♀, here designated, in Dresden. Distr. — Peru, Chile. N. COMB.
- MEXICANA (Duda), 1930: 118 (*Merectecephala*). Type locality: "Mexico", lectotype ♂, here designated, in Vienna. N. COMB.
- PULCHRIPES (Loew), 1872: 108 (Cent. 10, no. 96) (*Chlorops*). Type locality: U.S.A. (Texas). Distr. — Canada (Manitoba) to sw U.S.A., Mexico (Distrito Federal, Jalisco).

Genus PARECTECEPHALA Becker

- PARECTECEPHALA Becker, 1910: 105. Type species, *Oscinis longicornis* Fallén, 1820 (Duda, 1933b: 208).

The species are here retained in the traditional New World usage of *Parectecephala*, pending revisionary studies.

- BREVIS (Becker), 1916: 446 (*Chlorops*). Type locality: Colombia, Baranquilla. N. COMB.
- COSTARICANA (Duda), 1930: 126 (*Chlorops brevis* var.). Type locality: Costa Rica (inferred from name) [San José, presumably Farm La Caja, from presumed holotype in Hamburg, now destroyed]. N. COMB.
- DORSALIS (Becker), 1912a: 143 (*Anthracophaga*). Type locality: "Brazil" [São Paulo, from lectotype ♀, here designated, in Milan]. N. COMB.
- GUNDLACHI (Loew), 1872: 107 (Cent. 10, no. 93) (*Chlorops*). Type locality: Cuba. N. COMB.
- trivittata* Williston, 1896: 425 (*Chlorops*). Type locality: S. Vincent [Leeward side, from lectotype ♀, here designated, in London]. N. SYN.



HUMERALIS (Duda), 1930: 124 (*Chlorops*). Type locality: "Brazil" [Rio de Janeiro, Petrópolis, from lectotype ♂, here designated, in São Paulo]. N. COMB.

HYALIPENNIS Duda, 1930: 126. Type locality: "Brazil" [Rio Negro, from lectotype ♀, here designated, in Berlin].

INDISTINCTA Duda, 1930: 121. Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed holotype in Hamburg, now destroyed].

MACULIFRONS Becker, 1912a: 155. Type locality: Brazil, Santa Catarina. Distr. — Brazil (Rio de Janeiro, Santa Catarina).

MACULOSA (Loew), 1872: 111 (Cent. 10, no. 99) (*Chlorops*). Type locality: U.S.A., Texas, lectotype ♀, here designated, in Cambridge. Distr. — Southwestern U.S.A., Mexico (Chihuahua).

MICROPS Duda, 1930: 126. Type locality: "Brazil" [Rio de Janeiro, Petrópolis, from lectotype ♀, here designated, in Berlin].

OBSCURATA (Duda), 1930: 125 (*Chlorops*). Type locality: Not stated [Costa Rica, Suiza de Turrialba, from presumed holotype in Budapest].

SCHINERI Becker, 1912a: 154. Type locality: "South America" [Venezuela, from presumed holotype in Vienna].

transatlantica Schiner, 1868, in part (*Platycephala*)

Genus PSEUDOPACHYCHAETA Strobl

Diplotoxa, subg. PSEUDOPACHYCHAETA Strobl, 1902: 500. Type species, *pachycera* Strobl, 1902 (mon.).

APPROXIMATONERVIS (Zetterstedt), 1848: 2622 (*Oscinis*). Type locality: Scandinavia. Distr. — Holarctic, Mexico (Baja California).

microcera Loew, 1872: 108 (Cent. 10, no. 95) (*Chlorops*). Type locality: U.S.A., Texas.

Unidentified species. Colombia, Ecuador.

Genus PSILOCHLOROPS Duda

PSILOCHLOROPS Duda, 1930: 113. Type species: *clavitibia* Duda, 1931 (sub. des. and sub. mon., Duda, 1931: 167).

CLAVITIBIA Duda, 1931: 168. Type locality: Bolivia, Mapi, Lorenzopata.

Genus STELEOCERELLUS Frey

Mepachymerus, subgenus STELEOCERELLUS Frey, 1961: 35. Type species, *Steleocerus tenellus* Becker, 1910 (orig. des.).

Steleocerellus Frey; Andersson, 1977: 134. Generic status.

ABERRANS (Schiner), 1868: 244 (*Elachiptera*). Type locality: "Colombia" [Venezuela, from lectotype ♀, here designated, in Vienna]. N. COMB.



FLAVIFRONS (Frey), 1918: 19 (*Steleocerus neotropicus* var.). Type locality: Brazil, Rio de Janeiro. N. COMB.

NEOTROPICUS (Becker), 1912a: 178 (*Steleocerus*). Type locality: Peru, mouth of Río Pachitea. N. COMB.

RUFUS (Duda), 1930: 119 (*Steleocerus*). Type locality: Not stated [Peru, Río Urubamba, from presumed holotype in Dresden]. N. COMB.

Genus THAUMATOMYIA Zenker

THAUMATOMYIA Zenker, 1833: 344. Type species, *prodigiosa* Zenker, 1833 (mon.) = *Thaumatomyia notata* (Meigen).

Chloropisca Loew, 1866b: 79. Type species, *Chlorops glabra* Meigen, 1830 (Coquillett, 1910: 522).

Two species, *columbiana*, and *megacera*, are atypical but we leave them here for the present.

APACHE Sabrosky, 1943: 111. Type locality: U.S.A., Arizona, Chiricahua Mts. Distr. — U.S.A. (Arizona to Texas), Mexico (Chihuahua, Sonora, Puebla).

sulfurifrons, authors, not Duda.

APPROPINQUA (Adams), 1903: 39 (*Chlorops*). Type locality: U.S.A., Colorado, Kansas, and Wyoming. Distr. — southwest Canada, western U.S.A., Mexico (Chihuahua, Durango, San Luis Potosí, Zacatecas).

COLUMBIANA (Becker), 1912a: 149 (*Chlorops*). Type locality: Colombia, Fusagasugá. N. COMB.

strigata Duda, 1930: 122 (*Chloropisca*). Type locality: Not stated [Peru, Chanchamayo, from presumed holotype in Dresden]. N. SYN., N. COMB.

GEMINA (Becker), 1912a: 139 (*Chloropisca*). Type locality: Bolivia, Yungas de Coroico, 1200 m., and Brazil, São Paulo. Distr. — Bolivia, Brazil (Santa Catarina, São Paulo), Argentina (Misiones, Tucumán). N. COMB.

rufithorax Duda, 1930: 122 (*Chloropisca*). Type locality: "Bolivia" [Error? The only specimen found, and labeled new species by Duda, is from Peru, Meshagua, Río Urubamba, in Dresden]. N. SYN., N. COMB.

GLABRA (Meigen), 1830: 149 (*Chlorops*). Type locality: Europe. Distr. — Holarctic, Mexico (Aguascalientes, Distrito Federal, Hidalgo, México, Querétaro).

GLABRINA (Becker), 1912a: 138 (*Chloropisca*). Type locality: Peru, Cuzco (lectotype ♂, here designated, in Dresden), and near Urubamba, 3000 m. Distr. — Costa Rica, Colombia, Ecuador. Peru, Argentina (Tucumán). N. COMB.

MEGACERA (Becker), 1912b: 645 (*Chlorops*; nom. nov. for *grandicornis* Becker, 1912). Type locality: Bolivia, Mapi, Sarampiuní, 700 m. N. COMB.

- grandicornis* Becker, 1912a: 147 (*Chlorops*; preocc. Strobl, 1909). Type locality: Bolivia, Mapiri, Sarampiuní, 700 m. N. COMB.
- NIGRESCENS (Duda), 1930: 123 (*Chloropisca*). Type locality: "Mexico" [Puebla, from presumed holotype in Vienna]. Distr. — Mexico (México, Puebla). N. COMB.
- PULLA (Adams), 1904a: 303 (*Chlorops*). Type locality: U.S.A., Colorado. Distr. — S. Canada, U.S.A., Mexico (Veracruz).
- SULFURIFRONS (Duda), 1930: 124 (*Chloropisca*). Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed holotype ♂ (Sabrosky, 1943: 112), now to be interpreted as lectotype in view of a recently discovered presumed syntype in Berlin; the lectotype was in Hamburg, now destroyed]. Distr. — Mexico (Jalisco, Michoacán, San Luis Potosí, Veracruz), Costa Rica, Colombia, Ecuador.

Genus TRIGONOMMA Enderlein

- TRIGONOMMA Enderlein, 1911b: 124. Type species, *lippulum* Enderlein, 1911 (orig. des.).
- Beckerella* Enderlein, 1911b: 122. Type species, *speculifrons* Enderlein, 1911 (orig. des.). N. SYN.
- Epichlorops* Becker of Duda, 1931.
- ALBIPES (Wiedemann), 1830: 595 (*Chlorops*). Type locality: "South America". Distr. — Honduras, Costa Rica, Panama, Colombia, Guyana, Surinam, Trinidad, Peru. N. COMB.
- ATRICEPS (Duda), 1930: 127 (*Chlorops*). Type locality: Not stated [Costa Rica, Farm La Caja, near San José, from presumed type series in Hamburg, now destroyed, and lectotype ♀, here designated, in Berlin]. N. COMB.
- BISULCATUM (Enderlein), 1911a: 208 (*Tricimba*). Type locality Brazil, Santa Catarina. N. COMB.
- CINEREOMICANS (Frey), 1918: 21 (*Chlorops*). Type locality: Brazil, Rio de Janeiro. Distr. — Argentina (Tucumán), Brazil (Santa Catarina, Rio de Janeiro). N. COMB.
- COERULEIFRONS (Becker), 1912a: 150 (*Chlorops*). Type locality: Brazil, Santa Catarina. N. COMB.
- FLAVIFRONS (Duda), 1930: 128 (*Chlorops* or *Epichlorops*). Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed holotype in Hamburg, now destroyed]. N. COMB.
- FLAVIPES (Duda): 126 (*Chlorops*). Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed type series in Hamburg, now destroyed, and lectotype ♀, here designated, in Berlin]. N. COMB.
- FOLIATUM (Becker), 1912a: 219 (*Oscinella*). Type locality: Brazil, Santa Catarina, lectotype ♀, here designated, in Warsaw. N. COMB.

FOSSULATUM (Loew), 1863: 43 (Cent. 3, no. 82) (*Chlorops*). Type locality: "Cuba". Distr. — U.S.A. (Texas), Cuba, Jamaica, Puerto Rico, Virgin Islands, Panama.

atra Curran, 1926: 3 (*Chloropisca*). Type locality: Puerto Rico, Arceibo.

FOVEOLATUM (Duda), 1930: 128 (*Chlorops* or *Epichlorops*). Type locality: Not stated [Paraguay, San Bernardino, from presumed holotype in Vienna]. N. COMB.

LIPPULUM Enderlein, 1911b: 124. Type locality: Brazil, Santa Catarina. Distr. — Brazil (Rio de Janeiro, Santa Catarina).

NIGRUM (Becker), 1916: 447 (*Chlorops*). Type locality: Trinidad, Port of Spain. N. COMB.

PUBICOLLIS (Becker), 1912a: 148 (*Chlorops*). Type locality: Peru, Arequipa, lectotype ♀, here designated, in Dresden, and Chile, Quillota. N. COMB.

SPECULIFRONS (Enderlein), 1911b: 122 (*Beckerella*). Type locality: Brazil, Santa Catarina, lectotype ♀, here designated, in Berlin. N. COMB.

TENUIFRONS (Duda), 1930: 126 (*Chlorops*). Type locality: "Costa Rica" [Farm La Caja, near San José, from presumed syntypes in Hamburg, now destroyed]. N. COMB.

VERRUCOSUM (Duda), 1930: 128 (*Chlorops* or *Epichlorops*). Type locality: "Peru" [Pinipini, from presumed holotype in Dresden]. N. COMB.

Species *incertae sedis*, CHLOROPINAE

BRASILIENSIS Walker, 1853: 414 (*Chlorops*). Type locality: "Brazil". Placed by Duda (1930: 123) as *Chloropisca*, but the description is unrecognizable.

NIGELLUS Wiedemann, 1830: 595 (*Chlorops*). Type locality: "South America". Description unrecognizable and type lost. Possibly *Trigonomma*.

SPECIES TRANSFERRED TO OTHER FAMILIES


DIMIDIATUS Wiedemann, 1830: 597 (*Chlorops*). Type locality: Surinam. To *Milichiella* in Milichiidae (see Sabrosky, 1973, Family 75).

FLAVUS Wiedemann, 1830: 595 (*Chlorops*). Type locality: Uruguay, Montevideo. To Lauxaniidae, type species of *Physoclypeus* Hendel.

MELANOSOMA Brèthes, 1924: 110 (*Somillus*). Type locality: Chile, Isla Chiloé. The genus and species belong to Dolichopodidae.

PORTERI Brèthes, 1925: 35 (*Hippelates*). Type locality: Chile, San José de Maipo. To Ephyridae, synonym of *Discocerina obscurella* (Fallén) [ident. W. W. Wirth, W. M. Mathis].

VITTATUS Wiedemann, 1830: 594 (*Chlorops*). Type locality: "West Indies." Probably Lauxaniidae. Type lost.



LITERATURE CITED

ADAMS, C. F.

1903. Dipterological contributions. *Kans. Univ. Sci. Bul.* 2: 21-47.
- 1904a. Descriptions of new Oscinidae. *Ent. News* 15: 303-304.
- 1904b. On the North American species of *Siphonella*. *Psyche* 11: 103-104.
- 1905a. On the North American species of *Oscinis*. *Ent. News* 16: 108-111.
- 1905b. Diptera africana. I. *Kans. Univ. Sci. Bul.* 3: 149-208.
1908. Notes on North American species of *Crassiseta* v. Ros. *Jour. N. Y. Ent. Soc.* 16: 151-152.

ALDRICH, J. M.

1923. A new sugarcane miner. *Bull. Brooklyn Ent. Soc.* 18: 22-23.
1924. A new genus and species of two-winged flies of the family Chloropidae injuring manihot in Brazil. *Proc. U. S. Nat. Mus.* 65 (Art. 21): 1-2.
1929. Three new acalyptrate Diptera. *Proc. Ent. Soc. Wash.* 31: 89-91.
1931. Notes on *Hippelates* (Diptera: Chloropidae), with a new Brazilian species. *Proc. Ent. Soc. Wash.* 33: 69-72.

ANDERSSON, H.

1977. Taxonomic and phylogenetic studies on Chloropidae (Diptera) with special reference to Old World genera. *Ent. Scand., Suppl.* 8: 200 pp.

BECKER, T.

1909. Collections recueillies par M. Maurice de Rothschild dans l'Afrique orientale anglaise. Insectes: Diptères nouveaux. *Bul. Mus. Nat. d'Hist. Nat. (Paris)* 15: 113-121.
1910. Chloropidae. Eine monographische Studie. I. Teil. Paläarktische Region. *Arch. Zool. Budapest.* 1: 33-174, pls. 2-3.
- 1912a. Chloropidae. Eine monographische Studie. IV. Teil. Nearktische Region. V. Teil. Neotropische Region. *Ann. Hist. Nat. Magyar Nemzeti Muz.* 10: 21-256, 2 figs., 1 pl.
- 1912b. Berichtigungen zur Monographie der Chloropiden. *Ann. Hist. Nat. Magyar Nemzeti Muz.* 10: 645-646.
1916. Neue Chloropiden aus dem ungarischen National-Museum. *Ann. Hist. Nat. Magyar Nemzeti Muz.* 14: 423-453, 5 figs.
1919. Diptères. Brachycères. *Mission Serv. Géogr. de l'Armée, Arc de Méridien Équatorial en Amérique du Sud*, 10: 163-215, pls. 14-17.



BLANCHARD, E. E.

1941. Nuevos parasitos del bicho de cesto *Oeceticus kirbyi*, Guild. *Rev. Soc. Ent. Argentina* 11: 3-21, 8 figs.

BRÈTHES, J.

1924. Sur quelques Diptères Chiliens. *Rev. Chilena Hist. Nat.* 28: 104-111.
1925. Sur quelques insectes de San José de Maipo. *Rev. Chilena Hist. Nat.* 29: 34-35.

COQUILLETT, D. W.


1898. Notes and descriptions of Osecinidae. *Jour. N. Y. Ent. Soc.* 6: 44-49.
1900. Report on a collection of dipterous insects from Puerto Rico. *Proc. U.S. Natl. Mus.* 22: 249-270.
1904a. (New genera and new species). In Baker, C. F. Reports on Californian and Nevadan Diptera, I. *Invertebrata Pacifica* 1: 17-39.
1904b. New Diptera from Central America. *Proc. Ent. Soc. Wash.* 6: 90-98.
1910. The type-species of the North American genera of Diptera. *Proc. U.S. Natl. Mus.* 37: 499-647.

CURRAN, C. H.

1926. New Diptera from the West Indies. *Amer. Mus. Novitates* 220: 1-14.
1928. Insects of Porto Rico and the Virgin Islands. Diptera or two-winged flies. Pp. 1-118, 39 figs. In *New York Academy of Sciences, Scientific Survey of Porto Rico and the Virgin Islands*, Vol. XI, Part 1.
1931. First supplement to the 'Diptera of Porto Rico and the Virgin Islands'. *Amer. Mus. Novitates* 456: 1-23, 4 figs.
1932. The Norwegian Zoological Expedition to the Galapagos Islands 1925, conducted by Alf Wollebaek. IV Diptera (excl. of Tipulidae and Culicidae). *Medd. Zool. Mus. Oslo* 30: 347-366, 1 fig.

DUDA, O.

1921. *Fiebrigella* und *Archiborborus*, zwei neue südamerikanische Borboridengattungen (Dipteren). *Tijd. v. Ent.* 64: 119-146, 1 fig.
1929. Die Ausbeute der deutschen Chaco-Expedition 1925/26 (Diptera). X. Chloropidae. (Fortsetzung). *Konowia* 8: 165-169.
1930. Die neotropischen Chloropiden (Dipt.). *Folia Zool. et Hydrobiol.* 2: 46-128.
1931. Die neotropischen Chloropiden (Dipt.). 1. Fortsetzung: Nachtrag, Ergänzungen, Berichtigungen und Index. *Folia Zool. et Hydrobiol.* 3: 159-172.



- 1933a. Die neotropischen Chloropiden (Dipt.). 2. Fortsetzung. *Folia Zool. et Hydrobiol.* 5: 41-47.
- 1933b, 1934. Neue und bekannte neotropische Chloropidae (Dipt.) des U.S. National-Museums, Smithsonian Institution. *Konowia* (1933) 12: 192-209; (1934) 13: 58-69, 101-110.
1936. Ergebnisse einer zoologischen Sammelreise nach Brasilien, insbesondere in das Amazonasgebiet, ausgeführt von Dr. H. Zerny. XI. Teil. Diptera: Chloropidae. *Ann. Naturhist. Mus. Wien* 47: 107-113.

ENDERLEIN, G.

- 1911a. Klassifikation der Oscinosominen. *Sitzber. Gesell. Naturf. Freunde* 1911: 185-244, 11 figs.
- 1911b. Einige neue Gattungen und Arten aussereuropaischer Chloropinen. *Zool. Anz.* 38: 122-126, 1 fig.
1913. Dipterologische Studien. IV. Neues und altes über Chloropiden. *Zool. Anz.* 42: 352-374, 21 figs.
1938. Die Dipterenfauna der Juan-Fernandez-Inseln und der Oster-Insel. In *Skottsberg: Nat. Hist. Juan Fernandez & Easter Id. (Uppsala)*, 3 (Zool.) (Pt. V): 643-680.

FEDOSEEVA, L. I.

1971. (A revision of nearctic species of the genus *Meromyza* Mg. (Diptera, Chloropidae) (In Russian). *Ent. Oboz.* 50: 911-930, 26 figs. (In English, *Ent. Review* 50: 520-529, 26 figs.).
1978. (New data on the genus *Meromyza* Mg. (Diptera, Chloropidae) in North America). (In Russian). *Ent. Oboz.* 57: 910-914, 6 figs. (In English, *Ent. Review* 57: 630-633, 6 figs.).

FITCH, A.

1856. Report on the noxious, beneficial and other insects of the State of New York [II]. *Trans. N. Y. State Agric. Soc.* (1855) 15: 409-559.

FREY, R.

1918. Mitteilungen über südamerikanische Dipteren. *Förhandl. Finska Vetensk.-Soc.* 60, Afd. A, no. 14: 1-35, 1 pl.
1961. Neue orientalische Dipteren. *Notulae Ent.* 41: 33-35.

GIGLIO-TOS, E.

1893. Diagnosi di nuovi generi e di nuove specie di Ditteri. IX. *Boll. Mus. Zool. Anat. Compar. Torino* 8(158): 1-14.
1895. *Ditteri del Messico*, Parte IV. 74 pp., 1 pl.



GRIOT, M., R. SILBERMANN, A. ICART, & A. FREIBERG

1946. Revisión bibliográfica sobre el bicho de cesto. *Rev. Argentina de Agron.* 13: 310-330.

HALL, D. G.

1937. The North and Central American spider parasites of the genus *Pseudogaurax* (Diptera: Chloropidae). *Jour. Wash. Acad. Sci.* 27: 255-261, 7 figs.

HENDEL, F.

1907. Nomina nova für mehrere Gattungen der acalyptraten Musciden. *Wien Ent. Ztg.* 26: 98.

ICHES, L.

1906. Sobre cinco Dipteros nuevos del Chaco Austral. *Bol. Min. Agric., Argentina* 6: 262-273.

I. C. Z. N. (INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE)

1955. Opinion 348. Suppression, under the Plenary Powers, of the generic name *Titania* Meigen, 1800, for the purpose of validating the generic name *Chlorops* Meigen, 1803 (Class Insecta, Order Diptera). *Opinions and Declarations rendered by the International Commission on Zoological Nomenclature* 10: 421-436.
1978. Opinion 1100. Designation of *Musca frit* Linnaeus, 1758, as type-species of *Oscinella* Becker, 1909 (Diptera, Chloropidae). *Bull. Zool. Nomencl.* 34: 203-204.

KUMM, H. W.

1936. The Jamaican species of *Hippelates* and *Oscinella* (Diptera, Chloropidae). *Bul. Ent. Research* 27: 307-329, 1 pl., 6 figs.

LINNAEUS, C.

1758. *Systema naturae* (ed. X). 824 pp.


LIOY, P.

1864. I ditteri distribuiti secondo un nuovo metodo di classificazione naturale. (part). *Atti I. R. Ist. Veneto di Sci., Let. ed Arti*, ser. 3, 9: 1087-1126.

LOEW, H.

1863. *Diptera Americae septentrionalis indigena*. Centuria tertia. *Berl. Ent. Ztschr.* 7: 1-55.

81.46



- 1866a. Ibid. Centuria sexta. *Berl. Ent. Ztschr.* 9: 127-186.
- 1866b. Ueber die bisher in Schlesien aufgefundenen Arten der Gattung *Chlorops* Macq. *Ztschr. f. Ent. Breslau* 15: 3-96.
- 1869a. *Diptera Americae septentrionalis indigena*. Centuria octava. *Berl. Ent. Ztschr.* 13: 1-52.
- 1869b. Ibid. Centuria nona. *Berl. Ent. Ztschr.* 13: 129-186.
1872. Ibid. Centuria decima. *Berl. Ent. Ztschr.* 16: 49-115.
1876. Beschreibungen neuer amerikanischer Dipteren. *Ztschr. f. die Gesam. Naturw.* 48: 317-340.

MACQUART, J.


1835. *Histoire naturelle des insectes. Diptères*. Vol. 2: 703 pp., pls. 13-24.
1851. Diptères exotiques nouveaux ou peu connus. Suite du 4^e supplément. *Mem. Soc. Natl. des Sci., de l'Agr. et des Arts, Lille*, 1850: 134-294, pls. 15-28 (also as separate work).

MALLOCH, J. R.

- 1913a. Four new species of North American Chloropidae. *Insecutor Inscitiae Menstruus* 1: 60-64.
- 1913b. New North American Diptera. *Canad. Ent.* 45: 282-284.
- 1913c. The genera of flies in the subfamily Botanobiinae with hind tibial spur. *Proc. U. S. Natl. Mus.* 46: 239-266, 2 pls.
- 1913d. A new genus and two new species of Chloropidae (Diptera). *Insecutor Inscitiae Menstruus* 1: 46-48.
1914. A synopsis of the genera in Chloropidae, for North America. *Canad. Ent.* 46: 113-120.
1915. Notes on North American Chloropidae (Diptera). *Proc. Ent. Soc. Wash.* 17: 158-162.
1916. A new genus and species of North American Chloropidae (Diptera). *Bul. Brooklyn Ent. Soc.* 11: 86-87.
1918. Notes on Chloropidae, with descriptions (Diptera). *Bul. Brooklyn Ent. Soc.* 13: 19-21.
1934. Acalyptrata (concluded). pp. 393-489, pl. 8, figs. 69-84. In *British Museum (Nat. Hist.), Diptera of Patagonia and South Chile*, Part VI, Fasc. 5.

MEIGEN, J. W.

1803. Versuch einer neuen Gattungseintheilung der europäischen zweiflügeligen Insekten. *Mag. f. Insektenkunde* (Illiger's Magazin) 2: 259-281.



1830. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten*. Vol. VI: iv — 401 pp., pls. 55-66.

PAPAVERO, N.

1973. Essays on the history of Neotropical Dipterology, with special reference to collectors (1750-1905). Vol. II: iii — 217-446. São Paulo, Museu de Zoologia, Universidade de São Paulo.

PAZOS, J. H.

1914. Catalogo de los Dípteros de la Isla de Cuba. *An. Acad. Cien. Habana* 1: 990-1003.

ROBERTS, R. A.

1934. Some insects collected in Mexico, mostly in association with man and animals or animal products. *Jour. N. Y. Ent. Soc.* 42: 249-262.

RONDANI, C.

1856. *Dipterologiae Italicae prodromus*. Vol. 1: 228 pp.

SABROSKY, C. W.

1936. Some synonymy in the chloropid genera *Hippelates* and *Diplotoxa* (Diptera). *Ent. News* 47: 245-248.

1938. Taxonomic notes on the dipterous family Chloropidae. I. *Jour. N. Y. Ent. Soc.* 46: 417-434, 4 figs.

1939. Chloropidae (Diptera). pp. 177-179. In *Résultats scientifiques des croisières du Navire-École Belge "Mercator", Vol. II (Mém. Mus. Roy. d'Hist. Nat. de Belgique, ser. 2, fasc. 15)*.

1940. Twelve new North American species of *Oscinella* (Diptera, Chloropidae). *Canad. Ent.* 72: 214-230.

1941a. The *Hippelates* flies or eye gnats: preliminary notes. *Canad. Ent.* 73: 23-27.

1941b. An annotated list of genotypes of the Chloropidae of the world (Diptera). *Ann. Ent. Soc. Amer.* 34: 735-765.

1943. A revised synopsis of nearctic *Thaumatomyia* (= *Chloropisca*). (Diptera, Chloropidae). *Canad. Ent.* 75: 109-117.

1947. *Rhodesiella*: a genus new to the Western Hemisphere (Diptera: Chloropidae). *Ent. News* 58: 174-177.

1949. A new species of "*Pseudogaurax*" from Brazil (Diptera, Chloropidae). *Rev. Brasil. Biol.* 9: 33-34.

1950a. A new species of *Pseudogaurax* from Florida (Diptera, Chloropidae). *Bul. Brooklyn Ent. Soc.* 45: 33-34.

- 1950b. A new species of *Eribolus* from California (Diptera, Chloropidae). *Pan-Pacific Ent.* 26: 91-92.
- 1950c. A synopsis of the chloropid genera *Chaetochlorops* and *Eugaurax* (Diptera). *Jour. Wash. Acad. Sci.* 40: 183-188.
1951. A review of the nearctic species of *Lasiopleura* (Diptera, Chloropidae). *Canad. Ent.* 83: 336-343, 4 figs.
1955. Los insectos de las islas Juan Fernandez. 19. Chloropidae (Diptera). *Rev. Chilena de Ent.* 4: 45-49.
1961. A new nearctic species of *Stenoscinis*, with key to the species of the Western Hemisphere (Diptera, Chloropidae). *Ent. News* 72: 19-23, 1 fig.
1966. Three new Brazilian species of *Pseudogaurax* with a synopsis of the genus in the Western Hemisphere (Diptera, Chloropidae). *Papéis Avulsos Dept. Zool., Sec. Agric. São Paulo* 19: 117-127, 24 figs.
1970. The identity of *Fiebrigella* Duda (Diptera: Chloropidae). *Jour. Kans. Ent. Soc.* 43: 182-184.
1973. Family Milichiidae. Family 75: 12 pp., in *A Catalogue of the Diptera of the Americas South of the United States*. São Paulo.
1974. *Eugaurax setigena* (Diptera: Chloropidae), a new stem miner in water hyacinth. *Fla. Ent.* 57: 347-348.
- 1980a. Unexpected synonymy in Chloropidae, from the family Ephydriidae (Diptera). *Austral. Ent. Mag.* 6(6): 101-102.
- 1980b. New genera and new combinations in nearctic Chloropidae (Diptera). *Proc. Ent. Soc. Wash.* 82: 412-429, 12 figs.

SCHINER, I. R.

1868. Diptera. 388 pp., 4 pls. In *Reise der österreichischen Fregatte Novara, Zool.*, Vol. 2, Abt. 1, B.

SÉGUY, E.

1934. Étude sur quelques Muscides de l'Amérique Latine. *Rev. Soc. Ent. Argentina* 6: 9-16, 3 figs.
1940. Étude sur les Diptères Hippelatoïdes pathogènes (mouches des yeux) de la région néotropicale (Diptera Chloropidae). *Mém. Mus. Natl. d'Hist. Nat.*, n. ser., 13: 331-357, 14 figs.

STROBL, G.

1902. [New contributions on the dipterous fauna of the Balkan Peninsula] (In Serbian). *Glasnik Zemalj. Muz. Bosni Herz.* 14: 461-517.



THOMSON, C. G.

1869. *Diptera. Species novas descripsit.* pp. 443-614, pl. 9. In K. Svenska Vetenskaps-Akademien, *Kongliga svenska fregatten Eugénies resa omkring jorden*, Part 2: Zoologie, (Sec.) 1: Insekter.

TOWNSEND, C. H. T.

1895. On the Diptera of Baja California, including some species from adjacent regions. *Proc. Calif. Acad. Sci.*, ser. 2, 4: 593-620.

TUCKER, E. S.

1908. Incidental studies of new species of *Oscinis*. *Ent. News* 19: 272-274.

WALKER, F.

1853. *Insecta Saundersiana, Diptera*, Vol. 1 (Pt. 4): 253-414.
1859. Catalogue of dipterous insects collected at Makassar in Celebes, by Mr. A. R. Wallace, with descriptions of new species. *Jour. of Proc. Linn. Soc. (London)*, Zool. 4: 90-144 (cont.).

WENDT, H.

1975. Die Chloropiden-Typen der Dipteren-Sammlung des Zoologischen Museums in Berlin. *Mitt. Zool. Mus. Berlin* 51: 83-137.
1978. Die Chloropiden-Typen der Dipteren-Sammlung des Zoologischen Museums in Berlin (Nachtrag). *Mitt. Zool. Mus. Berlin* 54: 369-376.

WIEDEMANN, C. R. W.

1830. *Aussereuropäische zweiflügelige Insekten*. Vol. 2: xii-684 pp., pls. 7-10b.

WILLISTON, S. W.

1896. On the Diptera of St. Vincent (West Indies). *Trans. Ent. Soc. London* 1896: 253-446, pls. 8-14.

WOLLASTON, T. V.

1858. Brief diagnostic characters of undescribed Madeiran insects. *Ann. and Mag. Nat. Hist.*, ser. 3, 1: 18-28, 113-125, 2 pls.


ZENKER, J. C.

1833. *Miscellen. Not. aus dem Geb. der Natur- u. Heilk.* (Froriep's Notizen) 35: 344.

ZETTERSTEDT, J. W.

1848. *Diptera Scandinaviae*. Vol. 7: 2581-2934.

81.50




INDEX

aberrans (Schiner), <i>Steleocerellus</i>	81.39
<i>Acanthopeltastes</i> Enderlein	81.4
<i>aenea</i> (Duda), <i>Apallates</i>	81.5
<i>aeneiventris</i> (Becker), <i>Liohippclates</i>	81.19
<i>aeneus</i> (Duda), <i>Apallates</i>	81.5
<i>aequatorialis</i> (Becker), <i>Liohippelates</i>	81.19
<i>aequisecta</i> (Enderlein), <i>Conioscinella</i>	81.9
<i>albifacies</i> (Duda), <i>Fiebrigella</i>	81.15
<i>albipes</i> (Wiedemann), <i>Trigonomma</i>	81.41
<i>albiseta</i> (Becker), <i>Neodiptotxa</i>	81.38
<i>albistyla</i> Duda, <i>Ectecephala</i>	81.36
<i>albistylum</i> Macquart, <i>Ectecephala</i>	81.36
<i>albohirtum</i> Enderlein, <i>Coryphisoptron</i>	81.11
<i>angulicornis</i> Duda, <i>Conioscinella</i>	81.9
<i>angulicornis</i> (Malloch), <i>Hippelates</i>	81.16
<i>angustata</i> (Enderlein), <i>Notaulacella</i>	81.24
<i>angustibucca</i> (Duda), <i>Olcella</i>	81.25
<i>angustibuccus</i> (Duda), <i>Liohippelates</i>	81.20
<i>annotatus</i> Becker, <i>Hippelates</i>	81.17
<i>annulatus</i> Enderlein, <i>Hippelates</i>	81.17
<i>annulicrus</i> (Duda), <i>Oscinoides</i>	81.28
<i>annulifemur</i> Duda, <i>Conioscinella</i>	81.9
<i>anonyma</i> (Williston), <i>Monochaetoscinella</i>	81.24
<i>Anoscinella</i> Becker	81.13
<i>Anthrachophaga</i> Loew	81.34
<i>antiguanus</i> (Duda), <i>Apallates</i>	81.5
<i>antiguensis</i> (Duda), <i>Stenoscinis</i>	81.32
<i>apache</i> Sabrosky, <i>Thaumatomyia</i>	81.40
<i>Apallates</i> Sabrosky	81.5
<i>apicale</i> (Williston), <i>Onychaspidium</i>	81.27
<i>apicatus</i> (Malloch), <i>Liohippelates</i>	81.19
<i>Apotropina</i> Hendel	81.3
<i>appropinqua</i> (Adams), <i>Thaumatomyia</i>	81.40
<i>approximatonervis</i> (Zetterstedt), <i>Pseudopachychaeta</i>	81.39
<i>arcuata</i> Duda, <i>Conioscinella</i>	81.9
<i>argentina</i> (Malloch), <i>Ectecephala</i>	81.36
<i>argentinica</i> (Duda), <i>Conioscinella</i>	81.10
<i>armillata</i> Enderlein, <i>Notaulacella</i>	81.24
<i>atra</i> (Curran), <i>Trigonomma</i>	81.42
<i>atriceps</i> (Duda), <i>Trigonomma</i>	81.41
<i>attenuata</i> (Adams), <i>Elachiptera</i>	81.13
<i>attingens</i> (Duda), <i>Monochaetoscinella</i>	81.24
<i>australis</i> (Malloch), <i>Olcella</i>	81.25
<i>australis</i> (Sabrosky), <i>Liohippelates</i>	81.19
<i>Baseoneura</i> Duda	81.24
<i>Beckerella</i> Enderlein	81.41
<i>beckeri</i> (Enderlein), <i>Conioscinella</i>	81.9
<i>benoisti</i> (Séguy), <i>Hippclates</i>	81.17
<i>bicolor</i> (Coquillett), <i>Liohippelates</i>	81.19




bicolor Malloch, Eugaurax	81.14
Biorbitella Sabrosky	81.8
bipunctatus Malloch, Eugaurax	81.14
bishoppi (Sabrosky), Liohippелates	81.19
bisulcatum (Enderlein), Trigonomma	81.41
blanda Becker, Oscinella	81.28
<i>blanda</i> group, Oscinella	81.28
boliviensis (Becker), Ectecephala	81.36
boliviensis Duda, Lioscinella	81.22
boliviensis Sabrosky, Fiebrigella	81.15
borgmeieri Duda, Ectecephala	81.36
<i>Botanobia</i> Lioy	81.15
bracata (Duda), Oscinella	81.28
bracatus (Duda), Hippelates	81.17
<i>brasilienis</i> (Aldrich), Liohippелates	81.20
brasilienis Becker, Ectecephala	81.36
<i>brasilienis</i> Duda, Hippelates	81.17
brasilienis Walker, Chlorops (<i>incertae sedis</i>)	81.42
brevipila Duda, Tricimba	81.32
brevis (Becker), Parectecephala	81.38
brevistylus (Malloch), Liohippелates	81.19
breviventris (Becker), Fiebrigella	81.15
breviventris Malloch, Tricimba	81.33
brumpti Séguy, Hippelates	81.17
<i>buccalis</i> (Duda), Apallates	81.7
Cadrema Walker	81.9
calcaratus (Enderlein), Olcanabates	81.25
californicus Sabrosky, Eribolus	81.14
capax (Coquillett), Apotropina	81.3
capillata (Coquillett), Ectecephala	81.36
carbonaria (Loew), Oscinella	81.27
catharinae Sabrosky & Paganelli, Hippelates	81.17
catharinensis Enderlein, Oscinosoma (<i>incertae sedis</i>)	81.34
caviventris (Enderlein), Tricimba	81.33
Chaethippus Duda	81.9
Chaetochlorops Malloch	81.9
chamaeleon (Becker), Homaluroides	81.37
chilenica Duda, Conioscinella	81.10
CHLOROPINAE	81.34
<i>Chloropisca</i> Loew	81.40
Chlorops Meigen	81.35
cinerea (Loew), Olcella	81.25
cinereomicans (Frey), Trigonomma	81.41
cingulatus Sabrosky, Pseudogaurax	81.29
circularis (Becker), Acanthopeltastes	81.4
<i>circumdata</i> (Becker), Liohippелates	81.21
circumdata (Duda), Apotropina	81.3
clavitibia Duda, Psilochlorops	81.39
coeruleifrons (Becker), Trigonomma	81.41
coeruleifrons (Duda), Malloewia	81.23
coeruleifrons (Frey), Ectecephala	81.36
<i>collusor</i> (Curran), Liohippелates	81.20
collusor (Townsend), Liohippелates	81.20



columbiana (Becker), Thaumatomyia	81.40
columbianus (Enderlein), Apallates	81.5
communis Fedoseeva, Meromyza	81.38
compta (Duda), Notaulacella	81.24
concinna (Williston), Oscinella	81.28
confluens (Becker), Conioscinella	81.10
confluens Becker, Ectecephala	81.36
confusa (Malloch), Biorbitella	81.8
Conioscinella Duda	81.9
coniotrigona Duda, Elachiptera	81.13
convexa (Malloch), Olcella	81.25
convexiformis (Duda), Apallates	81.6
convexus (Loew), Apallates	81.5
Coryphisoptron Enderlein	81.11
costaricana (Duda), Biorbitella	81.8
costaricana (Duda), Parectecephala	81.38
costaricanus (Duda), Apallates	81.6
costaricanus (Duda), Gaurax	81.15
costaricanum (Duda), Coryphisoptron	81.11
costata (Loew), Elachiptera	81.13
coxendix (Fitch), Apallates	81.6
coxendix of Curran, Apallates	81.6
crassipes (Aldrich), Fiebrigella	81.15
crassisetia (Duda), Tricimba	81.32
cricopus (Enderlein), Conioscinella	81.10
cricopus Enderlein, part, Tricimba	81.32
cuncata (Becker), Anthracophaga	81.34
currani (Aldrich), Liohippelates	81.20
curvinervis (Duda), Acanthopeltastes	81.4
curvinervis Duda, Conioscinella (<i>incertae sedis</i>)	81.34

<i>Dactylothyrea</i> of Becker, Duda	81.4
dampfi Aldrich, Oscinella	81.27
delicata (Becker), Fiebrigella	81.15
dentigena (Enderlein), Parasiphonella	81.28
dentitibia (Enderlein), Lioscinella	81.22
dimidiata (Becker), Conioscinella	81.10
dimidiatus Wiedemann, Chlorops	81.42
Diplotoxa Loew	81.35
disciventre (Enderlein), Coryphisoptron	81.11
disciventris Enderlein, Homalura	81.37
Discogastrella Enderlein	81.12
dissidens (Tucker), Apallates	81.6
dissidens group, Apallates	81.5
distinctus (Becker), Homaluroides	81.37
diversipes (Curran), Apallates	81.6
dorsalis (Becker), Parectecephala	81.38
dorsalis Loew, Hippelates	81.17
dorsata (Loew), Incertella	81.18
dorsatus Williston, Hippelates	81.17
dubiosa (Becker), Elachiptera	81.13
dubium (Duda), Coryphisoptron	81.12



<i>Ectecephala</i> Macquart	81.35
<i>Ectropa</i> Schiner	81.3
<i>Elachiptera</i> Macquart	81.12
<i>Elliponeura</i> Loew	81.37
<i>enderleini</i> Duda, <i>Tricimba</i>	81.32
<i>Enderleiniella</i> Becker	81.13
<i>Epichlorops</i> Becker of Duda	81.41
<i>equalis</i> (Williston), <i>Goniaspis</i>	81.16
<i>Eribolus</i> Becker	81.14
<i>escomeli</i> (Séguy), <i>Liohippелates</i>	81.20
<i>Eugaurax</i> Malloch	81.14
<i>eunota</i> (Loew), <i>Melanochaeta</i>	81.23
<i>exlineatus</i> Becker, <i>Hippelates</i>	81.18
<i>fasciata</i> (Duda), <i>Olcella</i>	81.25
<i>fasciventris</i> (Duda), <i>Eugaurax</i>	81.14
<i>femoralis</i> (Duda), <i>Liohippелates</i>	81.20
<i>femorata</i> (Duda), <i>Liohippелates</i>	81.20
<i>festina</i> (Becker), <i>Elachiptera</i>	81.13
<i>ferrugineus</i> Duda, <i>Chlorops</i>	81.35
<i>Fiebrigella</i> Duda	81.14
<i>fiebrigi</i> (Duda), <i>Notaulacella</i>	81.24
<i>flava</i> Duda, <i>Strandimyia</i>	81.32
<i>flavescens</i> (Duda), <i>Ectecephala</i>	81.36
<i>flavibucca</i> Duda, <i>Liohippелates</i>	81.20
<i>flaviceps</i> (Enderlein), <i>Coryphisoptron</i>	81.12
<i>flaviceps</i> (Loew), <i>Liohippелates</i>	81.20
<i>flavicorne</i> Duda, <i>Rhopalopterum</i>	81.31
<i>flavida</i> Williston, <i>Elachiptera</i>	81.13
<i>flavidulus</i> (Malloch), <i>Olcanabates</i>	81.25
<i>flavifrons</i> (Duda), <i>Coryphisoptron</i>	81.12
<i>flavifrons</i> (Duda), <i>Trigonomma</i>	81.41
<i>flavifrons</i> (Frey), <i>Steleocerellus</i>	81.40
<i>flavipes</i> (Duda), <i>Onychaspidium</i>	81.27
<i>flavipes</i> Duda, <i>Tricimba</i>	81.32
<i>flavipes</i> (Duda), <i>Trigonomma</i>	81.41
<i>flavipes</i> Enderlein, <i>Coryphisoptron</i>	81.12
<i>flavipes</i> (Loew), <i>Liohippелates</i>	81.20
<i>flaviscutellata</i> (Enderlein), <i>Conioscinella</i>	81.10
<i>flaviseta</i> (Malloch), <i>Olcella</i>	81.25
<i>flavitarse</i> (Enderlein), <i>Coryphisoptron</i>	81.12
<i>flavithorax</i> (Duda), <i>Gaurax</i>	81.15
<i>flavitibia</i> (Duda), <i>Coryphisoptron</i>	81.12
<i>flavitibia</i> (Duda), <i>Oscinella</i>	81.28
<i>flaviventre</i> (Malloch), <i>Coryphisoptron</i>	81.12
<i>flavolineata</i> (Duda), <i>Tricimba</i>	81.32
<i>flavotrigona</i> Duda, <i>Conioscinella</i>	81.10
<i>flavus</i> Séguy, <i>Hippelates</i>	81.17
<i>flavus</i> Wiedemann, <i>Chlorops</i>	81.42
<i>floridensis</i> Sabrosky, <i>Pseudogaurax</i>	81.29
<i>foliatum</i> (Becker), <i>Trigonomma</i>	81.41
<i>forbesi</i> (Curran), <i>Apallates</i>	81.6
<i>formosa</i> (Becker), <i>Conioscinella</i>	81.10
<i>fortis</i> (Becker), <i>Apotropina</i>	81.3



fossulatum (Loew), Trigonomma	81.42
foveolatum (Duda), Trigonomma	81.42
frit (Linnaeus), Oscinella	81.28
frontalis (Becker), Apallates	81.6
<i>frontella</i> (Fallén), Conioscinella	81.10
frontoorbitalis (Sabrosky), Biorbitella	81.8
fur (Williston), Apallates	81.6

galapagensis (Curran), Conioscinella	81.10
galapagensis (Curran), Liohippелates	81.21
<i>Gampsocera</i> Schiner of Becker, Enderlein	81.12
<i>Gauracisoma</i> Duda	81.33
Gaurax Loew	81.15
<i>Gaurax</i> Loew of Becker, Duda	81.29
gemina (Becker), Thaumatomyia	81.40
genalis Thomson, Hippelates	81.17
giganteus (Duda), Homaluroides	81.37
glabra (Meigen), Thaumatomyia	81.40
glabricollis (Thomson), Diplotoxa	81.35
glabrina (Becker), Thaumatomyia	81.40
glabrum (Duda), Rhopalopterum	81.31
Goniaspis Duda	81.16
<i>Goniopsis</i> Duda	81.14
<i>Goniopsita</i> Duda	81.14
gracilis (Malloch), Apotropina	81.3
<i>grandicornis</i> (Becker), Thaumatomyia	81.41
grandiusculus Becker, Hippelates	81.17
grisea (Malloch), Apotropina	81.4
<i>griseolineata</i> (Duda), Olcella	81.26
griseoscutellata (Duda), Olcella mendozana var.	81.25
gruenbergi Enderlein, Discogastrella	81.12
gundlachi (Loew), Parectecephala	81.38


<i>Hammaspis</i> Duda	81.32
hastatus (Duda), Acanthopeltastes	81.5
hemiptera Séguéy, Siphonellomyia	81.31
hermsi (Sabrosky), Apallates	81.6
hesperia (Sabrosky), Biorbitella	81.8
Hippelates Loew	81.16
<i>Hippelates</i> Loew, part	81.5
<i>Hippelatinus</i> Enderlein	81.25
hirta (Duda), Ectecephala	81.36
hirtiventris (Malloch), Apotropina	81.4
<i>Homalura</i> Meigen of Becker, etc.	81.37
Homaluroides Sabrosky	81.37
horvathi (Becker), Acanthopeltastes	81.5
humeralis (Duda), Parectecephala	81.39
hyalinipennis Enderlein, Acanthopeltastes	81.5
hyalipennis Duda, Parectecephala	81.39
<i>hyalipennis</i> , error, Acanthopeltastes	81.5
hystrix (Duda), Eugaurax	81.14

<i>ignobilis</i> Becker, Hippelates	81.17
<i>ignorata</i> (Duda), Lioscinella	81.22
<i>illicis</i> (Curran), Liohippelates	81.19
<i>immaculatus</i> (Becker), Pseudogaurax	81.29
<i>impressus</i> Becker, Hippelates	81.17
<i>improvisa</i> (Becker), Conioscinella	81.10
<i>Incertella</i> Sabrosky	81.18
<i>incipiens</i> Curran, Hippelates	81.17
<i>incipiens</i> (Williston), Conioscinella	81.10
<i>incolorata</i> (Becker), Acanthopeltastes	81.5
<i>incolumis</i> (Becker), Malloewia	81.23
<i>incompletus</i> (Becker), Homaluroides	81.37
<i>incompletus</i> (Becker), Liohippelates	81.21
<i>inconspicua</i> (Becker), Conioscinella	81.10
<i>inconstans</i> (Becker), Conioscinella	81.10
<i>indistincta</i> Duda, Parectecephala	81.39
<i>infumata</i> (Becker), Apotropina	81.4
<i>inquilinus</i> (Coquillett), Chaetochlorops	81.9
<i>insignis</i> Becker, Anthracophaga	81.34
<i>insularis</i> Malloch, Eugaurax	81.14
<i>intermedia</i> Duda, Ectecephala	81.36
<i>intermedius</i> Sabrosky, Eugaurax	81.14
<i>interruptus</i> (Becker), Pseudogaurax	81.29
<i>iota</i> (Malloch), Conioscinella	81.10

<i>Lacommetopa</i> Duda	81.15
<i>lacteihalterata</i> (Enderlein), Conioscinella	81.10
<i>laevifrons</i> Becker, Ectecephala	81.36
<i>laevis</i> (Becker), Homaluroides	81.37
<i>lancifer</i> (Coquillett), Pseudogaurax	81.29
<i>Lasiopleura</i> Becker	81.3
<i>latimaculatus</i> Hall, Pseudogaurax	81.29
<i>leprae</i> Linnaeus, Musca (<i>incertae sedis</i>)	81.34
<i>Leptopeltastes</i> Enderlein	81.27
<i>Leptotrigonum</i> Becker	81.35
<i>limitatum</i> (Becker), Rhopalopterum	81.31
<i>lindneri</i> Duda, Oscinella (<i>incertae sedis</i>)	81.34
<i>linearis</i> (Malloch), Hippelates	81.17
<i>Liohippelates</i> Duda	81.19
<i>Lioscinella</i> Duda	81.22
<i>lippulum</i> Enderlein, Trigonomma	81.42
<i>longiceps</i> Duda, Anthracophaga	81.34
<i>longilineatus</i> Sabrosky, Pseudogaurax	81.29
<i>longipalpus</i> Enderlein, Oscinosoma (<i>incertae sedis</i>)	81.34
<i>longipennis</i> (Becker), Apotropina	81.4
<i>longiscutellatum</i> (Enderlein), Onychaspidium	81.27
<i>longiventris</i> (Enderlein), Enderliniella	81.14
<i>longula</i> (Becker), Apotropina	81.4
<i>Loxobathmis</i> Enderlein	81.23
<i>luciscens</i> (Enderlein), Pseudogaurax	81.29
<i>luscens</i> , error, Pseudogaurax	81.29
<i>luederwaldti</i> Enderlein, Tropicodiscinis	81.33
<i>luteipes</i> (Becker), Apallates	81.6

lutzi (Curran), Apallates	81.6
lutzi (Curran), Liohippелates	81.21
lyricus Hall, Pseudogaurax	81.30
<i>maculata</i> Duda, Chlorops	81.35
<i>maculicornis</i> (Duda), Apallates	81.6
<i>maculifrons</i> Becker, Parectecephala	81.39
<i>maculiventris</i> (Malloch), Fiebrigella	81.15
<i>maculosa</i> Duda, Chlorops	81.35
<i>maculosa</i> (Loew), Parectecephala	81.39
<i>Madiza</i> Fallén, part	81.25
<i>magnicornis</i> (Duda), Apallates	81.6
<i>magnifica</i> Duda, Metasiphonella	81.23
<i>magnipalpoides</i> (Curran), Apallates	81.7
<i>major</i> (Duda), Stenoscini	81.32
<i>Malloewia</i> Sabrosky	81.23
<i>mars</i> (Curran), Conioscinella	81.10
<i>mattea</i> (Curran), Oscinella	81.28
<i>megacera</i> (Becker), Thaumatomyia	81.40
<i>melaena</i> (Becker), Elachiptera	81.13
<i>Melanochaeta</i> Bezzi	81.23
<i>Melanochaeta</i> , part	81.12
<i>melancholica</i> (Becker), Tricimba	81.32
<i>melanosoma</i> Brèthes, Somillus	81.42
<i>mellinus</i> (Becker), Chaethippus	81.9
<i>mendozæ</i> (Malloch), Homaluroides	81.37
<i>mendozaana</i> Enderlein, Olcella	81.25
<i>Merectecephala</i> Duda	81.38
<i>Meromyza</i> Meigen	81.38
<i>Meroscini</i> of Becker, Duda	81.27
<i>metallicus</i> (Becker), Liohippелates	81.20
<i>metallica</i> of Enderlein, Liohippелates	81.19
<i>Metasiphonella</i> Duda	81.23
<i>mexicana</i> (Duda), Neodiptotaxa	81.38
<i>mexicanus</i> (Duda), Homaluroides	81.37
<i>miccotrogus</i> (Séguy), Apallates	81.7
<i>miccotrogus</i> (Séguy), Apallates	81.7
<i>microcentrus</i> (Coquillett), Apallates	81.7
<i>microcera</i> (Loew), Pseudopachychaeta	81.39
<i>Microchaetaspis</i> Duda	81.32
<i>microneurinus</i> (Duda), Liohippелates	81.21
<i>microps</i> Duda, Parectecephala	81.39
<i>Mimogaurax</i> Hall	81.29
<i>Mimoscini</i> Malloch	81.15
<i>minor</i> (Adams), Incertella	81.18
<i>misceomaculatus</i> Hall, Pseudogaurax	81.29
<i>mitis</i> (Williston), Elachiptera	81.13
<i>mona</i> (Curran), Incertella	81.19
<i>Monochaetoscinella</i> Duda	81.24
<i>nabis</i> Séguy, Hippelates	81.17
<i>nana</i> (Williston), Olcella	81.26
<i>neglecta</i> (Becker), Malloewia	81.23
<i>neocoxendix</i> (Sabrosky), Apallates	81.7

Neodiplotoxa Malloch	81.38
Neohippelates Roberts	81.19
Neolcanabates Duda	81.33
neotropica (Becker), Olcella	81.26
neotropicus (Becker), Steleocerellus	81.40
nigellus Wiedemann, Chlorops (<i>incertae sedis</i>)	81.42
nigra Malloch, Tricimba	81.32
nigrescens (Duda), Thaumatomyia	81.41
nigribucca Duda, Liohippelates	81.21
nigriceps (Duda), Onychaspidium	81.27
nigricollis Becker, Hippelates	81.17
nigricornis (Duda), Apotropina	81.4
nigricoxa (Malloch), Apallates	81.7
nigricoxa of Malloch, Apallates	81.7
nigrifrons Duda, Liohippelates	81.21
nigrilineatus, error, Pseudogaurax	81.29
nigripes (Duda), Apallates	81.7
nigripes Duda, Conioscinella soluta var.	81.11
nigripes Duda, Monochaetosciniella	81.24
nigrirostra (Adams), Lioscinella	81.22
nigriseta (Malloch), Olcella	81.26
nigrithorax (Duda), Olcella	81.26
nigrivertex Duda, Liohippelates	81.21
nigrofemorata Duda, Tricimba	81.33
nigrohalterata (Duda), Conioscinella	81.10
nigrolineatus (Enderlein), Pseudogaurax	81.29
nigromaculatus (Duda), Pseudogaurax	81.29
nigronotata (Duda), Biorbitella	81.8
nigrum (Becker), Trigonomma	81.42
nitens Duda, Neolcanabates (<i>incertae sedis</i>)	81.34
nitida (Duda), Malloewia	81.23
nitidifrons Duda, Ectecephala	81.36
nocens (Becker), Apallates	81.7
Notaulacella Enderlein	81.24
nuda (Adams), Conioscinella	81.10
nudifrons (Malloch), Liohippelates	81.20
numerata (Becker), Discogastrella	81.12
obliquans Enderlein, Loxobathmis	81.23
obscura (Coquillett), Oscinella	81.28
obscura (Schiner), Ectecephala	81.36
obscurata Duda, Goniaspis	81.16
obscurata (Duda), Parectecephala	81.39
obscuricornis Loew, Chlorops	81.35
obscuripila (Duda), Olcella mendozana var.	81.25
ochripes Enderlein, Tricimba	81.33
oculatus (Becker), Pseudogaurax	81.29
oeceticola, error, Pseudogaurax	81.29
oecetiphagus (Blanchard), Pseudogaurax	81.29
Olcanabates Enderlein	81.25
Olcella Enderlein	81.25
Omochaeta Duda	81.3
Onychaspidium Enderlein	81.27
opacula Fedoseeva, Meromyza	81.38



Opetiophora Loew	81.27
opposita (Becker), Olcella	81.26
orbitalis (Duda), Biorbitella	81.8
Oscinella Becker	81.27
Oscinella Becker, part	81.5
OSCINELLINAE	81.4
Oscinoides Malloch	81.28

paessleri (Duda), Apallates	81.7
Palaeoconioscinella Duda	81.16
Palaeoenderleiniella Duda	81.16
Palaeogaurax Duda	81.9
palliata (Curran), Apotropina	81.4
pallida (Loew), Cadrema	81.9
pallidicornis (Duda), Gaurax	81.16
pallipes (Duda), Pseudogaurax	81.30
pallipes (Loew), Liohippелates	81.21
pallipes, authors, Liohippелates	81.20
pallipes of Malloch, Liohippелates	81.19
palpalis Becker, Tricimba	81.33
panamensis (Malloch), Apotropina	81.4
papillipes, error, Liohippелates	81.20
paraguayensis Sabrosky & Paganelli, Olcella	81.26
parallelinervis (Duda), Pseudogaurax	81.30
Parasiphonella Enderlein	81.28
Parectecephala Becker	81.38
particeps (Becker), Apallates	81.7
particeps (Becker), part, Apallates	81.7
partitus (Becker), Liohippелates	81.21
partitus (Becker), part, Liohippелates	81.19
parva (Adams), Olcella	81.26
parviseta (Malloch), Hippelates	81.18
parvulus (Malloch), Apallates	81.7
pehlkei Enderlein, Hippelates	81.18
pennatus (Becker), Pterogaurax	81.30
Pentanotaulax Enderlein	81.33
perlineata Duda, Conioscinella	81.10
peruana Duda, Goniaspis	81.16
peruanus (Becker), Liohippелates	81.21
peruanus Duda, Pterogaurax	81.30
picta Duda, Ectecephala	81.36
pilosa Duda, Elachiptera	81.13
pilosa Fedoseeva, Meromyza	81.38
plaumanni Sabrosky, Pseudogaurax	81.30
plebejus Loew, Hippelates	81.18
plebejus group, Hippelates	81.16
plcsia (Curran), Oscinella	81.28
plcuralis (Becker), Olcella	81.26
pleuralis (Duda), Olcella	81.26
pleuralis Malloch, Eugaurax	81.14
pleuriticus (Wiedemann), Hippelates	81.18
plumbella (Wiedemann), Siphunculina	81.31
pollinosa Sabrosky, Elachiptera	81.13
porteri Brèthes, Hippelates	81.42


<i>proboscideus</i> Williston, <i>Hippelates</i>	81.18
<i>Prohippелates</i> Malloch	81.9
<i>projecta</i> (Malloch), <i>Olcella</i>	81.26
<i>prosthionelas</i> (Enderlein), <i>Lioscinella</i>	81.22
<i>pruinosis</i> (Duda), <i>Apallates</i>	81.7
<i>Pseudochlorops</i> Duda	81.15
<i>pseudogauracica</i> Duda, <i>Conioscinella</i>	81.11
<i>Pseudogaurax</i> Duda	81.29
<i>Pseudogaurax</i> Malloch	81.29
<i>Pseudohippelates</i> Malloch	81.3
<i>Pseudopachychaeta</i> Strobl	81.39
<i>Pseudoscinella</i> Duda	81.15
<i>Psilochlorops</i> Duda	81.39
<i>Pterogaurax</i> Duda	81.30
<i>pubescens</i> (Becker), <i>Apotropina</i>	81.4
<i>pubicollis</i> (Becker), <i>Trigonomma</i>	81.42
<i>pubiseta</i> (Becker), <i>Tricimba</i>	81.33
<i>pulchripes</i> (Loew), <i>Neodiplotoxa</i>	81.38
<i>pulla</i> (Adams), <i>Thaumatomyia</i>	81.41
<i>pulverulenta</i> (Duda), <i>Conioscinella</i>	81.11
<i>punctatus</i> Becker, <i>Hippelates</i>	81.18
<i>punctatus</i> (Duda), <i>Homaluroides</i>	81.37
<i>punctifrons</i> (Becker), <i>Olcella</i>	81.26
<i>punctulatus</i> (Duda), <i>Apallates</i>	81.7
<i>pura</i> (Curran), <i>Monochaetoscinella</i>	81.24
<i>pusio</i> (Loew), <i>Liohippелates</i>	81.21
<i>pusio</i> group, <i>Liohippелates</i>	81.19
<i>pygmaea</i> Duda, <i>Tricimba</i>	81.33
<i>quadridentata</i> (Duda), <i>Onychaspidium</i>	81.27
<i>quadrilineata</i> Williston of Becker, <i>Conioscinella</i>	81.11
<i>quadrilineata</i> Williston of Coquillett, <i>Eugaurax</i>	81.14
<i>quadrilineatus</i> (Williston), <i>Eugaurax</i>	81.14
<i>quadrivittata</i> (Duda), <i>Conioscinella</i>	81.11
<i>quinquelineata</i> (Adams), authors, <i>Olcella</i>	81.26
<i>quinquepunctatus</i> (Loew), <i>Homaluroides</i>	81.37
<i>reticulata</i> (Loew), <i>Siphunculina</i>	81.31
<i>retractum</i> (Duda), <i>Coryphisopteron</i>	81.12
<i>Rhodesiella</i> Adams	81.30
<i>Rhopalopteron</i> Duda	81.31
<i>Rhopalopteronum</i> Duda	81.31
<i>rimata</i> Enderlein, <i>Tricimba</i>	81.33
<i>robertsoni</i> (Sabrosky), <i>Liohippелates</i>	81.21
<i>rubicunda</i> (Duda), <i>Biorbitella</i>	81.8
<i>rubida</i> Becker, <i>Elachiptera</i>	81.13
<i>rubra</i> (Becker), <i>Goniaspis</i>	81.16
<i>rubrolimbata</i> Duda, <i>Elachiptera</i>	81.13
<i>rufescens</i> (Duda), <i>Olcanabates</i>	81.25
<i>ruficollis</i> (Frey), <i>Elachiptera</i>	81.13
<i>rufipectus</i> (Becker), <i>Homaluroides</i>	81.37
<i>rufipleura</i> (Duda), <i>Incertella</i>	81.19
<i>rufithorax</i> (Duda), <i>Apotropina</i>	81.4
<i>rufithorax</i> Duda, <i>Conioscinella</i>	81.11


<i>rufithorax</i> (Duda), <i>Thaumatomyia</i>	81
<i>rufomarginata</i> Duda, <i>Diplotoxa</i>	81.11
<i>rufoscutellata</i> Duda, <i>Conioscinella soluta</i> var.	81.12
<i>rufoscutellatum</i> (Duda), <i>Coryphisoptron</i>	81.30
<i>rufus</i> (Duda), <i>Pseudogaurax</i>	81.40
<i>rufus</i> (Duda), <i>Steleocerellus</i>	
<i>sacculicornis</i> (Enderlein), <i>Elachiptera</i>	81.13
<i>salebrosum</i> (Enderlein), <i>Coryphisoptron</i>	81.12
<i>satanas</i> (Becker), <i>Liohippelates</i>	81.21
<i>saundersi</i> Kumm, <i>Hippelates</i>	81.18
<i>schildi</i> Duda, <i>Notaulacella</i>	81.24
<i>schineri</i> Becker, <i>Parectecephala</i>	81.39
<i>schnusei</i> (Becker), <i>Acanthopeltastes</i>	81.5
<i>scutellaris</i> (Becker), <i>Chaetochlorops</i>	81.9
<i>scutellaris</i> (Malloch), <i>Ocella mendozana</i> var.	81.26
<i>scutellaris</i> (Williston), <i>Goniaspis</i>	81.16
<i>selectus</i> Becker, <i>Hippelates</i>	81.18
<i>semicircularis</i> Duda, <i>Conioscinella soluta</i> var.	81.11
<i>senilis</i> (Duda), <i>Apotropina</i>	81.4
<i>setigena</i> Sabrosky, <i>Eugaurax</i>	81.14
<i>sexdentatum</i> Enderlein, <i>Onychaspidium</i>	81.27
<i>sicatrix</i> (Curran), <i>Apallates</i>	81.7
<i>signata</i> (Wollaston), <i>Siphunculina</i>	81.31
<i>signatus</i> (Loew), <i>Pseudogaurax</i>	81.30
<i>similis</i> Becker, <i>Ectecephala</i>	81.36
<i>similis</i> Enderlein, <i>Tricimba</i>	81.33
<i>simulator</i> (Séguy), <i>Apallates</i>	81.6
<i>Siphomyia</i> Williston	81.16
<i>Siphonella</i> Macquart, part	81.25
<i>Siphonellomyia</i> Séguy	81.31
SIPHONELLOPSINAE	81.3
<i>Siphunculina</i> Rondani	81.31
<i>soluta</i> (Becker), <i>Conioscinella</i>	81.11
<i>Somillus</i> Brèthes	81.42
<i>specularifrons</i> (Enderlein), <i>Apotropina</i>	81.4
<i>speculifrons</i> (Enderlein), <i>Trigonomma</i>	81.42
<i>speculiger</i> (Enderlein), <i>Lioscinella</i>	81.22
<i>spineus</i> Enderlein, <i>Acanthopeltastes</i>	81.5
<i>Steleocerellus</i> Frey	81.39
<i>Stenoprosopon</i> Duda	81.19
<i>Stenoscinis</i> Malloch	81.31
<i>stenrothi</i> (Frey), <i>Homaluroides</i>	81.37
<i>stigmaticus</i> (Duda), <i>Hippelates</i>	81.18
<i>straminea</i> Loew, <i>Opetiophora</i>	81.27
<i>strandi</i> (Duda), <i>Acanthopeltastes</i>	81.5
<i>strandi</i> (Duda), <i>Apallates</i>	81.6
<i>strandi</i> (Duda), <i>Liohippelates</i>	81.22
<i>Strandimyia</i> Duda	81.32
<i>Stratiomicroneurum</i> Duda	81.19
<i>strigata</i> (Duda), <i>Thaumatomyia</i>	81.40
<i>strigatus</i> Enderlein, <i>Olcenabates</i>	81.25
<i>striolata</i> (Wiedemann), <i>Siphunculina</i>	81.31
<i>subannulatus</i> (Malloch), <i>Hippelates</i>	81.18

subequalis (Malloch), Goniaspis	81.16
sublineata (Becker), Elachiptera	81.13
sublineata Duda, Conioscinella	81.11
subtomentosus (Enderlein), Apallates	81.8
sulcata Duda, Tricimba	81.33
<i>sulcatus</i> Enderlein, Olcanabates	81.25
sulfurifrons (Duda), Thaumatomyia	81.41
<i>sulfurifrons</i> (Duda), part, Thaumatomyia	81.40
sulfurihalterata (Enderlein), Lioscinella	81.22
tecta (Becker), Tricimba	81.33
tectus (Becker), Pseudogaurax	81.30
<i>Teleocoma</i> Aldrich	81.15
tenacissimus (Iches) Liohippелates	81.22
tener (Coquillett), Apallates	81.8
tenuifrons (Duda), Trigonomma	81.42
tenuiseta Duda, Ectecephala	81.36
testaceus (Enderlein), Pseudogaurax	81.30
<i>texana</i> (Malloch), Apallates	81.6
Thaumatomyia Zenker	81.40
tibialis Duda, Liohippелates	81.22
tibialis Malloch, Pseudogaurax	81.30
tincta Becker, Ectecephala	81.36
tinctipes (Becker), Gaurax	81.16
<i>tofinigastrea</i> , error, Tricimba	81.33
tofinistrigata (Enderlein), Tricimba	81.33
tomentosus (Enderlein), Apallates	81.8
trabeatus (Enderlein), Pseudogaurax	81.30
transatlantica (Schiner), Ectecephala	81.37
<i>transatlantica</i> , part, Parectecephala	81.39
trapezisopteron (Enderlein), Notaulacella	81.24
triangularis (Williston), Pseudogaurax	81.30
triangulinus (Duda), Apallates	81.8
Tricimba Lioy	81.32
tridens Sabrosky, Pseudogaurax	81.30
trifidus (Duda), Pseudogaurax	81.30
trigonalifrons (Enderlein), Coryphisopteron	81.12
Trigonomma Enderlein	81.41
trigramma (Loew), Olcella	81.26
trilineata (Duda), Olcella	81.26
trilineatus (Duda), Pseudogaurax	81.30
trimaculatus (Duda), Hippelates	81.18
tripunctata Aldrich, Ectecephala	81.37
tripunctata (Becker), Enderleiniella	81.14
tripunctata (Curran), Conioscinella	81.11
<i>trivittata</i> (Williston), Parectecephala	81.38
Tropidoscinis Enderlein	81.33
truncata (Malloch), Goniaspis	81.16
turbata (Duda), Olcella	81.27
turbida (Becker), Conioscinella	81.11
turneri Kumm, Hippelates	81.18
ujhelyi Duda, Oscinella	81.28
<i>umbrosa</i> (Loew), Oscinella	81.28



<i>umbrosa</i> of Coquillett, Rhopalopterum	81.31
unicolor Becker, Diplotoxa	81.35
unicornis Giglio-Tos, Chlorops	81.35
unilineatus Hall, Pseudogaurax	81.30
uniseriata (Duda), Conioscinella soluta var.	81.11
varipalpus (Curran), Apallates	81.8
verrucosa Duda, 1921, Fiebrigella	81.15
verrucosa (Duda, 1929), Fiebrigella	81.15
verrucosum (Duda), Trigonomma	81.42
versicolor (Locw), Diplotoxa	81.35
virgata of Coquillett, Biorbitella	81.8
virgulata (Enderlein), Tricimba	81.33
viridifrons (Duda), Fiebrigella	81.15
viridiniger (Enderlein), Apallates	81.8
vittatus Wiedemann, Chlorops	81.42
vulgarimonens Duda, Oscinella	81.28
vulgaris (Duda), Malloewia	81.23
yungas Sabrosky & Paganelli, Chlorops	81.35
zernyi Duda, Monochactoscincella	81.24





MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO

(Formerly Departamento de Zoologia, Secretaria da Agricultura, São Paulo)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS SOUTH
OF THE UNITED STATES

82

FAMILY ODINIIDAE

ANGELO PIRES DO PRADO

*Departamento de Zoologia, Setor de Parasitologia
Universidade Estadual de Campinas, São Paulo, Brasil*

This small family of small to medium-sized flies, usually dark brown in color, is poorly known in the Neotropical Region. Habits and immature stages of strictly Neotropical species are unknown. Palearctic species of *Odinia* Robineau-Desvoidy have been reared in association with wood boring beetles or from trees attacked by such beetles; less commonly, from larvae of cossid and other moths.

Ref. — Prado, 1973 (revision).

Subfamily ODINIINAE

Genus ODINIA Robineau-Desvoidy

ODINIA Robineau-Desvoidy, 1830: 648. Type-species, *trinitata* Robineau-Desvoidy (Rondani, 1875:167; 1875:2) = *maculata* (Meigen).

BIGUTTATA Sabrosky, 1959:225, pl. 1, fig. 1. Type-locality: Virgin Islands, St. Croix, Peter Res. Distr. — West Indies (Virgin Islands, Puerto Rico); U.S.A. (Florida).

BREVITIBIA Shewell, 1960:632, figs. 12-14. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Roraima, Pará, Santa Catarina). Ref. — Prado, 1973:484, figs. 1-9.

CORONATA Sabrosky, 1959:234, pl. 1, fig. 8. Type-locality: El Salvador, La Unión. Distr. — U.S.A. (Arizona, Texas), El Salvador, Panama.

PARVIPUNCTATA Sabrosky, 1959:235, pl. 1, fig. 9. Type-locality: Puerto Rico, Isabela, Guatajaca Forest. Distr. — West Indies (Virgin Islands, Puerto Rico).

SP. Hennig, 1969:617. — Jamaica.

SURUMUANA Prado, 1973:484, figs. 10-11. Type-locality: Brazil, Roraima, Surumu. Distr. — Brazil (Roraima).

WILLIAMSII Johnson, 1924:90. Type-locality: "Galápagos Islands". Distr. — Galápagos Islands. Ref. — Sabrosky, 1959:632, pl. 1, fig. 7.

82.1

Subfamily TRAGINOPINAE

Genus LOPESODINIA Prado

LOPESODINIA Prado, 1973:500. Type-species, *diversa* Prado (orig. des.).

ALVARENGAI Prado, 1973:503. Type-locality: Brazil, Pará, Jacareacanga. Distr. — Brazil (Pará).

DIVERSA Prado, 1973:502, figs. 51-55. Type-locality: Brazil, Guanabara. Distr. — Brazil (Guanabara).

lanei Prado, 1973:504 (*lapsus*).

Genus NEOTRAGINOPS Prado

NEOTRAGINOPS Prado, 1973:504. Type-species, *Traginops clathratus* Hendel (orig. des.).

CLATHRATUS (Hendel), 1909:52, 2 figs. (*Traginops*). Type-locality: Peru, Vilcanota, and "Paraguay". Distr. — Peru, Paraguay, Brazil (Mato Grosso, Goiás, Minas Gerais, Espírito Santo, Rio de Janeiro, Guanabara, São Paulo, and Santa Catarina), Colombia. Ref. — Prado, 1973:506, figs. 56-61.

Genus PARATRAGINOPS Hendel

PARATRAGINOPS Hendel, 1917:38. Type-species, *Traginops pilicornis* Cresson (orig. des.).

PILICORNIS (Cresson), 1912:395, pl. 19, figs. 3-4 (*Traginops*). Type-locality: Guyana, Bartica. Distr. — Guyana, Brazil (Mato Grosso, Goiás, São Paulo). Ref. — Prado, 1973:488, figs. 12-18.

PLAUMANNI Shewell, 1960:626, fig. 1. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Goiás, Minas Gerais, Espírito Santo, Rio de Janeiro, São Paulo, Santa Catarina). Ref. — Prado, 1973:490, figs. 19-23.

Genus SHEWELLIA Hennig

SHEWELLIA Hennig, 1969:620. Type-species, *agromyzina* Hennig (orig. des.).

AGROMYZINA Hennig, 1969:621, figs. 48-49. Type-locality: Peru, Madre de Dios, Avispas. Distr. — Peru.

Genus SCHILDOMYIA Malloch

SCHILDOMYIA Malloch, 1926:26. Type-species, *vittithorax* Malloch (orig. des.).

BREVIHIRTIA Malloch, 1926:27. Type-locality: Panama, Canal Zone, Gatún Lake, Cano Saddle. Distr. — Panama.

FLAVIDA Prado, 1973:499, figs. 48-50. Type-locality: Brazil, São Paulo, Capital (Cantareira). Distr. — Brazil (São Paulo).

- GOYANA Prado, 1973:498, figs. 41-47. Type-locality: Brazil, Goiás, Brasília, D.F., Parque Municipal. Distr. — Brasil (Goiás).
- LANEI Prado, 1973:492, figs. 25-31. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães). Distr. — Brazil (Amapá, Mato Grosso).
- PERUVIANA Hennig, 1969:618, fig. 46. Type-locality: Peru, Madre de Dios, Avispas. Distr. — Peru.
- PUNCTIFRONS Malloch, 1926:28, pl. 2, fig. 16. Type-locality: Costa Rica, San Mateo, Higuito. Distr. — Costa Rica.
- RETICULATA Prado, 1973:494, figs. 32-40. Type-locality: Brazil, São Paulo, Andes. Distr. — Brazil (São Paulo).
- TRINIDADENSIS Hennig, 1969:620, fig. 47. Type-locality: Trinidad, Arima Valley. Distr. — Trinidad.
- VITTITHORAX Malloch, 1926:27. Type-locality: Costa Rica, San Mateo, Higuito. Distr. — Costa Rica, Panama, Peru. Ref. — Hennig, 1969: 618.

LITERATURE CITED

- CRESSON, E. T., Jr.
1912. Descriptions of several new neotropical acalyptrate Diptera. *Ent. News* 23:389-396, 1 pl., 6 figs.
- HENDEL, F.
1909. Ueber eine Dipterengattung mit turmartigen Scheitel, *Tragionops* Coquillett. *Wien. ent. Ztg.* 28(2):49-52, 2 figs.
1917. Beiträge zur Kenntnis der acalyptraten Musciden. *Deutsche ent. Ztschr.* 1917:36-38.
- HENNIG, W.
1969. Neue Gattungen und Arten der Acalyptraten. *Canad. Ent.* 101 (6):589-633, 72 figs.
- JOHNSON, C. W.
1924. Diptera of the Williams Galapagos Expedition. *Zoologica*, New York 5:85-92, 2 figs.
- MALLOCH, J. R.
1926. New genera and species of acalyptrate flies in the United States National Museum. *Proc. U. S. Natl. Mus.* 68(21):1-35, pls. 1-2.
- PRADO, A. P. DO
1973. Contribuição ao conhecimento da família Odiniidae (Diptera, Acalyptratae). *Studia Ent.* 16(1-4):481-510, 61 figs.
- ROBINEAU-DESVOIDY, J. B.
1830. Essai sur les Myodaires. Institut de France, [Cl. des] Sci. Math. et Phys., Acad. Roy. des Sci., *Mém présentées par divers Savans* (2)2:1-813.
- RONDANI, C.
1875. Species Italicae ordinis dipterorum (Muscaria Rndn.) collectae et observatae. Stirps XXIII. Agromyzinae. *Bol. Soc. ent.*



Ital. 7:166-191. (Also published separately as "*Dipterologiae Italicae Prodromus*" 8 (Stirps 23 [= Pt 3]) : 23 pp., Firenze, 1875).

SABROSKY, C. W.

1959. Flies of the genus *Odinia* in the Western Hemisphere (Diptera: Odiniidae). *Proc. U. S. Natl. Mus.* 109 (= n.º 3414): 223-236, 1 pl.

SHEWELL, G. E.

1960. Notes on the family Odiniidae with a key to the genera and descriptions of new species. *Canad. Ent.* 92(8):625-633, 13 figs.

INDEX

agromyzina Hennig, 1969, <i>Shewellia</i>	82.2
alvarengai Prado, 1973, <i>Lopesodinia</i>	82.2
biguttata Sabrosky, 1959, <i>Odinia</i>	82.1
brevihirta Malloch, 1926, <i>Schildomyia</i>	82.2
brevitibia Shewell, 1960, <i>Odinia</i>	82.1
clathratus (Hendel), 1909 (Traginops), <i>Neotraginops</i>	82.2
coronata Sabrosky, 1959, <i>Odinia</i>	82.1
diversa Prado, 1973, <i>Lopesodinia</i>	82.2
flavida Prado, 1973, <i>Schildomyia</i>	82.2
goyana Prado, 1973, <i>Schildomyia</i>	82.3
lanei Prado, 1973, <i>Odinia</i> (<i>lapsus</i>)	82.2
lanei Prado, 1973, <i>Schildomyia</i>	82.3
Lopesodinia Prado, 1973	82.2
Neotraginops Prado, 1973	82.2
<i>Odinia</i> Robineau-Desvoidy, 1830	82.1
Odiniinae	82.1
Paratraginops Hendel, 1917	82.2
parvipunctata Sabrosky, 1959, <i>Odinia</i>	82.1
peruviana Hennig, 1969, <i>Schildomyia</i>	82.3
pilicornis (Cresson), 1912 (Traginops), <i>Paratraginops</i>	82.2
plaumanni Shewell, 1960, <i>Paratraginops</i>	82.2
punctifrons Malloch, 1926, <i>Schildomyia</i>	82.3
reticulata Prado, 1973, <i>Schildomyia</i>	82.3
<i>Schildomyia</i> Malloch, 1926	82.2
<i>Shewellia</i> Hennig, 1969	82.2
sp. Hennig, 1969, <i>Odinia</i>	82.1
surumuana Prado, 1973, <i>Odinia</i>	82.1
Traginopinae	82.2
trinidadensis Hennig, 1969	82.3
vittithorax Malloch, 1926. <i>Schildomyia</i>	82.3
williamsi Johnson, 1924, <i>Odinia</i>	82.1

DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

'83

FAMILY AGROMYZIDAE

KENNETH A. SPENCER

19, Redington Road, Hampstead
London, England

The Agromyzidae are known primarily as leaf-miners but recent advances in our knowledge of the family have shown that a significant proportion of species are internal stem-borers and seed-feeders and, to a lesser extent, root-feeders and gall-causers.

The family is well represented throughout the Neotropical region. In areas, such as around São Paulo, Caracas, and in the mountains below Bogotá, leaf-mines can be seen in abundance on much of the vegetation and this is in striking contrast to the obvious scarcity of species in tropical areas of other regions I have visited, as for example Ceylon, the Philippines, and Darwin, northern Australia. Dr. P. Wygodzinsky has discovered numerous species, as yet undescribed, around Tucumán in Argentina and many species occur in Chile. There has been insufficient collecting to indicate to what extent the family is represented in Patagonia.

The two largest genera, *Liriomyza* Mik and *Melanagromyza* Hendel, occur widely elsewhere in the world but the next largest genus, *Calucomyza* Hendel, has proliferated in a remarkable way in the Neotropical region and there is no doubt that many further species remain to be discovered. It is clear that the 141 species listed below represent only a fraction of those existing in the Neotropical region.

The following abbreviations have been used for the locations of type-specimens:

A.M.N.H. — American Museum of Natural History, New York.

B.M. — British Museum (Natural History), London.

Dep. Zool., São Paulo — Departamento de Zoologia da Secretaria da Agricultura do Estado de São Paulo.

M.C.Z. — Museum of Comparative Zoology, Cambridge, Massachusetts.

U.S.N.M. — United States National Museum, Washington, D.C.

83.1



Genus AGROMYZA Fallén

- AGROMYZA Fallén, 1810:21. Type-species, *reptans* Fallén (Rondani, 1875a: 177, 1875:12). [See note in the North American catalogue, p. 795].
- APPELBECKI Strobl, 1902:504. Type-locality: "Southern Spain". Distr. — Chile (Valparaiso, La Cruz); Southern Europe. Syntypes in coll. Strobl, Admont, Austria.
- andalusiaca* Strobl, 1906:380. Type-locality: "Southern Spain". Syn-
types in coll. Strobl, Admont, Austria.
- FUSCA Spencer, 1963:295, figs. 1a.b. Type-locality: Brazil, Santa Catarina.
Nova Teutônia. Distr. — Brazil. Holotype in B.M.
- INSOLENS Spencer, 1963:296, figs. 2a-c (male genitalia). Type-locality:
Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Holotype
in B.M.
- PLAUMANNI Spencer, 1963:297, figs. 3a-c (male genitalia). Type-locality:
Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Holotype
in B.M.
- SIMILLIMA Spencer, 1963:297, figs. 4a-b (male genitalia). Type-locality:
Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Holotype
in B.M.
- SPADIX Spencer, 1963:298, figs. 5a-b (male genitalia). Type-locality: British
Guiana, Tukeit. Distr. — British Guiana. Holotype in B.M.
- TEREBRANS Bezzi & Tavares, 1916:166. Type-locality: Brazil, Bahia. Distr.
— Brazil. Ref. — Spencer, 1963:300. Location of types not known.
- VENEZOLANA Spencer, 1963:299, figs. 6a-c (male genitalia). Type-locality:
Venezuela, Caracas, Botanical Gardens. Distr. — Venezuela. Ho-
lotype in coll. Spencer.

Genus JAPANAGROMYZA Sasakawa


- JAPANAGROMYZA Sasakawa, 1958:138. Type-species, *Agromyza duchesneae*
Sasakawa (orig. des.).
- ALDRICH (Frick), 1952:377 (*Melanagromyza*; nom. nov. for *tibialis* Frost).
Type-locality: Panama, Canal Zone, Barro Colorado I. Distr. —
Guatemala, Panama. Ref. — Spencer, 1963:301. Holotype in
U.S.N.M.
- tibialis* Frost, 1936:312 (*Agromyza*; preocc. Fallén, 1823).
- ARNAUDI Sasakawa, 1963:95. Type-locality: "El Salvador". Distr. — El
Salvador. Holotype in California Academy of Sciences.
- CENTROSEMAE (Frost), 1936:301 (*Agromyza*). Type-locality: Panama, Canal
Zone. Distr. — Panama. Ref. — Spencer, 1963:301, fig. 7. Holotype
in U.S.N.M.
- CURRANI (Frost), 1936:305 (*Agromyza*). Type-locality: Panama, Canal
Zone, Barro Colorado I. Distr. — Panama. Ref. — Spencer, 1963:
301, fig. 8. Holotype in A.M.N.H.
- FROSTI (Frick), 1952:373 (*Agromyza*; nom. nov. for *schmidtii* Frost). Type-
locality: Costa Rica, La Caja. Distr. — Costa Rica. Ref. —
Spencer, 1963:302. Holotype in U.S.N.M.
- schmidtii* Frost, 1936:302 (*Agromyza*; preocc. Aldrich, 1929).



- INAEQUALIS (Malloch), 1914:89 (*Agromyza*). Type-locality: "Puerto Rico"
Distr. — U.S.A. (Florida). Puerto Rico. Ref. — Spencer, 1963:302,
figs. 11a-b (male genitalia). Holotype in U.S.N.M.
- IRIDESCENS (Frost), 1936:303 (*Agromyza*). Type-locality: Panama, Canal
Zone, Barro Colorado I. Distr. — Panama (Canal Zone). Ref. —
Spencer, 1966c:5. Holotype in A.M.N.H.
- JAMAICENSIS Spencer, 1963:303. Type-locality: Jamaica. Rio Cobre Gorge.
Distr. — Jamaica. Holotype in coll. Spencer.
- ORBITALIS (Frost), 1936:314 (*Agromyza*). Type-locality: Panama, Canal
Zone, Barro Colorado I. Distr. — Panama. Ref. — Spencer, 1963:
303, figs. 12a-d (male genitalia). Holotype in A.M.N.H.

Genus MELANAGROMYZA Hendel

- MELANAGROMYZA Hendel, 1920:114, 126. Type-species, *Agromyza aeneoven-*
tris Fallén (orig. des.).
- Limnoagromyza* Malloch, 1921:147. Type-species, *diantherae* Malloch (orig.
des.).
Ref. — Spencer, 1966a:3-60.
- ALTERNANTHERAE Spencer, 1963:307, figs. 16a-d (male genitalia). Type-
locality: Argentina, Santa Fé. Distr. — Brazil, Paraguay, Argen-
tina. Holotype in U.S.N.M.
- ANGOLAE (Malloch), 1934:483 (*Agromyza*). Type-locality: Chile, Angol.
Distr. — Chile. Holotype in U.S.N.M.
- APPROXIMATA (Frost), 1936:316 (*Agromyza*). Type-locality: "Guatemala".
Distr. — Guatemala. Holotype in U.S.N.M.
- CAERULEA (Malloch), 1913:322 (*Agromyza*). Type-locality: "Mexico". Distr.
— Mexico, Cuba, Jamaica, Puerto Rico, Venezuela, Brazil, Peru.
Ref. — Spencer, 1963:307, figs. 17a-b (male genitalia). Holotype
in U.S.N.M.
- plumisetæ* Malloch, 1913:324 (*Agromyza*). Type-locality: "Puerto
Rico". Holotype in U.S.N.M.
- CHIENOPODI Spencer, 1963:308, figs. 18a-c (male genitalia). Type-locality:
Chile, Santiago. Distr. — Chile. Holotype in U.S.N.M.
- COLOMBIENSIS Spencer, 1963:309, figs. 20a-b (male genitalia). Type-locality:
Colombia, Bogotá. Distr. — Colombia. Holotype in U.S.N.M.
- COMPOSITOIDES Spencer, 1963:310, fig. 21. Type-locality: Jamaica, Rio Cobre
Gorge. Distr. — Jamaica. Holotype in coll. Spencer.
- CROTONIS (Frost), 1936:313 (*Agromyza*). Type-locality: Panama, Canal
Zone, Barro Colorado I. Distr. — Cuba, Panama. Ref. — Spencer,
1963:310, figs. 22a-c (male genitalia). Holotype in U.S.N.M.
- CUNCTANOIDES Blanchard, 1954:12. Type-locality: Argentina, General Pico,
Eva Perón (ex La Pampa). Distr. — Argentina. Holotype in coll.
Blanchard, Buenos Aires.
- DIADEMA (Melander), 1913:259 (*Agromyza*). Type-locality: "Haiti". Distr.
— Haiti, ?Brazil. Ref. — Spencer, 1963:311. Holotype in coll.
A. L. Melander.
- DISTINCTA Spencer, 1963:311, figs. 23a-b (male genitalia). Type-locality:
Brazil, Santa Catarina, Nova Teutônia. Holotype in B.M.



- ERIGERONIS Spencer, 1963:311, figs. 24a-b (male genitalia), 24c. Type-locality: Brazil, São Paulo, Capital, Água Funda. Distr. — Brazil. Holotype in coll. Spencer.
- FLORIDENSIS Spencer, 1963:312, figs. 25a-b (male genitalia), 25c. Type-locality: Dominican Republic, Ridgefield. Distr. — U.S.A. (Florida), Dominican Republic, British Guiana. Holotype in U.S.N.M.
- FLORIS Spencer, 1963:313, figs. 26a-c (male genitalia). Type-locality: Mexico, Cuernavaca. Distr. — U.S.A. (Florida), Mexico, Cuba, Costa Rica, Puerto Rico. Holotype in U.S.N.M.
- HALTERELLA Spencer, 1963:314, fig. 27 (male genitalia). Type-locality: Mexico, Cuernavaca. Distr. — Mexico. Holotype in U.S.N.M.
- HELIANTHI Spencer, 1963:314, fig. 28 (male genitalia). Type-locality: Cuba, Santiago de las Vegas. Distr. — U.S.A. (Florida), Mexico, Cuba, Trinidad. Holotype in U.S.N.M.
- LACUSTRIS (Malloch), 1934:482 (*Agromyza*). Type-locality: Argentina, Rio Negro, L. Nahuel Huapi. Distr. — Argentina. Ref. — Spencer, 1963:315. Holotype in B.M.
- LINI Spencer, 1963:315, figs. 29a-b (male genitalia). Type-locality: Peru, Barranca. Distr. — Peru. Holotype in U.S.N.M.
- LONGICAUDALIS Spencer, 1963:316, fig. 30. Type-locality: Jamaica, Mt. Diavolo. Distr. — U.S.A. (Florida), Jamaica, Puerto Rico, Virgin Is. (St. Croix). Holotype in coll. Spencer.
- LONGISETA (Malloch), 1913:326 (*Agromyza*). Type-locality: Mexico, Tabasco. Distr. — Mexico. Holotype in U.S.N.M.
- MALLOCHI (Hendel), 1923:145 (*Agromyza*; *nom. nov. for eupatoriae* Malloch). Type-locality: "Puerto Rico". Distr. — Puerto Rico, Cuba, Trinidad. Ref. — Spencer, 1963:316, figs. 31a, 31b-c (male genitalia). Holotype in U.S.N.M.
- eupatoriae* Malloch, 1915:107 (*Agromyza*; *preocc.* Kaltenbach, 1874). Type-locality: "Puerto Rico".
- MARELLI (Brèthes), 1920:282 (*Agromyza*). Type-locality: Argentina, Buenos Aires. Distr. — Brazil, Argentina. Ref. — Spencer, 1963:317, figs. 32a (male genitalia), 32b-c. Location of holotype not known.
- MINIMA (Malloch), 1913:328 (*Agromyza*). Type-locality: "Trinidad". Distr. — U.S.A. (Florida), Mexico, Jamaica, Trinidad, Guatemala, Puerto Rico, Virgin Is., Panama (Canal Zone), British Guiana, Venezuela. Ref. — Spencer, 1963:317, figs. 33a, 33b-d (male genitalia). Holotype in U.S.N.M.
- NEOTROPICA Spencer, 1963:319, figs. 34a-c (male genitalia). Type-locality: Mexico, Cuernavaca. Distr. — Mexico, Brazil. Holotype in U.S.N.M.
- NORILIS Spencer, 1963:320, fig. 35 (male genitalia). Type-locality: Colombia, Bogotá, near foot of Teleferico. Distr. — Colombia. Holotype in coll. Spencer.
- PFAFFIAE Spencer, 1963:320, figs. 36a-c (male genitalia), 36d-e. Type-locality: Venezuela, Caracas, Botanical Gardens. Distr. — Venezuela. Holotype in coll. Spencer.
- PRAECLARA Spencer, 1963:321, figs. 37a-b (male genitalia). Type-locality: Chile, Puerto Puyuhuapi. Distr. — Chile. Type in Deutsches Entomologisches Institut, Eberswalde, German Democratic Republic.



- SETIFERA Spencer, 1963:322, fig. 38 (male genitalia). Type-locality: British Guiana, Mazaruni. Distr. — British Guiana. Holotype in B.M.
- TETRAE (Malloch), 1934:483 (*Agromyza*). Type-locality: Chile, Peulla. Distr. — Chile. Ref. — Spencer, 1963:322, figs. 39a-c (male genitalia). Holotype in B.M.
- VIRENS (Loew), 1869:46 (*Agromyza*). Type-locality: U.S.A., Pennsylvania. Distr. — U.S.A., Canada, Hawaii; Jamaica. Ref. — Spencer, 1963:322, figs. 40a-c (male genitalia), 40d-e. Lectotype in M.C.Z.
- splendida* Frick, 1953:207. Type-locality: "Hawaii". Holotype in U.S.N.M.

Genus OPHIOMYIA Braschnikov

- Agromyza*, subg. OPHIOMYIA Braschnikov, 1897:40. Type-species, *Agromyza pulicaria* Meigen (mon.; misident.) = *maura* (Meigen).
- Ref. — Spenceer, 1964b:773-822.
- BUSCKI (Frost), 1936:315 (*Agromyza*). Type-locality: Panama, Canal Zone, Corozal. Distr. — Panama, Brazil, Argentina. Ref. — Spencer, 1963:325, figs. 41a-b (male genitalia), 41c. Holotype in U.S.N.M.
- CAMARAE Spenceer, 1963:326, figs. 42a, f, 42b-e (male genitalia). Type-locality: Trinidad, Curepe. Distr. — U.S.A. (Florida), Trinidad. Holotype in B.M.
- CURVIBRISATA (Frost), 1936:309 (*Agromyza*). Type-locality: Guatemala, Cobán, Alta Vera Paz. Distr. — Guatemala. Refs. — Frick, 1959:369, fig. 49; Spencer, 1963:327. Holotype in U.S.N.M.
- FASCICULATA (Malloch) 1934:479, fig. 22 (*Agromyza*). Type-locality: Argentina, Maquinhao. Distr. — Argentina. Ref. — Spencer, 1963:327. Holotype in B.M.
- HIRTICEPS (Malloch), 1934:481, fig. 23 (*Agromyza*). Type-locality: Argentina, Rio Negro, L. Gutierrez. Distr. — Argentina. Ref. — Spencer, 1963:328. Holotype in B.M.
- INSULARIS (Malloch), 1913:318 (*Agromyza*). Type-locality: "Cuba", Distr. — Cuba. Syntypes in U.S.N.M.
- LANTANAE (Froggatt), 1919:665 (*Agromyza*). Type-locality: Australia, New South Wales. Distr. — Mexico, Puerto Rico, Cuba, Honduras, Costa Rica, Panama, Trinidad; Australia, Hawaii, India, Kenya, South Africa. Ref. — Spencer, 1963:328, fig. 43. Types lost.
- longicauda* Curran, 1928:65 (*Agromyza*). Type-locality: "Puerto Rico". Holotype ♀ in A.M.N.H.
- ORALIS (Frost), 1936:309 (*Agromyza*). Type-locality: Guatemala, El Salto, Antigua. Distr. — Guatemala. Holotype in U.S.N.M.
- PUNCTOHALTERATA (Frost), 1936:311 (*Agromyza*). Type-locality: "Guatemala". Distr. — Guatemala, Costa Rica. Ref. — Spencer, 1963:328. Holotype in U.S.N.M.
- SPICATAE Spencer, 1963:329, fig. 44. Type-locality: Brazil, Rio Grande (?). Distr. — Brazil. Ref. — Spenceer, 1966b:141, figs. 1, 2 (male genitalia). Holotype in B.M.

Genus CEROBONTHA Rondani

- Odontocera* Macquart, 1835:614 (preocc. Serville, 1833). Type-species, *Chlorops denticornis* Panzer (orig. des.).



- CERODONTA Rondani, 1861:10 (nom. nov. for *Odontocera* Macquart).
Type-species, *Chlorops denticornis* Panzer (aut.).
Micromma Philippi, 1865:777. Type-species, ?*Dolichopus flavifrons* Philippi (orig. des.). N. SYN.
Ref. — Nowakowski, 1962:102.

Subgenus CERODONTA Rondani

- BRAZILIANA Spencer, 1963:330, figs. 45a, 45b-c (male genitalia). Type-locality: Brazil, Santa Catarina. Nova Teutônia. Distr. — Brazil. Holotype in B.M.
CRUCIATA (Blanchard), 1938:356 (*Triticomysa*). Type-locality: Argentina, La Pampa. Distr. — Argentina. Refs. — Frick, 1952:399; Spencer, 1963:331. Holotype in coll. Blanchard, Buenos Aires.
DORSALIS (Loew), 1863:54 (*Odontocera*). Type-locality: U.S.A., District of Columbia. Distr. — U.S.A., Puerto Rico, Guatemala, Brazil.
FLAVIFRONS (Philippi), 1865:777 (as ?*Dolichopus*, and at same time proposed as type of *Micromma*). Type-locality: Chile, Valdivia. Distr. — Chile, Juan Fernandez Is. Ref. — Spencer, 1963:332, figs. 46a-b (male genitalia). Syntypes believed in Museo Nacional, Santiago.
FULVITHORAX Malloch, 1934:484. Type-locality: Argentina, Buenos Aires. Distr. — Argentina. Ref. — Spencer, 1963:332. Holotype in B.M.
NIGRICORNIS Becker, 1919:212. Type-locality: "Ecuador". Distr. — Ecuador. Ref. — Spencer, 1963:332. Holotype in Musée d'Histoire Naturelle, Paris.

Subgenus DIZYGOMYZA Hendel

- DIZYGOMYZA Hendel, 1920:114, 130 (as genus). Type-species, *Agromyza morosa* Meigen (orig. des.).
ANDENSIS (Spencer), 1963:334, figs. 48a-b (*Phytobia*). Type-locality: Colombia, Tequendama Falls. Distr. — Colombia. Ref. — Spencer, 1966b:142. Holotype in coll. Spencer.
ANGULATA (Loew), 1869:47 (*Agromyza*). Type-locality: U.S.A., Pennsylvania. Distr. — U.S.A., British Guiana, Brazil; Europe. Ref. — Spencer, 1963:335; 1966b:142. Lectotype in M.C.Z.
semiposticata Hendel, 1920:131. Type-locality: "Austria". Ref. — Frick, 1957:203. Holotype in Naturhistorisches Museum, Vienna.
INEPTA (Spencer), 1963:335 (*Phytobia*, subg. *Poemyza*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Ref. — Spencer, 1966b:142. Holotype in B.M.
NITIDIVENTRIS (Malloch), 1934:477 (*Agromyza*). Type-locality: Argentina, Rio Negro, L. Nahuel Huapi. Distr. — Argentina. Refs. — Spencer, 1963:335, 1966b:142. Holotype in B.M.

Subgenus ICTEROMYZA Hendel

- Dizygomyza*, subg. ICTEROMYZA Hendel, 1931:51. Type-species, *Agromyza geniculata* Fallén (orig. des.).



LONGIPENNIS (Loew), 1869:48 (*Agromyza*). Type-locality: U.S.A., District of Columbia. Distr. — U.S.A., Brazil (São Paulo), Chile (Angol). Ref. — Spencer, 1963a: fig. 49 (male genitalia), 1966b:142. Holotype in M.C.Z.

Genus PHYTOBIA Lioy

PHYTOBIA Lioy, 1864:1313. Type-species, *Agromyza errans* Meigen (mon.). *Dizygomyza*, subg. *Dendromyza* Hendel, 1931:22. Type-species, *Agromyza carbonaria* Zetterstedt (orig. des.).

Refs. — Nowakowski, 1962:97; Spencer, 1966b:143 (key to spp.). KALLIMA (Frost), 1936:299 (*Agromyza*). Type-locality: Panama, Canal Zone, Barro Colorado I. Distr. — Panama. Refs. — Frick, 1959: 375, fig. 60; Spencer, 1963:333, 1966b:142, 143 (key). Holotype in U.S.N.M.

KUHLMANNI Spencer, 1966b:143, figs. 3-5 (male genitalia). Type-locality: Brazil, São Paulo, Capital, Cantareira. Distr. — Brazil. Holotype in Dep. Zool., São Paulo.

LANEI Spencer, 1966b:144, figs. 6, 7-8 (male genitalia). Type-locality: Brazil, Minas Gerais, Sapucaí-Mirim, Cidade Azul. Distr. — Brazil. Holotype in Dep. Zool., São Paulo.

RABELLOI Spencer, 1966b:145, figs. 9, 10-11 (male genitalia). Type-locality: Brazil, Mato Grosso, Rio Paraná. Distr. — Brazil. Holotype in Dep. Zool., São Paulo.

XANTHOPHORA (Schiner), 1868:291 (*Agromyza*). Type-locality: "South America". Distr. — Mexico, St. Vincent Is., Honduras, Costa Rica, Brazil. Refs. — Frick, 1957:202, 1959:376; Spencer, 1963:333, fig. 47 (male genitalia), 1966b:146, fig. 12 (male genitalia).

picta Coquillett, 1902:188 (*Agromyza*). Type-locality: Mexico, Tabasco, Frontera. Holotype in U.S.N.M.

dorsocentralis Frost, 1936:307 (*Agromyza*). Type-locality: Panama, Patilla Point. Holotype in U.S.N.M.

Genus PRASPEDOMYZA Hendel

Dizygomyza, subg. PRASPEDOMYZA Hendel, 1931:177. Type-species, *approximata* Hendel (orig. des.).

Praspedomyza Hendel; Spencer, 1966b:146.

FLAVIANTENNATA Spencer, 1966b:147, fig. 15 (male genitalia). Type-locality: Brazil, São Paulo, Barueri. Distr. — Brazil. Holotype in Dep. Zool., São Paulo.

PEULLAE (Malloch), 1934:476 (*Agromyza*). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Chile, Argentina. Ref. — Spencer, 1963:337, fig. 51 (male genitalia). Holotype in B.M.

Genus AMAUROMYZA Hendel

Dizygomyza, subg. AMAUROMYZA Hendel, 1931:59. Type-species, *Agromyza lamii* Kaltenbach (orig. des.).

Amauromyza Hendel; Nowakowski, 1962:97.

FUSCIBASIS (Malloch), 1934:476 (*Agromyza*; as *maculosa* var.). Type-locality: Argentina, Rio Negro, L. Nahuel Huapi (eastern end), Bariloche. Distr. — Brazil, Argentina. Ref. — Spencer, 1963:336. Holotype in B.M.

MACULOSA (Malloch), 1913:302 (*Agromyza*). Type-locality: U.S.A., New York; Jamaica. Distr. — U.S.A. (New York, Florida, Hawaii), Jamaica, Venezuela, Brazil, Uruguay, Argentina. Ref. — Spencer, 1963:336, fig. 50. Holotype in B.M.

guaranitica Brèthes, 1920:283 (*Agromyza*). Type-locality: Brazil, Rio Grande do Sul.

Genus CALYCOMYZA Hendel

Dizygomyza, subg. CALYCOMYZA Hendel, 1931:65. Type-species, *Agromyza artemisiae* Kaltenbach (orig. des.).

Calycomyza Hendel; Frick, 1957:59-70 (as subgenus); Nowakewski, 1962:97.

ALLECTA (Melander), 1913:257 (*Agromyza*; as *platyptera* var.; nom. nov. for *lateralis* Williston). Type-locality: West Indies, St. Vincent. Distr. — U.S.A. (east of Rocky Mountains), Haiti, Puerto Rico, St. Vincent, Brazil. Refs. — Frick, 1956:298; Spencer, 1963:333, figs. 52a-b, 53a-b (male genitalia). Lectotype in Snow Entomological Collections, University of Kansas.

lateralis Williston, 1896:428 (*Agromyza*; preocc. Macquart, 1835). Type-locality: West Indies, St. Vincent.

ARTEMISIAE (Kaltenbach), 1856:236 (*Phytomyza*). Type-locality: "Germany". Distr. — Europe, Canada, U.S.A., Guatemala, Colombia. Refs. — Frick, 1956:294; Spencer, 1963:339, figs. 54a-b (male genitalia), 54c. Types lost.

CASSIAE (Frost), 1936:306 (*Agromyza*). Type-locality: Panama, Canal Zone, Barro Colorado I. Distr. — Panama. Ref. — Spencer, 1963:340, figs. 55a-c (male genitalia). Holotype in U.S.N.M.

CROTALARIIVORA Spencer, 1963:340, figs. 56a-c (male genitalia), 56d. Type-locality: Venezuela, Caracas, hill above Hotel Avila. Distr. — Venezuela. Holotype in coll. Spencer.

ECLIPTAE Spencer, 1963:341, figs. 57a (male genitalia), 57b. Type-locality: Brazil, Guanabara, Rio de Janeiro, Jardim Botânico. Distr. — Brazil. Holotype in coll. Spencer.

ECREGIA Spencer, 1963:341, figs. 58a, 58b (male genitalia). Type-locality: Brazil, São Paulo, Capital. Distr. — Brazil. Holotype in coll. Spencer.

EXQUISITA Spencer, 1963:342, figs. 59a, e, 59b-d (male genitalia). Type-locality: Brazil, São Paulo, Capital. Distr. — Brazil. Holotype in coll. Spencer.

FLAVINOTUM Frick, 1956:297, fig. 1c. Type-locality: U.S.A., New York, Elma. Distr. — U.S.A., Jamaica, Venezuela, Brazil. Ref. — Spencer, 1963:343, figs. 60a-d (male genitalia), 60c-f. Holotype in Science Museum, San Francisco.

HUMERALIS (Roser), 1840:63 (*Agromyza*). Type-locality: "Germany". Distr. — Europe, Canada, U.S.A., Hawaii, Japan, Africa, India, Australia,

- Fiji; Jamaica, Brazil, Argentina. Refs. — Frick, 1956:290, figs. 1b, 4a-e; Spencer, 1963:344, figs. 61a, 61b-d (male genitalia). Holotype in Museum für Naturkunde, Stuttgart, Germany.
- IPOMAEAE (Frost), 1931:74 (*Agromyza*). Type-locality: "Puerto Rico". Distr. — Puerto Rico, Dominican Republic, Jamaica, Brazil. Ref. — Spencer, 1963:345, figs. 62a-b, 62c (male genitalia). Holotype in U.S.N.M.
- IRREPERTA Spencer, 1963:345, fig. 63a (male genitalia). Type-locality: Brazil, São Paulo, Capital, Água Funda. Distr. — Brazil. Holotype in coll. Spencer.
- JUCUNDACEA (Blanchard), 1954:22 (*Dizygomyza*). Type-locality: Argentina, Mendoza. Distr. — Argentina. Ref. — Spencer, 1963:351. Holotype in coll. Blanchard, Buenos Aires.
- LANTANAE Frick, 1956:287. Type-locality: "Trinidad". Distr. — U.S.A. (Texas), Mexico, Puerto Rico, Grenada, Trinidad, Venezuela. Ref. — Spencer, 1963:346, figs. 64a-b (male genitalia), 64c. Holotype in U.S.N.M.
- LONGICAUDA (Blanchard), 1954:24 (*Dizygomyza*). Type-locality: Argentina, Buenos Aires. Distr. — Argentina. Ref. — Spencer, 1963:346, fig. 65 (male genitalia). Holotype in coll. Blanchard, Buenos Aires.
- MALVAE (Burgess), in Comstock, 1880b:202 (*Oscinis*). Type-locality: U.S.A., District of Columbia. Distr. — U.S.A., Panama, Brazil. Refs. — Frick, 1956:298; Spencer, 1963:347, figs. 66a-c (male genitalia), 66d. Lectotype in U.S.N.M.
- MERIDIANA (Hendel), 1923:123 (*Agromyza*; nom. nov. for *meridionalis* Malloch). Type-locality: "Costa Rica". Distr. — Mexico, Costa Rica, British Guiana, Brazil. Refs. — Spencer, 1963:348, fig. 68 (male genitalia); 1966b:146.
- meridionalis* Malloch, 1914:89 (*Agromyza*; preocc. Strobl, 1900). Type-locality: "Costa Rica".
- RICHARDSI Spencer, 1963:348, figs. 69a, 69b-d (male genitalia). Type-locality: British Guiana, Mazaruni. Distr. — British Guiana. Holotype in B.M.
- VERBENIVORA Spencer, 1963:350, figs. 71a-b (male genitalia). Type-locality: Venezuela, Caracas, hill behind Hotel Avila. Distr. — Venezuela. Holotype in coll. Spencer.
- VERBESINAE Spencer, 1963:351, figs. 72a, c, 72b (male genitalia). Type-locality: Brazil, São Paulo, Capital, Água Funda. Distr. — Brazil. Holotype in coll. Spencer.

Genus LIRIOMYZA Mik

- LIRIOMYZA Mik, 1894:289. Type-species, *urophorina* Mik (mon.).
- ANDINA (Malloch), 1934:471 (*Agromyza*). Type-locality: Chilc, Los Andes. Distr. — Chilc. Ref. — Spencer, 1963:353. Holotype in B.M.
- BACCHARIDIS Spencer, 1963:354, figs. 74a-b (male genitalia), 74c. Type-locality: Colombia, Tequendama Falls. Distr. — Colombia. Holotype in coll. Spencer.
- BARROCOLORADENSIS (Frost), 1936:301 (*Agromyza*). Type-locality: Panama, Barro Colorado I. Distr. — Panama. Ref. — Spencer, 1963:355, figs. 75a-b (male genitalia).



- BIFORMATA (Becker), 1919:213 (*Agromyza*). Type-locality: "Ecuador".
Distr. — Ecuador. Ref. — Spencer, 1963:372. Holotype in Musée d'Histoire Naturelle, Paris.
- BRASSICAE (Riley), 1885:322 (*Oscinis*). Type-locality: U.S.A., Montana, St. Louis. Distr. — Europe, U.S.A., Hawaii; British Guiana (cosmopolitan). Ref. — Spencer, 1963:356.
mitis Curran, 1931:97 (*Phytomyza*). Type-locality: U.S.A., Manitoba.
hawaiiensis Frick, 1952:513. Type-locality: "Hawaii". Holotype (♀) in coll. Hawaii Sugar Planters Association.
- BRAZILIAE Spencer, 1963:356, figs. 76a-b (male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Holotype in B.M.
- BRAZILIENSIS (Frost), 1939:97 (*Agromyza*). Type-locality: Brazil, São Paulo, Campinas. Distr. — Colombia, Ecuador, Brazil. Ref. — Spencer, 1963:357, figs. 77a-c (male genitalia), 77d. Holotype in U.S.N.M.
ecuadorensis Frost, 1939:99 (*Agromyza*). Type-locality: Ecuador, Los Baños. Holotype in U.S.N.M.
- EULNESIAE Spencer, 1963:360, figs. 79a-b (male genitalia), 79c-d. Type-locality: Venezuela, Caracas, Botanical Gardens. Distr. — Venezuela. Holotype in coll. Spencer.
- CHILENSIS (Malloch), 1934:469 (*Agromyza*). Type-locality: Chile, Llanquihue, Puerto Varas, Casa Pangue. Distr. — Chile. Ref. — Spencer, 1963:361. Holotype in B.M.
- COMMELINAE (Frost), 1931:72 (*Agromyza*). Type-locality: West Indies, St. Vincent. Distr. — U.S.A. (Florida), Jamaica, St. Vincent, Brazil, Argentina. Refs. — Silva & Oliveira, 1952: figs. 1-22; Spencer, 1963:361, figs. 80a-b (male genitalia).
bahamondesi Blanchard, 1954:28. Type-locality: Argentina, Córdoba. Holotype in coll. Blanchard, Buenos Aires.
- CUCUMIFOLIAE Blanchard, 1938:352. Type-locality: Argentina, Buenos Aires. Distr. — Argentina. Ref. — Spencer, 1963:372. Holotype in coll. Blanchard, Buenos Aires.
- GAYI (Porter), 1915:56 (*Agromyza*). Type-locality: "Chile". Distr. — Chile. Ref. — Spencer, 1963:372. Location of types not known.
- GRANDIS Spencer, 1963:361, figs. 81a-d (male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Holotype in B.M.
- HUIDORENSIS Blanchard, 1926:10. Type-locality: Argentina, Buenos Aires. Distr. — Argentina. Ref. — Spencer, 1963:372. Holotype in coll. Blanchard, Buenos Aires.
- INSIGNIS Spencer, 1963:363, fig. 83. Type-locality: Costa Rica, La Suiza. Distr. — Costa Rica, Brazil. Ref. — Spencer, 1966b:147. Holotype in B.M.
- KUSCHELI Spencer, 1964a:253, figs. 1a-c (male genitalia). Type-locality: Juan Fernandez Is., Masatierra I. Distr. — Juan Fernandez Is. (Masatierra I.). Holotype in Museo Nacional, Santiago.
- LANGEI Frick, 1951:81. Type-locality: U.S.A., California. Distr. — U.S.A., Brazil, Argentina, Juan Fernandez Is. Refs. — Spencer, 1963:359 (as *bryoniae* Kaltcnbach, misident.), figs. 78a-c (male genitalia), 1965c:108. Holotype in Science Museum, San Francisco.
decora Blanchard, 1954:31. Type-locality: Argentina, Buenos Aires. Holotype in coll. Blanchard, Buenos Aires.

- MICROGLOSSAE Spencer, 1963:363, figs. 84a, d, 84b-c (male genitalia). Type-locality: Brazil, São Paulo, Capital. Distr. — Brazil. Holotype in coll. Spencer.
- MUNDA Frick, 1957:61. Type-locality: U.S.A., California. Distr. — U.S.A., Jamaica. Refs. — Steyskal, 1964:388, figs. 1a-f (male genitalia); Spencer, 1965a:36, figs. 4a-b (male genitalia). Holotype in California Academy of Sciences.
- guytona* Freeman, 1958:344. Type-locality: U.S.A., Alabama. Ref. — Spencer, 1963:362, figs. 82a-b (male genitalia), 82c. Holotype in U.S.N.M.
- PAGANA (Malloch), 1934:475 (*Agromyza*). Type-locality: Argentina, Rio Negro, L. Nahuel Huapi. Distr. — Argentina. Ref. — Spencer, 1963:364. Holotype in B.M.
- PARANAENSIS Spencer, 1966b:148, figs. 16, 17-19 (male genitalia). Type-locality: Brazil, São Paulo, Pôrto Cabral. Distr. — Brazil. Holotype in Dep. Zool., São Paulo.
- PATAGONICA (Malloch), 1934:472 (*Agromyza*). Type-locality: Argentina, Rio Negro, L. Gutierrez. Distr. — Chile, Argentina. Ref. — Spencer, 1963:364, figs. 85a-d (male genitalia). Holotype in B.M.
- POLITELLA (Malloch), 1934:473 (*Agromyza*). Type-locality: Argentina, Vidma. Distr. — Chile, Argentina. Ref. — Spencer, 1963:365. Holotype in B.M.
- QUADRATA (Malloch), 1934:471 (*Agromyza*). Type-locality: Chile, Concepción. Distr. — Colombia, Chile. Ref. — Spencer, 1963:365, figs. 86a, 86b-d (male genitalia). Holotype in B.M.
- subandina* Blanchard, 1954:34. Type-locality: Argentina, Mendoza, Tunuyán. Holotype in coll. Blanchard, Buenos Aires.
- SABAZIAE Spencer, 1963:366, figs. 87a-b (male genitalia), 87c. Type-locality: Venezuela, Caracas, near Hotel Humboldt. Distr. — Venezuela, Colombia. Holotype in coll. Spencer.
- SATIVAE Blanchard, 1938:354. Type-locality: Argentina, La Pampa. Distr. — Argentina. Ref. — Spencer, 1963:372. Holotype in coll. Blanchard, Buenos Aires.
- SCHMIDTI (Aldrich), 1929:89 (*Agromyza*). Type-locality: "Costa Rica". Distr. — U.S.A. (Florida), Costa Rica. Ref. — Spencer, 1963:367, figs. 88a-b (male genitalia). Holotype in U.S.N.M.
- SCHWABEI Spencer, 1963:368, fig. 89 (male genitalia). Type-locality: Chile, Viña. Distr. — Chile. Holotype in Deutsches Entomologisches Institut.
- SENECIELLA Spencer, 1963:368, figs. 90a-c (male genitalia), 90d. Type-locality: Brazil, São Paulo, Capital. Distr. — Brazil. Holotype in coll. Spencer.
- SIMULATOR (Malloch), 1934:474 (*Agromyza*). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile, Argentina. Ref. — Spencer, 1963:369. Holotype in B.M.
- SOLANITA Spencer, 1963:369, figs. 91a-c (male genitalia), 91d-e. Type-locality: Colombia, Bogotá, Ciudad Universitaria. Distr. — Colombia. Holotype in coll. Spencer.

- SOROSIS (Williston), 1896:429 (*Agromyza*). Type-locality: West Indies, St. Vincent, Distr. — Southern U.S.A., Jamaica, St. Vincent, Venezuela, British Guiana. Ref. — Spencer, 1963:370, figs. 92a-c (male genitalia), 92d-e. Lectotype in B.M.
- marginalis* Malloch, 1913:283 (*Agromyza*; as *melanopyga* var.). Type-locality: U.S.A., South Carolina. Holotype in U.S.N.M.
- STRIGOSA Spencer, 1963:371, figs. 93a-b (male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Holotype in B.M.
- TRIFOLI (Burgess) in Comstock, 1880a:201 (*Oscinis*). Type-locality: U.S.A., District of Columbia. Distr. — Southern U.S.A., Barbados. Ref. — Spencer, 1963:354 (as *archboldi* Frost, misident.), figs. 73a-c (male genitalia), 1965a:37, figs. 7a-c (male genitalia). Neotype in U.S.N.M.
- TUBIFER (Melander), 1913:266 (*Agromyza*). Type-locality: "Haiti". Distr. — Haiti. Ref. — Spencer, 1963:372. Holotype in U.S.N.M.

Genus METOPOMYZA Enderlein


- METOPOMYZA Enderlein, 1936:180. Type-species, *Agromyza flavonotata* Haliday (mon.).
- TEQUENDAMAE Spencer, 1963:373. Type-locality: Colombia, Tequendama Falls. Distr. — Colombia. Holotype in coll. Spencer.

Genus HAPLOMYZA Hendel

- Antineura* Melander, 1913:149 (preocc. Osten Sacken, 1881). Type-species, *togata* Melander (orig. des.).
- HAPLOMYZA Hendel, 1914:73 (nom. nov. for *Antineura* Melander). Type-species, *Antineura togata* Melander (aut.).
- Ref. — Spencer, 1963:373.
- BULLATI Spencer, 1963:374, figs. 95a, c, 95b-d (male genitalia). Type-locality: Brazil, São Paulo, Capital, Água Funda. Distr. — Brazil. Holotype in coll. Spencer.
- LOPESI Oliveira & Silva, 1954:25, figs. 1-2, 8-18, 3-7 (male genitalia). Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil. Ref. — Spencer, 1963:375. Holotype in Instituto Oswaldo Cruz, Guanabara.
- TIGRENSIS Spencer, 1963:375, figs. 96a, 96b-c (male genitalia). Type-locality: Argentina, Buenos Aires, Tigre. Distr. — Argentina. Holotype in coll. Spencer.
- VERBASCIFOLI Spencer, 1963:376, figs. 97a-c (male genitalia), 97d. Type-locality: Brazil, São Paulo, Capital, Água Funda. Distr. — Brazil. Holotype in coll. Spencer.

Genus PHYTOLIRIOMYZA Hendel

- Liriomyza*, subg. PHYTOLIRIOMYZA Hendel, 1931:203. Type-species, *Agromyza perpusilla* Meigen (mon.).
- Ref. — Spencer, 1965b:657-663.



- ARCTICA (Lundbeck), 1901:304 (*Agromyza*). Type-locality: "Greenland". Distr. — Europe, N. America, Brazil, Paraguay, Chile; semi-cosmopolitan. Ref. — Spencer, 1963:377, figs. 98a-e (male genitalia)., Leetotype in Universitetets Zoologiska Museum, Copenhagen.
- immaculata* Coquillett, 1902:185 (*Odinia*). Type-locality: U.S.A., New Hampshire. Holotype in U.S.N.M.
- halterata* Beeker, 1908:169 (*Agromyza*). Type-locality: "Canary Is.". Holotype in Zoologisches Museum, Berlin.
- formosensis* Malloch, 1914a:315 (*Agromyza*). Type-locality: "Formosa". Holotype in Hungarian National Museum, Budapest.
- IMPERFECTA (Malloch), 1934:474 (*Agromyza*). Type-locality: Chile, Llanquihue, Puerto Varas. Distr. — U.S.A. (California), Chile. Ref. — Spencer, 1963:379, figs. 99a-c (male genitalia). Holotype in U.S.N.M.
- LONGIPENIS Spencer, 1964a:253, figs. 2a-d (male genitalia). Type-locality: Juan Fernandez Is., Masafuera I. Distr. — Juan Fernandez Is. Holotype in Museo Nacional, Santiago.
- LURIDA Spencer, 1963:379, figs. 100a, 100b-c (male genitalia). Type-locality: Brazil. Santa Catarina, Nova Teutônia. Distr. — Brazil. Holotype in B.M.
- PICEA Spencer, 1963:380, figs. 101a-e (male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Holotype in B.M.

Genus PHYTOMYZA Fallén

- PHYTOMYZA Fallén, 1810:21, 26. Type-species, *flaveola* Fallén (mon.).
- CAESALPINIAE Blanchard, 1954:17. Type-locality: Argentina, Mendoza. Distr. — Argentina. Ref. — Spencer, 1963:381. Holotype in coll. Blanchard, Buenos Aires.
- CENTRALIS Frost, 1936:317. Type-locality: "Costa Rica". Distr. — Costa Rica. Ref. — Spencer, 1963:381. Holotype in U.S.N.M.
- ENIGMA Malloch, 1934:485. Type-locality: Chile, Llanquihue, Aneud. Distr. — Chile. Ref. — Spencer, 1963:382. Holotype in B.M.
- MELANOGASTER Thomson, 1869:610. Type-locality: "Patagonia". Distr. — Chile. Ref. — Spencer, 1963:382, figs. 102a, 102b (male genitalia).
- PAMPEANA Blanchard, 1954:18. Type-locality: Argentina, La Pampa. Distr. — Argentina. Ref. — Spencer, 1963:383. Holotype in coll. Blanchard, Buenos Aires.
- PLATENSIS Brèthes, 1923:154. Type-locality: Argentina, Buenos Aires. Distr. — Argentina. Ref. — Spencer, 1963:383. Location of types not known.
- TUEUMANA Blanchard, 1954:20. Type-locality: "Argentina". Distr. — Argentina. Ref. — Spencer, 1963:383. Holotype in coll. Blanchard, Buenos Aires.
- WILLIAMSONI Blanchard, 1954:20. Type-locality: Argentina, Tucumán. Distr. — Argentina. Ref. — Spencer, 1963:383. Holotype in coll. Blanchard, Buenos Aires.



LITERATURE CITED

ALDRICH, J. M.

1929. Three new acalyptate Diptera. *Proc. Ent. Soc. Washington* 31:89-91.

BECKER, T.

1908. Dipteren der Kanarischen Inseln. *Mitt. Zool. Mus. Berlin* 4:1-180, 4 pls.
1919. Diptères, Brachycères. *Mission Arc Méridien Amérique du Sud*, Paris 10:163-215.

EEZZI, M. & J. S. TAVARES

1916. Alguns Muscideos cecidogênicos do Brazil. *Brotéria, Zool.*, 14:166-169.

BLANCHARD, E.

1926. A dipterous leaf-miner on *Cineraria*, new to science. *Rev. Soc. Ent. Argent.* 1:10-11.
1938. Descripciones y anotaciones de dipteros argentinos. *An. Soc. Cient. Argent.* 126:345-386.
1954. Sinopsis de los Agromizidos argentinos (Diptera, Agromyzidae). *Publ. Inst. San. Veg. (A)* 10(56):50 pp., figs.

BRASCHNIKOV, W. C.

1897. Zur Biologie und Systematik einiger Arten minierender Dipteren. *Izv. Moskva Selsk. Khoz. Inst.* 3(2):19-43, 7 figs., pl. 2.

BRÉTHES, J.

1920. Insectos útiles y daños de Rio Grande do Sul y de la Plata. *An. Soc. Rur. Argent.* 54:281-290.
1923. Sur un Diptère mineur de feuilles de *Salvia splendens* et deux Hymenoptères parasites. *Rev. Zool. Agr., App.* 22:153-158.

COMSTOCK, J. H.

- [1880a]. *Report upon cotton insects*. 511 pp., 77 figs., 3 pls. Washington, D. C., 1879.
[1880b]. Report of the entomologist. *U. S. Dept. Agric. Comnr. Agr. Rept.* 1879:185-248, 16 pls.

COQUILLETT, D. W.

1902. New acalyptate Diptera from North America. *Jour. N. Y. Ent. Soc.* 10:177-191.

CURRAN, C. H.

1928. Diptera or two-winged flies. *Scient. Surv. Porto Rico & Virgin Is.*, New York 11:1-118, 39 figs.
1931. Four new Diptera in the Canadian National Collection. *Canad. Ent.* 63:93-98.



ENDERLEIN, G.

1936. Zweiflügler, Diptera, in Brohmer et. al.: *Tierwelt Mitteleuropas* 6 (Lief. 2, Insekt. Teil 3): 179-183.

FALLÉN, C. F.

1810. *Specimen entomologicum novam Diptera disponendi methodum exhibens*: 1-26. Lund.

FREEMAN, C. C.

1958. *Liriomyza guytona*: A new species of Agromyzid leaf-miner. *Ann. Ent. Soc. America* 51:344-345, 1 fig.

FRICK, K. E.

1951. *Liriomyza langei*, a new species of leaf-miner of economic importance in California (Diptera: Agromyzidae). *Pan-Pacific Ent.* 21:81-88, 2 figs.
1952. A generic revision of the family Agromyzidae (Diptera) with a catalogue of New World species. *Univ. Calif. Publ. Ent.* 8:339-452, 34 figs.
1953. Further studies on Hawaiian Agromyzidae (Diptera) with descriptions of four new species. *Proc. Hawaii. Ent. Soc.* 15:207-215.
1956. Revision of the North American *Calycomyza* species north of Mexico (*Phytobia*: Agromyzidae: Diptera). *Ann. Ent. Soc. Amer.* 49:284-300, 1 fig., 2 pls.
1957. Nearctic species in the *Liriomyza pusilla* complex. N° 2. *L. munda* and two other species attacking crops in California (Diptera: Agromyzidae). *Pan-Pacific Ent.* 33:59-70.
1959. Synopsis of the species of agromyzid leaf-miners described from North America (Diptera). *Proc. U. S. Nat. Mus.* 108:347-465, 170 figs.

FROGGATT, W. W.

1919. The lantana fly (*Agromyza lantanae*). *Agr. Gaz. N. S. Wales* 30:665-668, 1 pl.

FROST, S. W.

1931. New species of West Indian Agromyzidae (Diptera). *Ent. News* 42:72-76.
1936. New Central American Agromyzidae. *Ann. Ent. Soc. Amer.* 29:298-318.
1939. Two new species of *Agromyza* from South America (Dipt.: Agromyzidae). *Ent. News* 50:97-100.

HENDEL, F.

1914. Namensänderungen (Dipt.). *Ent. Mitt.* 3:73.
1920. Die paläarktischen Agromyziden (Dipt.). (Prodromus einer Monographic). *Arch. f. Naturg.* (1918) 84 A (7):109-174, 4 figs.



1923. Neue europäische *Melanagromyza*-Arten (Dipt.). *Konowia* 2: 142-145.

1931. Agromyzidae, in E. Lindner: *Fliegen der paläarktischen Region* 6:1-570.

KALTENBACH, J. H.

1856. Die deutschen Phytophagen aus der Klasse der Insekten. *Verh. naturhist. Vereins preuss. Rheinl. u. Westfalens* 26:106-224.

LIOY, P.

1864. I ditteri distribuiti secondo un nuovo metodo di classificazione naturale. *Atti. Ist. Veneto Sci., Lett. ed Arti* (3)9:1311-1317.

LOEW, H.

1863. Diptera Americae septentrionalis indigena. Centuria tertia. *Berlin. ent. Zeitschr.* 7:1-55.

1869. Idem. Centuria octava. *Ibidem*, 13:1-52.

LUNDBECK, W.

1901. Diptera Groenlandica. *Videnskab. Medd. Naturh. For., Copenhagen* 5:236-314.

MACQUART, J.

1835. *Histoire Naturelle des Insectes. Diptères* 2:703 pp., 12 pls. Paris.

MALLOCH, J. R.

1913. A revision of the species in *Agromyza* Fallén, and *Cerodontha* Rondani (Diptera). *Ann. Ent. Soc. Amer.* 6:269-336, 4 pls.

1914. Descriptions of new species of *Agromyza* from Porto Rico. *Proc. Ent. Soc. Washington* 16:89-90.

1914a. Formosan Agromyzidae. *Ann. hist.-nat. Mus. Hung.* 12:306-336.

1915. Flies of the genus *Agromyza*, related to *A. virens*. *Proc. U.S. Nat. Mus.* 49:103-108, pl. 36.

1921. A new genus of Agromyzidae (Diptera). *Bull. Brooklyn Ent. Soc.* (1920)15:147-148.

1934. Agromyzidae, in *Diptera of Patagonia and South Chile* 6(5): 465-486. British Museum (Natural History), London.

MARELLI, C.

1926. La *Agromyza productora* de agallas nodicolas en la lagunilla. *Rev. Soc. Ent. Argent.* 1:13-21.

MELANDER, A. L.

1913. A synopsis of the dipterous groups Agromyzinae, Millehiinae, Oethiophilinae and Geomyzinae. *Jour. N. Y. Ent. Soc.* 21:219-273, 283-300, 1 pl.



MEENDES, L. O. T.

1940. O minador da batatinha. *J. Agron.*, Piracicaba 3:207-220.

MIK, J.

1894. Über eine neue *Agromyza*, deren Larven in den Blütenknospen von *Lilium martagon* leben. *Wien. ent. Z.* 13:284-290, 1 pl.

NOWAKOWSKI, J. T.

1962. Introduction to a systematic revision of the family Agromyzidae (Diptera). *Ann. Zool.*, Warsaw 20:67-183.

OLIVEIRA, S. J. & G. A. DA SILVA

1954. Sobre uma nova espécie de Agromyzidae (Diptera) cujas larvas minam folhas de *Solanum argenteum*. *Rev. Brasil. Entom.* 1:25-30, 18 figs.

PHILIPPI, R. A.

1865. Aufzählung der chilnischen Dipteren. *Verh. zool-bot. Ges. Wien* 15:595-782, 7 pls.

PORTER, C. E.

1915. Descripción de un nuevo Diptero chileno. *Bol. Mus. Nac. Chile* 8:56-57.

RILEY, C. V.

1885. The cabbage *Oscinis*, *Oscinis brassicae* n. sp., in *U. S. Dept. Agric. Ann. Rept.*: p. 322.

RONDANI, C.

1861. *Dipterologiae Italicae Prodomus* 4:10. Parmae.

1875a. Species Italicae ordinis Dipteriorum (Muscaria Rndn.) collectae et observatae. Stirps XXIII. Agromyzinae. *Boll. Soc. Ent. Ital.* 7:166-191.

1875. *Dipterologiae Italicae Prodomus* 8 (Stirps 23 = Pt. 3):26 pp. Firenze.

ROSER, C. VON

1840. Erster Nachtrag zu dem im Jahre 1834 bekannt gemachten Verzeichnisse in Württemberg vorkommender zweiflügliger Insekten. *Correspondenzbl. K. Württemb. Landw. Ver.*, Stuttgart 37 (n. s., 17) (1):49-64.

SASAKAWA, M.

1958. The female terminalia of the Agromyzidae, with description of a new genus. *Sci. Rep. Saikyo Univ. Agric.* 10:133-150, 5 pls.

1963. A new species of *Japanagromyza* from Central America (Diptera: Agromyzidae). *Wasmann J. Biol.* 21:95-96.



SCHINER, J. R.

1868. *Reise der österreichischen Fregatte Novara um die Erde*, Zool. Theil, Diptera 2(1)b:290-291.

SILVA, G. A. DA & S. J. DE OLIVEIRA

1952. Sobre um "Agromyzidae" (Diptera) e suas larvas minam folhas de trapoeiraba (Commelinaceae). *Rev. Brasil. Biol.* 12(3):293-299, 22 figs.

SPENCER, K. A.

1963. A synopsis of the neotropical Agromyzidae (Diptera). *Trans. R. Ent. Soc. London* 115:291-389, 104 figs.
- 1963a. The Australian Agromyzidae (Diptera, Insecta). *Rec. Austr. Mus., Sydney* 25(15):305-354.
- 1964a. Some Agromyzidae (Diptera) from the Juan Fernandez Islands. *Pacific Insects* 6(2):251-255.
- 1964b. A revision of the palaearctic species of the genus *Ophiomyia* Braschnikov. *Beitr. Ent.* 14(7/8):773-822.
- 1965a. A clarification of the status of *Liriomyza trifolii* (Burgess) and some related species. *Proc. Ent. Soc. Washington* 67:32-40, 7 figs.
- 1965b. The genus *Phytoliriomyza* Hendel (Agromyzidae: Diptera). *Ann. Mag. Nat. Hist.* (13)7:657(1964).
- 1965c. Agromyzidae (Diptera) from the Azores and Madeira. *Bol. Mus. Municipal Funchal, Madeira* 19:104-110, 3 figs.
- 1966a. A revision of European species of the genera *Melanagromyza* Hendel and *Hexomyza* Enderlein, with a supplement on the genus *Ophiomyia* Braschnikov (Diptera: Agromyzidae). *Beitr. Ent.* 16(1/2):3-60.
- 1966b. Notes on Neotropical Agromyzidae (Diptera). *Papéis Avulsos Dep. Zool., São Paulo* 19(11):141-150, 19 figs.
- 1966c. New and interesting Agromyzidae (Diptera) from Florida. *Stuttgart. Beitr. Ent.* 158:1-20, 54 figs.

STEYSKAL, G. C.

1964. Descriptive and synonymical notes on *Liriomyza munda* (Diptera: Agromyzidae). *Ann. Ent. Soc. Amer.* 57(3):388-389, 1 fig.

STROBL, G.

1902. [Studies on the Dipteran fauna of Bosnia-Herzegovina]. *Glasnik Zemaljski Muscj u Bosni i Hercegovini* 14:461-517.
1906. Spanische Dipteren. II. Beitrag. *Mem. Soc. Esp. Hist. Nat., Madrid* 3:271-422.



THOMSON, C. G.

1869. *K. Svenska fregatten Eugenie resa omkring jorden 2* (Zoologie). Sec. 1: Insekter, 617 pp., 9 pls. Stockholm, "1868".

WILLISTON, S. W.

1896. On the Diptera of St. Vincent (West Indies). *Trans. Ent. Soc. London* (1896):253-449.

INDEX

Agromyza Fallén, 1810	83.2
aldrichi (Frick), 1952 (Melanagromyza), Japanagromyza	83.2
allecta (Melander), 1913 (Agromyza), Calycomyza	83.8
alternantherae Spencer, 1963, Melanagromyza	83.3
Amauromyza Hendel, 1931	83.7
andalusiaca Strobl, 1906, Agromyza	83.2
andensis (Spencer), 1963 (Phytobia), Cerodontha	83.6
andina (Malloch), 1934 (Agromyza), Liriomyza	83.9
angolae (Malloch), 1934 (Agromyza), Melanagromyza	83.3
angulata (Loew), 1869 (Agromyza), Cerodontha	83.6
Antineura Melander, 1913	83.12
apfelbecki Strobl, 1902, Agromyza	83.2
approximata (Frost), 1936 (Agromyza), Melanagromyza	83.3
arctica (Lundbeck), 1901 (Agromyza), Phytoliriomyza	83.13
arnaudi Sasakawa, 1963, Japanagromyza	83.2
artemisiae (Kaltenbach), 1856 (Phytomyza), Calycomyza	83.8
baccharidis Spencer, 1963, Liriomyza	83.9
bahamondesi Blanchard, 1954, Liriomyza	83.10
barrocoloradensis (Frost), 1936 (Agromyza), Liriomyza	83.9
biformata (Becker), 1919 (Agromyza), Liriomyza	83.10
brassicae (Riley), 1885 (Oscinis), Liriomyza	83.10
braziliae Spencer, 1963, Liriomyza	83.10
braziliana Spencer, 1963, Cerodontha	83.6
braziliensis (Frost), 1939 (Agromyza), Liriomyza	83.10
bullati Spencer, 1963, Haplomyza	83.12
bulposiae Spencer, 1963, Liriomyza	83.10
buscki (Frost), 1936 (Agromyza), Ophiomyia	83.5
caerulea (Malloch), 1913 (Agromyza), Melanagromyza	83.3
cacsalpiniae Blanchard, 1954, Phytomyza	83.13
Calycomyza Hendel, 1931	83.8
camarae Spencer, 1963, Ophiomyia	83.5
cassiae (Frost), 1936 (Agromyza), Calycomyza	83.8
centralis Frost, 1936, Phytomyza	83.13
centrosemae (Frost), 1936 (Agromyza), Japanagromyza	83.2
Cerodontha Rondani, 1861	83.6
chenopodii Spencer, 1963, Melanagromyza	83.3
chilensis (Malloch), 1934 (Agromyza), Liriomyza	83.10
colombiensis Spencer, 1963, Melanagromyza	83.3




commelinae (Frost), 1931 (Agromyza), Liriomyza	83.10
compositoides Spencer, 1963, Melanagromyza	83.3
crotalarivora Spencer, 1963, Calycomyza	83.8
crotonis (Frost), 1936 (Agromyza), Melanagromyza	83.3
cruciata (Blanchard), 1938 (Triticomyza), Cerodontha	83.6
cucumifoliae Blanchard, 1938, Liriomyza	83.10
cunctanoides Blanchard, 1954, Melanagromyza	83.3
currani (Frost), 1936 (Agromyza), Japanagromyza	83.2
curvibrissata (Frost), 1936 (Agromyza), Ophiomyia	83.5
<i>decora</i> Blanchard, 1954, Liriomyza	83.10
<i>Dendromyza</i> Hendel, 1931	83.7
diadema (Melandar), 1913 (Agromyza), Melanagromyza	83.3
distincta Spencer, 1963, Melanagromyza	83.3
Dizygomyza Hendel, 1920	83.6
dorsalis (Loew), 1863 (Odontocera), Cerodontha	83.6
dorsocentralis (Frost), 1936 (Agromyza), Phytobia	83.7
ecliptae Spencer, 1963, Calycomyza	83.8
ecuadorensis Frost, 1939 (Agromyza), Liriomyza	83.10
egregia Spencer, 1963, Calycomyza	83.8
enigma Malloch, 1934, Phytomyza	83.13
erigeronis Spencer, 1963, Melanagromyza	83.4
eupatoriae (Malloch), 1915 (Agromyza), Melanagromyza	83.4
exquisita Spencer, 1963, Calycomyza	83.8
fasciculata (Malloch), 1934 (Agromyza), Ophiomyia	83.5
flaviantennata Spencer, 1966, Praspodomyza	83.7
flavifrons (Philippi), 1865 (Micromma), Cerodontha	83.6
flavinotum Frick, 1956, Calycomyza	83.8
floridensis Spencer, 1963, Melanagromyza	83.4
floris Spencer, 1963, Melanagromyza	83.4
formosensis (Malloch), 1914 (Agromyza), Phytoliriomyza	83.13
frosti (Frick), 1952 (Agromyza), Japanagromyza	83.2
fulvithorax Malloch, 1934, Cerodontha	83.6
fusca Spencer, 1963, Agromyza	83.2
fuscibasis (Malloch), 1934 (Agromyza), Amauromyza	83.8
gayi (Porter), 1915 (Agromyza), Liriomyza	83.10
grandis Spencer, 1963, Liriomyza	83.10
guaranitica (Brèthes), 1920 (Agromyza), Amauromyza	83.8
guytona Freeman, 1958, Liriomyza	83.11
halterata (Becker), 1919 (Agromyza), Phytoliriomyza	83.13
halterella Spencer, 1963, Melanagromyza	83.4
Haplomyza Hendel, 1914	83.12
hawaiiensis Frick, 1952, Liriomyza	83.10
helianthi Spencer, 1963, Melanagromyza	83.4
hirticeps (Malloch), 1934 (Agromyza), Ophiomyia	83.5
huidobrensis Blanchard, 1926, Liriomyza	83.10
humeralis (Roser), 1840 (Agromyza), Calycomyza	83.8




<i>Icteromyza</i> Hendel, 1931	83.6
<i>immaculata</i> (Coquillett), 1902 (Odinia), <i>Phytoliriomyza</i>	83.13
<i>imperfecta</i> (Malloch), 1934 (<i>Agromyza</i>), <i>Phytoliriomyza</i>	83.13
<i>inaequalis</i> (Malloch), 1914 (<i>Agromyza</i>), <i>Japanagromyza</i>	83.3
<i>inepta</i> (Spencer), 1963 (<i>Phytobia</i>), <i>Cerodontha</i>	83.6
<i>insignis</i> Spencer, 1963, <i>Liriomyza</i>	83.10
<i>insolens</i> Spencer, 1963, <i>Agromyza</i>	83.2
<i>insularis</i> (Malloch), 1913 (<i>Agromyza</i>), <i>Ophiomyia</i>	83.5
<i>ipomeae</i> (Frost), 1931 (<i>Agromyza</i>), <i>Calycomyza</i>	83.9
<i>iridescens</i> (Frost), 1936 (<i>Agromyza</i>), <i>Japanagromyza</i>	83.3
<i>irreperita</i> Spencer, 1963, <i>Calycomyza</i>	83.9
<i>jamaicensis</i> Spencer, 1963, <i>Japanagromyza</i>	83.3
<i>Japanagromyza</i> Sasakawa, 1958	83.2
<i>jacundacea</i> (Blanchard), 1954 (<i>Dizygomyza</i>), <i>Calycomyza</i>	83.9
<i>kallima</i> (Frost), 1936 (<i>Agromyza</i>), <i>Phytobia</i>	83.7
<i>kuhlmanni</i> Spencer, 1966, <i>Phytobia</i>	83.6
<i>kuhlmanni</i> Spencer, 1966, <i>Phytobia</i>	83.7
<i>lacustris</i> (Malloch), 1934 (<i>Agromyza</i>), <i>Melanagromyza</i>	83.4
<i>lanci</i> Spencer, 1966, <i>Phytobia</i>	83.7
<i>langei</i> Frick, 1951, <i>Liriomyza</i>	83.10
<i>lantanae</i> Frick, 1951, <i>Calycomyza</i>	83.9
<i>lantanae</i> (Froggatt), 1919 (<i>Agromyza</i>), <i>Ophiomyia</i>	83.5
<i>lateralis</i> (Williston), 1896 (<i>Agromyza</i>), <i>Calycomyza</i>	83.8
<i>limnoagromyza</i> Malloch, 1921	83.3
<i>lini</i> Spencer, 1963, <i>Melanagromyza</i>	83.4
<i>Liriomyza</i> Mik, 1894	83.9
<i>longicauda</i> (Blanchard), 1954 (<i>Dizygomyza</i>), <i>Calycomyza</i>	83.9
<i>longicauda</i> (Curran), 1928 (<i>Agromyza</i>), <i>Ophiomyia</i>	83.5
<i>longicaudalis</i> Spencer, 1963, <i>Melanagromyza</i>	83.4
<i>longipennis</i> Spencer, 1964, <i>Phytoliriomyza</i>	83.13
<i>longipennis</i> (Loew), 1869 (<i>Agromyza</i>), <i>Cerodontha</i>	83.7
<i>longisetæ</i> (Malloch), 1913 (<i>Agromyza</i>), <i>Melanagromyza</i>	83.4
<i>lopesi</i> Oliveira & Silva, 1954, <i>Haplomyza</i>	83.12
<i>lurida</i> Spencer, 1963, <i>Phytoliriomyza</i>	83.13
<i>maculosa</i> (Malloch), 1913 (<i>Agromyza</i>), <i>Amauromyza</i>	83.8
<i>mallochi</i> (Hendel), 1923 (<i>Agromyza</i>), <i>Melanagromyza</i>	83.4
<i>malvae</i> (Burgess), 1880 (<i>Oscinis</i>), <i>Calycomyza</i>	83.9
<i>marcellii</i> (Brèthes), 1920 (<i>Agromyza</i>), <i>Melanagromyza</i>	83.4
<i>marginalis</i> (Malloch), 1913 (<i>Agromyza</i>), <i>Liriomyza</i>	83.12
<i>Melanagromyza</i> Hendel, 1920	83.3
<i>melanogaster</i> Thomson, 1869, <i>Phytomyza</i>	83.13
<i>meridiana</i> (Hendel), 1923 (<i>Agromyza</i>), <i>Calycomyza</i>	83.9
<i>meridionalis</i> (Malloch), 1914 (<i>Agromyza</i>), <i>Calycomyza</i>	83.9
<i>Metopomyza</i> Enderlein, 1936	83.12
<i>microglossae</i> Spencer, 1963, <i>Liriomyza</i>	83.11
<i>Micromma</i> Philippi, 1865	83.6
<i>minima</i> (Malloch), 1913 (<i>Agromyza</i>), <i>Melanagromyza</i>	83.4
<i>mitis</i> (Curran), 1931 (<i>Phytomyza</i>), <i>Liriomyza</i>	83.10
<i>munda</i> Frick, 1957, <i>Liriomyza</i>	83.11

neotropica Spencer, 1963, <i>Melanagromyza</i>	83.4
nigricornis Becker, 1919, <i>Cerodontha</i>	83.6
nitidiventris (Malloch), 1934 (<i>Agromyza</i>), <i>Cerodontha</i>	83.6
nobilis Spencer, 1963, <i>Melanagromyza</i>	83.4
<i>Odontocera</i> Macquart, 1835	83.5
Ophiomyia Braschnikov, 1897	83.5
oralis (Frost), 1936 (<i>Agromyza</i>), <i>Ophiomyia</i>	83.5
orbitalis (Frost), 1936 (<i>Agromyza</i>), <i>Japanagromyza</i>	83.3
pagana (Malloch), 1934 (<i>Agromyza</i>), <i>Liriomyza</i>	83.11
pampeana Blanchard, 1954, <i>Phytomyza</i>	83.13
paranaensis Spencer, 1966, <i>Liriomyza</i>	83.11
patagonica (Malloch), 1934 (<i>Agromyza</i>), <i>Liriomyza</i>	83.11
peullae (Malloch), 1934 (<i>Agromyza</i>), <i>Praspedomyza</i>	83.7
pfaffiae Spencer, 1963, <i>Melanagromyza</i>	83.4
Phytobia Lioy, 1864	83.7
Phytoliriomyza Hendel, 1931	83.12
Phytomyza Fallén, 1810	83.13
picca Spencer, 1963, <i>Phytoliriomyza</i>	83.13
picta (Coquillett), 1902 (<i>Agromyza</i>), <i>Phytobia</i>	83.7
platensis Brèthes, 1923, <i>Phytomyza</i>	83.13
plaumanni Spencer, 1963, <i>Agromyza</i>	83.2
plumisetæ (Malloch), 1913 (<i>Agromyza</i>), <i>Melanagromyza</i>	83.3
politella (Malloch), 1934 (<i>Agromyza</i>), <i>Liriomyza</i>	83.11
praeclara Spencer, 1963, <i>Melanagromyza</i>	83.4
Praspedomyza Hendel, 1931	83.7
punctohalterata (Frost), 1936 (<i>Agromyza</i>), <i>Ophiomyia</i>	83.5
quadrata (Malloch), 1934 (<i>Agromyza</i>), <i>Liriomyza</i>	83.11
rabelloi Spencer, 1966, <i>Phytobia</i>	83.7
richardsi Spencer, 1963, <i>Calycomyza</i>	83.9
sabaziae Spencer, 1963, <i>Liriomyza</i>	83.11
sativæ Blanchard, 1938, <i>Liriomyza</i>	83.11
schmidtii (Aldrich), 1929 (<i>Agromyza</i>), <i>Liriomyza</i>	83.11
schmidtii (Frost), 1936 (<i>Agromyza</i>), <i>Japanagromyza</i>	83.2
schwabeii Spencer, 1963, <i>Liriomyza</i>	83.11
semiposticata Hendel, 1920, <i>Cerodontha</i>	83.6
seneciella Spencer, 1963, <i>Liriomyza</i>	83.11
setifera Spencer, 1963, <i>Melanagromyza</i>	83.5
simillima Spencer, 1963, <i>Agromyza</i>	83.2
simulator (Malloch), 1934 (<i>Agromyza</i>), <i>Liriomyza</i>	83.11
solanita Spencer, 1963, <i>Liriomyza</i>	83.11
sorosis (Williston), 1891 (<i>Agromyza</i>), <i>Liriomyza</i>	83.12
spadix Spencer, 1963, <i>Agromyza</i>	83.2
spicatae Spencer, 1963, <i>Ophiomyia</i>	83.5
splendida Frick, 1953, <i>Melanagromyza</i>	83.5
strigosa Spencer, 1963, <i>Liriomyza</i>	83.12
subandina Blanchard, 1954, <i>Liriomyza</i>	83.11



tequendamac Spencer, 1963, <i>Metopomyza</i>	83.12
terebrans Bezzi & Tavares, 1911, <i>Agromyza</i>	83.2
tetrac (Malloch), 1934 (<i>Agromyza</i>), <i>Melanagromyza</i>	83.5
<i>tibialis</i> (Frost), 1936 (<i>Agromyza</i>), <i>Japanagromyza</i>	83.2
tigrensis Spencer, 1963, <i>Haplomyza</i>	83.12
trifolii (Burgess), 1880 (<i>Oscinis</i>), <i>Liriomyza</i>	83.12
tubifer (Melander), 1913 (<i>Agromyza</i>), <i>Liriomyza</i>	83.12
tucumana Blanchard, 1954, <i>Phytomyza</i>	83.13
venezolana Spencer, 1963, <i>Agromyza</i>	83.2
verbascifolii Spencer, 1963, <i>Haplomyza</i>	83.12
verbenivora Spencer, 1963, <i>Calycomyza</i>	83.9
verbesinac Spencer, 1963, <i>Calycomyza</i>	83.9
virens (Loew), 1869 (<i>Agromyza</i>), <i>Melanagromyza</i>	83.5
xanthophora (Schiner), 1868 (<i>Agromyza</i>), <i>Phytobia</i>	83.7
williamsoni Blanchard, 1954, <i>Phytomyza</i>	83.13





SciELO₉

DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

84

FAMILY CLUSIIDAE

Á. Soós

*Zoological Department of the Hungarian Natural History
Museum, Budapest, Hungary*

The minute (1.5–7.5 mm, but usually 3–4 mm), black, black-yellow, or entirely yellowish, hyaline-, fumosely clouded-, or spotted-winged Clusiids are among the rarest flies, with hardly any big population known up to now. Their biology and immature stages are very little known. Adults can be found in wet, shaded sites, in woods, mainly around decaying trees and wooden artifacts; they are absent in arid regions. The larvae and pupae of only a few European, North American, and Australian species have been described. They have been found in burrows and galleries made by other insects either under the bark of living trees, or, in the majority, among soft rotting fibers and almost desintegrating tree-trunks.

Clusiid species occur in all parts of the world, except continental Africa. To date, disregarding varieties, 22 genera, 199 species, and 5 subspecies are known. Of these, 10 genera, 95 species, and 4 subspecies were heretofore found to occur in the Neotropical Region (within the limits of this Catalogue), that is, about half of the known species inhabit this area.

Czerny (1903), Kertész (1903), Melander & Argo (1924), Frey (1960), McAlpine (1960), and Soós (1964) have published revisions of the family and keys to world genera, respectively.

Genus ALLOCLUSIA Hendel

Peratochaetus Rondani, 1863:43 (*nee* Rondani, 1856). Type-species, *philippii* Rondani (mon.).

Heteroehraa Schiner, 1868:236 (preocc. Boisduval, 1835). Type-species, *pieta* Schiner (orig. des.) = *philippii* (Rondani).

ALLOCLUSIA Hendel, 1917:36 (nom. nov. for *Peratochaetus* Rondani). Type-species, *Peratochaetus philippii* Rondani (orig. des.).

Ref. — Malloch, 1933:235 (key to species).

CHILOENSIS Malloch, 1933:238, text-fig. 50a (hypopygium), pl. III, fig. 3 (wing). Type-locality: Chile, Chiloé Island, Castro. Distr. — Chile (Chiloé Island, Meehuque Island).



- COSTALIS** Malloch, 1933:239, text-fig. 50e (forceps). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina (Bariloche).
- EXCEPTA** Malloch, 1933:241, text-figs. 50b-e (forceps from side and from below), pl. III, fig. 6 (wing). Type-locality: Chile, Ancud. Distr. — Chile (Ancud).
- LIMBIPENNIS** (Rondani), 1863:43 (*Peratochaetus*). Type-locality: "Chile". Distr. — Argentina (Rio Negro: Bariloche, L. Correntoso), Chile (Angol, Casa Pangue, Limache). Refs. — Czerny, 1903:99 (*Apiochaeta*); Melander & Argo, 1924:7 (*Apiochaeta*); Malloch, 1933:238, text-fig. 49b (hypopygium) pl. III, fig. 4 (wing).
- PHILIPPIN** (Rondani), 1863:43 (*Peratochaetus*). Type-locality: "Chile". Distr. — Chile (Ancud, Angol, Cholchol, Puerto Montt). Refs. — Czerny, 1903:94 (*Peratochaetus*), pl. II, figs. 4, 5 (wing, head); Melander & Argo, 1924:32; Malloch, 1933:235, text-fig. 49a (hypopygium), pl. III, fig. 1 (wing).
- picta* Schiner, 1868:236 (*Heterochroa*). Type-locality: "Chile".
- PICTIPENNIS** (Wulp), 1882:91 (*Heterochroa*). Type-locality: "Chile". Distr. — Chile (Puerto Varas). Refs. — Czerny, 1903:96; Melander & Argo, 1924:33; Malloch, 1933:236, text-fig. 49c (hypopygium), pl. III, fig. 2 (wing).
- SIMILIS** Malloch, 1933:237, text-fig. 49d (hypopygium). Type-locality: Chile, Puerto Varas. Distr. — Chile (Puerto Varas).
- TINCTICORNIS** Malloch, 1933:242, text-fig. 50f (forceps). Type-locality: Chile, Puntra. Distr. — Chile (Ancud, Casa Pangue, Peulla, Puntra).
- VARIA** Malloch, 1933:241, text-fig. 50d (forceps), pl. III, fig. 5 (wing). Type-locality: Chile, Puerto Varas. Distr. — Chile (Peulla, Puerto Varas, Puntra).
- var. **CLARIPENNIS** Malloch, 1933:242. Type-locality: Chile, Puerto Varas. Distr. — Chile (Ancud, Castro, Puerto Varas, Puntra).

Genus APIOCHAETA Czerny

- APIOCHAETA** Czerny, 1903:98. Type-species, *Heterochroa bicolor* Schiner (by present des.).
Ref. — Malloch, 1933:243 (key to species).
- AEQUALIS** Malloch, 1933:245. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina (Bariloche).
- ATERRIMA** Malloch, 1933:244. Type-locality: Chile, Casa Pangue. Distr. — Chile (Casa Pangue).
- BICOLOR** (Schiner), 1868:237 (*Heterochroa*). Type-locality: "Chile". Distr. — Chile. Refs. — Czerny, 1903:98, pl. II, figs. 6, 7 (wing, head); Melander & Argo, 1924:6; Malloch, 1933:243.
- CONNEXA** Malloch, 1933:244. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina (Bariloche).
- VITICOLLIS** (Blanchard), 1852:452 (*Helomyza*). Type-locality: Chile, "en las provincias centrales de la Republica". Distr. — Chile.

Genus CHAETOCLUSIA Coquillett

- CHAETOCLUSIA** Coquillett, 1904:93. Type-species, *bakeri* Coquillett (orig. des.).
Refs. — Malloch, 1918:5; Melander & Argo, 1924:7 (key to species).

- BAKERI Coquillett, 1904:94. Type-locality: Nicaragua, Chinandega and Granada. Distr. — Nicaragua (Chinandega and Granada), Costa Rica. Refs. — Malloch, 1918:5; Melander & Argo, 1924:8, pl. 2, fig. 16 (photo, lateral aspect).
- ssp. PERUANA Hennig, 1938:128. Type-locality: Peru, Meshagua, Urbamba River. Distr. — Peru (Meshagua).
- LONGEFILATA Melander & Argo, 1924:8, pl. 2, fig. 12 (photo, lateral aspect). Type-locality: Panama, Alhajuela. Distr. — Panama (Alhajuela).
- NIGROMACULATA Melander & Argo, 1924:9. Type-locality: "Haiti". Distr. — Haiti.
- QUADRIVITTATA Melander & Argo, 1924:9, pl. 2, fig. 15 (photo, lateral aspect). Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica (La Suiza de Turrialba).
- XANTHOPS (Williston), 1896:386 (*Heteroneura*) (in part). Type-locality: "St. Vincent". Distr. — West Indies (St. Vincent). [See also *Sobarocephala bistrigata*].

Genus CHAETOCLUSIELLA Soós

- CHAETOCLUSIELLA Soós, 1962a:138. Type-species, *richardfreyi* Soós (orig. des.).
- RICHARDFREYI Soós, 1962a:138, figs. 3A, B (head, notum). Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil (Rio de Janeiro).

Genus CZERNYOLA Bezzi

- Craspedochacta* Czerny, 1903:103 (preocc. Macquart, 1851). Type-species, *transversa* Czerny (by present des.).
- CZERNYOLA Bezzi, 1907:52 (nom. nov. for *Craspedochaeta* Czerny). Type-species, *Craspedochaeta transversa* Czerny (aut.).
- Ref. — Soós, 1962c:449 (review and key of the species).
- ALBOHALTERATA Soós, 1962a:145. Type-locality: Bolivia, Cachuela, Esperanza. Distr. — Bolivia (Cachuela, Esperanza).
- ATRA (Kertész), 1903:570 (*Craspedochaeta*). Type-locality: Bolivia, Songo. Distr. — Costa Rica (La Suiza de Turrialba), Bolivia (Songo). Ref. — Melander & Argo, 1924:25, pl. 2, fig. 11 (photo, lateral aspect).
- BASALIS (Czerny), 1903:104 (*Craspedochacta*). Type-locality: "Peru". Distr. — Peru (Vilcanota). Ref. — Melander & Argo, 1924:26.
- ssp. BRASILIENSIS Soós, 1961:399. Type-locality: Brazil, Mato Grosso, Maracaju. Distr. — Brazil (Mato Grosso).
- CLARIPENNIS Melander & Argo, 1924:26. Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica (La Suiza de Turrialba).
- FASCIPENNIS Melander & Argo, 1924:26, pl. 2, fig. 13 (photo, lateral aspect). Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica (La Suiza de Turrialba), Bolivia (Mapiri, S. Antonio).
- FESTIVA Soós, 1962a:142. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil (Guanabara).

- MINUTA Soós, 1962a:148. Type-locality: Costa Rica, Higuito, San Mateo. Distr. — Costa Rica (Higuito, San Mateo).
- NIGRITHORAX Hennig, 1938:133. Type-locality: Bolivia, Mapiri, S. Carlos. Distr. — Bolivia (Mapiri, S. Carlos, Sarampiuní, S. Antonio).
- ORNATA Hennig, 1938:132. Type-locality: Bolivia, Mapiri, S. Carlos. Distr. — Bolivia (Mapiri, Yungas de Coroico).
- PICEOFLAVA Soós, 1962a:146. Type-locality: Dominican Republic, Trujillo, Colonia Rafnis. Distr. — Dominican Republic (Trujillo).
- SYNNEURA Hennig, 1938:132. Type-locality: Bolivia, Mapiri, S. Carlos. Distr. — Bolivia (Mapiri).
- TRANSVERSA (Czerny), 1903:103 (*Craspedochaeta*). Type-locality: "Peru". Distr. — Peru, Bolivia (Mapiri), Paraguay. Ref. — Melander & Argo, 1924:27.
- TRIVITTATA Soós, 1962a:144. Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil (Rio de Janeiro).

Genus HETEROMERINGIA Czerny

- HETEROMERINGIA Czerny, 1903:72. Type-species, *Heteroneura nigrimana* Loew (mon.). Ref. — Melander & Argo, 1924:28 (revision).
- CZERNYI Kertész, 1903:568. Type-locality: Peru, Callanga. Distr. — Costa Rica (San Mateo), Panama (Cabima), Peru (Callanga), Paraguay.
- DIMIDIATA Hennig, 1938:134. Type-locality: Peru, Unini, Ucayali River. Distr. — Peru (Unini).
- FLAVIFRONS Hennig, 1938:134. Type-locality: Bolivia, Mapiri, S. Carlos. Distr. — Bolivia (Mapiri).
- FLAVIPES (Williston), 1896:387, pl. XIII, fig. 135 (wing) (*Heteroneura*). Type-locality: "St. Vincent". Distr. — U.S.A. (Florida), Nicaragua (Chinandega), West Indies (St. Vincent). Ref. — Melander & Argo, 1924:29.
- FUCATA Hendel, 1936:89. Type-locality: Brazil, Pará, Santarém, Farm Taperinha. Distr. — Brazil (Pará).
- FUMIPENNIS Melander & Argo, 1924:30, pl. 2, fig. 19 (photo, lateral aspect). Type-locality: Costa Rica, Turrialba. Distr. — Costa Rica (Turrialba, Juan Vinas).
- NIGRIFRONS Kertész, 1903:568. Type-locality: Peru, Vilcanota. Distr. — Peru (Vilcanota). Ref. — Melander & Argo, 1924:30.

Genus LABOMYIA Frey

- LABOMYIA Frey, 1928:14. Type-species, *mirabilis* Frey (orig. des.).
- MIRABILIS Frey, 1928:16, fig. 1 (head). Type-locality: Peru, Ostanden, Roque. Distr. — Peru (Ostanden).

Genus SOBAROCEPHALA Czerny

- SOBAROCEPHALA Czerny, 1903:85. Type-species, *ruebsaameni* Czerny (mon.).
- Meriza Kertész, 1903:571. Type-species, *dorsata* Kertész (by present des.).
- Monorrhhexa Kertész, 1903:572. Type-species, *pictipennis* Kertész (mon.).



Rcfs. — Melander & Argo, 1924:33 (revision); Soós, 1963:391 (key to the *plumata*-group).

ALBITARSIS Czerny, 1929:10. Type-locality: "Paraguay". Distr. — Paraguay.

ALBOMACULATA Soós, 1965:355. Type-locality: Panama, Canal Zone, Ft. Randolph. Distr. — Costa Rica (Higuito, San Mateo), Panama (Canal Zone).

ANNULATA Melander & Argo, 1924:36, pl. 4, fig. 34 (photo, lateral aspect). Type-locality: Costa Rica, Turrialba. Distr. — Costa Rica (Turrialba), Panama (Barro Colorado Island, Erwin Island, Porto Bello), Peru, Bolivia.

ssp. ALBIVENTRIS Soós, 1962b:391, figs. 4A-B (hypopygium, head). Type-locality: Ecuador, Guayas, Chilcales. Distr. — Ecuador (Guayas, Chilcales).

BECKERI Hennig, 1956:146, fig. 1 (hypopygium). Type-locality: Guatemala, Los Trojados. Distr. — Guatemala (Los Trojados).

BISTRIGATA (Kertész), 1903:572 (*Meriza*). Type-locality: Peru, Callanga. Distr. — Costa Rica, Jamaica, West Indies (St. Vincent), Peru (Callanga). Ref. — Melander & Argo, 1924:36, pl. 4, fig. 33 (photo, lateral aspect).

xanthops Williston, 1896:386 (*Heteroneura*) (in part).

BIVITTATA Melander & Argo, 1924:37, pl. 4, fig. 36 (photo, lateral aspect). Type-locality: Costa Rica, Turrialba. Distr. — Costa Rica (Turrialba).

BOLIVIANA Soós, 1963:395. Type-locality: Bolivia, Songo. Distr. — Bolivia (Songo).

COLUMBIENSIS Soós, 1962a:387, figs. 3A-D (head, abdomen, wing, hypopygium). Type-locality: Colombia, Aracataca. Distr. — Colombia (Aracataca).

CONCINNA (Williston), 1896:387 (*Heteroneura*). Type-locality: St. Vincent". Distr. — West Indies (St. Vincent).

DIVERSIPES Curran, 1939:2. Type-locality: Panama, Canal Zone, Barro Colorado Island. Distr. — Panama (Canal Zone).

DORSATA Czerny, 1903:105. Type-locality: "Peru". Distr. — Costa Rica, Peru (Callanga). Refs. — Kertész, 1903:571 (*Meriza*); Melander & Argo, 1924:38, pl. 4, fig. 38 (photo, lateral aspect).

DUDICHI Soós, 1963:392. Type-locality: "Costa Rica". Distr. — Costa Rica.

ELEGANS Czerny, 1929:9. Type-locality: "Paraguay". Distr. — Paraguay.

FASCIPENNIS Melander & Argo, 1924:39, pl. 3, fig. 29 (photo, lateral aspect). Type-locality: Panama, Alhajuela. Distr. — Panama (Alhajuela).

FERRUGINEA Czerny, 1903:105. Type-locality: "Peru". Distr. — Peru. Ref. — Melander & Argo, 1924:39.

FESTIVA Czerny, 1929:9. Type-locality: "Paraguay". Distr. — Paraguay.

FINNILAEI Frey, 1919:32, fig. 5 (wing). Type-locality: Brazil, Rio de Janeiro, Petrópolis. Distr. — Venezuela (Akuriman, Gran Sabana), Brazil (Guanabara; Rio de Janeiro, Petrópolis, Teresópolis), Paraguay (S. Bernardino). Ref. — Melander & Argo, 1924:39.

ornata Czerny, 1929:12. Type-locality: Paraguay, S. Bernardino. Distr. — Paraguay (S. Bernardino).

- FUMPENNIS Soós, 1963:393. Type-locality: "West Indies". Distr. — West Indies.
- GUIANICA Curran, 1934:442. Type-locality: Guyana, Kartabo. Distr. — Guyana (Kartabo).
- HUMERALIS Melander & Argo, 1924:40, pl. 3, fig. 25 (photo, lateral aspect). Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica (La Suiza de Turrialba).
- var. DIVES Czerny, 1929:11. Type-locality: Paraguay, S. Bernardino. Distr. — Paraguay (S. Bernardino).
- var. DISTINCTA Soós, 1965:354. Type-locality: Brazil, Rio de Janeiro, Tinguá. Distr. — Brazil (Rio de Janeiro).
- var. MELANDERI Soós, 1965:354. Type-locality: Costa Rica, Juan Vinas. Distr. — Costa Rica (Juan Vinas).
- var. DIVERSA Soós, 1965:354. Type-locality: Panama, Canal Zone, Cloc's, Cano Saddle. Distr. — Panama (Canal Zone).
- var. BRASILIENSIS Soós, 1965:354. Type-locality: Brazil, Mato Grosso, Maracaju. Distr. — Brazil (Mato Grosso).
- HYPOPYGALIS Melander & Argo, 1924:41, pl. 4, fig. 41 (photo, lateral aspect). Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica (La Suiza de Turrialba).
- IMITANS Curran, 1934:443. Type-locality: Guyana, Kartabo. Distr. — Guyana (Kartabo).
- ISLA Curran, 1939:1. Type-locality: Panama, Canal Zone, Barro Colorado Island. Distr. — Panama (Canal Zone).
- LANEI Curran, 1939:1. Type-locality: Brazil, Mato Grosso, Ponc. Distr. — Brazil (Mato Grosso).
- LATIPENNIS Melander & Argo, 1924:43, pl. 3, fig. 24 (photo, lateral aspect). Type-locality: Costa Rica, Turrialba. Distr. — Costa Rica (Turrialba), Panama (Porto Bello).
- var. LATIPENNODIS Soós, 1965:361. Type-locality: Panama, Canal Zone, Barro Colorado Island. Distr. — Panama (Canal Zone).
- LITURATA Melander & Argo, 1924:43, pl. 4, fig. 42 (photo, lateral aspect). Type-locality: Costa Rica, Turrialba. Distr. — Costa Rica (Turrialba), Panama (Canal Zone), Peru (Iquitos). Ref. — Soós, 1965:361.
- LUMBALIS (Williston), 1896:388 (*Heteroneura*). Type-locality: "St. Vincent". Distr. — West Indies (St. Vincent, Grenada), Panama (Alhajueta, Paraiso). Ref. — Melander & Argo, 1924:44, pl. 4, fig. 37 (photo, lateral aspect).
- MACALPINEI Soós, 1964a:453, fig. 2B (head). Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil (Rio de Janeiro).
- MELANOPYGA Czerny, 1929:12. Type-locality: "Paraguay". Distr. — Paraguay.
- NIGROANTENNATA Soós, 1962a:389, fig. 4A (head). Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Panama (Ft. Gulick), Brazil (Guanabara). Ref. — Soós, 1965:357.
- NIGROFACIES Soós, 1962b:383, figs. 1A-C (head, thorax, wing). Type-locality: Brazil, Rio de Janeiro, Tinguá. Distr. — Brazil (Rio de Janeiro).
- NIGROHUMERALIS Curran, 1929:38. Type-locality: Jamaica, Cinchona. Distr. — Jamaica (Cinchona).

NIGRONOTA Melander & Argo, 1924:45, pl. 3, fig. 28 (photo, lateral aspect).
Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica (La Suiza de Turrialba).

nigronotum, error.

ssp. PERUANA Soós, 1965:358. Type-locality: "Peru". Distr. — Peru
PANAMAENSIS Soós, 1965:356. Type-locality: Panama, Garachine. Distr. —
Panama (Aguadulce, Garachine, Patino Pt. Darion).

PICTA Hennig, 1938:135. Type-locality Paraguay, "Hohenau". Distr. —
Paraguay ("Hohenau").

PICTIPENNIS (Kertész), 1903:573 (*Monorrhexa*). Type-locality: "Surinam".
Distr. — Surinam. Ref. — Melander & Argo, 1924:45.

PLEURALIS (Williston), 1896:387 (*Heteroneura*). Type-locality: "St. Vincent".
Distr. — West Indies (St. Vincent). Ref. — Melander & Argo, 1924:45.

PRUINOSA Soós, 1962a:385, figs. 2A-C (head, surstylus, hypopygium)
Type-locality: Panama, San Juan. Distr. — Panama (Mayaguez San Juan).

PLUMATELLA Melander & Argo, 1924:46, pl. 4, fig. 39 (photo, lateral aspect).
Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica (La Suiza de Turrialba). Ref. — Soós, 1965:359.

PRUINOSA Soós, 1962a:385, figs. 2A-C (head, surstylus, hypopygium)
Type-locality: Panama, San Juan. Distr. — Panama (San Juan, Mayaguez).

QUADRIVITTATA Czerny, 1929:12. Type-locality: Brazil, Rio Grande do Sul.
Distr. — Brazil (Rio Grande do Sul).

RUEBSAAMENI Czerny, 1903:86, pl. III, figs. 3, 4, 5 (head, wing). Type-locality: "Peru". Distr. — Salvador (Rosario Cuzertlan), Costa Rica (San Mateo, Surrubes), Panama (El Cermeño, Jas. Zetek, Mindi Dairy, Trinidad River), Peru. Refs. — Melander & Argo, 1924:47, pl. 3, fig. 23 (photo, lateral aspect); Soós, 1965:351.

var. REDUCTA Soós, 1965:351. Type-locality: Panama, Canal Zone, Mojinga Swamp. Distr. — Costa Rica, Panama (Canal Zone), Brazil (Magé, R. Janeiro).

SABROSKYI Soós, 1963:395. Type-locality: Paraguay, S. Bernardino. Distr. — Paraguay (S. Bernardino).

SEXVITTATA Soós, 1964a:451, fig. 2A (head). Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil (Guanabara).

STEYSKALI Soós, 1963:395. Type-locality: "Brazil". Distr. — Brazil.

styskali, error.

STRIGATA Hennig, 1938:136. Type-locality: Peru, Pachitea. Distr. — Peru (Pachitea).

SUBFASCIATA Curran, 1939:2. Type-locality: Panama, Canal Zone, Barro Colorado Island. Distr. — Panama (Canal Zone).

VALIDA (Williston), 1896:388, pl. XIII, fig. 136 (wing) (*Heteroneura*).
Type-locality: "St. Vincent". Distr. — West Indies (St. Vincent). Ref. — Melander & Argo, 1924:48, pl. 3, fig. 30 (photo, lateral aspect).

- VARIEGATA Melander & Argo, 1924:48, pl. 4, fig. 40 (photo, lateral aspect). Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica (La Suiza de Turrialba), Panama (Canal Zone, Corozal).
- VITTATIFRONS Hennig, 1938:137. Type-locality: Peru, Meshagua, Urubamba River. Distr. — Peru (Meshagua).
- XANTHOMELANA Melander & Argo, 1924:49, pl. 3, fig. 27 (photo, lateral aspect). Type-locality: Costa Rica, Turrialba. Distr. — Costa Rica (Turrialba). Ref. — Soós, 1965:360.
- ZEUGMA Hennig, 1938:136. Type-locality: Bolivia, Yungas de Coroico. Distr. — Bolivia (Yungas de Coroico).

Genus SOBAROCEPHALOIDES Soós


- SOBAROCEPHALOIDES Soós, 1964b:1. Type-species, *discolor* Soós (orig. des.).
- DISCOLOR Soós, 1964b:2, figs. A-B (head, thorax). Type-locality: Costa Rica, La Suiza. Distr. — Costa Rica (La Suiza).

Genus TRICHOCCLUSIA Soós

- TRICHOCCLUSIA Soós, 1962a:135. Type-species, *sabroskyi* Soós (orig. des.).
- SABROSKYI Soós, 1962a:135, fig. 1, 2 (antenna, head, notum, wing). Type-locality: Cuba, Las Animas, Sierra Rangol. Distr. — Cuba (Las Animas, Pinar del Rio).

LITERATURE CITED

- BEZZI, M.
1907. Nomenklatorisches über Dipteren. *Wien. ent. Ztg.* 26:51-56.
- BLANCHARD, E.
1852. Orden IX. Dipteros, in C. Gay. *Historia física y política de Chile, Zoología* 7:327-468.
- COQUILLET, D. W.
1904. New Diptera from Central America. *Proc. Entomol. Soc. Washington* 6:90-98.
- CURRAN, C. H.
1928. Records and descriptions of Diptera, mostly from Jamaica, in C. C. Gowdey: *Catalogus insectorum jamaicensis. Entomol. Bull. Dept. Agric. Jamaica* 4:29-45.
1934. The Diptera of Kartabo, Bartica District, British Guiana, with descriptions of new species from other British Guiana localities. *Bull. Amer. Mus. Nat. Hist.* 66(Art.III):287-532.
1939. Four new species of *Sabarocephala* Diptera: Clusiidae. *Amer. Mus. Nat. Hist., Amer. Mus. Novitates* 1029:1-2.
- CZERNY, L.
1903. Revision der Heteroneuriden. *Wien. ent. Ztg.* 22:61-107.
1929. Neue Clusiiden aus Paraguay und Rio Grande do Sul. *Konowia* 8:9-13.



FREY, R.

1919. Mitteilungen über südamerikanische Dipteren. *Öfvers. Finska Vet. Soc. Förh.* 60(A,14):1-35.
1928. Eine neue Heteroneuriden-Gattung mit stiel förmigen Kopf-
fortsätzen (Dipt., Haplost.). *Notulae Ent.* 8(1):14-16.
1960. Studien über indoaustralische Clusiiden (Dipt.), nebst Katalog
der Clusiiden. *Comment. Biol.* 22(1):1-31.

HENDEL, F.

1917. Beiträge zur Kenntnis der acalyptraten Musciden. *Deutsch. Ent.
Ztschr.* 1917(I/II):33-47.
1936. Ergebnisse einer zoologischen Sammelreise nach Brasilien,
insbesondere in das Amazonasgebiet, ausgeführt von Dr. H. Zerny.
X. Diptera: Muscidae acalyptratae (excl. Chloropidae). *Ann.
Naturhist. Mus. Wien* 47:61-106.

HENNIG, W.

1938. Beiträge zur Kenntnis der Clusiiden und ihres Kopulations-
apparates (Dipt., Acalypt.). *Encycl. Entomol.* (B) II. *Diptera*
9:121-138.
1956. Neue neotropische Acalyptrata aus dem Deutschen Entomolo-
gischen Institut (Diptera: Acalyptrata) *Beitr. z. Ent.*, Berlin,
6(1/2):146-154.

KERTÉSZ, K.

1903. Beiträge zur Kenntniss der Heteroneuriden. *Ann. Mus. Nat.
Hung.* 1:566-573.

MALLOCH, J. R.

1918. A revision of the dipterous family Clusiodidae (Heteroneu-
ridae). *Proc. Entomol. Soc. Washington* 20(1):2-8.
1933. Acalyptrata, in: *Diptera of Patagonia and South Chilc.* British
Museum (Nat. Hist.), London, Part VI, Fasc. 4:177-391.

MCALPINE, D. K.

1960. A review of the Australian species of Clusiidae (Diptera,
Acalyptrata). *Rec. Austr. Mus.* 25(4):63-94.

MELANDER, A. L. & N. G. ARGO

1924. Revision of the two-winged flies of the family Clusiidae. *Proc.
U. S. Nat. Mus.* 64 (Art. 11):1-54.

RONDANI, C.

1863. *Diptera exotica revista et annotata.* Modena, 99 pp.

SCHINER, I. R.

1868. Diptera, in: *Reise der österreichischen Fregatte Novara um die
Erde in den Jahren 1857, 1858, 1859.* Wien, Zool. Vol. 2, Abt. 1B:
1-388.

SÓOS, Á.

1961. A new subspecies of *Czernyola basalis* Czerny, 1903 (Diptera,
Clusiidae). *Folia Entomol. Hung.* (n. s.) 14(25):399-402.
1962a. New genera and species of Clusiidae (Diptera: Acalyptrata).
Acta Zool. Hung. 8(1-2):135-149.
1962b. New species of *Sobaroccephala* Czerny (Diptera: Clusiidae).
Ann. Hist.-nat. Mus. Nat. Hung. 54:383-392.

- 1962e. A review of the species of *Czernyola* Bezzi, 1907 (Diptera: Clusiidae). *Acta Zool. Hung.* 8(3-4):449-457.
1963. Identification key to the species of the "plumata-group" of the genus *Sobarocephala* Czerny (Diptera: Clusiidae). *Acta Zool. Hung.* 9(3-4):391-396.
- 1964a. New *Sobarocephala*-species from the "plumata-group" (Diptera, Clusiidae). *Ann. Hist.-nat. Mus. Nat. Hung.* 56:449-455.
- 1964b. *Sobarocephaloides discolor* gen. nov., spec. nov., eine neue Clusiide aus Costa-Rica, nebst einer Bestimmungstabelle der Gattungen der Familie Clusiidae (Diptera: Acalyptrata). *Reichenbachia* 4(1):1-8.
1965. On some new and known species of *Sobarocephala* Czerny, 1903 (Diptera: Clusiidae). *Ann. Hist.-nat. Mus. Nat. Hung.* 57:349-362.
- WILLISTON, S. W.
1896. On the Diptera of St. Vincent (West Indies). *Trans. Entomol. Soc. London, 1896* (Part III):253-446.
- WULP, F. M. VAN DER
1882. Remarks on certain American Diptera in the Leyden Museum and description of nine new species. *Not. Leyden Mus.* 4(2):73-92.

INDEX


<i>aequalis</i> Malloch, 1933, <i>Apiochaeta</i>	84.2
<i>albitarsis</i> Czerny, 1929, <i>Sobarocephala</i>	84.5
<i>albiventris</i> Soós, 1962, <i>Sobarocephala</i>	84.5
<i>albohalterata</i> Soós, 1962, <i>Czernyola</i>	84.3
<i>albomaculata</i> Soós, 1965, <i>Sobarocephala</i>	84.5
<i>Alloclusia</i> Hendel, 1917	84.1
<i>annulata</i> Melander & Argo, 1924, <i>Sobarocephala</i>	84.5
<i>Apiochaeta</i> Czerny, 1903	84.2
<i>aterrima</i> Malloch, 1933, <i>Apiochaeta</i>	84.2
<i>atra</i> (Kertész), 1903 (<i>Craspedochaeta</i>), <i>Czernyola</i>	84.3
<i>bakeri</i> Coquillett, 1904, <i>Chaetoclusia</i>	84.3
<i>basalis</i> (Czerny), 1903 (<i>Craspedochaeta</i>), <i>Czernyola</i>	84.3
<i>beckeri</i> Hennig, 1956, <i>Sobarocephala</i>	84.5
<i>bicolor</i> (Schiner), 1868 (<i>Heterochroa</i>), <i>Apiochaeta</i>	84.2
<i>bistrigata</i> (Kertész), 1903 (<i>Meriza</i>), <i>Sobarocephala</i>	84.5
<i>bivittata</i> Melander & Argo, 1924, <i>Sobarocephala</i>	84.5
<i>boliviana</i> Soós, 1963, <i>Sobarocephala</i>	84.5
<i>brasiliensis</i> Soós, 1961, <i>Czernyola</i>	84.3
<i>brasiliensis</i> Soós, 1965, <i>Sobarocephala</i>	84.6
<i>Chaetoclusia</i> Coquillett, 1904	84.2
<i>Chaetoclusiella</i> Soós, 1962	84.3
<i>chiloensis</i> Malloch, 1933, <i>Alloclusia</i>	84.1
<i>claripennis</i> Malloch, 1933, <i>Alloclusia</i>	84.2
<i>claripennis</i> Melander & Argo, 1924, <i>Czernyola</i>	84.3




columbiensis Soós, 1962, Sobarocephala	84.5
concinna (Williston), 1896 (Heteroneura), Sobarocephala	84.5
connexa Malloch, 1933, Apiochaeta	84.2
costalis Malloch, 1933, Alloclusia	84.2
<i>Craspedochaeta</i> Czerny, 1903	84.2
czernyi Kertész, 1903, Heteromeriingia	84.4
Czernyola Bezzi, 1907	84.3
dimidiata Hennig, 1938, Heteromeriingia	84.4
discolor Soós, 1964, Sobarocephaloides	84.8
distincta Soós, 1965, Sobarocephala	84.6
diversa Soós, 1965, Sobarocephala	84.6
diversipes Curran, 1939, Sobarocephala	84.5
dives Czerny, 1929, Sobarocephala	84.6
dorsata Czerny, 1903, Sobarocephala	84.5
dudichi Soós, 1963, Sobarocephala	84.5
elegans Czerny, 1929, Sobarocephala	84.5
excepta Malloch, 1933, Alloclusia	84.2
fascipennis Melander & Argo, 1924, Czernyola	84.3
fascipennis Melander & Argo, 1924, Sobarocephala	84.5
ferruginea Czerny, 1903, Sobarocephala	84.5
festiva Soós, 1962, Czernyola	84.3
festiva Czerny, 1929, Sobarocephala	84.5
finnilaei Frey, 1919, Sobarocephala	84.5
flavifrons Hennig, 1938, Heteromeriingia	84.4
flavipes (Williston), 1896 (Heteroneura), Heteromeriingia	84.4
fucata Hendel, 1936, Heteromeriingia	84.4
fumipennis Melander & Argo, 1924, Heteromeriingia	84.4
fumipennis Soós, 1963, Sobarocephala	84.6
guianica Curran, 1934, Sobarocephala	84.6
<i>Heterochroa</i> Schiner, 1868	84.1
Heteromeriingia Czerny, 1903	84.4
humeralis Melander & Argo, 1924, Sobarocephala	84.6
hypopygialis Melander & Argo, 1924, Sobarocephala	84.6
imitans Curran, 1934, Sobarocephala	84.6
isla Curran, 1939, Sobarocephala	84.6
Labomyia Frey, 1928	84.4
lanci Curran, 1939, Sobarocephala	84.6
latipennis Melander & Argo, 1924, Sobarocephala	84.6
latipennoides Soós, 1965, Sobarocephala	84.6
limbipennis (Rondani), 1863 (Peratochaetus), Alloclusia	84.2
liturata Melander & Argo, 1924, Sobarocephala	84.6
longefilata Melander & Argo, 1924, Chaetoclusia	84.3
lumbalis (Williston), 1896 (Heteroneura), Sobarocephala	84.6





macalpinei Soós, 1964, Sobarocephala	84.6
melanderi Soós, 1965, Sobarocephala	84.6
melanopyga Czerny, 1929, Sobarocephala	84.6
<i>Meriza</i> Kertész, 1903	84.4
minuta Soós, 1962, Czernyola	84.4
mirabilis Frey, 1928, Labomyia	84.4
<i>Monorrhexa</i> Kertész, 1903	84.4
nigrifrons Kertész, 1903, Heteromeringia	84.4
nigrithorax Hennig, 1938, Czernyola	84.4
nigroantennata Soós, 1962, Sobarocephala	84.6
nigrofacies Soós, 1962, Sobarocephala	84.6
nigrohumeralis Curran, 1929, Sobarocephala	84.6
nigromaculata Melander & Argo, 1924, Chaetoclusia	84.3
nigronota Melander & Argo, 1924, Sobarocephala	84.7
<i>nigronotum</i> , error	84.7
ornata Hennig, 1938, Czernyola	84.4
<i>ornata</i> Czerny, 1929, Sobarocephala	84.5
panamaensis Soós, 1965, Sobarocephala	84.7
<i>Peratochaetus</i> Rondani, 1863	84.1
peruana Hennig, 1938, Chaetoclusia	84.3
peruana Soós, 1965, Sobarocephala	84.7
philippii (Rondani), 1863 (<i>Peratochaetus</i>), Alloclusia	84.2
piccoflava Soós, 1962, Czernyola	84.4
<i>picta</i> (Schiner), 1868 (<i>Heterochroa</i>), Alloclusia	84.2
<i>picta</i> Hennig, 1938, Sobarocephala	84.7
pictipennis (Wulp), 1882 (<i>Heterochroa</i>), Alloclusia	84.2
pictipennis (Kertész), 1903 (<i>Monorrhexa</i>), Sobarocephala	84.7
pleuralis (Williston), 1896 (<i>Heteroneura</i>), Sobarocephala	84.7
plumata Melander & Argo, 1924, Sobarocephala	84.7
plumatella Melander & Argo, 1924, Sobarocephala	84.7
pruinosa Soós, 1962, Sobarocephala	84.7
quadrivittata Czerny, 1929, Sobarocephala	84.7
quadrivittata Melander & Argo, 1924, Chaetoclusia	84.3
reducta Soós, 1965, Sobarocephala	84.7
richardfreyi Soós, 1962, Chaetoclusicla	84.3
ruebsaameni Czerny, 1903, Sobarocephala	84.7
sabroskyi Soós, 1963, Sobarocephala	84.7
sabroskyi Soós, 1962, Trichoclusia	84.8
sexvittata Soós, 1964, Sobarocephala	84.7
similis Malloch, 1933, Alloclusia	84.2
Sobarocephala Czerny, 1903	84.4
Sobarocephaloides Soós, 1964	84.8
steyskali Soós, 1963, Sobarocephala	84.7
strigata Hennig, 1938, Sobarocephala	84.7




<i>styskali</i> , error	84.7
<i>subfasciata</i> Curran, 1939, <i>Sobarocephala</i>	84.7
<i>synneura</i> Hennig, 1938, <i>Czernyola</i>	84.4
<i>tincticornis</i> Malloch, 1933, <i>Alloclusia</i>	84.2
<i>transversa</i> (Czerny), 1903 (<i>Craspedochaeta</i>), <i>Czernyola</i>	84.4
<i>Trichoclusia</i> Soós, 1962	84.8
<i>trivittata</i> Soós, 1962, <i>Czernyola</i>	84.4
<i>valida</i> (Williston), 1896 (<i>Heteroneura</i>), <i>Sobarocephala</i>	84.7
<i>varia</i> Malloch, 1933, <i>Alloclusia</i>	84.2
<i>variegata</i> Melander & Argo, 1924, <i>Sobarocephala</i>	84.8
<i>vittatifrons</i> Hennig, 1938, <i>Sobarocephala</i>	84.8
<i>vitticollis</i> (Blanchard), 1852 (<i>Hclomyza</i>), <i>Apiochaeta</i>	84.2
<i>xanthomelana</i> Melander & Argo, 1924, <i>Sobarocephala</i>	84.8
<i>xanthops</i> (Williston), 1896 (<i>Heteroneura</i>), <i>Chaetoclusia</i>	84.3
<i>xanthops</i> (Williston), 1896 (<i>Heteroneura</i>), <i>Sobarocephala</i>	84.5
<i>zeugma</i> Hennig, 1938, <i>Sobarocephala</i>	84.8









SciELO₉

DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

85

FAMILY HELEOMYZIDAE

(HELOMYZIDAE)

INCLUDING THE TRIXOSCELIDIDAE
(TRICHOSCELIDAE) (86)

GORDON D. GILL

*Department of Biology, Northern Michigan University
Marquette, Michigan, U. S. A.*

When considering only the Nearectic, Palaearctic or Holarctic fauna, it is not difficult to establish distinctions between the Heleomyzidae and the Trixoscelididae. The two groups are considered as separate families in *A Catalog of the Diptera of America North of Mexico* (Stone *et al.*, 1965), although European dipterists have sometimes treated the Trixoscelididae as a subfamily of the Heleomyzidae. Malloch (1933, pp. 178-179) maintained that distinctions between the two groups do not hold when the fauna of the entire world are considered, and he made no designation of subfamilies in his treatment of the Heleomyzidae of Patagonia and South Chile. This point of view is supported by Harrison (1959, pp. 111-113) in his work on the New Zealand acalyptate Diptera. Although I am reluctant to deviate from the classification followed in the North American Catalog, I am unable at this time to arrange the Neotropical genera into families and subfamilies consistent with the treatment of North American forms. It therefore seems advisable to unite the Heleomyzidae and Trixoscelididae in this report and to present all genera and species alphabetically, rather than according to supposed phylogeny. Further studies of the entire complex on a world-wide basis are required before the problems of family and subfamily limits can be resolved.

Relatively little is known about the biology of the Neotropical Heleomyzidae. The immature stages of several species of *Snilla* have been reared from fungi. Members of several other genera, including *Aecothea* and *Pseudolertia*, are associated with birds' nests, mammal burrows and caves. Virtually nothing is known regarding the immature stages and habits of members of the other genera known from the Neotropical Region.

I am indebted to Dr. J. R. Vockeroth for several helpful suggestions and for providing distribution records from specimens in the Canadian



National Collection. The literature search for this report covers *Zoological Record* through 1963 and *Biological Abstracts* through September 15, 1967.

General references: Czerny's (1924) monograph and subsequent *Ergänzungen* (1926, 1927, 1929) include Neotropical genera and species. Malloch (1933) treats the family for Patagonia and South Chile.

Genus AECOTHEA Haliday

AECOTHEA Haliday, 1838:187. Type-species, *Helomyza fenestralis* Fallén (mon.).

Oecothea, emend.

FENESTRALIS (Fallén), 1820:5 (*Helomyza*). Type-locality: "Sweden". Distr. — Holarctic Region; Argentina (Rio Negro Prov., Lake Gutiérrez). Refs. — Gorodkov, 1959:915, figs. 42, 43, 44-47; Gill, 1962:519, figs. 21, 22.

Genus ANASTOMYZA Malloch

ANASTOMYZA Malloch, 1933:216. Type-species, *discalis* Malloch (orig. des.).

DELETA Edwards, *in* Malloch, 1933:224, text-figs. 47c, 47d, 47e, 47f. Type-locality: Argentina, Rio Negro Prov., Bariloche.

DISCALIS Malloch, 1933:220. Type-locality: Argentina, Rio Negro Prov., Lake Nahuel Huapi. Distr. — Argentina (SW Rio Negro Prov.).

FRACTURA Malloch, 1933:218, text-fig. 46; pl. 2, fig. 10. Type-locality: Argentina, Rio Negro Prov., Bariloche.

FULVITHORAX Malloch, 1933:219. Type-locality: Argentina, Rio Negro Prov., Bariloche.

HYALIPENNIS Malloch, 1933:220. Type-locality: Argentina, Rio Negro Prov., Lake Correntoso.

LATERALIS Malloch, 1933:221, text-figs. 47a, 47b, 47d; pl. 2, fig. 12. Type-locality: Argentina, Rio Negro Prov., Lake Correntoso. Distr. — Argentina (SW Rio Negro Prov.).

MULTIPUNCTATA Malloch, 1933:218, pl. 2, fig. 11. Type-locality: Argentina, Rio Negro Prov., Lake Gutiérrez. Distr. — Argentina (SW Rio Negro Prov.).


NEGLECTA Edwards, *in* Malloch, 1933:223, text-figs. 47e, 47g, 47h. Type-locality: Argentina, Rio Negro Prov., Lake Gutiérrez. Distr. — Argentina (SW Rio Negro Prov.), Chile (Chiloé Prov., Castro).

SPECULARIS Malloch, 1933:221. Type-locality: Argentina, Rio Negro Prov., Lake Gutiérrez. Distr. — Chile (Malleco Prov., Angol), Argentina (SW Rio Negro Prov.).

Genus APOPHONEURA Malloch

APOPHONEURA Malloch, 1933:225. Type-species, *exquisita* Malloch (orig. des.).

EXQUISITA Malloch, 1933:229, pl. 2, fig. 3. Type-locality: Argentina, Rio Negro Prov., Lake Correntoso. Distr. — Argentina (SW Rio Negro Prov.), Chile (Aysén Prov.).



- INCONSPICUA Malloch, 1933:231, pl. 2, fig. 7. Type-locality: Argentina, Rio Negro Prov., Bariloche. Distr. — Chile (Malleco and Chiloé Prov.), Argentina (SW Rio Negro Prov.).
- ineonspicua* Malloch, 1933:226 (misspelling, in key to species).
- PICTA Malloch, 1933:231, pl. 2, fig. 5. Type-locality: Chile, Chiloé Prov., Castro.
- PUNCTIPENNIS Malloch, 1933:227, pl. 2, fig. 9. Type-locality: Argentina, Rio Negro Prov., Lake Gutiérrez. Distr. — Argentina (SW Rio Negro Prov.).
- RECURRENS Malloch, 1933:226, text-fig. 48. Type-locality: Argentina, Rio Negro Prov., Bariloche. Distr. — Chile (Ñuble Prov.), Argentina (SW Rio Negro Prov.).
- SUPERBA Malloch, 1933:228, pl. 2, fig. 8. Type-locality: Argentina, Rio Negro Prov., Lake Correntoso. Distr. — Argentina (SW Rio Negro Prov.), Chile (Aysén Prov.).
- UNIVITTATA Malloch, 1933:230, pl. 2, fig. 4. Type-locality: Chile, Osorno Prov., San Pablo.
- VITTIGERA Malloch, 1933:232, pl. 2, fig. 6. Type-locality: Argentina, Rio Negro Prov., Bariloche. Distr. — Argentina (SW Rio Negro Prov.).

Genus BLAESOCHIAETOPHORA Czerny

- BLAESOCHIAETOPHORA Czerny, 1904:206. Type-species, *Leria pieticornis* Bigot (mon.).
- Camaterus* Aldrich, 1926:97. Type-species, *dichaeta* Aldrich (orig. des.).
- PICTICORNIS (Bigot), 1888:35 (*Leria*). Type-locality: Chile, Magallanes Prov., Tierra del Fuego, Orange Bay. Distr. — Chile (Malleco, Llanquihue, Chiloé, Aysén and Magallanes Prov., Juan Fernández Islands), Argentina (SW Rio Negro Prov.). Ref.: Malloch, 1933:191.
- bezzii* Brèthes, 1924:108 (*Chiromyia*). Type-locality: Chile, Chiloé Prov., Island of Chiloé.
- dichaeta* Aldrich, 1926:98 (*Camaterus*). Type-locality: Chile, Puyen [possibly Purén, Bio-Bio Prov.].

Genus CEPHODAPEDON Malloch

- CEPHODAPEDON Malloch, 1933:212. Type-species, *fulvicornis* Malloch (orig. des.).
- FULVICORNIS Malloch, 1933:212. Type-locality: Argentina, Rio Negro Prov., Bariloche. Distr. — Chile (Santiago, O' Higgins, Curicó and Aysén Prov.), Argentina (Rio Negro and Santa Cruz Prov.).

Genus DIACIA Wiedemann

- DIACIA Wiedemann, 1830:448. Type-species, *Seatophaga diadema* Wiedemann (Aldrich, 1926:101).
- Thyreophorella* Hendel, 1910:111. Type-species, *Seatophaga diadema* Wiedemann (orig. des.). Arguments have been presented to oppose

the treatment of *Thyreophorella* Hendel as a junior synonym of *Diacia* Wiedemann (e. g., Czcrny, 1927, pp. 42-43; Hendel, 1928, pp. 216-217). However, the use of *Diacia* Wiedemann follows that of Malloch (1933, p. 190) in his work on the Diptera of Patagonia and South Chile.

DIADEMA (Wiedemann), 1830:448 (*Scatophaga*). Type-locality: Uruguay, Montevideo. Distr. — Brazil, Uruguay. Ref. — Hendel, 1910:116.

Genus DIHOPLOPYGA Malloch

DIHOPLOPYGA Malloch, 1933:192. Type-species, *polita* Malloch (orig. des.).

NORMA Malloch, 1933:194. Type-locality: Chile, Llanquihue Prov., Casa Pangue. Distr. — Chile (Valdivia, Osorno, Llanquihue, Chiloé and Aysén Prov.).

POLITA Malloch, 1933:193, text-fig. 40. Type-locality: Chile, Llanquihue Prov., Puerto Montt. Distr. — Chile (Llanquihue, Chiloé and Aysén Prov.).

Genus EPISTOMYIA Malloch

EPISTOMYIA Malloch, 1933:198. Type-species, *aurifrons* Malloch (orig. des.).

AURIFRONS Malloch, 1933:198. Type-locality: Argentina, Rio Negro Prov., Bariloche. Distr. — Argentina (SW Rio Negro Prov.).

Genus GEPHYROMYZA Malloch

GEPHYROMYZA Malloch, 1933:208. Type-species, *testaceipes* Malloch (orig. des.).

GRANDIS Edwards in Malloch, 1933:210. Type-locality: Chile, Valdivia Prov., Lake Panguipulli. Distr. — Chile (Valdivia and Aysén Prov.).

PENICILLIPES (Enderlein), 1912:95, fig. 19 (*Leria*). Type-locality: Chile, Magallanes Prov., Tierra del Fuego, Orange Bay; Argentina, Tierra del Fuego Prov., Ushuaia. Distr. — Argentina (Rio Negro and Tierra del Fuego Prov.), Chile (Curicó, Arauco, Malleco, Aysén and Magallanes Prov.). Ref. — Malloch, 1933:211.

TESTACEIPES Malloch, 1933:209, text-fig. 44. Type-locality: Argentina, Rio Negro Prov., Lake Correntoso. Distr. — Chile (Curicó, Talca and Nuble Prov.), Argentina (SW Rio Negro Prov.).

VARIPES Malloch, 1933:212. Type-locality: Argentina, Rio Negro Prov., Lake Correntoso.

Genus HELOCLUSIA Malloch

HELOCLUSIA Malloch, 1933:184. Type-species, *imperfecta* Malloch (orig. des.).

IMPERFECTA Malloch, 1933:186; text-figs. 36, 37; pl. 2, fig. 13. Type-locality: Chile, Concepción Prov., Concepción. Distr. — Chile (Concepción and Aysén Prov.), Argentina (SW Rio Negro Prov.).

Genus NEPHONEURA Malloch

- NEPHONEURA Malloch, 1933:214. Type-species, *dendrophila* Malloch (orig. des.).
- DENDROPHILA Malloch, 1933:214, text-fig. 45. Type-locality: Argentina, Rio Negro Prov., Lake Gutiérrez. Distr. — Argentina (SW Rio Negro Prov.).

Genus NOTOMYZA Malloch

- NOTOMYZA Malloch, 1933:195. Type-species, *abrupta* Malloch (orig. des.).
- ABRUPTA Malloch, 1933:195, text-figs. 41a, 41d. Type-locality: Argentina, Rio Negro Prov., Bariloche. Distr. — Chile (Llanquihue Prov.), Argentina (SW Rio Negro Prov.).
- EDWARDSI Malloch, 1933:197, text-figs. 41a, 41b. Type-locality: Argentina, Rio Negro Prov., Lake Correntoso. Distr. — Chile (Llanquihue Prov.), Argentina (Rio Negro Prov.).
- MICROPYGA Malloch, 1933:197, text-figs. 41c, 41f. Type-locality: Argentina, Rio Negro Prov., Lake Correntoso. Distr. — Chile (Llanquihue Prov.), Argentina (SW Rio Negro Prov.).
- SPHAEROCERA (Thomson), 1869:597 (*Drosophila*). Type-locality: Argentina, Patagonia. Stuardo Ortiz (1946, p. 189) places this species under "Especies presuntivamente de Chile".

Genus PROSOPANTRUM Enderlein

- PROSOPANTRUM Enderlein, 1912:135. Type-species, *austrinum* Enderlein (orig. des.) = *diadematum* (Bigot).
- ACROSTICHALIA Tonnoir & Malloch, 1927:86. Type-species, *flavifrons* Tonnoir & Malloch (orig. des.).
- CNEMOSPATHIS Enderlein, 1940:655. Type-species, *baeckstroemi* Enderlein (orig. des.).
- DIDYMOCHAETA Czerny, 1929:87. Type-species, *peruana* Czerny (mon.).
- AEQUISETA Malloch, 1933:204, text-fig. 42c. Type-locality: Argentina, Rio Negro Prov., Lake Correntoso. Distr. — Chile (Llanquihue Prov., Peulla), Argentina (Rio Negro Prov., Lake Correntoso), Falkland Islands (?).
- DASYURA Edwards in Malloch, 1933:207, text-fig. 43c. Type-locality: Argentina, Rio Negro Prov., Lake Correntoso. Distr. — Argentina (SW Rio Negro Prov.), Chile (Magallanes Prov.).
- DIADEMATUM (Bigot), 1888:40, pl. 4, fig. 6 (*Agromyza*). Type-locality: Chile, Magallanes Prov., Tierra del Fuego, Orange Bay. Distr. — Chile (Magallanes Prov.), South Georgia Island. Ref. — Malloch, 1933:202, text-fig. 42b.
- austrinum* Enderlein, 1912:135, fig. 33. Type-locality: South Georgia Island, Bay of Islets.
- FLAVIFRONS (Tonnoir & Malloch), 1927:87 (*Acrostichalia*). Type-locality: New Zealand, Christchurch. Distr. — Argentina (Rio Negro Prov.), Chile (Llanquihue, Chiloé, Osorno and Magallanes Prov.,

- Juan Fernández Islands); New Zealand, South Africa. Ref. — Malloch, 1933:200, text-fig. 42a.
- baeckstroemi* Enderlein, 1940:656 (*Cnemospathis*). Type-locality: Juan Fernández Islands.
- schoenemanni* Enderlein, 1940:657, fig. 20 (*Cnemospathis*). Type-locality: Juan Fernández Islands.
- INERMALLOCH Malloch, 1933:207, text-fig. 42d. Type-locality: Argentina, Rio Negro Prov., Bariloche. Distr. — Chile (Llanquihue Prov.), Argentina (SW Rio Negro Prov.).
- MAGELLANICUM Edwards in Malloch, 1933:202. Type-locality: Chile, Magallanes Prov. Distr. — Argentina (Rio Negro Prov., Lake Correntoso), Chile (Magallanes Prov.).
- PAUCISETA (Enderlein), 1940:657 (*Cnemospathis*). Type-locality: Juan Fernández Islands. The two other species described by Enderlein from the Juan Fernández Islands, *baeckstroemi* and *schoenemanni*, were subsequently found by Hennig (1951, 1955) to be synonymous with *Prosopanthrum flavifrons* Tonnoir & Malloch. This species presumably also belongs in the genus *Prosopanthrum*.
- PERUANUM (Czerny), 1929:87 (*Didymochaeta*). Type-locality: "Peru". Malloch (1933, p. 208) considered this species very close to *P. flavifrons* (Tonnoir & Malloch). Soukup (1959, p. 213) retained *Didymochaeta* Czerny for this species.
- PRISTIS Edwards in Malloch, 1933:205, text-fig. 43b. Type-locality: Argentina, Rio Negro Prov., Lake Gutiérrez. Distr. — Chile (Llanquihue Prov.), Argentina (SW Rio Negro Prov.).
- RUFICORNE Edwards in Malloch, 1933:205, text-fig. 43a. Type-locality: Argentina, Rio Negro Prov., Bariloche.
- UNISETA Malloch, 1933:208. Type-locality: Argentina, Rio Negro Prov., Bariloche.
- VARIFRON Edwards in Malloch, 1933:202. Type-locality: Chile, Magallanes Prov.

Genus PSEUDOLERIA Garrett

- PSEUDOLERIA Garrett, 1921:128. Type-species, *Blepharoptera pectinata* Loew (orig. des.).
- PECTINATA (Loew), 1872:99 (*Blepharoptera*). Type-locality: U.S.A., Texas. Distr. — North America, Mexico (Guadalupe Island), Chile (Arauco, Malleco and Llanquihue Prov.); New Zealand, Australia. Ref. — Gill, 1962:538, fig. 36. The difficulty in identifying species of *Pseudoleria* without careful examination of male terminalia makes the Chile, New Zealand, and Australia records questionable; however, I personally examined male specimens from Guadalupe Island and am confident they are *P. pectinata* (Loew).

Genus SPILOCHROA Williston

- SPILOCHROA Williston, 1907:2. Type-species, *Heterochroa ornata* Johnson (mon.). Malloch (1933, p. 214) considered *Spilochroa* a subgenus of *Diastata* auct. nec Meigen (= *Trixoscelis* Rondani). *Diastata* Meigen is in the Diastatidae and is a genus quite distinct from *Trixoscelis* Rondani.

GEMINATA Sabrosky, 1961:233. Type-locality: U.S.A., Arizona, Maricopa County, Buckeye. Distr. — U.S.A. (Arizona), Mexico (Sonora, Imuris).

GUTTATA (Macquart), 1843:418, pl. 35, fig. 3 (*Opomyza*). Type-locality: "Chile". Distr. — Chile (Coquimbo, Aconcagua, Valparaíso, Santiago and Malleco Prov.), Argentina (Rio Negro and Santa Cruz Prov.).

chilensis Schiner, 1868:235 (*Diastata*). Type-locality: "Chile".

ORNATA (Johnson), 1895:306 (*Heterochroa*). Type-locality: U.S.A., Florida. Distr. — Southern U.S.A., West Indies.

PERUVIANA (Malloch), 1931:31 (*Diastata*, subg. *Spilochroa*). Type-locality: Peru, Lima, Verrugas Canyon.

Genus SUILLIA Robineau-Desvoidy

SUILLIA Robineau-Desvoidy, 1830:642. Type-species, *fungorum* Robineau-Desvoidy (Coquillett, 1910a: 610), probably = *rufa* (Fallén).

Helomyza, authors, not Fallén.

DISTIGMA (Wulp), 1897:352, pl. 9, fig. 5 (*Helomyza*). Type-locality: Mexico, Guerrero.

INIENS (Giglio-Tos), 1893:8 (*Helomyza*). Type-locality: "Mexico".

POLYSTIGMA (Wulp), 1897:353, pl. 9, fig. 6 (*Helomyza*). Type-locality: Mexico, Guerrero.

PUNCTULATA (Wulp), 1897:352, pl. 9, fig. 4 (*Helomyza*). Type-locality: Mexico, Guerrero.

Genus TRIXOSCELIS Rondani

TRIXOSCELIS Rondani, 1856:134. Type-species, *Gcomyza obscurella* Fallén (orig. des.).

Trichoscelis, emend.

Parodina Coquillett, 1902:186. Type-species, *cinerea* Coquillett (orig. des.).

Diastata, authors, not Meigen.

COSTALIS (Coquillett), 1901:378 (*Rhienoessa*). Type-locality: Ecuador, Galápagos Islands, Albemarle, Tagus Cove. Distr. — Ecuador (Galápagos Islands).

NUDA (Coquillett), 1910b:130 (*Leria*). Type-locality: U.S.A., California. Distr. — Western U.S.A., Mexico (Guadalupe Island). Ref. — Melander, 1952:48.

prima Hendel, 1911:43. Type-locality: U.S.A., California, Claremont.

TRIPLEX Melander, 1952:50. Type-locality: U.S.A., Arizona, Organpipe Cactus National Monument. Distr. — U.S.A. (California, Arizona), Mexico.

Genus ZACHAETOMYIA Malloch

ZACHAETOMYIA Malloch, 1933:187. Type-species, *atriventris* Malloch (orig. des.).



ATRIVENTRIS Malloch, 1933:188, text-fig. 38. Type-locality: Argentina, Rio Negro Prov., Bariloche. Distr. — Argentina (SW Rio Negro Prov.), Central Chile ("Cajón de Longave, Baños").

SPECIES OF UNCERTAIN POSITION

- APICALIS Schiner, 1868:232 (*Heteromyza*). Type-locality: "Süd Amerika".
- CHILENSIS Macquart, 1851:252, pl. 25, fig. 16 (*Helomyza*). Type-locality: "Chile." Stuardo Ortiz (1946, p. 154) lists this species among "species incertae sedis", although Czerny (1924, p. 56) considered it to be a *Suillia*.
- CHILOENSIS Brèthes, 1924:108 (*Anorostoma*). Type-locality: Chile, Chiloé Prov., Chiloé Is. Czerny (1927, p. 38) states this species cannot be an *Anorostoma*. Stuardo Ortiz (1946, p. 154) lists it among "species incertae sedis".
- ELONGATA Blanchard in Gay, 1852:451 (*Helomyza*). Type-locality: "Central Chile". Czerny (1904, p. 285; 1925, p. 156), Malloch (1933, p. 182) and Stuardo Ortiz (1946, p. 154) were unable to place this species.
- PALLIDICEPS Blanchard in Gay, 1852:451 (*Helomyza*). Type-localit*: Chile, "Coquimbo, Serena, etc." Czerny (1904, p. 285; 1924, p. 156), Malloch (1933, p. 183) and Stuardo Ortiz (1946, p. 154) were unable to place this species.

LITERATURE CITED

ALDRICH, J. M.

1926. A new genus of Heleomyzidae from Chile with key to genera. *Insecutor Inscitiae Menstruus* 14(4-6):96-102.

BIGOT, J. M. F.

1888. *Mission Sci. du Cap Horn. Diptères* 6:1-45, 4 pls.

BLANCHARD, E.

1852. Dipteros. In Gay, C. (ed.). *Historia fisica y politica de Chile. Zoologia (Dipteros)* 7:327-468.

BRÈTHES, J.

1924. Sur quelques Diptères chiliens. *Rev. Chilena Hist. Nat.* 28:104-111.

COQUILLET, D. W.

1901. Papers from the Hopkins Stanford Galapagos Expedition, 1898-1899. II. Entomological Results (2): Diptera. *Proc. Washington Acad. Sci.* 3:371-379.
1902. New acalyptrate Diptera from North America. *J. New York Entomol. Soc.* 10:177-191.
- 1910a. The type-species of the North American genera of Diptera. *Proc. U. S. Natl. Mus.* 37:499-647.
- 1910b. New genera and species of North American Diptera. *Proc. Entomol. Soc. Washington* 12:124-131.



CZERNY, L.

1904. Revision der Helomyziden. *Wiener Entomol. Ztg.* 23:199-244, 263-286, 1 pl.
1924. Monographie der Helomyziden (Dipteren). *Abhandl. Zool. Botan. Ges. Wien* 15(1):1-166, 1 pl.
1926. Ergänzungen und Berichtigungen zu meiner Monographie der Helomyziden. *Konowia* 5(1):53-56.
1927. Ergänzungen und Berichtigungen zu meiner Monographie der Helomyziden, II. *Konowia* 6(1):35-49, 2 figs.
1929. Ergänzungen zu meiner Monographie der Helomyziden, IV. *Konowia* 8:87-92.

ENDERLEIN, G.

1912. Die Insekten des Antarkto-Archipelago-Gebietes (Feuerland, Falklands-Inseln, Süd-Georgien). *K. Svenska Vetensk. Akad. Handl.* 48(3):1-170, 4 pls.
1940. Die Dipterenfauna der Juan-Fernández-Inseln und der Oster-Inseln. In Skottsberg, Nat. Hist. Juan Fernández and Easter Isl., Zool. 3:643-680, 49 figs.

FALLÉN, C. F.

1810. *Specimen entomologicum novum Diptera disponendi methodum exhibens*, 26 pp., 1 pl. Lund.
1820. *Heteromyzides Sveciae*. 10 pp., Lund.

GARRETT, C. B. D.

1921. Notes on Helomyzidae and descriptions of new species (Diptera). *Insector Inscitiae Menstruus* 9:119-132.

GIGLIO-TOS, E.

1893. Diagnosi di nuovi generi e di nuove specie di Ditteri. IX. *Boll. Mus. Zool. ed Anat. Comp.* [Torino] 8(158):1-14.

GILL, G. D.

1962. The helomyzid flies of America north or Mexico (Diptera: Helomyzidae). *Proc. U. S. Natl. Mus.* 113:495-603, 96 figs.

GORODKOV, K. B.

1959. Revision of the Palaearctic species of the genus *Occotheca* Hal. *Entomologicheskoe Obozrenie* 38(4):905-922, 64 figs. (Pages refer to the original, in Russian; in the English translation of this journal, *Entomological Review*, the pages are 398-416).

HALIDAY, A. H.

1838. Descriptions of new British insects indicated in Mr. Curtis's Guide. *Ann. Nat. Hist.* (1839) 2:183-190.

HARRISON, R. A.

1959. Family Helomyzidae, pp. 110-169, figs. 108-200, in Acalypterate Diptera of New Zealand. *New Zealand Dept. Sci. Indust. Res., Bull.* 128:382 pp., 438 figs.

HENDEL, F.

1910. Über acalypterate Museiden. *Wiener Entomol. Ztg.* 29:101-127.
1911. Über von Professor J. M. Aldrich erhaltene und einige andere amerikanische Dipteren. *Wiener Entomol. Ztg.* 30:19-46.



1928. Über zwei meiner Dipterengattungen die Doktor J. M. Aldrich nicht anerkennen will. *Konowia* 7(3):214-217.

HENNIG, W.

1948. Ueber einige verkannte Dipteren-Gattungen. *Acta Zool. Lilloana* 6:169-170.

1951. Neue acalyptraten aus Europa und Südafrika (Diptera: Pionilidae, Helomyzidae, Lauxanidae). *Beitr. z. Entomol.* [Berlin] 1 (1):70-76.

1955. Los insectos de las islas Juan Fernández, 16. *Rev. Chilena Entomol.* 4:21-34.

JOHNSON, C. W.

1895. Diptera of Florida. *Proc. Acad. Nat. Sci. Philadelphia* 1895: 303-340.

LOEW, H.

1872. Diptera Americae septentrionalis indigena. *Berliner Entomol. Zeitschr.* 16:49-115.

MACQUART, J.

1843. Diptères exotiques nouveaux ou peu connus. *Mém. Soc. Roy. des Sci., de l'Agr. et des Arts (Lille)* 1842:162-460, 36 pls.

1851. Diptères exotiques nouveau ou peu connus. Suite du 4^e supplément. *Mém. Soc. Natl. des Sci., de l'Agr. et des Arts [Lille]* 1850:134-294, pls. 15-28.

MALLOCH, J. R.

1931. Notes on some acalyptrate flies in the United States National Museum. *Proc. U. S. Natl. Mus.* 78:1-32, 10 figs.

1933. Acalyptrata, pp. 177-391, figs. 36-68, pls. 2-7, in *British Museum (Natural History), Diptera of Patagonia and South Chile*, Pt. 6, Fasc. 4, 499 pp., 84 figs., 8 pls. London.

MELANDER, A. L.

1952. The American species of Trixoseclidae. *J. New York Entomol. Soc.* 60:37-52.

ROBINEAU-DESVOIDY, J. B.

1830. Essai sur les Myodaires. *Mem. Acad. Roy. des Sci.* [Paris]: 2:1-813.

FONDANI, C.

1856. *Dipterologiae Italicae prodromus*. Vol. 1, 228 pp. Parma.

SABROSKY, C. W.

1961. Three new nearctic acalyptrate Diptera. *Entomol. News* 72:229-234, 1 fig.

SCHINER, J. R.

1868. Diptera, 388 pp., 4 pls., in *Reise der österreichischen Fregatte Novara um die Erde*. Zool., Vol. 2, Abt. 1, B. Wien.

SOUKUP, J.

1959. Segunda lista de Dipteros Peruanos. *Biota* 2(18):201-223.

- STONE, A., C. W. SABROSKY, W. W. WIRTH, R. H. FOOTE, AND J. R. COULSON (editors).
1965. *A Catalog of the Diptera of America North of Mexico*. U. S. Dept. Agr., Res. Scr., Agr. Handb. No. 276. 1696 pp.
- STUARDO ORTIZ, C.
1946. *Catalogo de los dipteros de Chilc*. Minist. Agr. 250 pp. Santiago.
- THOMSON, C. G.
(1869). Diptera. Species nova descripsit. pp. 443-614, pl. 9. In K. Svenska Vetenskaps-Akademien, *Kongliga svenska fregatten Eugenie res a omkring jorden*. Pt. 2: Zoologic, 1: Insekter, 617 pp., 9 pls. Stockholm (1868).
- TONNOIR, A. L. & J. R. MALLOCH
1927. New Zealand Muscidae Acalyptratae. Helomyzidae. *Rec. Canterbury Mus.* 3(2):83-100, 3 pls.
- WESTWOOD, J. O.
1840. Diptera, in *An introduction to the modern classification of insects*. 158 pp. London.
- WIEDEMANN, C. R. W.
1830. *Aussereuropäische zweiflügelige Insekten*. Vol. 2. xii + 684 pp., 5 pls. Hamm.
- WILLISTON, S. W.
1907. Dipterological notes. *J. New York Entomol. Soc.* 15:1-2.
- WULP, F. M. VAN DER
1897. Fam. Muscidae. Pp. 345-376, pl. 9. In Godman, F. D. and Salvin, O. (editors). *Biologia Centrali-Americana*. Zoologia-Insecta-Diptera, Vol. 2, 489 pp., 11 figs., 13 pls. London.

INDEX

abrupta Malloch, 1933, Notomyza	85.5
Acrostichalia Tonnoir & Malloch, 1927	85.5
Accothes Haliday, 1838	85.2
aequiseta Malloch, 1933, Prosopantrum	85.5
Anastomyza Malloch, 1933	85.2
apicalis Sehiner, 1868, Heteromyza	85.8
Apophoneura Malloch, 1933	85.2
atriventris Malloch, 1933, Zachatomyia	85.8
aurifrons Malloch, 1933, Epistomyia	85.4
austrinum Enderlein, 1912, Prosopantrum	85.5
baeckstroemi (Enderlein), 1940 (Cnemospathis), Prosopantrum ...	85.6
buzzi (Brèthes), 1924 (Chiromyia), Blaesochactophora	85.3
Blaesochactophora Czerny, 1904	85.3
Camaterus Aldrich, 1926	85.3
Cephodapedon Malloch, 1933	85.3
chilensis Macquart, 1851 (Helomyza)	85.8





<i>chilensis</i> (Schiner), 1868 (Diastata), <i>Spilochroa</i>	85.7
<i>chilocensis</i> Brèthes, 1924 (Anorostoma)	85.8
<i>Cnemospithis</i> Enderlein, 1940	85.5
<i>costalis</i> (Coquillett), 1901 (Rhicnocssa), <i>Trixoscelis</i>	85.7
<i>dasyura</i> Edwards, 1933, <i>Prossopantrum</i>	85.5
<i>deleta</i> Edwards, 1933, <i>Anastomyza</i>	85.2
<i>dendrophila</i> Malloch, 1933, <i>Nephoneura</i>	85.5
<i>Diacia</i> Wiedemann, 1830	85.3
<i>diadema</i> (Wiedemann), 1830 (Scatophaga), <i>Diacia</i>	85.4
<i>diadematum</i> (Bigot), 1888 (Agromyza), <i>Prosopantrum</i>	85.5
<i>Diastata</i> , authors, not Meigen	85.7
<i>dichaeta</i> (Aldrich), 1926 (Camaterus), <i>Blaesochaetophora</i>	85.3
<i>Didymochaeta</i> Czerny, 1929	85.5
<i>Dihoplopyga</i> Malloch, 1933	85.4
<i>discalis</i> Malloch, 1933, <i>Anastomyza</i>	85.2
<i>distigma</i> (Wulp), 1897 (Helomyza), <i>Suillia</i>	85.7
<i>edwardsi</i> Malloch, 1933, <i>Notomyza</i>	85.5
<i>elongata</i> Blanchard, 1852 (Helomyza)	85.8
<i>Epistomyia</i> Malloch, 1933	85.4
<i>exquisita</i> Malloch, 1933, <i>Apophoneura</i>	85.2
<i>fenestralis</i> (Fallén), 1820 (Helomyza), <i>Accothea</i>	85.2
<i>flavifrons</i> (Tonnoir & Malloch), 1927 (Acrostichalia), <i>Prosopantrum</i>	85.5
<i>fractura</i> Malloch, 1933, <i>Anastomyza</i>	85.2
<i>fulvicornis</i> Malloch, 1933, <i>Cephodapedon</i>	85.3
<i>fulvithorax</i> Malloch, 1933, <i>Anastomyza</i>	85.2
<i>geminata</i> Sabrosky, 1961, <i>Spilochroa</i>	85.7
<i>Gephyromyza</i> Malloch, 1933	85.4
<i>grandis</i> Edwards, 1933, <i>Gephyromyza</i>	85.4
<i>guttata</i> (Macquart), 1843 (Opomyza), <i>Spilochroa</i>	85.7
<i>Heleomyza</i> , authors	85.7
<i>Heloclusia</i> Malloch, 1933	85.4
<i>hyalipennis</i> Malloch, 1933, <i>Anastomyza</i>	85.2
<i>imperfecta</i> Malloch, 1933, <i>Heloclusia</i>	85.4
<i>inconspicua</i> Malloch, 1933, <i>Apophoneura</i>	85.3
<i>inconspicua</i> Malloch, 1933, <i>Apophoneura</i>	85.3
<i>inermis</i> Malloch, 1933, <i>Prosopantrum</i>	85.6
<i>iniens</i> (Giglio-Tos), 1893 (Helomyza), <i>Suillia</i>	85.7
<i>latralis</i> Malloch, 1933, <i>Anastomyza</i>	85.2
<i>magellanicum</i> Edwards, 1933, <i>Prosopantrum</i>	85.6
<i>micropyga</i> Malloch, 1933, <i>Notomyza</i>	85.5
<i>multipunctata</i> Malloch, 1933, <i>Anastomyza</i>	85.2




<i>neglecta</i> Edwards, 1933, <i>Anastomyza</i>	85.2
<i>Nephoneura</i> Malloch, 1933	85.5
<i>norma</i> Malloch, 1933, <i>Dihoplopyga</i>	85.4
<i>Notomyza</i> Malloch, 1933	85.5
<i>nuda</i> (Coquillett), 1910 (<i>Leria</i>), <i>Trixoscelis</i>	85.7
<i>Oecothoa</i> , emend. of <i>Accothoa</i> Haliday, 1838	85.2
<i>ornata</i> (Johnson), 1895 (<i>Heterochroa</i>), <i>Spilochroa</i>	85.7
<i>pallidiceps</i> Blanchard, 1852 (<i>Helomyza</i>)	85.8
<i>Parodina</i> Coquillett, 1902	85.7
<i>pauciseta</i> (Enderlein), 1940 (<i>Cnemospathis</i>), <i>Prosopantrum</i>	85.6
<i>pectinata</i> (Locw), 1872 (<i>Blepharoptera</i>), <i>Pseudoleria</i>	85.6
<i>penicillipes</i> (Enderlein), 1912 (<i>Leria</i>), <i>Gephyromyza</i>	85.4
<i>peruanum</i> (Czerny), 1929 (<i>Didymochaeta</i>), <i>Prosopantrum</i>	85.6
<i>peruviana</i> (Malloch), 1931 (<i>Diastata</i> , subg. <i>Spilochroa</i>), <i>Spilochroa</i>	85.7
<i>picta</i> Malloch, 1933, <i>Apophoneura</i>	85.3
<i>picticornis</i> (Bigot), 1888 (<i>Leria</i>), <i>Blaesochaetophora</i>	85.3
<i>polita</i> Malloch, 1933, <i>Dihoplopyga</i>	85.4
<i>polystigma</i> (Wulp), 1897 (<i>Helomyza</i>), <i>Suillia</i>	85.7
<i>prima</i> Hendel, 1911, <i>Trixoseclis</i>	85.7
<i>pristis</i> Edwards, 1933, <i>Prosopantrum</i>	85.6
<i>Prosopantrum</i> Enderlein, 1912	85.5
<i>Pseudoleria</i> Garrett, 1921	85.6
<i>punctipennis</i> Malloch, 1933, <i>Apophoneura</i>	85.3
<i>punctulata</i> (Wulp), 1897 (<i>Helomyza</i>), <i>Suillia</i>	85.7
<i>recurrens</i> Malloch, 1933, <i>Apophoneura</i>	85.3
<i>ruficorne</i> Edwards, 1933, <i>Prosopantrum</i>	85.6
<i>shoenemannii</i> (Enderlein), 1940 (<i>Cnemospathis</i>), <i>Prosopantrum</i> ...	85.6
<i>specularis</i> Malloch, 1933, <i>Anastomyza</i>	85.2
<i>sphaerocera</i> (Thomson), 1869 (<i>Drosophila</i>), <i>Notomyza</i>	85.5
<i>Spilochroa</i> Williston, 1907	85.6
<i>Suillia</i> Robineau-Desvoidy, 1830	85.7
<i>superba</i> Malloch, 1933, <i>Apophoneura</i>	85.3
<i>testaceipes</i> Malloch, 1933, <i>Gephyromyza</i>	85.4
<i>Thyreophorella</i> Hendel, 1910	85.3
<i>Trichoseclis</i> , emend. of <i>Trixoscelis</i> Rondani, 1858	85.7
<i>triplex</i> Melander, 1952, <i>Trixoseclis</i>	85.7
<i>Trixoscelis</i> Rondani, 1856	85.7
<i>uniseta</i> Malloch, 1933, <i>Prosopantrum</i>	85.6
<i>univittata</i> Malloch, 1933, <i>Apophoneura</i>	85.3
<i>varifrons</i> Edwards, 1933, <i>Prosopantrum</i>	85.6
<i>varipes</i> Malloch, 1933, <i>Gephyromyza</i>	85.4
<i>vittigera</i> Malloch, 1933, <i>Apophoneura</i>	85.3
<i>Zachatomyia</i> Malloch, 1933	85.7









MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO
(Formerly Departamento de Zoologia, Secretaria da Agricultura, São Paulo)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

85A

FAMILY NOTOMYZIDAE

ANGELO PIRES DO PRADO

*Departamento de Zoologia, Setor de Parasitologia
Universidade Estadual de Campinas, São Paulo, Brasil*

This monobasic family has until fairly recently been included with the Heleomyzidae, where it was catalogued by Gill (1968). It was elevated to familial rank by Griffiths (1972:260).

Genus NOTOMYZA Malloch

- NOTOMYZA Malloch, 1933:195. Type-species, *abrupta* Malloch (orig. des.).
ABRUPTA Malloch, 1933:195, text-figs. 41a, 41d. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Chile (Llanquihue), Argentina (sw. Rio Negro prov.).
EDWARDSI Malloch, 1933:197, text-figs. 41b, 41e. Type-locality: Argentina, Rio Negro, Lake Correntoso. Distr. — Chile (Llanquihue), Argentina (sw. Rio Negro prov.).
MICROPYGA Malloch, 1933:197, text-figs. 41c, 41f. Type-locality: Argentina, Rio Negro, Lake Correntoso. Distr. — Chile (Llanquihue), Argentina (sw. Rio Negro prov.).
SPHAEROCERA (Thomson), 1869:597 (*Drosophila*). Type-locality: Argentina, Patagonia. Stuardo Ortiz (1946:189) places this species under "Especies presuntivamente de Chile".

LITERATURE CITED

- GILL, G. D.
1968. Family Heleomyzidae (Helomyzidae), including the Trixoscelididae (Trichoscelidae) (86), in Departamento de Zoologia, Secretaria da Agricultura, *A Catalogue of the Diptera of the Americas South of the United States* 85:1-13. São Paulo.
- GRIFFITHS, G. C. D.
1972. *The phylogenetic classification of Diptera Cyclorrhapha, with special reference to the structure of the male postabdomen*, 341 pp., 154 figs., 2 pls. [In W. Junk Publ., Series Entomologica, Vol. 8]. The Hague.



MALLOCH, J. R.


1933. Acalyptrata, pp. 177-391, figs. 36-38, pls. 2-7, in British Museum (Natural History), *Diptera of Patagonia and South Chile* 6(4) : 499 pp., 84 figs., 8 pls. London.

STUARDO ORTIZ, C.

1946. *Catalogo de los dipteros de Chile*, 250 pp. Minist. Agric., Santiago de Chile.

THOMSON, C. G.

1869. Diptera. Species novas descripsit, pp. 443-614, pl. 9, in K. Svenska Vetenskaps-Akademien, *Konglika svenska fregatten Eugenie resa omkring jorden 2* (Zool.) (1; Insekter) : 617 p., 9 pls. Stockholm, "1868".



MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO
(FORMERLY DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

85B

FAMILY CNEMOSPATHIDAE

ANGELO PIRES DO PRADO

*Departamento de Parasitologia,
Universidade Estadual de Campinas, São Paulo*

This monobasic family of South American flies was reinstated by Griffiths (1972). Until recently it has been included with the Heleomyzidae (Gill, 1968). The type-genus *Cnemospathis* was synonymized with *Prosopanthrum* by Hennig (1948). Because of Enderleins's proposal of family rank, the name Cnemospathidae is yet available. One species, *P. flavifrons* (Tonnoir & Malloch), 1927, also occurs in New Zealand and South Africa (Hennig, 1951; Cogan, 1971).

Genus PROSOPANTHRUM Enderlein

PROSOPANTHRUM Enderlein, 1912:135. Type-species, *austrinum* Enderlein (orig. des.) = *diadematum* (Bigot).

Acrostichalia Tonnoir & Malloch, 1927:86. Type-species, *flavifrons* Tonnoir & Malloch (orig. des.).

Cnemospathis Enderlein, 1940:655. Type-species, *baeckstroemi* Enderlein (orig. des.).

Didymochaeta Czerny, 1929:87. Type-species, *peruana* Czerny (mon.).

AEQUISETA Malloch, 1933:204, text-fig. 42c. Type-locality: Argentina, Rio Negro Prov., Lake Correntoso. Distr. - Chile (Llanquihue Prov., Peulla); Argentina (Rio Negro Prov.); Falkland Islands ?

DASYURA Edwards *in* Malloch, 1933:207, text-fig. 43c. Type-locality: Argentina, Rio Negro Prov., Lake Correntoso. Distr. - Chile (Magallanes Prov.); Argentina (SW Rio Negro Prov.).

DIADEMATUM (Bigot), 1888:40, pl. 4, fig. 6 (*Agromyza*). Type-locality: Chile, Magallanes Prov.; South Georgia Island. Ref.: Malloch, 1933:202, text-fig. 42b.

austrinum Enderlein, 1912:135, fig. 33. Type-locality: South Georgia Island, Bay of Islets.

FLAVIFRONS (Tonnoir & Malloch), 1927:87 (*Acrostichalia*). Type-locality: New Zealand, Christchurch. Distr. - Chile (Llanquihue, Chiloé, Osorno and Magallanes Prov.); Argentina (Rio Negro Prov.); Juan Fernandez Islands; New Zealand (South Island); South Africa. Refs.: Hennig, 1951; Cogan, 1971:693, fig. 101.

baeckstroemi Enderlein, 1940:656 (*Cnemospathis*). Type-locality: Juan Fernandez Islands.

schoenemanni Enderlein, 1940:657, fig. 20 (*Cnemospathis*). Type-locality: Juan Fernandez Islands.

INERME Malloch, 1933:207, text-fig. 42d. Type-locality: Argentina, Rio Negro Prov., Bariloche. Distr. - Chile (Magallanes Prov.); Argentina (SW Rio Negro Prov.).

MAGELLANICUM Edwards *in* Malloch, 1933:202. Type-locality: Chile, Magallanes Prov. Distr. - Chile; Argentina (Rio Negro Prov., Lake Correntoso).



- PAUCISETA (Enderlein), 1940:657 (*Cnemospathis*). Type-locality: Juan Fernandez Islands. Distr. - Juan Fernandez Islands.
- PERUANUM (Czerny), 1929:87 (*Didymochaeta*). Type-locality: "Peru". Distr. Peru.
- PRISTIS Edwards in Malloch, 1933:205, text-fig. 43b. Type-locality: Argentina, Rio Negro Prov., Lake Gutiérrez. Distr. - Chile (Llanquihue Prov.); Argentina (SW Rio Negro Prov.).
- RUFICORNE Edwards in Malloch, 1933:205, text-fig. 43a. Type-locality: Argentina, Rio Negro Prov., Bariloche. Distr. - Argentina.
- UNISETA Malloch, 1933:208. Type-locality: Argentina, Rio Negro Prov., Bariloche. Distr. - Argentina.
- VARIFRONS Edwards in Malloch, 1933:202. Type-locality: Chile, Magallanes Prov. Distr. - Chile.

LITERATURE CITED

BIGOT, J.M.F.

1888. *Mission Sci. du Cap Horn. Diptères* 6: 1-45, 4 pls.

CZERNY, L.

1929. Ergänzungen zu meiner Monographie der Helomyziden. IV. *Konowia* 8: 87-92

COGAN, B.H.

1971. The Heleomyzidae of the Ethiopian Region (Diptera). *Ann. Natal Mus.* 20 (3): 627-696, 101 figs.

ENDERLEIN, G.

1912. Die Insekten des Antarkto-Archipelago-Gebietes (Feuerland, Falklands-Inseln, Süd-Georgien). *K. svenska Vetensk-Akad. Handl.* 48 (3): 1-170, 4 pls.
1940. Die Dipterenfauna der Juan-Fernández-Inseln und der Oster-Inseln. In Skottsberg, *Nat. Hist. Juan Fernández and Easter Isl.*, Zool. 3: 643-680, 49 figs.

GILL, D.G.

1968. Family Heleomyzidae (Helomyzidae), including the Trixoscelididae (Trichoscelidae) (86), in Departamento de Zoologia, Secretaria da Agricultura, *A Catalogue of the Diptera of the Americas South of the United States* 85: 1-13, São Paulo.

GRIFFITHS, G.C.D.

1972. *The phylogenetic classification of Diptera Cyclorrhapha, with special reference to the structure of the male postabdomen*, 341 pp., 154 figs., 2 pls. (in W. Junk Publ., Series Entomologica, vol. 8). The Hague.

HENNIG, W.


1948. Ueber einige verkannte Dipteren-Gattungen. *Acta Zool. Lilloana* 6: 169-170.
1951. Neue Acalypraten aus Europa und Südafrika (Diptera: Piophilidae, Helomyzidae, Lauxaniidae). *Beitr. Ent.* 1: 70-76.

MALLOCH, J.R.

1933. Acalyprata, pp. 177-391, figs. 36-68, pls. 2-7, in British Museum (Natural History), *Diptera of Patagonia and South of Chile* 6 (4): 499 pp., 84 figs., 8 pls. London.

TONNOIR, A.L. & MALLOCH, J.R.

1927. New Zealand Muscidae Acalypratae. Helomyzidae. *Rec. Canterbury Mus.* 3 (2): 83-100, 3 pls.



DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

87

FAMILY RHINOTORIDAE

- NELSON PAPAVERO

*Departamento de Zoologia, Secretaria da Agricultura
São Paulo, Brasil*

A small family of forest-inhabiting flies, ranging from the south-western United States to southern Brazil. Nothing is known about their biology, except that the adults live on the bark of trees, licking the sap exuded from the trunks. The family is still poorly known systematically and many more species are expected to occur in the Neotropical region.

We are greatly indebted to Dr. George Steyskal, for his suggestion of a new generic arrangement. According to him the genera *Rhinotora* and *Neorhinotora* should be distinguished on the basis of the shape of the scutellum. In *Rhinotora*, the scutellum has one apical and two lateral spurs, while in *Neorhinotora* the scutellum has no spurs. The character used before, the presence of crossveins between C and Sc is variable, and of no generic importance. We have accepted the redefinition of the above cited genera, and consequently two new combinations were made.

Genus NEORHINOTORA Lopes

NEORHINOTORA Lopes, 1934:517. Type-species, *Rhinotora mutica* Schiner (orig. des.).

AMAPAENSIS Guimarães & Papavero, 1966:263, figs. 1-2 (male genitalia).
Type-locality: Brazil, Território do Amapá, Serra do Navio. Distr. — Brazil (Território do Amapá, Serra do Navio).

ARISTALIS (Fischer), 1932:446, figs. 11, 22 (scutellum, wing) (*Rhinotora*).
Type-locality: Brazil, Mato Grosso, Fazenda Murtinho. Distr. — Brazil (Mato Grosso, Guanabara, São Paulo). Ref. — Lopes, 1935: 25, figs. 3, 13-16 (egg, male genitalia).

Published under the auspices of the "Fundação de Amparo à Pesquisa do Estado de São Paulo."

87.1

DEPARTAMENTO DE ZOOLOGIA
BIBLIOTECA
SECRETARIA DA AGRICULTURA

- DIVERSA (Giglio-Tos), 1893:13 (*Rhinotora*). Type-locality: "Mexico". Distr. — U.S.A. (Arizona, New Mexico), Mexico. Refs. — Steyskal, 1957:70, figs. 5A-C (male genitalia); Wheeler, 1954:35, figs. 1-2 (male genitalia). N. comb.
- FONSECAI (Lopes), 1934:517 (*Rhinotora*). Type-locality: Brazil, São Paulo, Capital, Cantareira. Distr. — Brazil (São Paulo). Ref. — Lopes, 1935:22, fig. 8, pl. 1, fig. 2 (scutellum, wing). N. comb.
- MUTICA (Schiner), 1868:234 (*Rhinotora*). Type-locality: "Brazil". Distr. — Brazil (Guanabara, São Paulo, Paraná, Santa Catarina, Rio Grande do Sul). Ref. — Lopes, 1935:25, figs. 2, 11-12, 17 (egg, male genitalia).

Genus RHINOTORA Schiner

- RHINOTORA Schiner, 1868:233. Type-species, *pluricellata* Schiner (orig. des.).
- DIVERSIPENNIS Lopes, 1936:107, figs. 3-4, 7-9 (male genitalia, crossveins between C and Sc, wing). Type-locality: Brazil, São Paulo, Capital, Cantareira. Distr. — Brazil (São Paulo, Rio Grande do Sul).
- LOPESI Guimarães & Papavero, 1966:264, figs. 3-4 (male genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão, Alto da Boa Vista, Fazenda da Guarda. Distr. — Brazil (São Paulo, Rio Grande do Sul).
- PERUANA Kertész, 1901:415, pl. 20, figs. 8-9 (scutellum, wing). Type-locality: Peru, Callanga. Distr. — Peru (Callanga).
- PLURICELLATA Schiner, 1868:233. Type-locality: "Brazil". Distr. — Brazil (Guanabara, São Paulo). Refs. — Fischer, 1932:425, figs. 8-9, 20 (antenna, scutellum, wing); Lopes, 1935:20, figs. 1, 4-7 (egg, male genitalia, ovipositor).
- SALESOPOLITANA Guimarães & Papavero, 1966:266, figs. 5-7 (male genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica do Boracéia. Distr. — Brazil (São Paulo, Salesópolis).
- sp.* Wulp, 1898:380, pl. 10, figs. 4, 4a (head). Locality: Volcán de Irazu, 6000 feet, Costa Rica.
- SPILOPTERA Guimarães & Papavero, 1966:266, figs. 8-10 (male genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica do Boracéia. Distr. — Brazil (São Paulo, Salesópolis).
- TRAVASSOSI Lopcs, 1934:517. Type-locality: Brazil, São Paulo, Capital, Cantareira. Distr. — Brazil (Minas Gerais, São Paulo). Refs. — Lopes, 1935:20, pl. 1, fig. 1 (wing); 1936:106, figs. 1-2, 5-6 (male genitalia, crossveins between C and Sc).

Genus RHINOTOROIDES Lopes

- RHINOTOROIDES Lopes, 1934:517. Type-species, *bifurcata* Lopes (orig. des.).
- BIFURCATA Lopes, 1934:517. Type-locality: Brazil, São Paulo, Capital, Cantareira. Distr. — Brazil (São Paulo, Paraná). Ref. — Lopes, 1935:24, figs. 9-10, pl. 1, fig. 3 (male genitalia, wing).



LITERATURE CITED

FISCHER, C. P.

1932. Um genero e duas especies novas de Rhopalomeridae do Brasil o o pupario de *Willistonella pleuropunctata* Wied. (Dipt.). *Rev. Entom.*, Rio de Janeiro 2:441-450, 23 figs.

GIGLIO-TOS, E.

1893. Diagnosi di nuovi generi e nuove specie di ditteri. *Boll. Mus. Torino* 8(158):1-14.

GUIMARÃES, J. H. & N. PAPAVERO

1966. A contribution to the knowledge of the Neotropical Rhinotoridae (Diptera, Acalyptratae). *Papéis Avulsos Dep. Zool. S. Paulo* 18:261-269, 13 figs.

KERTÉSZ, K.

1901. Neue und bekannte Dipteren in den Sammlung des Ungarischen Nationalmuseums. *Természetrázi Füzetek* 24:403-432, pl. 20.

LOPES, H. S.

1934. Nota previa sobre alguns Rhopalomeridae (Dipt.). *Rev. Entom.*, Rio de Janeiro 4:517.
1935. A sub-familia Rhinotorinae Williston (Dipt. — Rhopalomeridae). *Arch. Inst. Biol. Veget.*, Rio de Janeiro 2(1):19-26, 17 figs.
1936. Sobre duas especies brasileiras de *Rhinotora* Schiner (Dipt. Rhopalomeridae). *Rev. Entom.*, Rio de Janeiro 6:106-109, 9 figs.

SCHINER, J. R.

1868. *Reise der österreichischen Fregatte Novara um die Erde*. Zool. Theil, Diptera: 388 pp., 4 pls. K. K. Hof- und Staatsdruckerei. Wien.

STEYSKAL, G. C.

1957. The postabdomen of male acalyptrate Diptera. *Ann. Entom. Soc. Amer.* 50:66-73, 5 figs.

WHEELER, M. R.

1954. *Rhinotora diversa* Giglio-Tos from the southwestern United States (Diptera, Rhinotoridae). *Wasmann Journ. Biol.* 12:35-39, 2 figs.

WULP, F. M. VAN DER


1898. *Biologia Centrali-Americana, Diptera* 2: ix + 490 pp., 13 pls.

INDEX

amapaensis Guimarães & Papavero, 1966, Neorhinotora	87.1
aristalis (Fischer), 1932 (Rhinotora), Neorhinotora	87.1



bifurcata Lopes, 1934, Rhinotoroides	87.2
diversa (Giglio-Tos), 1893 (Rhinotora), Neorhinotora	87.2
diversipennis Lopes, 1936, Rhinotora	87.2
fonsecai (Lopes), 1934 (Rhinotora), Neorhinotora	87.2
lopesi Guimarães & Papavero, 1966, Rhinotora	87.2
mutica (Schiner), 1868 (Rhinotora), Neorhinotora	87.2
Neorhinotora Lopes, 1934	87.1
peruana Kertész, 1901, Rhinotora	87.2
pluricellata Schiner, 1868, Rhinotora	87.2
Rhinotora Schiner, 1868	87.2
Rhinotoroides Lopes, 1934	87.2
salepolitana Guimarães & Papavero, 1966, Rhinotora	87.2
sp. Wulp, 1898, Rhinotora	87.2
spiloptera Guimarães & Papavero, 1966, Rhinotora	87.2
travassosi Lopes, 1934, Rhinotora	87.2



MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO
(Formerly Departamento de Zoologia, Secretaria da Agricultura, São Paulo)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

89A

FAMILY TERATOMYZIDAE

ANGELO PIRES DO PRADO

*Departamento de Zoologia, Setor de Parasitologia
Universidade Estadual de Campinas, São Paulo, Brasil*

This is a family of rarely encountered flies with a discontinuous distribution, occurring in New Zealand, Australia, Nepal, and South America (Chile and Brazil). Members of this group were formerly included in the Anthomyzidae or Opomyzidae, only recently having been given family rank (Colless & McAlpine, 1970). Many of the Neotropical species in collections remain undescribed.


Genus TERATOPTERA Malloch

- Teratomyza*, subg. TERATOPTERA Malloch, 1933:387. Type-species, *chilensis* Malloch (orig. des.).
CHILENSIS Malloch, 1933:388, pl. 7, figs. 12, 14. Type-locality: Chile, Ancud. Distr. — Chile.
LATIPENNIS Malloch, 1933:388, pl. 7, figs. 13, 15. Type-locality: Chile, Mechuque Island. Distr. — Chile.
sp. Hennig, 1971:37, figs. 13, 43, 45 ("*Teratomyza*-Gruppe"). — Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

LITERATURE CITED

- COLLESS, D. H. & D. K. MCALPINE
1970. [Fam. Teratomyzidae] Diptera (Flies), pp. 656-740, in *The Insects of Australia*. Melbourne Univ. Press.
HENNIG, W.
1971. Neue Untersuchungen über die Familien der Diptera Schizophora (Diptera: Cyclorrhapha). *Stuttgart. Beitr. z. Naturk.* 226:1-76, 108 figs.
MALLOCH, J. R.
1933. *Diptera of Patagonia and South Chile* 4(4) (Acalyptatae) pp. 176-391, pls. 2-7, figs.





MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO

(FORMERLY DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

90

FAMILY CHYROMYIIDAE

ANGELO PIRES DO PRADO

*Departamento de Parasitologia,
Universidade Estadual de Campinas, São Paulo*

One Nearctic species recorded from Mexico, and one indetermined species from West Indies (Bahamas), are the only records of this small family in the Neotropical Region. Nothing is known about the habits or immature stages of the Neotropical forms.

Genus APHANIOSOMA Becker

APHANIOSOMA Becker, 1903:186. Type-species, *approximatum* Becker (mon.).

FRONTATUM Wheeler, 1961:89. Type-locality: U.S.A., California. Distr. - U.S.A. - (California; Arizona; Utah); Mexico.

SP. Melander, 1913:300. West Indies, Bahamas.

LITERATURE CITED

BECKER, T.


1903. Aegyptische Dipteren. *Berlin. Zool. Mus. Mitt.* 2(H.3):67-195, 4 pls.

WHEELER, M.R.

1961. New species of southwestern acalyptrate Diptera. *Southwest Nat.* 6:86-91, 1 fig.

MELANDER, A.L.

1913. A synopsis of the Dipterous groups Agromyzinae, Milichiinae, Ochthiphiinae and Geomyzinae. *Jour. N.Y. Ent. Soc.* 21(4): 283-300, pl. 8, 8 figs.



SciELO

DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

91

FAMILY AULACIGASTRIDAE

NELSON PAPAVERO

*Departamento de Zoologia, Secretaria da Agricultura
São Paulo, Brasil*

This family is represented in the Neotropical region by two genera and two species. The biology and immature stages of *Aulacigaster leucopeza* have been studied by Dufour (1864), Malloch & McAtee (1924), and Robinson (1953). Hennig (1958, 1965) has studied the phylogenetic relationships of the family and its fossil representatives.

Genus AULACIGASTER Macquart

AULACIGASTER Macquart, 1835:579. Type-species, *rufitarsis* Macquart (mon.)
= *leucopeza* (Meigen).

Aulacogaster, emend.

LEUCOPEZA (Meigen), 1830:100 (*Diastata*). Type-locality: "Europe". Distr.
— Europe, North America (Saskatchewan to Quebec, south to
Florida), Mexico.

Genus SCHIZOCHROA Hennig

SCHIZOCHROA Hennig, 1956:151. Type-species, *melanoleuca* Hennig (orig.
des.).

MELANOLEUCA Hennig, 1956:152, figs. 5-8. Type-locality: Costa Rica,
Raventa, Farm Hamburg. Distr. — Costa Rica.

Published under the auspices of the "Fundação de Amparo à Pesquisa do
Estado de São Paulo."

91.1



LITERATURE CITED

DUFOUR, L.

1846. Histoire des métamorphoses de l'*Aulacigaster rufitarsis* et observations critiques sur ce genre de Muscides Acalypterécs. *Ann. Soc. Ent. France* (2)4:455-463, pl. 11, no. 1, figs. 1-14.

HENNIG, W.

1956. Neue neotropische Acalyptrata im Deutschen Entomologischen Institut. *Beitr. z. Ent.* 6:146-154.
1958. Die Familien der Diptera Schizophora und ihre phylogenetischen Verwandtschaftsbeziehungen. *Beitr. z. Ent.* 8:505-688, 365 figs.
1965. Die Acalyptratae des baltischen Bernsteins. *Stuttgarter Beitr. z. Naturkunde* 145:1-215, 519 figs.

MALLOCH, J. R. & W. L. MCATEE

1924. Flies of the family Drosophilidae of the District of Columbia region, with keys to genera, and other notes, of broader application. *Proc. Biol. Soc. Washington* 37:25-41, pls. 8-9.

MACQUART, J.

1835. *Histoire naturelle des insectes diptères* 2: 703 pp., 12 pls.

MEIGEN, J. W.

1830. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten* 6: iv + 401 pp., pls. 55-66, Hamm.

ROBINSON, I.

1953. The postembryonic stages in the life cycle of *Aulacigaster leucopera* (Meigen) (Diptera Cyclorrhapha: Aulacigasteridae). *Proc. Roy. Ent. Soc. London* (A) 28:77-84, 10 figs.



DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

92

FAMILY ASTEIIDAE

CURTIS W. SABROSKY

Entomology Research Division, Agric. Res. Serv.
U.S. Department of Agriculture
Washington, D.C.

The small family Asteiidae contains small, delicate, seldom-collected flies. Virtually nothing is known of their biology. Sabrosky (1956, 1957) published a key to the genera of the world and a synopsis of the New World species. The family is best developed in tropical regions, and many more Neotropical species may be found.

Subfamily SIGALOESSINAE

Genus PHLEBOSOTERA Duda

PHLEBOSOTERA Duda, 1927:119, 125. Type-species, *mollis* Duda (orig. des., as gen. n., sp. n.).

PECULIARIS Sabrosky, 1957:51. Type-locality: Virgin Is., St. Thomas. Distr. — Virgin Is. (St. Thomas, St. Croix, St. John), Puerto Rico, Antigua.

Genus BAHAMIA Sabrosky

BAHAMIA Sabrosky, 1956:219 (unavailable, no type-species); 1957:52. Type-species, *longiceps* Sabrosky (orig. des.).

LONGICEPS Sabrosky, 1957:52. Type-locality: Bahamas, Andros Is. Distr. — Bahamas.

Genus ASTIOSOMA Duda

ASTIOSOMA Duda, 1927:119, 127. Type-species, *rufifrons* Duda (orig. des., as gen. n., sp. n.).

Unnamed sp. Distr. — Puerto Rico, San Juan

Genus SIGALOESSA Loew

SIGALOESSA Loew, 1866:186 (Cent. 6, no. 100). Type-species, *bicolor* Loew (mon.).

92.1

DEPARTAMENTO DE ZOOLOGIA
BIBLIOTECA

- Crepidohamma* Enderlein, 1915:185. Type-species, *brasiliense* Enderlein (orig. des.).
- BICOLOR** Loew, 1866:186 (Cent. 6, no. 100). Type-locality: "Cuba". Distr. — Uncertain, a species complex ranging from Illinois and Texas to s. Brazil.
- BLANTONI** Sabrosky, 1957:55. Type-locality: Panama, Darien, Jaqué. Distr. — El Salvador, Panama, Antigua.
- BRASILIENSIS** (Enderlein), 1915:186 (*Crepidohamma*). Type-locality: Brazil, Santa Catarina. Unrecognized.
- CINCTIPES** (Sabrosky), 1943:509 (*Crepidohamma*). Type-locality: Panama, Gatun Lake, Cano Saddle. Distr. — Panama.
- FLAVIFRONS** Sabrosky, 1957:56. Type-locality: U.S.A., Florida, Miami. Distr. — U.S.A. (Florida), Bahamas.
- FRONTALIS** Aldrich, 1915:98. Type-locality: Peru, Lima. Distr. — Panama (Canal Zone), Ecuador, Peru.
- HILLIFERA** Sabrosky, 1957:55. Type-locality: Costa Rica, San Mateo, Higuító. Distr. — Costa Rica.
- NIGRA** Sabrosky, 1957:55. Type-locality: Argentina, Tucumán, La Toma, Quebrada. Distr. — Argentina (Tucumán).
- NIGRIFRONS** Sabrosky, 1957:56. Type-locality: Costa Rica, San José. Distr. — Costa Rica.
- PSEUDOCINCTIPES** (Sabrosky), 1943:510 (*Crepidohamma*). Type-locality: Costa Rica, San Mateo, Higuító. Distr. — Costa Rica, Panama (Canal Zone), Ecuador.
- RICA** Curran, 1934:328 (n. name for *insularis* Curran, 1931). Type-locality: Puerto Rico, Vieques Is., Puerto Real. Distr. — Puerto Rico.
- insularis* Curran, 1931:13 (preoce. Malloch, 1930). Type-locality: Puerto Rico, Vieques Is., Puerto Real.
- SCHULDI** Sabrosky, 1957:54. Type-locality: Costa Rica, San Mateo, Higuító. Distr. — Costa Rica.

Genus TUCUMYIA Sabrosky

- TUCUMYIA** Sabrosky, 1956:219, 240; 1957:58. Type-species, *aczei* Sabrosky (orig. des.).
- ACZEI** Sabrosky, 1956:240; 1957:59. Type-locality: Argentina, Tucumán, Quebrada. Distr. — Argentina (Tucumán).
- ANGUSTIGENA** Sabrosky, 1957:59. Type-locality: Argentina, Tucumán, Burruyacu. Distr. — Argentina (Tucumán).
- LINEARIS** Sabrosky, 1957:59. Type-locality: Panama, Colón, Cativa. Distr. — Panama, Trinidad (?).
- POLLINOSA** Sabrosky, 1957:59. Type-locality: Argentina, Tucumán, Quebrada. Distr. — Brazil, (Santa Catarina), Argentina (Tucumán).

Subfamily ASTEINAE

Genus ASTEIA Meigen

- ASTEIA** Meigen, 1830:88. Type-species, *amoena* Meigen (Westwood, 1840:152).
- Asteimya* Sabrosky, 1943:505. Type-species, *spinosa* Sabrosky (orig. des.).



- ALBOVARIA Aldrich, 1915:95. Type-locality: "Peru". Distr. — Peru.
- ANTENNATA (Sabrosky), 1943:506 (*Asteimyia*). Type-locality: Costa Rica, San Mateo, Higuito. Distr. — Costa Rica, Panama (Canal Zone).
- ARGENTINICA Sabrosky, 1957:46. Type-locality: Argentina, Formosa, Clorinda. Distr. — Argentina (Formosa, Chaco).
- EXPANSA Sabrosky, 1957:46. Type-locality: Costa Rica, San José. Distr. — U.S.A. (Florida), Haiti, Costa Rica, Colombia, Ecuador.
- FASCIATA Sabrosky, 1957:44. Type-locality: El Salvador, Santa Tecla. Distr. — El Salvador.
- LINEIFACIES Sabrosky, 1957:46. Type-locality: Argentina, Tucumán, Quebrada. Distr. — Argentina (Tucumán).
- PLAUMANNI Sabrosky, 1957:46. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Santa Catarina).
- SEMILUNATA Sabrosky, 1957:46. Type-locality: Venezuela, Merida, Tovar. Distr. — Venezuela.
- SPINOSA (Sabrosky), 1943:505 (*Asteimyia*). Type-locality: Costa Rica, San Mateo, Higuito. Distr. — Costa Rica.
- STRIATIFRONS Malloch, 1930:323. Type-locality: Peru, Chanchamayo, Colonia Perenc. Distr. — Peru.

Genus LOEWIMYIA Sabrosky

- LOEWIMYIA Sabrosky, 1943:503. Type-species, *bifurcata* Sabrosky (orig. des.).
- BIFURCATA Sabrosky, 1943:504. Type-locality: Panama, Sabanas. Distr. — Panama, Ecuador.

LITERATURE CITED

ALDRICH, J. M.

1915. New American species of *Asteia* and *Sigaloesa* [sic]. *Psyche* 22:94-98, 2 figs. [Corrected to *Sigaloessa* in reprint].

CURRAN, C. H.

1931. First Supplement to the 'Diptera of Porto Rico and the Virgin Islands.' *Amer. Mus. Nov.* 456:1-23, 4 figs.
1934. *The families and genera of North American Diptera*. 512 pp., 235 figs., 2 pls. The Author, New York.

DUDA, O.


1927. Revision der altweltischen Astiidae (Dipt.). *Deut. ent. Ztschr.* (1927):113-147, 3 figs., 2 pls.

ENDERLEIN, G.

1915. Dipterologische Studien XV. *Crepidohamma*, eine neue Drosophilidengattung aus Brasilien. *Wien. Ent. Ztg.* 34:185-186, 1 fig.

LOEW, H.

1866. *Diptera Americae septentrionalis indigena. Centuria sexta.* *Berlin. ent. Ztschr.* 9(1865):127-186. [Also published separately,



1872, with Centuriae 6-10 bound together as *Diptera Americae septentrionalis indigena*, Vol. 2:300 ppl.

MALLOCH, J. R.

1930. Exotic Muscaridae (Diptera). — XXX. *Ann. & Mag. Nat. Hist.* (10)6:321-334, 3 figs.

MEIGEN, J. W.

1830. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten* 6(4):401 pp., pls. 55-66. Hamm.

SABROSKY, C. W.

1943. New genera and species of Asteiidae (Diptera), with a review of the family in the Americas. *Ann. Ent. Soc. Amer.* 36:501-514, 5 figs.

1956. Additions to the knowledge of Old World Asteiidae (Diptera). *Rev. Franc. d'Ent.* 23:216-243, 9 figs.

1957. Synopsis of the New World species of the Dipterous family Asteiidae. *Ann. Ent. Soc. Amer.* 50:43-61, 15 figs.

WESTWOOD, J. O.

1838-1840. *An introduction to the modern classification of insects. Synopsis of the genera of British insects.* 158 pp., London. [Diptera, 1840, pp. 125-158].


INDEX

aczeli Sabrosky, 1956, Tucumyia	92.2
albovaria Aldrich, 1915, Asteia	92.3
angustigena Sabrosky, 1957, Tucumyia	92.2
antennata (Sabrosky), 1943 (Asteimyia), Asteia	92.3
argentinica Sabrosky, 1957, Asteia	92.3
Asteia Meigen, 1830	92.2
Asteiinae	92.2
Asteimyia Sabrosky, 1943	92.2
Astiosoma Dura, 1927	92.1
Bahamia Sabrosky, 1957	92.1
bicolor Loew, 1866, Sigaloessa	92.2
bifurcata Sabrosky, 1943, Locwimyia	92.3
blantoni Sabrosky, 1957, Sigaloessa	92.2
brasilensis (Enderlein), 1915 (Crepidohamma), Sigaloessa	92.2
cinctipes (Sabrosky), 1943 (Crepidohamma), Sigaloessa	92.2
Crepidohamma Enderlein, 1915	92.2
expansa Sabrosky, 1957, Asteia	92.3
fasciata Sabrosky, 1957, Asteia	92.3
flavifrons Sabrosky, 1957, Sigaloessa	92.2
frontalis Aldrich, 1915, Sigaloessa	92.2
hillifera Sabrosky, 1957, Sigaloessa	92.2





<i>insularis</i> Curran, 1931, Sigaloessa	92.2
<i>linearis</i> Sabrosky, 1957, Tucumyia	92.2
<i>lineifacies</i> Sabrosky, 1957, Asteia	92.3
<i>Loewimyia</i> Sabrosky, 1943	92.3
<i>longiceps</i> Sabrosky, 1957, Bahamia	92.1
<i>nigra</i> Sabrosky, 1957, Sigaloessa	92.2
<i>nigrifrons</i> Sabrosky, 1957, Sigaloessa	92.2
<i>peculiaris</i> Sabrosky, 1957, Phlebosotera	92.1
<i>Phlebosotera</i> Duda, 1927	92.1
<i>plaumanni</i> Sabrosky, 1957, Asteia	92.3
<i>pollinosa</i> Sabrosky, 1957, Tucumyia	92.2
<i>pseudocinctipes</i> (Sabrosky), 1943 (Crepidohamma), Sigaloessa ..	92.2
<i>rica</i> Curran, 1934, Sigaloessa	92.2
<i>schildi</i> Sabrosky, 1957, Sigaloessa	92.2
<i>scmilunata</i> Sabrosky, 1957, Asteia	92.3
<i>Sigaloessa</i> Loew, 1866	92.1
<i>Sigaloessinae</i>	92.1
<i>spinosa</i> (Sabrosky), 1943 (Asteimyia), Asteia	92.3
<i>striatifrons</i> Malloch, 1930, Asteia	92.3
<i>Tucumyia</i> Sabrosky, 1956	92.2





SciELO





SciELO

MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO
(Formerly Departamento de Zoologia, Secretaria da Agricultura, São Paulo)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

93

FAMILY CRYPTOCHETIDAE

ANGELO PIRES DO PRADO

*Departamento de Zoologia, Setor de Parasitologia
Universidade Estadual de Campinas, São Paulo, Brasil*

One Australian species of *Cryptochetum* Rondani has been successfully introduced in Chile, via California. This is one of the natural enemies of the cottony-cushion scale, *Pericleria purchasi* (Maskell) (Homoptera, Margarodidae).

Ref. — Graf & Cortés, 1940:352.

Genus CRYPTOCHETUM Rondani

CRYPTOCHETUM Rondani, 1875:172 (as *Cryptochoetum*, p. 167). Type-species, *grandicorne* Rondani (orig. des.).

Cryptochaetum, emend.

Lestophonus Williston, 1888:21. Type-species, *iccryae* Williston (mon.).

ICERYAE (Williston), 1888:21 (*Lestophonus*). Type-locality: Australia, Crawford of Adelaide. Distr. — Australia; U.S.A. (California; introduced); Chile (introduced).

LITERATURE CITED

GRAF M., A. & R. CORTÉS P.


1940. Introducción de hiperparásitos en Chile: Resumen de las importaciones hechas y de sus resultados. *Proc. VI Pacif. Sci. Congr.* 4 (1939) : 351-353.

RONDANI, C.

1875. Species Italicae ordines Dipteriorum (Muscaria Rndn.) collectae et observatae. Stirps XXIII. Agromyzinae. *Bol. Soc. Ent. Ital.* 7:166-191.

WILLISTON, S. W.

1888. An Australian parasite of *Icerya purchasi*. *Insect Life* 1: 21-22, 1 fig.



PAPAVERO, N.

1964. Sobre o gênero *Somatia* Schiner, 1868 (Diptera, Somatiidae).
Papéis Avulsos Dept. Zool. Secr. Agr., São Paulo, Brazil 16:57-61.

SCHINER, I. R.

1868. Diptera, in B. von Wüllersdorff-Urbair: *Reise der österreichischen Fregatte Novara*. Zool. 2(1B):1-388 pp., pls. 1-4. Wien.

STEYSKAL, G. C.

1958. The genus *Somatia* Schiner (Diptera, Somatiidae). *Rev. Brasil. Ent.* 8:69-74.
1968. The genus *Somatia* Schiner, II. (Diptera: Somatiidae). *Proc. Ent. Soc. Wash.* 70:255-257.

Index

<i>aestiva</i> (Fabricius), 1805 (<i>Tephritis</i>), <i>Somatia</i>	94.1
<i>australis</i> Steyskal, 1958, <i>Somatia</i>	94.1
<i>carrerae</i> Papavero, 1964, <i>Somatia</i>	94.1
<i>lanei</i> Papavero, 1964, <i>Somatia</i>	94.1
<i>papaveri</i> Steyskal, 1968, <i>Somatia</i>	94.1
<i>schildi</i> Steyskal, 1968, <i>Somatia</i>	94.1
<i>Somatia</i> Schiner, 1868	94.1
<i>sophiston</i> Steyskal, 1958, <i>Somatia</i>	94.1
<i>xanthomelas</i> Schiner, 1868, <i>Somatia</i>	94.1



MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO
(FORMERLY DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS SOUTH OF THE UNITED STATES

96a

FAMILY ANTHOMYIIDAE

ADRIAN C. PONT

British Museum (Natural History), London, England

The family Anthomyiidae contains species that are generally small in size, drab in appearance and retiring in habit. The greatest development of the family has taken place in the Holarctic region, and in South America, as in Africa, the number of species is comparatively small. Most of the species are restricted to the temperate southern areas and to the mountain ranges and are clearly comparatively recent derivatives from elements of the northern hemisphere, with little endemism at the generic level.

Adults generally feed on nectar and pollen, but a few genera contain predaceous species. Larvae are mainly phytophagous, and the family contains a number of species of considerable economic importance such as Beet Fly, Cabbage-Root Fly, Onion Fly, etc. A few larvae are saprophagous or coprophagous.

There is no modern revisionary work on Neotropical Anthomyiidae, either at the generic or the specific level, and this catalogue can only be regarded as a provisional step, as a compilation of the available names. For this reason, the genera are listed alphabetically. A number of types has been examined, but it is not usually possible to be certain of generic assignments in the Anthomyiidae without studying the male genitalia, and types are better dissected in the context of revisionary work rather than as a matter of routine.

The catalogues of Anthomyiidae and Muscidae have been prepared simultaneously, and the bibliography contains all taxonomic and faunistic papers that have been consulted during the preparation of this catalogue.

I am grateful to the following colleagues for the loan of material from the collections in their charge: Dr. A. Kaltenbach (Vienna), Dr. R. Krause (Dresden), Dr. C. Leonardi (Milan), Prof. U. Parenti (Turin), and Dr. H. Schumann (Berlin). I am grateful to Professor H. C. Hockett and Mr. D. M. Ackland for advice on the generic assignment of a number of species, and also to Mr. Ackland for a great deal of help and discussion of South American Anthomyiidae.

Genus ANTHOMYIA Meigen

ANTHOMYIA Meigen, 1803:281. Type-species, *Musca pluvialis* Linnaeus (Westwood, 1840:143).

OCULIFERA Bigot, 1885:299. Type-locality: United States, Baltimore. Distr. — Venezuela; Nearctic region.

96a.1

PLUVIALIS (Linnaeus), 1758:597 (*Musca*). Type-locality: "Europe". Distr. — Mexico; Nearctic and Palaearectic regions.

Genus CALYTHEA Schnabl & Dziedzicki

Pegomyia, subg. CALYTHEA Schnabl & Dziedzicki, 1911:274 (1911:59). Type-species, *Musca albicincta* Fallén (mon.) = *nigricans* (Robineau-Desvoidy).

COMIS (Stein), 1911:155 (*Anthomyia*). Type-locality: Chile, Tacna. Distr. — Brazil, Argentina, Chile.

COSTANA (Séguy), 1934:9, fig. 1 (*Chortophila*). Type-locality: Argentina, La Plata. Distr. — Argentina.

CRENATA (Bigot), 1885:282 (*Trichopticus*). Type-locality: "Mexico". Distr. — Mexico, Puerto Rico, Colombia, Ecuador, Peru.

mexicana Giglio-Tos, 1893:10; 1896:28 (*Lasiops*). Type-locality: Mexico, Orizaba.

MICROPTERYX (Thomson), 1869:555 (*Anthomyia*). Type-locality: United States, California. Distr. — Virgin Islands; Nearctic region.

NIGRICANS (Robineau-Desvoidy), 1830:584 (*Anthomyia*). Type-locality: France, Saint-Sauveur. Distr. — Jamaica, Venezuela, Brazil, Bolivia; Europe, ?Asia, ?North Africa. This species is erroneously recorded from the New World, but it is not known to which species the above records refer.

albicincta Fallén, 1825:73 (*Musca*; preocc. Rossi, 1794). Type-locality: Sweden, Scania, Westrogothia and Ostrogothia.

Genus CRASPEDOCHAETA Macquart

CRASPEDOCHAETA Macquart, 1851a:241; 1851b:268 (as *Craspedochaeta*). Type-species, *Anthomyia punctipennis* Wiedemann (mon.).

AENEA (Malloch), 1934:186 (*Hylemyia*). Type-locality: Argentina, Lake Correntoso. Distr. — Argentina. N. comb.

ALBILAMELLATA (Stein), 1911:154 (*Chortophila*). Type-locality: Peru, Lake Titicaca, and Bolivia, Island in Lake Titicaca. Distr. — Peru, Bolivia. N. comb.

ALBOSTRIATA (Wulp), 1883:46 (*Chortophila*). Type-locality: "Argentina". Distr. — Peru, Venezuela, Argentina. N. comb.

CHILENSIS (Macquart), 1851a:238, pl. 24, fig. 9; 1851b:265, pl. 24, fig. 9 (*Chortophila*). Type-locality: Chile, Coquimbó. Distr. — Chile. N. comb.

CHIROSUNA Hennig, 1969:3, figs. 1-4 and 6-7. Type-locality: Chile, Andes slopes on the Santiago-Farellones road. Distr. — Chile.

CITREBASIS (Malloch), 1934:186, fig. 24d (*Hylemyia*). Type-locality: Argentina, River Pichilcufu. Distr. — Argentina. N. comb. *citribasis*, error.


CONFUSA (Albuquerque), 1959b:19, figs. 30-36 [*Hylemyia* (*Craspedochaeta*)]. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil. N. comb.



- EDWARDSI (Malloch), 1934:185, fig. 24a-c (*Hylemyia*). Type-locality: Argentina, Lake Nahuel Huapi. Distr. — Argentina. N. comb.
- LIMBINERVIS (Macquart), 1843a:326, pl. 22, fig. 2; 1843b:169, pl. 22, fig. 2 (*Chortophila*). Type-locality: "Chile". Distr. — Argentina, Chile, Juan Fernandez Islands.
- chilensis* Jaennicke, 1867a:373; 1867b:65 (*Anthomyia*; preocc. Macquart, 1851). Type-locality: "Chile".
- LINDIGI (Schiner), 1868:298 (*Anthomyia*). Type-locality: "South America". Distr. — Colombia, Peru, Brazil.
- lindgi*, error.
- lingindii*, error.
- MACULIPENNIS (Stein), 1918:241 (*Pegomyia*). Type-locality: Argentina, Tucuman. Distr. — Argentina. N. comb.
- PALLIDITHORAX (Albuquerque), 1959b:35, fig. 50 [*Hylemyia* (*Craspedochaeta*)]. Type-locality: Brazil, Rio de Janeiro, Pctrópolis, Alto da Mosela, Le Vallon. Distr. — Brazil. N. comb.
- PLURIPUNCTATA (Albuquerque), 1959b:30, figs. 43-49 [*Hylemyia* (*Craspedochaeta*)]. Type-locality: Brazil, São Paulo, Eugenio Lefevre. Distr. — Brazil. N. comb.
- PULLULA (Zetterstedt), 1845:1449 (*Aricia*). Type-locality: "In Scandinavia meridionali et media, Scania, Ostrogothia, Dania". Distr. — Mexico; Nearctic and Palearctic regions.
- PUNCTIPENNIS (Wiedemann), 1830:435 (*Anthomyia*). Type-locality: Uruguay, Montevideo. Distr. — Bolivia, Brazil, Uruguay, Argentina; South Africa.
- punctipennis* Shannon & Del Ponte, 1926:570 (*Hylemyia*; preocc. Wiedemann, 1830). Type-locality: Argentina, San Isidro.
- RECURRENTS (Malloch), 1934:180 (*Hylemyia*). Type-locality: Argentina, Bariloche. Distr. — Argentina. N. comb.
- SPILOTA (Wulp), 1883:46 (*Chortophila*). Type-locality: "Argentina". Distr. — Argentina. N. comb.
- VIDEO (Gaminara), 1930:1252, fig. 7; 1931:1251, fig. 7 (*Hylemyia*). Type-locality: Uruguay, Montevideo, Pocitos and Malvín. Distr. — Uruguay. N. comb.
- VIRGATA (Stein), 1911:153 (*Chortophila*). Type-locality: Peru, Sicuani, Cuzco and Tarma; Bolivia, Sorata; Chile, Santiago, Coquimbo and Guayacán. Distr. — Peru, Bolivia, Argentina, Chile. N. comb.
- VITIFERA (Stein), 1911:151 (*Pegomyia*). Type-locality: Peru, Lake Titicaca, Sicuani, Cuzco and Tarma; Bolivia, Lake Titicaca; Chile, Palca and Talca. Distr. — Peru, Bolivia, Chile. N. comb.
- XANTHOPYGA (Albuquerque), 1959b:39 [*Hylemyia* (*Craspedochaeta*)]. Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil. N. comb.

Genus DELIA Robineau-Desvoidy

- DELIA Robineau-Desvoidy, 1830:571. Type-species, *floricola* Robineau-Desvoidy (Coquillett, 1910:531) = *cardui* (Meigen).



- ANTIQUA (Meigen), 1826:166 (*Anthomyia*). Type-locality: "Germany". Distr. — Colombia, Brazil; Nearctic region, Europe.
- FLORILEGA (Zetterstedt), 1845:1555 (*Aricia*). Type-locality: "Sweden and Norway". Distr. — ?Argentina; Nearctic region, Europe. Probably an erroneous record.
- liturata* Meigen, 1838:329 (*Anthomyia*; preocc. Robineau-Desvoidy, 1830). Type-locality: "Germany".
- trichodactyla* Rondani, 1866:164 (*Chorthophila*). Type-locality: "Italy".
- LATIFRONS (Stein), 1918:236 (*Hylemyia*). Type-locality: Mexico, Maltrata. Distr. — Mexico. N. COMB.
- PLATURA (Meigen), 1826:171 (*Anthomyia*). Type-locality: "Germany". Distr. — Bermuda, Mexico, Venezuela, Colombia, Peru, Bolivia, Brazil, Uruguay, Argentina, Chile; cosmopolitan.
- diversa* Wiedemann, 1830:436 (*Anthomyia*). Type-locality: Uruguay, Montevideo.
- cana* Macquart, 1835:340 (*Anthomyia*). Type-locality: "North France".
- fusciceps* Zetterstedt, 1845:1552 (*Aricia*). Type-locality: "Denmark".
- fuscipes*, error.
- corella* Walker, 1849:953 (*Anthomyia*). Type-locality: Uruguay, Montevideo.
- cilicrura* Rondani, 1866:165 (*Chorthophila*). Type-locality: "Italy".
- lanicrus* Thomson, 1869:556 (*Anthomyia*). Type-locality: Argentina, Buenos Aires.
- rubrifrons* Bigot, 1885:279 (*Chorthophila*). Type-locality: Argentina, Buenos Aires. N. SYN.
- rubifrons* Bigot, 1885:297 (?*Anthomyia*). Type-locality: "Chile".
- rufifrons*, error.
- trifilis* Giglio-Tos, 1893:10; 1896:28 (*Anthomyia*). Type-locality: Mexico, Coscomatepec]. N. SYN.
- ceparum* (Meigen) of Johnson, 1904:79. Misidentification.
- PRISCA (Wulp), 1896:340 (*Phorbia*). Type-locality: Mexico, Ciudad in Durango, Pátzcuaro and Jalisco. Distr. — Mexico. N. COMB.
- SANCTIJACOBI (Bigot), 1885:296 (*Anthomyia*). Type-locality: "Chile". Distr. — Uruguay, Argentina, Chile, Juan Fernandez Islands.
- sanctojacobi*, error.
- SEGMENTATA (Wulp), 1896:336 (*Anthomyia*). Type-locality: Mexico, Guerrero, Omilteme and Sierra de las Aguas Escondidas. Distr. — Mexico. N. COMB.

Genus EMMESOMYIA Malloch

- EMMESOMYIA Malloch, 1917b:114. Type-species, *unica* Malloch (orig. des.) = *socialis* (Stein).
- ARGENTINA Snyder, 1957b:443. Type-locality: Argentina, Tucumán, Queb. La Toma. Distr. — Argentina.
- FLAVIPALPIS (Wulp), 1896:334 (*Hydrophoria*). Type-locality: Mexico, Guerrero, Venta de Zopilote, Chilpancingo, Amula and Sierra de las Aguas Escondidas. Distr. — Mexico.



- SEMPIPELLUCIDA (Stein), 1918:238 (*Taeniomymia*). Type-locality: Mexico, Orizaba and Tapachula. Distr. — Mexico. N. comb.
- SPADIBASIS Snyder, 1957b:442. Type-locality: Argentina, Tucumán, Queb. La Toma. Distr. — Argentina.

Genus ERIOSCHIA Lioy

- ERIOSCHIA Lioy, 1864:991. Type-species, *Chortophila floccosa* Macquart (mon.) = *brassicae* (Bouché).
- BRASSICAE (Bouché), 1834:74, pl. 5, fig. 34 (*Anthomyia*). Type-locality: not stated, presumably Germany. Distr. — Brazil; Nearctic and Palaearctic regions.

Genus FUCELLIA Robineau-Desvoidy

- FUCELLIA Robineau-Desvoidy, 1842:269. Type-species, *arenaria* Robineau-Desvoidy (mon.) = ?*maritima* (Haliday).
- CALCOERATA (Macquart), 1851a:246, pl. 25, fig. 5; 1851b:273, pl. 25, fig. 5 (*Scatophaga*). Type-locality: "Chile". Distr. — Peru, Chile, *calcarata*, error.
- ruficoxa* Bigot, 1885:280 (*Parachortophila*). Type-locality: "Chile".
- funifera* Stein, 1910a:82. Type-locality: Peru, Barranca, and Chile, Caldera.
- fumifera*, error.
- PACIFICA Malloch, 1923:427 [ssp. of *maritima* (Haliday)]. Type-locality: Gulf of California, San Esteban Island. Distr. — Mexico, Gulf of California Islands; Nearctic region.
- REJECTA Aldrich, 1918:171. Type-locality: United States, California, Ocean Beach near San Diego. Distr. — Mexico, Gulf of California Islands; Nearctic region.
- TERGINA (Zetterstedt), 1845:1690 (*Anthomyza*). Type-locality: Sweden, Scania. Distr. — Bermuda, Puerto Rico, Peru, Argentina, Chile, Juan Fernandez Islands; Nearctic and western Palaearctic regions, and Atlantic Ocean islands.
- fucorum* Fallén, 1819:5 (*Scatomyza*), of South American authors [misidentification]. Type-locality: "Sweden".
- maritima* Haliday, 1838:186 [*Scatophaga* (*Halithea*)], of New World authors [misidentification]. Type-locality: "British Isles".
- marina* Macquart, 1838:424, pl. 11, fig. 3 (*Scatophaga*), of South American authors [misidentification]. Type-locality: Belgium, Dunkerque.
- intermedia* Lundbeck, 1901:291, fig. 1b. Type-locality: Greenland, west coast, several localities.
- masafuerana* Enderlein, 1940:647 (*Thrixina*). Type-locality: Juan Fernandez Islands, Masafuera Island.
- masatierrana* Enderlein, 1940:648 (*Thrixina*). Type-locality: Juan Fernandez Islands, Masatierra Island.

Genus HYDROPHORIA Robineau-Desvoidy

- HYDROPHORIA Robineau-Desvoidy, 1830:503. Type-species, *littoralis* Robineau-Desvoidy (Coquillett, 1910:554) = *conica* (Wiedemann).
PLUMOSA Wulp, 1896:332. Type-locality: Mexico, Amula in Guerrero. Distr. — Mexico, Venezuela.
RURALIS (Meigen), 1826:101 (*Anthomyia*). Type-locality: "Germany". Distr. — Mexico, Bolivia; Nearctic and Palaearctic regions. Probably an erroneous record.

Genus HYLEMYA Robineau-Desvoidy


- HYLEMYA Robineau-Desvoidy, 1830:550. Type-species, *strenua* Robineau-Desvoidy (Rondani, 1860:74 and 184).
Hylemyia, emendation.
URBICA Wulp, 1896:338. Type-locality: Mexico, Mexico city. Distr. — Mexico.

Genus HYLEMYIOIDE Albuquerque

- HYLEMYIOIDE Albuquerque, 1949:419. Type-species, *brasiliensis* Albuquerque (orig. des.).
Hylemyioides, error or emendation.
Hylemyoides, error or emendation.
AURIFACIES Albuquerque, 1952:27, figs. 6-10. Type-locality: Brazil, São Paulo, São José dos Campos. Distr. — Brazil.
BRASILIENSIS Albuquerque, 1949:419, figs. 1-11. Type-locality: Brazil, São Paulo, São José dos Campos. Distr. — Brazil.
CRASSINERVIS Albuquerque, 1959a:1, figs. 1-8. Type-locality: Brazil, Rio de Janeiro, Le Vallon, Alto da Mosela. Distr. — Brazil.
LATIMACULATA Albuquerque, 1959a:6, figs. 9-15. Type-locality: Brazil, Rio de Janeiro, Petrópolis, Le Vallon, Alto da Mosela. Distr. — Brazil.
PAULISTENSIS Albuquerque, 1952:29, figs. 11-19. Type-locality: Brazil, São Paulo. Distr. — Brazil.
PLURINERVIS Albuquerque, 1958:341, figs. 1-6. Type-locality: Brazil, Petrópolis, Rio de Janeiro, Le Vallon, Alto da Mosela. Distr. — Brazil.

Genus LEUCOPHORA Robineau-Desvoidy

- LEUCOPHORA Robineau-Desvoidy, 1830:562. Type-species, *cinerea* Robineau-Desvoidy (Coquillett, 1901:138).
Hammomyia Rondani, 1877:13. Type-species, *Aricia albescens* Zetterstedt (orig. des.) = *cinerea* Robineau-Desvoidy.
Ammomyia, error.
Hylephila Rondani, 1877:13 (preocc. Billberg, 1820). Type-species, *Musca buccata* Fallén (orig. des.) = *sericca* Robineau-Desvoidy.



- ANDICOLA (Bigot), 1885:300 (*Hylemyia*). Type-locality: "Chile". Distr. — Argentina, Chile. N. comb.
- ARGENTINA (Malloch), 1934:193, fig. 27e-f (*Hammomyia*). Type-locality: Argentina, Mendoza Province. Distr. — Argentina. N. comb.
- CHILENSIS (Malloch), 1934:190, figs. 26 and 27a-b (*Hammomyia*). Type-locality: Argentina, Bariloche. Distr. — Argentina. N. comb.
- FLAVIPES (Stein), 1918:239 (*Ammomyia*). Type-locality: Peru, Piura. Distr. — Peru. N. comb.
- JOHNSONI (Stein), 1898:215 (*Hylemyia*). Type-locality: United States, Pennsylvania. Distr. — Bolivia; Nearctic region.
- MACULIPENNIS (Albuquerque), 1953:535, figs. 1-12 (*Hammomyia*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil. N. comb.
- PEULLAE (Malloch), 1934:192, fig. 27c-d (*Hammomyia*). Type-locality: Chile, Peulla. Distr. — Chile. N. comb.
- PLUMISETA (Malloch), 1934:194 (*Hammomyia*). Type-locality: Peru, Lima, Verrugas Canyon. Distr. — Peru. N. comb.
- RUFITIBIA (Stein), 1918:239 (*Ammomyia*). Type-locality: Peru, Piura. Distr. — Peru. N. comb.
- SUBVITTATA (Malloch), 1934:192, fig. 27g (*Hammomyia*). Type-locality: Argentina, Bariloche. Distr. — Argentina, Chile. N. comb.

Genus MYOPELLA Malloch


- MYOPELLA Malloch, 1934:196. Type-species, *Myopina appendiculata* Stein (orig. des.).
- ANGUSTIFRONS Malloch, 1934:193, fig. 31. Type-locality: Argentina, Lake Correntoso. Distr. — Argentina.
- APPENDICULATA (Stein), 1904:473 (*Myopina*). Type-locality: Peru, Callanga. Distr. — Peru, Bolivia, Argentina, Tierra del Fuego.

Genus NEOHYLEMYIA Malloch

- NEOHYLEMYIA Malloch, 1917a:37. Type-species, *proboscidalis* Malloch (orig. des.).
- NIGRICAUDA (Bigot), 1885:276 (*Hydrophoria*). Type-locality: United States, Rocky Mountains. Distr. — Nearctic region. Possibly a Mexican species and not Nearctic.
- TENUIROSTRIS (Wulp), 1896:337, pl. 8, figs. 15 and 15a (*Hylemyia*). Type-locality: Mexico, Guerrero, Tepetlapa and Amula. Distr. — Mexico. N. comb.

Genus PEGOMYA Robineau-Desvoidy

- PEGOMYA Robineau-Desvoidy, 1830:598. Type-species, *Musca hyoscyami* Panzer (Coquillett, 1901:140).
- Pegomyia*, emendation.



- AFFINIS Stein, 1898:286. Type-locality: United States, Louisiana. Distr. — Mexico; Nearctic region.
- BELLA Stein, 1911:152. Type-locality: Chile, Palca. Distr. — Chile.
- BRUCHI (Shannon & Del Ponte), 1926:571 (*Hylemyia*). Type-locality: Argentina, Buenos Aires Province. Distr. — Brazil, Uruguay, Argentina.
- poeciloptera* (Malloch) of Malloch, 1934:182. Misidentification.
- CARRERA Albuquerque, 1959b:6, figs. 9-13. Type-locality: Brazil, São Paulo, São Paulo (Ipiranga). Distr. — Brazil.
- poeciloptera* (Malloch) of Carrera & Travassos, 1947:49 ff. Misidentification.
- CINCTINERVIS Stein, 1918:239. Type-locality: Mexico, Orizaba and Cordoba. Distr. — Mexico.
- COGNATA Stein, 1920:67. Type-locality: United States, Stanford. Distr. — Guadalupe Island; Nearctic region.
- DOLOSA Stein, 1918:240. Type-locality: Mexico, Oaxaca. Distr. — Mexico.
- DORSIMACULATA (Wulp), 1896:336 (*Anthomyia*). Type-locality: Mexico, Omilteme in Guerrero. Distr. — Mexico.
- DUPLICOIDES Snyder, 1952:420. Type-locality: Mexico, Cuernavaca. Distr. — Mexico.
- ELONGATA (Wulp), 1896:335, pl. 8, figs. 14 and 14a (*Anthomyia*). Type-locality: Mexico, Sierra de las Aguas Escondidas in Guerrero. Distr. — Mexico. N. comb.
- HERNANDEZ Snyder in Olmedo, 1957a:126 and 129. Type-locality: Mexico, Chapingo. Distr. — Mexico.
- NEOPALPOSA Snyder, 1957b:444. Type-locality: Argentina, Tucumán, Queb. la Toma. Distr. — Argentina.
- neopalosa*, error.
- NERVICINCTA (Stein), 1911:154 (*Chortophila*). Type-locality: Bolivia, Sorata. Distr. — Bolivia.
- NIGRACAERULEA Snyder, 1952:418. Type-locality: Mexico, Amecameca. Distr. — Mexico.
- nigrocaerulea*, error.
- NIGRIFEMUR Stein, 1918:241. Type-locality: Mexico, mountains near Cuernavaca. Distr. — Mexico.
- NOTATA (Albuquerque), 1957a:119, figs. 1-8 (*Pegomyza*). Type-locality: Argentina, Tucumán, Sierra de San Xaxier. Distr. — Argentina.
- POECILOPTERA Malloch, 1921b:430. Type-locality: Argentina, La Plata city. Distr. — Brazil, Argentina.
- RUBRICEPS Hockett, 1965:861 (nom. nov. for *ruficeps* Stein, 1898). Type-locality: United States, Louisiana. Distr. — Mexico; Nearctic region.
- ruficeps* Stein, 1898:286 (preocc. Zetterstedt, 1838). Type-locality: North America, Louisiana.



- SPINISOIDES Snyder, 1952:417. Type-locality: Mexico, Cuernavaca. Distr. — Mexico.
- STLEMBIA (Giglio-Tos), 1893:10; 1896:29 (*Chortophila*). Type-locality: "Mexico". Distr. — Mexico.
- TARSADMSCA Snyder, 1952:415. Type-locality: Mexico, Cuernavaca. Distr. — Mexico.
- TRIFUMOSA Snyder, 1957b:445. Type-locality: Argentina, Jujuy, Termas de Reyes. Distr. — Argentina.
- VERTICALA Snyder, 1952:419. Type-locality: Mexico, Mexico City. Distr. — Mexico.

Genus PEGOPLATA Schnabl & Dziedzicki

- Pegomya, subg. PEGOPLATA Schnabl & Dziedzicki, 1911:108. Type-species, *Hydrophoria palposa* Stein (as *palpata* Stein, error) (Séguy, 1937:50).
- MERACULA (Giglio-Tos), 1893:9; 1896:24 (*Spilogaster*). Type-locality: "Mexico". Distr. — Mexico. N. comb.
- MONTICOLA (Wulp), 1896:338 (*Hylemyia*). Type-locality: Mexico, Guerrero, Sierra de las Aguas Escondidas. Distr. — Mexico. N. comb.
- RHODINA (Giglio-Tos), 1893:10; 1896:31 (*Hylemyia*). Type-locality: Mexico, Tuxpango. Distr. — Mexico. N. comb.

Genus PETRMYIA Albuquerque

- PETRMYIA Albuquerque, 1954:493. Type-species, *wygodzinskyi* Albuquerque (orig. des.).
- WYGODZINSKYI Albuquerque, 1954:493, figs. 1-10. Type-locality: Argentina, Tucumán. Distr. — Argentina.

Genus TAENIOMYIA Stein

- TAENIOMYIA Stein, 1918:237. Type-species, *auricollis* Stein (Séguy, 1937:141).
- AURICOLLIS Stein, 1918:237. Type-locality: Brazil, Rio Grande do Sul, Teresópolis. Distr. — Brazil.
- COLLARIS (Wulp), 1896:333, pl. 8, figs. 11 and 11a (*Hydrophoria*). Type-locality: Mexico, Guerrero, Amula, Xucumanatlán, Sierra de las Aguas Escondidas, Omilteme. Distr. — Mexico, Brazil.
- DEXIARIA (Stein), 1904:475 (*Hydrophoria*). Type-locality: "Colombian Cordillera". Distr. — Colombia.
- FLAVIBASIS (Stein), 1911:149 (*Hydrophoria*). Type-locality: Peru, Larcs Valley, and Bolivia, Lorenzopata. Distr. — Peru, Bolivia.
- PICTIPES (Bigot), 1885:275 (*Hydrophoria*). Type-locality: "Mexico". Distr. — Mexico.
- transversalis* Wulp, 1896:334, pl. 8, fig. 12 (*Hydrophoria*). Type-locality: Mexico, Guerrero, Xucumanatlán, Omilteme and Sierra de las Aguas Escondidas.



SCUTELLATA (Stein), 1911:149 (*Hydrophoria*). Type-locality: Bolivia, Sorata. Distr. — Bolivia. N. COMB.

TRIMACULATA (Stein), 1911:148 (*Hydrophoria*). Type-locality: Peru, Pachitea estuary, and Bolivia, Chimatc, San Carlos and Sarampiuni. Distr. — Peru, Bolivia. N. COMB.

SPLCIES OF DOUBTFUL GENERIC ASSIGNMENT

ABREPTA Giglio-Tos, 1893:10; 1896:30 (*Hylemyia*). Type-locality: "Mexico". Distr. — Mexico.

CHRYSOSTOMA Rondani, 1864:33 (*Anthomyia*). Type-locality: "Chile". Distr. — Chile.

DEVIA Cova Garcia, 1964: no page (*Pegomyia*). Type-locality: "Venezuela". Distr. — Venezuela. Cova Garcia attributes this name to Stein, but Stein described no Anthomyiid of this name.

LEUCOTELUS Walker, 1852:361 (*Anthomyia*). Type-locality: "Brazil". Distr. — Brazil. Possibly belongs to *Taeniomyia* Stein (see Stein, 1919: 150).

LEVIPES Giglio-Tos, 1893:10; 1896:30 (*Hylemyia*). Type-locality: "Mexico". Distr. — Mexico.

LITURATA Rondani, 1868:28 (*Chortophila*). Type-locality: Argentina, Buenos Aires. Distr. — Argentina.

NIGRIPES Robineau-Desvoidy, 1830:553 (*Hylemya*). Type-locality: French Guiana, Cayenne. Distr. — Guiana.

LITERATURE CITED

ACKLAND, D. M.

1967. Diptera from Nepal. Anthomyiidae. *Bull. Br. Mus. nat. Hist. (Ent.)*, 20 (4):105-139, 83 figs.

1968. The World Species of *Calythea* Schnabl and Dzied. (Dipt., Anthomyiidae) with notes on Bigot's Types. *Entomologist's mon. Mag.*, 104:135-144, 23 figs.

ALBUQUERQUE, D. DE O.

1949. Sobre um novo gênero neotrópico de "Anthomyiinae" (Diptera, Muscidae). *Rev. bras. Biol.* 9:419-423, 11 figs.

1952. Sobre "*Hylemyioide*" Albuquerque, 1949, com descrição de espécies novas (Diptera, Muscidae, Anthomyiinae). *Rev. bras. Biol.* 12:25-31, 19 figs.

1953. Contribuição ao conhecimento das espécies neotropicais de *Hammomyia* RDI. e *Calythea* Schnabl et Dziedzicki, com descrição de uma espécie nova (Diptera-Muscidae). *Anais Acad. bras. Ciênc.* 25:535-543, 19 figs.

1954. Um novo gênero e uma espécie nova de "Anthomyiinae" proveniente da Argentina (Dipt., Muscidae). *Rev. bras. Biol.* 14:493-496, 10 figs.



- 1957a. Uma espécie nova de "*Pegomyza*" Schnable [sic] & Dziedzicki, 1911 (Diptera, Muscidae). *Rev. bras. Biol.* 17:119-122, 8 figs.
- 1957b. Fauna do Distrito Federal, XLI. Notas sobre alguns Muscideos, com descrição de dois alótipos (Diptera-Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, No. 164:13 pp., 17 figs.
1958. Uma nova espécie de "*Hylemyioides*" Albuquerque, 1949 (Diptera, Muscidae, Anthomyiinae). *Rev. bras. Biol.* 18:341-344, 6 figs.
- 1959a. Contribuição ao conhecimento de *Hylemyioides* Albuquerque, 1949, com descrição de duas espécies novas (Diptera-Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, No. 194:12 pp., 18 figs.
- 1959b. Contribuição ao conhecimento de *Pegomyia* R.-D., 1830 e *Hylemyia* (*Craspedochaeta*) Macquart, 1850, na América do Sul, com descrições de espécies novas (Diptera-Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, No. 210:50 pp., 50 figs.
- ALDRICH, J. M.
1918. The Kelp-Flies of North America (Genus *Fucellia*, Family Anthomyidae). *Proc. Calif. Acad. Sci.* (4) 8:157-179, 10 figs.
- ANON.
1968. Catálogo de insectos en cultivos de importancia económica en Colombia. *Publ. Asoc. latinoam. Ent.* 1:156 pp.
- D'ARAÚJO E SILVA, A. G., ET AL.
1968. *Quarto catálogo dos insetos que vivem nas plantas do Brasil, seus parasitos e predadores. Parte II*, 1º Tomo. Min. Agric., Rio de Janeiro, xvi + 622 pp.
- ARNAUD, P. H.
1959. Records of Diptera from Guadalupe Island, Mexico. *Ent. News* 70:182-185.
1963. Records of Diptera from Guadalupe Island, Mexico. Second paper. *Ent. News* 74:117-129, 7 figs.
- BERG, C.
1899. Apuntes dipterológicos. *Comun. Mus. nac. B. Aires* 1 (4):124-130.
- BERRY, P. A., & VAQUERO, M. S.
1957. Lista de Insectos Clasificados de El Salvador. *Bol. técn. Min. Ganad. El Salvador* 21:134 pp.
- BIGOT, J.-M.-F.
1885. Diptères nouveaux ou peu connus. 25e partie. XXXIII. Anthomyzides nouvelles. *Ann. Soc. ent. Fr.* (6) 4 [1884]:263-304.
- BLANCHARD, E. E.
1933. Parasitos animales de la langosta. *Bol. Minist. Agric., B. Aires* 34:247-266, 5 pls.



BOUCHÉ, P. F.

1834. *Naturgeschichte der Insekten, besonders in Hinsicht ihrer ersten Zustände als Larven und Puppen. Erste Lieferung.* Berlin, v + 216 pp., 10 pls.

BRÉTHIES, J.

1908. Catálogo de los Dípteros de las Repúblicas del Plata. *An. Mus. nac. B. Aires* (3) 9:277-305.
1918. Quelques Díptères du Chili. *Rev. chil. Hist. nat.* 22:49-50.
1919. Cueillette d'Insectes au Río Blanco. III. Díptères. *Rev. chil. Hist. nat.* 23:40-44.

CARRERA, M., & TRAVASSOS, L.

1947. Dados morfológicos e bionômicos sobre *Hylemyia poeciloptera* (Malloch, 1921) (Diptera: Anthomyiidae), minadora das folhas de beterraba (*Beta vulgaris* L.). *Papéis Avulsos Dep. Zool. S. Paulo* 8:49-62, 12 figs., 2 pls.

COLLIN, J. E.

1921. A contribution towards the knowledge of the Anthomyid genera *Hammomyia* and *Hylephila* of Rondani (Diptera). *Trans. ent. Soc. Lond.* 53 [1920]:305-326, pls. 7-10.

COQUILLET, D. W.

1901. Types of Anthomyid Genera. *Jour. N. Y. ent. Soc.* 9:134-146.
1910. The type-species of the North American Genera of Diptera. *Proc. U. S. natn. Mus.* 37:499-647.

COVA GARCIA, P.

1964. *Moscas de Venezuela*, 17 unnumbered pages, 30 pls. Caracas.

CURRAN, C. H.

1928. Insects of Porto Rico and the Virgin Islands. Diptera or Two-winged Flies. In *Scientific Survey of Porto Rico and the Virgin Islands*, Vol. XI, part 1:118 pp., 39 figs. New York.

ENDERLEIN, G.

1940. *Die Dipterenfauna der Juan-Fernandez-Inseln und der Oster-Insel.* *Nat. Hist. Juan Fernandez and Easter Island*, III, Zoology, part 5:643-680, 49 figs. Uppsala.

ENGEL, E. O.

1931. Die Ausbeute der deutschen Chaco-Expedition 1925/26. Diptera. XXVI. Anthomyiidae, XXVII. Muscidae und XXVIII. Sarcophagidac. *Konowia* 10:133-154, pls. 1-6.

FALLÉN, C. F.

1819. *Scatomyzides Sueciae.* Lundae, 10 pp.
1825. *Monographia Muscidum Sueciae.* Part VIII, pages 73-80. Part IX, pages 81-94. Lundae.



GAMINARA, A.

1930. Clasificación de algunos Muscoideos Uruguayos (Muscidae y Calliphoridae). *An. Fac. Med. Univ. Montevideo* 14 [1929]:1235-1282, 19 figs.

1931. Clasificación de algunos Muscoideos Uruguayos (Muscidae y Calliphoridae). *Arch. Soc. Biol. Montev.*, Suppl. 5:1237-1280, 19 figs.

GAY, C.

1852. *Historia física y política de Chile. Zoología*, Vol. 7. Insectos. Paris and Santiago, 471 pp.

GIGLIO-TOS, E.

1893. Diagnosi di nuovi generi e di nuove specie di Ditteri. VIII. *Boll. Musei Zool. Anat. comp. R. Univ. Torino*, 7 (No. 147):11 pp.

1896. Ditteri del Messico. Parte Quarta. Muscidae calypteratae: Muscinae, Anthomyiinae. Muscidae acalypteratae: Scatophaginae, Helomyzinae, Tetanocerinae, Ortalinae, Ulidinae, Sapromyzinae, Trypetinae, Sepsinae, Tanypezinae, Psilinae, Chloropinae, Ephydrinae, Drosophilinae. *Memorie Accad. Sci. Torino* (2) 45:1-74, pl. 1.

GOWDEY, G. C.

1926. Catalogus Insectorum Jamaicensis. *Ent. Bull. Dep. Agric. Jamaica* 4 (1-2):114 + xiv and 10 + ii pp.

GUNDLACH, J.

1887. Apuntes para la Fauna Puerto-Riqueña. VIII. Insectos. *An. Soc. esp. Hist. nat.* 16:137-199.

HALIDAY, A. H.

1838. New British Insects indicated in Mr. Curtis' Guide. *Ann. nat. Hist.* 2:112-121 and 183-190.

HENNIG, W.

1955. Los Insectos de las Islas Juan Fernandez. 16. Phryneidae, Helomyzidae, Lonchaeidae, Piophilidae, Anthomyzidae und Muscidae (Diptera). *Rev. chil. Ent.* 4:21-34.

1957. Los Insectos de las Islas Juan Fernandez. 34. Threvidae, Helomyzidae, Piophilidae, Milchiidae und Muscidae (Diptera). (Nachtrag). *Rev. chil. Ent.* 5:409-412.


1966-. 63a, Anthomyiidae. In Lindner, E., *Fliegen palaearkt. Reg.* Stuttgart.

1969. Neue oder bisher ungedeutete Arten der Gattungen *Chirosta*, *Paraprosalpia* und *Craspedochaeta* (Diptera: Anthomyiidae). *Stuttg. Beitr. Naturk.* 208:12 pp., 19 figs.

JAENNICKÉ, F.

1867a. Neue exotische Dipteren. *Abh. senckenb. naturforsch. Ges.* 6: 311-408, pls. 43-44.

1867b. Ibidem, reprinted with pagination 3-99, 2 plates.



JAMES, M. T.

1948. The Flies that Cause Myiasis in Man. *Misc. Publ. U. S. Dep. Agric.* 631 [1947]:175 pp., 98 figs., 3 tables.

JOHNSON, C.

1904. A revised list of the Diptera of Bermuda. *Psyche*, Camb 11: 76-80.
1913. The Dipteran Fauna of Bermuda. *Ann. ent. Soc. Am.* 6:443-452, 2 figs.
1919. A Revised List of the Diptera of Jamaica. Phoridae by Charles T. Brues. *Bull. Am. Mus. nat. Hist.* 41 (8):421-449.

LAHILLE, F.

1907. La langosta y sus moscas parasitarias. *An. Minist. Agric. (Zool.)*, 3 (4):1-136, 29 figs., 6 pls.

LINNAEUS, C.

1758. *Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum caracteribus, differentiis, synonymis, locis.* 10th Edition. Volume 1. Holmiae, 824 pp.

LINS DE ALMEIDA, J.

1933. Nouveaux agents de transmission de la Berne [*Dermatobia hominis* (L. junior, 1781)] au Brésil. *C. r. Séanc. Soc. Biol.* 113, no. 27:1274-1275.

LIOY, P.

1864. I Ditteri distribuiti secondo un nuovo metodo di classificazione naturale. [Part]. *Atti R. Ist. veneto Sci.* (3) 9:989-1027.

LUNDBECK, W.

1901. Diptera groenlandica. *Vidensk. Medd. dansk. naturh. Foren.* (6), 2 [1900]:281-316, 5 figs.

MACQUART, J.

1835. *Histoire Naturelle des Insectes. Diptères, Tome deuxième.* Paris, 703 pp., 12 pls.
1838. Notice sur trois Diptères nouveaux des genres *Medeterus* et *Scatophaga*. *Ann. Soc. ent. Fr.* 7:421-424, pl. 11, figs. 1-3.
1843a. Diptères Exotiques nouveaux ou peu connus. Tome deuxième. 3e partie. *Mém. Soc. Sci. Agric. Lille* 1843:162-460, 36 pls.
1843b. Ibidem, reprinted with pagination 5-304.
1851a. Diptères Exotiques nouveaux ou peu connus. 4e Supplément (part). *Mém. Soc. Sci. Agric. Lille* 1850:134-294, pls. 15-28.
1851b. Ibidem, reprinted with pagination 161-336.

MALLOCH, J. R.

- 1917a. A new genus and species of Anthomyiidae (Diptera). *Bull. Brooklyn ent. Soc.* 12:37-38.

- 1917b. A new genus of Anthomyiidae (Diptera). *Bull. Brooklyn ent. Soc.* 12:113-115.
- 1921a. Notes on some of van der Wulp's Species of North American Anthomyiidae (Diptera). *Ent. News* 32:40-45.
- 1921b. Exotic Muscaridae (Diptera), II. *Ann. Mag. nat. Hist.* (9) 7: 420-431.
1923. Expedition of the California Academy of Sciences to the Gulf of California in 1921. Anthomyidae and Lonchaeidae (Kelp Flies and their allies). *Proc. Calif. Acad. Sci.* (4) 12:425-428.
1934. Muscidac. In *Diptera of Patagonia and South Chile, VII* (2): 171-346, figs. 22-60. London.
- MEIGEN, J. W.
1803. Versuch einer neuen Gattungseintheilung der europäischen zweiflügeligen Insekten. *Illig. Mag. Ins.* 2:259-281.
1826. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten.* 5. xii + 412 pp., pls. 42-54, Hamm.
1830. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten.* 6. xi + 401 pp., pls. 55-66, Hamm.
1838. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten.* 7. xi + 434 pp., pls. 67-74, Hamm.
- OCHILVIE, L.
1928. *The Insects of Bermuda.* Dept. of Agriculture, Bermuda, 52 pp.
- OLMEDO, I. H.
1957. Investigaciones sobre el Minador de la hoja del Huauzontle. *Revista Chapingo*, No. 64:117-131, 8 figs.
- OSTEN-SACKEN, C. R.
1877. Note on some Diptera from the Island Guadalupe (Pacific Ocean), collected by Mr. E. Palmer. *Proc. Boston Soc. nat. Hist.* 18:133-134.
- REED, E. C.
1888. Catalogo de los Insectos Dipteros de Chile. *An. Univ. Chile* 73: v + 46 pp.
- ROBERTS, R. A.
1934. Some Insects collected in Mexico, mostly in association with Man and Animals or animal Products. *Jour. N. Y. ent. Soc.* 42 (3): 249-262, 5 tables.
- ROBINEAU-DESVOIDY, A. J. B.
1830. Essai sur les Myodaires. *Mém. prés. div. Sav. Acad. Sci. Inst. Fr.*, 2:813 pp.
1842. Sur le genre Fucellie, *Fucellia*, R. D., et en particulier sur la *Fucellia arenaria*. *Ann. Soc. ent. Fr.* 10 [1841]:269-272.



RONDANI, C.

1864. Dipterorum species et genera aliqua exotica revisa et annotata novis nonnullis descriptis. *Arch. Zool. Anat. Fis.*, Modena, 3 [1863]: 1-99, pl. 5.
1866. Anthomyinae Italicae collectae distinctae et in ordinem dispositae. *Atti Soc. ital. Sci. nat.* 9:68-217.
1868. Diptera aliqua in America meridionali lecta a Prof. P. Strobel annis 1866-67. *Annuaire Soc. Nat. Modena* 3:24-40, pl. 4.
1877. *Dipterologiae Italicae Prodromus*. 7:304 pp., Parmae.

RUFFINELLI, A., & CARBONELL, C. S.

1954. Segunda lista de insectos y otros artrópodos de importancia económica en el Uruguay. *Rev. Asoc. Ing. agron.*, Montev., 94: 33-82.

SCHINER, J. R.

1868. Diptera. In *Reise der österreichischen Fregatte Novara. Zoologischer Theil*, Band II. vi + 388 pp., 4 plates, Wien.

SCHNABL, J., & DZIEDZICKI, H.

1911. Die Anthomyiden. *Nova Acta Acad. Caesar. Leop. Carol.* 95: 55-358, pls. 3-37.

SÉGUY, E.

1934. Étude sur Quelques Muscides de l'Amérique Latine. *Rev. Soc. ent. argent.* 6:9-16, 3 figs.
1937. Diptera, Family Muscidae. In Wytzman, P., *Genera Insect.*, Fasc. 205:604 pp., 9 plates. Bruxelles.

SHANNON, R. C., & DEL PONTE, E.

1926. Sinopsis parcial de los Muscoideos Argentinos. *Rev. Inst. bact.*, B. Aires 4 (5):549-590, 4 pls.
1928. Sinopsis parcial de los Muscoideos Argentinos (Addenda et Corrigenda). *Rev. Inst. bact.*, B. Aires 5 (1):141-147.

SNYDER, F. M.

1952. New Species of *Pegomyia* from Mexico (Diptera: Muscidae). *Ann. ent. Soc. Am.* 45:415-422.
1957a. A New *Pegomyia* Leaf Miner in *Chenopodium* From Mexico (Diptera: Muscidae). In Olmedo, I. H., Investigaciones sobre el Minador de la Hoja del Huauzontle. *Revista Chapingo* 64:126-131.
1957b. Notes and Descriptions of some Neotropical Muscidae (Diptera). *Bull. Am. Mus. nat. Hist.* 113 (6):437-490.

STEIN, P.

1898. Nordamerikanische Anthomyiden. Beitrag zur Dipterenfauna der Vereinigten Staaten. *Berl. ent. Z.* 42 [1897]:161-288.
1901. Die Walker'schen aussereuropäischen Anthomyiden in der Sammlung des British Museum zu London (Dipt.). *Z. syst. Hymenopt. Dipterol.* 1:185-221.



1902. Einige Wiedemann'sche u. Jaennicke'sche aussereuropäische Anthomyiden. (Dipt.). *Z. syst. Hymenopt. Dipterol.* 2:129-137.
 1904. Die Amerikanischen Anthomyiden des Königlichen Museums für Naturkunde zu Berlin und des Ungarischen National-Museums zu Budapest. *Ann. hist.-nat. Mus. natn. hung.* 2:414-495.
 1907. Revision der Bigot'schen und einiger von Macquart beschriebenen aussereuropäischen Anthomyiden (Dipt.). *Z. syst. Hymenopt. Dipterol.* 7:209-217 and 273-293.
 - 1910a. Zur Kenntnis der Gattung *Fucellia* Rob. Desv. *Wien. ent. Ztg.* 29:11-27.
 - 1910b. Die von Thomson beschriebenen Anthomyiden der *Eugenies* Resa. *Wien. ent. Ztg.* 29:66-78.
 1911. Die von Schuse in Südamerika gefangenen Anthomyiden. *Arch. Naturgesch.* 77, 1 (1):61-189.
 1918. Zur weitem Kenntnis aussereuropäischer Anthomyiden. *Ann. hist.-nat. Mus. natn. hung.* 16:147-244.
 1919. Die Anthomyidengattungen der Welt, analytisch bearbeitet, nebst einem kritisch-systematischen Verzeichnis aller aussereuropäischen Arten. *Arch. Naturgesch.* 83 A 1 [1917]:85-178.
 1920. Nordamerikanische Anthomyiden. 2. Beitrag. *Arch. Naturgesch.* 84 A 9 [1918]:1-106.
- STONE, A., SABROSKY, C. W., WIRTH, W. W., FOOTE, R. H., & COULSON, J. R.
1965.. A Catalog of the Diptera of America North of Mexico. U. S. Dep. Agric., *Agric. Handbook* 276:1696 pp.
- STUARDO ORTIZ, C.
1946. *Catalogo de los dipteros de Chile*, 250 pp. Ministerio de Agricultura, Santiago, Chile.
- THOMSON, C. G.
1869. Diptera. In *Kongliga svenska fregatten Eugenies Resa omkring jorden*, part 2, Zoologie, 1, Insekter: 443-614, pl. 9, Stockholm.
- WALKER, F.
1849. *List of the Specimens of Dipterous Insects in the collection of the British Museum.* 4:689-1172. London.
1852. *Insecta Saundersiana: or characters of undescribed insects in the collection of William Wilson Saunders, Esq., F. R. S., F. L. S., &c.* 1 Diptera, part 4:253-414. London.
- WESTWOOD, J. O.
1840. *An Introduction to the Modern Classification of Insects. Synopsis of the Genera of British Insects.* 158 pp. London.
- WIEDEMANN, C. R. W.
1830. *Aussereuropäische zweiflügelige Insekten* 2: xii + 684 pp., Hamm.
- WOLCOTT, G. N.
1924. "Insectae Portoricensis." A preliminary annotated check-list of the insects of Porto Rico, with descriptions of some new species. *J. Dep. Agric. P. Rico* 7 (1):1-313, 2 pls.



1936. "Insectae Borinquenses." A Revised Annotated Check-List of the Insects of Puerto Rico. *J. Agric. Univ. P. Rico* 20 (1):627 + 3 pp., illustr.
1948. The Insects of Puerto Rico. Diptera. *J. Agric. Univ. P. Rico* 32 (3):417-532, figs.

WULP, F. M. VAN DER

1883. Amerikaansche Diptera (III). *Tijdschr. Ent.* 26:1-60, pls. 1-2.
- 1888-1903. Insecta, Diptera. In Godman, F. D., and Salvin, O., *Biologia Centrali-Americana* 2: x + 489 pp., 11 figs., 13 pls. London.

ZETTERSTEDT, J. W.

1845. *Diptera Scandinaviae. Disposita et descripta.* 4:1281-1738. Lundae.

INDEX

abrepta Giglio-Tos, 1893 (Hylemyia)	96a.10
aenea (Malloch), 1934 (Hylemyia), Craspedochoeta	96a.2
affinis Stein, 1898, Pegomya	96a.8
albicincta (Fallén), 1825 (Musca), Calythia	96a.2
albilamellata (Stein), 1911 (Chortophila), Craspedochoeta	96a.2
albostriata (Wulp), 1883 (Chortophila), Craspedochoeta	96a.2
ammomyia, error	96a.6
andicola (Bigot), 1885 (Hylemyia), Leucophora	96a.7
angustifrons Malloch, 1934, Myopella	96a.7
Anthomyia Meigen, 1803	96a.1
antiqua (Meigen), 1826 (Anthomyia), Delia	96a.4
appendiculata (Stein), 1904 (Myopina), Myopella	96a.7
argentina (Malloch), 1934 (Hammomyia), Leucophora	96a.7
argentina Snyder, 1957, Emmesomyia	96a.4
auricollis Stein, 1918, Taeniomyia	96a.9
aurifacies Albuquerque, 1952, Hylemyioide	96a.6
bella Stein, 1911, Pegomya	96a.8
brasiliensis Albuquerque, 1949, Hylemyioide	96a.6
brassicae (Bouché), 1834 (Anthomyia), Eriopsis	96a.5
bruchii (Shannon & Del Ponte), 1926 (Hylemyia), Pegomya	96a.8
calcarata, Fucellia, error	96a.5
calcoerata (Macquart), 1851 (Scatophaga), Fucellia	96a.5
Calythia Schnabl & Dziedzicki, 1911	96a.2
cana (Macquart), 1835 (Anthomyia), Delia	96a.4
carrerae Albuquerque, 1959, Pegomya	96a.8
ceparum (Meigen), of Johnson, Delia	96a.4
chilensis (Jaenicke), 1867 (Anthomyia), Craspedochoeta	96a.3
chilensis Macquart, 1851 (Chortophila), Craspedochoeta	96a.2
chilensis (Malloch), 1934 (Hammomyia), Leucophora	96a.7
chirotilina Hennig, 1969, Craspedochoeta	96a.2
chrysostoma Rondani, 1864 (Anthomyia)	96a.10
cilicrura (Rondani), 1866 (Chortophila), Delia	96a.4





<i>cinctinervis</i> Stein, 1918, <i>Pegomya</i>	96a.8
<i>citreibasis</i> (Malloch), 1934 (<i>Hylemyia</i>), <i>Craspedochocta</i>	96a.2
<i>citribasis</i> , error	96a.2
<i>cognata</i> Stein, 1920, <i>Pegomya</i>	96a.8
<i>collaris</i> (Wulp), 1896 (<i>Hydrophoria</i>), <i>Taeniomyia</i>	96a.9
<i>comis</i> (Stein), 1911 (<i>Anthomyia</i>), <i>Calythea</i>	96a.2
<i>confusa</i> (Albuquerque), 1959 (<i>Hylemyia</i>), <i>Craspedochocta</i>	96a.2
<i>ecelesia</i> (Walker), 1949 (<i>Anthomyia</i>), <i>Delia</i>	96a.4
<i>costana</i> (Séguy), 1934 (<i>Chortophila</i>), <i>Calythea</i>	96a.2
<i>Craspedochaeta</i> , error	96a.2
<i>Craspedochocta</i> Macquart, 1851	96a.2
<i>crassinervis</i> Albuquerque, 1959, <i>Hylemyioide</i>	96a.6
<i>crenata</i> (Bigot), 1885 (<i>Trichopticus</i>), <i>Calythea</i>	96a.2
<i>Delia</i> Robineau-Desvoidy, 1830	96a.3
<i>devia</i> Cova Garcia, 1964 (<i>Pegomyia</i>)	96a.10
<i>dexiaria</i> (Stein), 1904 (<i>Hydrophoria</i>), <i>Taeniomyia</i>	96a.9
<i>diversa</i> (Wiedemann), 1830 (<i>Anthomyia</i>), <i>Delia</i>	96a.4
<i>dolosa</i> Stein, 1918, <i>Pegomya</i>	96a.8
<i>dorsimaculata</i> (Wulp), 1896 (<i>Anthomyia</i>), <i>Pegomya</i>	96a.8
<i>duplicoides</i> Snyder, 1952, <i>Pegomya</i>	96a.8
<i>edwardsi</i> (Malloch), 1934 (<i>Hylemyia</i>), <i>Craspedochocta</i>	96a.3
<i>elongata</i> (Wulp), 1896 (<i>Anthomyia</i>), <i>Pegomya</i>	96a.8
<i>Emmesomyia</i> Malloch, 1917	96a.4
<i>Erioschia</i> Lioy, 1864	96a.5
<i>flavibasis</i> (Stein), 1911 (<i>Hydrophoria</i>), <i>Taeniomyia</i>	96a.9
<i>flavipalpis</i> (Wulp), 1896 (<i>Hydrophoria</i>), <i>Emmesomyia</i>	96a.4
<i>flavipes</i> (Stein), 1918 (<i>Ammomyia</i>), <i>Leucophora</i>	96a.7
<i>florilega</i> (Zetterstedt), 1845 (<i>Aricia</i>), <i>Delia</i>	96a.4
<i>Fucellia</i> Robineau-Desvoidy, 1842	96a.5
<i>fucorum</i> (Fallén), 1819 (<i>Scatomyza</i>), <i>Fucellia</i>	96a.5
<i>fumifera</i> , <i>Fucellia</i> , error	96a.5
<i>funifera</i> Stein, 1910, <i>Fucellia</i>	96a.5
<i>fusciceps</i> (Zetterstedt), 1845 (<i>Aricia</i>), <i>Delia</i>	96a.4
<i>fuscipes</i> , <i>Delia</i> , error	96a.4
<i>Hammomyia</i> Rondani, 1877	96a.6
<i>hernandesi</i> Snyder, 1957, <i>Pegomya</i>	96a.8
<i>Hydrophoria</i> Robineau-Desvoidy, 1830	96a.6
<i>Hylemya</i> Robineau-Desvoidy, 1830	96a.6
<i>Hylemyia</i> , error	96a.6
<i>Hylemyioide</i> Albuquerque, 1949	96a.6
<i>Hylemyioides</i> , error	96a.6
<i>Hylemyoides</i> , error	96a.6
<i>Hylephila</i> Rondani, 1877	96a.6
<i>intermedia</i> Lundbeck, 1901, <i>Fucellia</i>	96a.5
<i>johnsoni</i> (Stein), 1898 (<i>Hylemyia</i>), <i>Leucophora</i>	96a.7



<i>lanicrus</i> (Thomson), 1869 (Anthomyia), Delia	96a.4
<i>latifrons</i> (Stein), 1918 (Hylemyia), Delia	96a.4
<i>latimaculata</i> Albuquerque, 1959, Hylemyoide	96a.6
<i>Leucophora</i> Robineau-Desvoidy, 1830	96a.6
<i>leucotelus</i> Walker, 1852 (Anthomyia)	96a.10
<i>levipes</i> Giglio-Tos, 1893 (Hylemyia)	96a.10
<i>limbinervis</i> (Macquart), 1843 (Chortophila), Craspedochoeta	96a.3
<i>lindgi</i> , Craspedochoeta, error	96a.3
<i>lindigii</i> (Schiner), 1868 (Anthomyia), Craspedochoeta	96a.3
<i>lingindii</i> , Craspedochoeta, error	96a.3
<i>liturata</i> (Meigen), 1838 (Anthomyia), Delia	96a.4
<i>liturata</i> Rondani, 1868 (Chortophila)	96a.10
<i>maculipennis</i> (Albuquerque), 1953 (Hammomyia), Leucophora ..	96a.7
<i>maculipennis</i> (Stein), 1918 (Pegomyia), Craspedochoeta	96a.3
<i>marina</i> (Macquart), 1838 (Scatophaga), Fucellia	96a.5
<i>maritima</i> (Haliday), 1838 (Scatophaga), Fucellia	96a.5
<i>masafuerana</i> (Enderlein), 1940 (Thrixina), Fucellia	96a.5
<i>masatierrana</i> (Enderlein), 1940 (Thrixina), Fucellia	96a.5
<i>meracula</i> (Giglio-Tos), 1893 (Spilogaster), Pegoplata	96a.9
<i>mexicana</i> (Giglio-Tos), 1893 (Lasiops), Calythea	96a.2
<i>micropteryx</i> (Thomson), 1869 (Anthomyia), Calythea	96a.2
<i>monticola</i> (Wulp), 1896 (Hylemyia), Pegoplata	96a.9
<i>Myopella</i> Malloch, 1934	96a.7
<i>Neohylemyia</i> Malloch, 1917	96a.7
<i>neopalosa</i> , Pegomya, error	96a.8
<i>neopalposa</i> Snyder, 1957, Pegomya	96a.8
<i>nervicincta</i> (Stein), 1911 (Chortophila), Pegomya	96a.8
<i>nigracaerulea</i> Snyder, 1952, Pegomya	96a.8
<i>nigricans</i> (Robineau-Desvoidy), 1830 (Anthomyia), Calythea	96a.2
<i>nigricauda</i> (Bigot), 1885 (Hydrophoria), Neohylemyia	96a.7
<i>nigrifemur</i> Stein, 1918, Pegomyia	96a.8
<i>nigripes</i> Robineau-Desvoidy, 1830 (Hylemyia)	96a.10
<i>nigrocaerulea</i> , Pegomya, error	96a.8
<i>notata</i> (Albuquerque), 1957 (Pegomyza), Pegomya	96a.8
<i>oculifera</i> Bigot, 1885, Anthomyia	96a.1
<i>pacifica</i> Malloch, 1923, Fucellia	96a.5
<i>pallidithorax</i> (Albuquerque), 1959 (Hylemyia), Craspedochoeta ..	96a.3
<i>paulistensis</i> Albuquerque, 1952, Hylemyioide ...	96a.6
<i>Pegomya</i> Robineau-Desvoidy, 1830	96a.7
<i>Pegomyia</i> , emendation	96a.7
<i>Pegoplata</i> Schnabl & Dziedzicki, 1911	96a.9
<i>Petrmyia</i> Albuquerque, 1954	96a.9
<i>peullae</i> (Malloch), 1934 (Hammomyia), Leucophora	96a.7
<i>pectipes</i> (Bigot), 1885 (?Hydrophoria), Taeniomyia	96a.9
<i>platura</i> (Meigen), 1826 (Anthomyia), Delia	96a.4
<i>plumiseta</i> (Malloch), 1934 (Hammomyia), Leucophora	96a.7
<i>plumosa</i> Wulp, 1896, Hydrophoria	96a.6
<i>plurinervis</i> Albuquerque, 1958, Hylemyioide	96a.6
<i>pluripunctata</i> (Albuquerque), 1959 (Hylemyia), Craspedochoeta	96a.3


<i>pluvialis</i> (Linnaeus), 1758 (<i>Musca</i>), <i>Anthomyia</i>	96a.2
<i>poeciloptera</i> Malloch, 1921, <i>Pegomya</i>	96a.8
<i>poeciloptera</i> Malloch of Malloch, 1934, <i>Pegomya</i>	96a.8
<i>poeciloptera</i> Malloch of Carrera & Travassos Filho, 1947, <i>Pegomya</i>	96a.8
<i>prisca</i> (Wulp), 1896 (<i>Phorbia</i>), <i>Delia</i>	96a.4
<i>pullula</i> (Zetterstedt), 1845 (<i>Aricia</i>), <i>Craspedochoeta</i>	96a.3
<i>punctipennis</i> (Shannon & Del Ponte), 1926 (<i>Hylemyia</i>), <i>Craspe-</i> <i>dochoeta</i>	96a.3
<i>punctipennis</i> (Wiedemann), 1830 (<i>Anthomyia</i>), <i>Craspedochoeta</i>	96a.3
<i>recurrens</i> (Malloch), 1934 (<i>Hylemyia</i>), <i>Craspedochoeta</i>	96a.3
<i>rejecta</i> Aldrich, 1918, <i>Fucellia</i>	96a.5
<i>rhodina</i> (Giglio-Tos), 1893 (<i>Hylemyia</i>), <i>Pegoplata</i>	96a.9
<i>rubifrons</i> (Bigot), 1885 (<i>Anthomyia</i>), <i>Delia</i>	96a.4
<i>rubriceps</i> Hockett, 1965, <i>Pegomya</i>	96a.8
<i>rubifrons</i> (Bigot), 1885 (<i>Chortophila</i>), <i>Delia</i>	96a.4
<i>ruficeps</i> Stein, 1898, <i>Pegomya</i>	96a.8
<i>ruficoxa</i> (Bigot), 1885 (<i>Parachortophila</i>), <i>Fucellia</i>	96a.5
<i>rufifrons</i> , <i>Delia</i> , error	96a.4
<i>rufitibia</i> (Stein), 1918 (<i>Ammomyia</i>), <i>Leucophora</i>	96a.7
<i>ruralis</i> (Meigen), 1826 (<i>Anthomyia</i>), <i>Hydrophoria</i>	96a.6
<i>sanetijacobi</i> (Bigot), 1885 (<i>Anthomyia</i>), <i>Delia</i>	96a.4
<i>sanctojacobi</i> , <i>Delia</i> , error	96a.4
<i>scutellata</i> (Stein), 1911 (<i>Hydrophoria</i>), <i>Taeniomyia</i>	96a.10
<i>segmentata</i> (Wulp), 1896 (<i>Anthomyia</i>), <i>Delia</i>	96a.4
<i>scnipellucida</i> (Stein), 1918 (<i>Taeniomyia</i>), <i>Emmesomyia</i>	96a.5
<i>spadibasis</i> Snyder, 1957, <i>Emmesomyia</i>	96a.5
<i>spilota</i> (Wulp), 1883 (<i>Chortophila</i>), <i>Craspedochoeta</i>	96a.3
<i>spinisoides</i> Snyder, 1952, <i>Pegomya</i>	96a.9
<i>stlemba</i> (Giglio-Tos), 1893 (<i>Chortophila</i>), <i>Pegomya</i>	96a.9
<i>subvittata</i> (Malloch), 1934 (<i>Hammomyia</i>), <i>Leucophora</i>	96a.7
<i>Taeniomyia</i> Stein, 1918	96a.9
<i>tarsadisca</i> Snyder, 1952, <i>Pegomya</i>	96a.9
<i>tenuirostris</i> (Wulp), 1896 (<i>Hylemyia</i>), <i>Neohylemyia</i>	96a.7
<i>tergina</i> (Zetterstedt), 1845 (<i>Anthomyza</i>), <i>Fucellia</i>	96a.5
<i>transversalis</i> (Wulp), 1896 (<i>Hydrophoria</i>), <i>Taeniomyia</i>	96a.9
<i>trichodactyla</i> (Rondani), 1866 (<i>Chortophila</i>), <i>Delia</i>	96a.4
<i>trifilis</i> (Giglio-Tos), 1893 (<i>Anthomyia</i>), <i>Delia</i>	96a.4
<i>trifumosa</i> Snyder, 1957, <i>Pegomya</i>	96a.9
<i>trimaculata</i> (Stein), 1911 (<i>Hydrophoria</i>), <i>Taeniomyia</i>	96a.10
<i>urbica</i> Wulp, 1896, <i>Hylemyia</i>	96a.6
<i>verticala</i> Snyder, 1952, <i>Pegomya</i>	96a.9
<i>video</i> (Gaminara), 1930 (<i>Hylemyia</i>), <i>Craspedochoeta</i>	96a.3
<i>virgata</i> (Stein), 1911 (<i>Chortophila</i>), <i>Craspedochoeta</i>	96a.3
<i>vittifera</i> (Stein), 1911 (<i>Pegomyia</i>), <i>Craspedochoeta</i>	96a.3
<i>wygodzinskyi</i> Albuquerque, 1954, <i>Petrmyia</i>	96a.9
<i>xanthopyga</i> (Albuquerque), 1959 (<i>Hylemyia</i>), <i>Craspedochoeta</i>	96a.3








Composto e impresso em 1974, nas oficinas da
EMPRESA GRÁFICA DA REVISTA DOS TRIBUNAIS S.A.
R. Conde de Sarzedas, 38, fone 33-4181, São Paulo, S.P., Brasil





SciELO

MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO

(FORMERLY DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

96B

FAMILY SCATOPHAGIDAE
(Scatomyzidae, Scopeumatidae, Cordiluridae)

† DALCY DE OLIVEIRA ALBUQUERQUE

Museu Nacional
Universidade Federal do Rio de Janeiro

This family, often treated as a part of the Anthomyiidae, is represented in the area covered by this Catalogue by 5 species in the genus *Scatophaga* Meigen, and by a new genus from Ecuador. The former is widely distributed in the holarctic region and contains the best known and commonest species. Larvae of several species of *Scatophaga* feed on dung.

The genera of Scatophagidae have been revised by Ségué (1952).

Genus SCATOGERA, gen. n.

Interfrontal space not depressed. Frontal bristles 4 pairs, the upper pair recumbent and divaricate. Parafacialia with 1 pair of proclinate and divaricate bristles. Face straight: arista weakly plumose (Fig. 1). Antenna long, apex rounded. Dorsocentral bristles 1:3. Propleura bare. Sternopleurals 1:0. Tibia I with 1 anteroventral bristle on the apical third and 1 dorsal immediately below. Tibia II with 2 posterodorsal, 1 posterior and 2 anterior bristles. Tibia III with 2 anterodorsal, 2 posterodorsal and 1 anteroventral bristles.

Type-species, *Scatogera primogenita*, sp.n.

Scatogera, gen. n., is related to *Spaziphora* Rondani, differing by the plumose arista.

Scatogera primogenita, sp. n.

♂ - Total length: 7 mm.

Yellowish-brown; thorax, scutellum, metanotum, ocellar triangle and abdomen darker. Antennae, tarsi and claws black. Wing with costal margin, apex and transverse veins darkened. Calypttrae dark-brown. Pulvilli yellow.

Head - Lozenge-shaped in profile, slightly inflated posteriorly, with eyes bare and separated by a parallel-sided space which measures, at the anterior ocellus level, circa 0.23 of head width. Ocellar triangle long, reaching the apex of the front, which does not show cruciate bristles. Frontal bristles 4 pairs, beginning at the antennal bases and extending to the level of the anterior ocellus. First and second pairs of frontals convergent; third and fourth (upper) pairs divaricate and recumbent, the pair nearest to the vertex very robust. Parafrontalia showing, between the 2 pairs of upper and lower bristles, 1 proclinate bristle, as strong as the anterior, convergent, frontal bristles and situated beyond the middle of the space separating the last pairs of frontals. Parafrontalia with a series of 1 or 2 cilia along its entire extension. Antennae with rounded apex, 2nd article measuring almost 3 times the length of the first. Arista weakly plumose, the branches spaced (Fig. 1). Vibrissae strong, with 1-3 additional bristles at base. Posterior surface of head with relatively strong, erect black cilia, which are not dense. Proboscis small, larger than palpi; these spatulate, leaf-shaped.


Thorax - With a few cilia in the vestiture. Dorsocentrals 1:3, the first post-sutural pair small. Achrosticals absent. 2 humeral, 1 post-humeral, 1 pre-sutural, 1 pre-alar, 2 intra-



alar, 2 supra-alar and 2 notopleural bristles, the posterior notopleural smaller. Mesopleura with a series of 5 bristles. Sternopleura with 1 strong bristle. Pteropleura with a few cilia. Prothorax and prothoracic stigma with 1 strong bristle and some satellite cilia at base. Prosternum and hypopleura bare. Scutellum naked, infrascutellum slightly inflated. Posterior spiracle circular, its opening smaller than the capitellum of the halter. Calyptrae reduced, the thoracic calyptra vestigial, the alar with long fringe. Scutellum with 2 pairs of marginal bristles, the apical small.

Wing (Fig. 2) - Covered with microtrichia; veins bare. Third and fourth veins parallel. Anal vein reaching wing apex. Anterior transverse vein situated beyond the middle of the discal cell. Costal spine reduced.

Legs - Elongate. Femur I with anterodorsal and ventral surfaces with a series of strong and spaced bristles. Tibia I with a mediocre, submedian bristle; dorsal surface with 1 strong bristle situated slightly below the level of the anteroventral bristle and more or less at the apical half; dorsal and anteroventral surfaces, apically, with 1 bristle, the latter stronger; pretarsus subequal to the sum of the other tarsomeres, with a few sensory cilia on the ventral surface, longer at the base; pulvilli and claws elongate. Femur II with a series of



Scatogera primogenita, gen. n., sp. n., ♂. Fig. 1: Arista; fig. 2: wing; fig. 3: 5th sternite; fig. 4: cercal plate, dorsal view.

bristles and 1 preapical bristle on the anterior surface; anteroventral and posteroventral surfaces with 1-2 bristles, longer than strong, inserted approximately at the level of the basal half; posterior surface with 2 bristles inserted pre-apically and 1 posteroventral situated at the same transverse level of femur. Tibia II with 2 bristles on the anterior surface, the one nearer to the apex robust; posterior surface with 2 bristles inserted at the limits of the third; posteroventral surface, in the limit of the inferior third, with 1 mediocre bristle; dorsal, anterodorsal, posterodorsal, anteroventral and ventral surfaces each with 1 apical bristle; that of the ventral surface very robust, pretarsus, tarsomeres, pulvilli and claws as in the anterior pair. Femur III with a series of strong, spaced bristles on the anterodorsal and anteroventral surfaces. Tibia III with 2 bristles (limiting the third) on the dorsal surface, the lower one very robust; anterodorsal surface with 4 bristles, the one nearer to the apex slightly smaller than the pre-apical posterodorsal; anteroventral surface with 2 small bristles on the apical half; anterodorsal and ventral surfaces each with 1 apical bristle, the anterodorsal more robust. Tibia III with closely set cilia on the posterior surface, not forming however any special aspect.

Abdomen - Long, with 5 visible segments and the genitalia discretely voluminous. All segments with strong apical bristles, including the genital segment. First sternite bare.

Genitalia - As in figures 3-4.

Holotype ♂, Ecuador, E. Papallacta (Quito-Baeza), 2.900 m, i.1971 (L.E. Peña), in the Museu de Zoologia da Universidade de São Paulo.

Genus SCATOPHAGA Meigen

Scopeuma Meigen, 1800: 36. Type-species, *Musca merdaria* Fabricius (Coquillett, 1910: 604) = *stercoraria* (Linnaeus). Suppressed by ICZN, 1963: 339.

SCATOPHAGA Meigen, 1803: 277 (as *Scathophaga*). Type-species, *Musca merdaria* Fabricius (mon.) = *stercoraria* (Linnaeus). Ref. - Malloch, 1935: 244 (key).

CHILENSIS Albuquerque, 1953: 1, figs. 1-7. Type-locality: Chile, Achacachi, 3850 m.

FURCATA (Say), 1823: 98 (1859: 85) (*Pyropa*). Type-locality: U.S.A., Missouri. Distr. - U.S.A., México (D.F.).

RESES Giglio-Tos, 1893: 7. Type-locality: "Mexico". Distr. U.S.A., s. to Belize and Guatemala.

STERCORARIA (Linnaeus), 1758: 599 (*Musca*). Type-locality: "Sweden". Distr. - Asia, Europe, North America, Mexico, Haiti.

TROPICALIS Malloch, 1931: 443 (nom. nov. for *lannata* Szilády). Type-locality: Bolivia, Coroico, Cillunticara and Rinconada. Distr. - Peru, Bolívia, Ref. - Malloch, 1935: 244, 251, fig. 2.

lannata Szilády, 1926: 595 (preocc. Lundbeck, 1895). Type-locality: Bolivia, Coroico, Cillunticara and Rinconada.

LITERATURE CITED

ALBUQUERQUE, D. de O.


1953. Contribuição ao conhecimento dos Scatophagidae neotropicais, com descrição de uma espécie nova (Diptera: Scatophagidae). *Bolm Mus. nac. R. de Janeiro, Zool.* (n.s.) 121: 1-7, 7 figs.

COQUILLET, D.W.

1910. The type-species of the North American genera of Diptera. *Proc. U.S. natl Mus.* 37: 499-647.

GIGLIO-TOS, E.

1893. Diagnosi di nuovi generi e di nuove specie di ditteri. IX. *Bol. Mus. Zool. Anat. comp., Torino* 8 (158): 1-14.



INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE

1963. *Opinion 678. The suppression under the Plenary Powers of the pamphlet published by Meigen, 1800. Bull. Zool. Nomencl. 20: 339-342.*

LINNAEUS, C.

1758. *Systema naturae per regna tria naturae*. Ed. X., 578 pp., 2 pls. Stockholm.

MALLOCH, J.R.

1931. Exotic Muscaridae (Diptera). XXXV. *Ann. Mag. nat. Hist.* (10) 8: 425-446, 6 figs.
1935. Exotic Muscaridae (Diptera). XXXVIII. *Ann. Mag. nat. Hist.* (10) 15: 242-266, 22 figs.

MEIGEN, J.W.

1800. *Nouvelle classification des mouches à deux ailes (Diptera L.) d'après un plan tout nouveau*, 40 pp. Paris.
1803. Versuch einer neuen Gattungseintheilung der europäischen zweiflügeligen Insekten. *Mag. f. Insektenkde* 2: 259-281.

SAY, T.

1823. Descriptions of dipterous insects of the United States. *J. Acad. nat. Sci. Phila.* 3: 9-54, 73-104.
1859. (In J.L. LeConte, ed.), *The complete writings of Thomas Say on the entomology of North America*, 1: 412 pp., 54 pls. New York.

SÉGUY, E.

1952. Diptera. Fam. Scatophagidae, in P. Wytsman, ed., *Genera Insectorum* 209: 1-107, 44 figs. Bruxelles.

SZILÁDY, Z.

1926. Dipterenstudien. *Ann. Mus. nat. Hung.* 24: 586-611, 18 figs.

MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO
(Formerly Departamento de Zoologia, Secretaria da Agricultura,
São Paulo)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS SOUTH OF THE UNITED STATES

97

FAMILY MUSCIDAE

ADRIAN C. PONT

British Museum (Natural History), London, England

Adult Muscidae are generally small to medium-sized flies, and seldom exhibit any striking developments of colour or form. They are found throughout the South American continent, from the temperate south to the equatorial north, and in almost all habitats except for the most dry, from sea-level up to 16,000 feet. Adults of Coenosiinae and Limnophorinae are predaceous upon other insects, whilst those of the other subfamilies feed on nectar and pollen, or are attracted to excrement, carrion and garbage for both feeding and oviposition. Some species suck blood, sweat, mucous secretions, etc., from man and his domestic animals and are well-known for their menace to good health.

Larval habitats are equally varied, and larvae may be coprophagous, saprophagous or carnivorous in a wide variety of habitats containing decaying animal or vegetable matter, or may even be aquatic. Few are of any medical or agricultural importance in South America.

In this catalogue, an attempt has been made to review the types of all species that had not been placed in their modern genera by Malloch, Snyder or Albuquerque, and many others were studied in order to confirm generic assignments. Even so, it has not always been possible to place species in existing genera, and it is clear that as much work needs to be done on generic structure as on species taxonomy on a South American basis. Where doubt exists as to generic assignment, this is indicated in the text. Brèthes' types and certain Wiedemann types could not be studied, to my great regret, and a number of Stein's types are known to have been destroyed. For various reasons, however, this catalogue should only be regarded as preliminary, for I am only too aware of its short-comings. A number of small or monotypic genera had to be placed in tribes without first-hand knowledge of the flies, and the group of genera around *Muscina* is only temporarily assigned to the Hydrataeini. Some subfamilies have a workable generic structure, but almost everywhere there is a need for further work, particularly in the Coenosiinae and Limnophorinae. Genera have often been defined quite arbitrarily in the past, and the assignment of species to genera is therefore sometimes a reflection of this and is not always based on any evident natural relationship.

The bibliography includes all taxonomic and faunistic papers that have been consulted during the compilation of this catalogue. Where

97.1

authors have published several papers in one year, the papers are listed chronologically. The pioneer work on South American Muscids was done by Paul Stein, and his papers are still useful. Amongst subsequent workers, who have been lamentably few, the papers of Malloch, Snyder and Albuquerque are most useful. Regarding supra-generic classification, I have followed the suggestions of Hennig (1965), with some minor modifications, which differs in certain important respects from that of earlier authors.

I am grateful to the following colleagues for the loan of material from the collections in their charge, and to several of them for additional help and information: Mr. D. M. Ackland (Oxford), Dr. G. W. Byers (Lawrence, Kansas), Mr. W. N. Ellis (Amsterdam), Dr. R. J. Gagné (Washington), Dr. J. H. Guimarães (São Paulo), Dr. P. J. Van Helsdingen (Leiden), Prof. W. Hennig (Stuttgart), Dr. J. Jezek (Prague), Dr. A. Kaltenbach (Vienna), Dr. R. Krause (Dresden), Dr. C. Leonardi (Milan), Dr. L. Lyneborg (Copenhagen), Mr. L. Matile (Paris), Dr. J. Moucha (Prague), Prof. U. Parenti (Turin), Mr. P. I. Persson (Stockholm), Dr. L. de Santis (La Plata), Dr. H. Schumann (Berlin), Mr. E. Taylor (Oxford), Prof. G. C. Varley (Oxford), and Dr. P. Wygodzinsky (New York). For advice on the nomenclature of *Syllimnophora* I am grateful to Dr. R. V. Melville (London) and Dr. C. W. Sabrosky (Washington). I am particularly grateful to Dr. R. W. Crosskey (London), for much discussion and for advice on numerous problems arising from this work, and to Dr. N. Papavero (São Paulo), for his continual help in obtaining literature and information for me.

Subfamily FANNIINAE

Genus EURYOMMA Stein

EURYOMMA Stein, 1899:19. Type-species, *hispaniense* Stein (mon.) = *peregrinum* (Meigen).

CARIOCA Albuquerque, 1956g:2, figs. 1-7. Type-locality: Brazil, Guanabara, Rio de Janeiro, Grajaú. Distr. — Brazil.

ERYTHROGASTER Séguay, 1941:54, figs. 1-3. Type-locality: "Chile". Distr. — Chile.

LONGICORNE Stein, 1911:111. Type-locality: Chile, Guayacán. Distr. — Chile.

NIGRIFEMUR Stein, 1911:110. Type-locality: Chile, Palca and Tacna; Peru, Arequipa, Tarma and Cuzco; Bolivia, Sorata. Distr. — Venezuela, Peru, Bolivia, Chile.

PANAMENSIS Chillcott, 1958:730, figs. 4-6, 8, 17-20. Type-locality: Panama Canal Zone, Barro Colorado Island. Distr. — Panama.

PEREGRINUM (Meigen), 1826:187 (*Anthomyia*). Type-locality: Germany, Hamburg, on a ship from America. Distr. — Ecuador, Peru, Chile, Tierra del Fuego, Guadalupe Is., Juan Fernandez Is.; tropicopolitan. Ref. — Hennig, 1955:14, pl. 1, fig. 3, pl. 3, fig. 61; Chillcott, 1961:224, figs. 149, 149-A-B, 217, 259.

brevipalpis Thomson, 1869:557 (*Anthomyia*). Type-locality: Ecuador, Guayaquil.

hispaniense Stein, 1899:20. Type-locality: Spain, Algeciras.

RETTENMEYERI Chillcott, 1958:725, figs. 1-3, 7, 9-16, 21. Type-locality: Panama Canal Zone, Barro Colorado Island. Distr. — Panama.

RUFFRONS Stein, 1911:110. Type-locality: Chile, Guayacán. Distr. — Peru, Chile.


Genus FANNIA Robineau-Desvoidy

- FANNIA Robineau-Desvoidy, 1830:567. Type-species, *saltatrix* Robineau-Desvoidy (mon.) = *scalaris* (Fabricius).
- Homalomyia* Bouché, 1834:89. Type-species, *Musca canicularis* Linnaeus (Westwood, 1840:143).
- Dasyphyma* Bigot, 1883:clxxxviii. Type-species, *armata* Bigot (mon.) = *setosa* (Bigot).
- Profannia* Séguy, 1937:163 (as subgenus). Type-species, *parasitica* Séguy (orig. des.).
- Mesazelia* Blanchard, 1942:57. Type-species, *trichopoda* Blanchard (orig. des.). N. SYN.
- ABNORMIS (Stein), 1900b:210 (*Homalomyia*). Type-locality: Bolivia, S. Antonio and Songo. Distr. — Venezuela, Colombia, Peru, Bolivia.
- ADMIRABILIS Albuquerque, 1958b:21, 7 figs. Type-locality: Brazil, Rio de Janeiro, Itatiaia, Lago Azul. Distr. — Brazil.
- ALBITARSIS Stein, 1911:105. Type-locality: Peru, Juliaca, Sicuani and Cuzco; Bolivia, Lake Titicaca; Chile, Guayacán, Coquimbo and Caldera. Distr. — Peru, Bolivia, Chile, Argentina; South Africa.
- ANNOSA Chillcott, 1961:60. Type-locality: Mexico, west side of Cortés Pass. Distr. — Mexico.
- ANTENNATA Stein, 1911:103. Type-locality: Peru, Umuhuankiali. Distr. — Mexico, Peru.
- ANTHRACINA (Walker), 1837:356 (*Anthomyia*). Type-locality: Port Famine. Distr. — Chile, Chiloé Is., Juan Fernández Is., Tierra del Fuego, Argentina. Ref. — Pont, 1966:427, figs. 1-12.
- albibasis* Malloch, 1934:202. In key only.
- ARCUATA Chillcott, 1961:152, figs. 89, 96, 192, 288. Type-locality: Mexico, Coapa. Distr. — Mexico.
- BAHIENSIS Albuquerque, 1954m:288, figs. 10-16. Type-locality: Brazil, Bahia, Irará. Distr. — Brazil.
- BELLA Albuquerque, 1957g:26, figs. 34-39. Type-locality: Brazil, Minas Gerais, Lassance. Distr. — Brazil.
- bela*, error.
- BENJAMINI Malloch, 1913:625, fig. 9. Type-locality: U.S.A., California, Claremont and Santa Clara County; Arkansas, Grand Canyon; Cuba, Cayamas and Baracoa. Distr. — Mexico, Cuba; North America.
- BIGOTI (Stein), 1907:287 (*Homalomyia*). Type-locality: "Chile". Distr. — Chile.
- BINOTATA Chillcott, 1961:203, figs. 139, 139A-B. Type-locality: Mexico, Atlacomulco. Distr. — Mexico; North America.
- CANICULARIS (Linnaeus), 1761:454 (*Musca*). Type-locality: "Sweden". Distr. — Mexico, Costa Rica, Ecuador, Brazil, Uruguay, Chile, Chiloé Is., Tierra del Fuego, Argentina, Juan Fernández Is., Galápagos Is.; cosmopolitan. Ref. — Hennig, 1955:32, figs. 3, 4, 8, 15, pl. 1, fig. 7, pl. 4, fig. 66; Chillcott, 1961:188, figs. 129, 129A-D, 203, 261, 271, 272, 283.



- lateralis* Linnaeus, 1758:597 (*Musca*). Type-locality: Europe.
Suppressed by I.C.Z.N. Opinion 884.
- chilensis* Macquart., 1843a:328, pl. 23, fig. 4; 1843b:171, pl. 23, fig. 4 (*Anthomyia*). Type-locality: "Chile".
- introducata* Walker, 1852:362 (*Anthomyia*). Type-locality: "Brazil".
- mexicana* Bigot, 1885:284 (*Homalomyia*). Type-locality: "Mexico".
- CLAVATA Chillcott, 1961:163, figs. 101, 108, 249. Type-locality: Mexico, Tamaulipas, Antiguo Morelos. Distr. — Mexico.
- COXATA Shannon & Del Ponte, 1926:574. Type-locality: Argentina, San Pedro de Jujuy. Distr. — Argentina.
- DASYOPS (Stein), 1900b:209 (*Homalomyia*). Type-locality: Bolivia, Cillutin-cara. Distr. — Bolivia.
- DODGEI Seago, 1954:4, figs. 3 and 4. Type-locality: Panama, David. Distr. — Panama, Cuba, Colombia.
- ERYTHROPSIS (Bigot), 1888c:30, pl. 4, fig. 1 (*Homalomyia*). Type-locality: Tierra del Fuego, Orange Bay. Distr. — Tierra del Fuego. Ref. — Albuquerque, 1953b:496, figs. 1-5.
- erythroopsis* Schnabl, 1902:160 (*Coelomyia*). Type-locality: Cape Horn. Nec (Bigot), 1888.
- EXILIS (Williston), 1896:369 (*Limnophora*). Type-locality: St. Vincent Is. Distr. — St. Vincent Is.
- FEMORALIS (Stein), 1898:282 (*Homalomyia*). Type-locality: U.S.A., Louisiana. Distr. — Mexico, Cuba, Virgin Is., Puerto Rico, Haiti, Dominican Republic, Guyana, Peru, Bolivia, Argentina; North America. Ref. — Chillcott, 1961:214, figs. 145, 145A, 210, 281.
- FLAVICINCTA (Stein), 1904:453 (*Homalomyia*). Type-locality: Peru, Vilcanota and Colombia, Cordillera. Distr. — Colombia, Peru, Bolivia, Brazil. Ref. — Albuquerque, 1954i:71, figs. 1-8.
- FLAVICORNIS Stein, 1911:102. Type-locality: Peru, Umuhuankiali and Paso de Lares and Bolivia, S. Ernesto and S. Carlos. Distr. — Peru, Bolivia, Argentina.
- FLAVIPALPIS Stein, 1911:103. Type-locality: Chile, Palca. Distr. — Brazil, Chile. Ref. — Albuquerque, 1957g:21, figs. 27-33.
- FLAVIPES Stein, 1918:234. Type-locality: Paraguay, Asunción. Distr. — Paraguay.
- FUSCONOTATA (Rondani), 1868:27 (*Myantha*). Type-locality: Argentina, Mendoza. Distr. — Argentina.
- GILVITARSIS Chillcott, 1961:153. Type-locality: Mexico, 4 miles west of Rio Frio. Distr. — Mexico.
- GRANDIS Malloch, 1912:3. Type-locality: Panama, Porto Bello. Distr. — Panama.
- HEYDENII (Wiedemann), 1830:429 (*Anthomyia*). Type-locality: "Brazil". Distr. — Peru, Bolivia, Brazil, Uruguay, Paraguay, Argentina. Ref. — Albuquerque, 1957g:6, figs. 7-13.
- platensis* (Brèthes), 1908:303 (*Homalomyia*). Type-locality: Argentina, Buenos Aires.
- heydeni*, error.
- heideni*, error.
- platense*, error.

- HIRTIFEMUR (Stein), 1904:457 (*Homalomyia*). Type-locality: Colombia, Cordillera. Distr. — Colombia, Peru.
- HYALINIPENNIS (Czerny), 1903:240 (*Homalomyia*). Type-locality: "Bolivia". Distr. — Bolivia.
- INCISURATA (Zetterstedt), 1838:679 (*Anthomyza*). Type-locality: Sweden ("in arenosis Lapponiae Tornensis"). Distr. — Mexico, Uruguay, Argentina; North America, Europe, Asia. Ref. — Hennig, 1955:51, fig. 12, pl. 1, fig. 19; Chillcott, 1961:57, figs. 1, 1A, 155, 285.
- ? *prostrata* Scopoli, 1763:328 (*Musca*). Type-locality: Carniolia.
- pruinosa* Shannon & Del Ponte, 1926:572. Type-locality: Argentina, San Isidro.
- INDUCTA (Walker), 1852:360 (*Anthomyia*). Type-locality: "Brazil". Distr. — Brazil.
- OBSCURINERVIS (Stein), 1900b:207 (*Homalomyia*). Type-locality: Bolivia, Mato Grosso, Salobra. Distr. — Brazil.
- INTENSICA Curran in Gowdey, 1928b:40. Type-locality: Jamaica, Cinchona. Distr. — Jamaica.
- ITATIAIENSIS Albuquerque, 1956b:33, figs. 1-4. Type-locality: Brazil, Rio de Janeiro, Itatiaia, Lago Azul. Distr. — Brazil.
- LONGIPILA Albuquerque, 1954m:385, figs. 1-9. Type-locality: Brazil, São Paulo, Ilha Sêca. Distr. — Brazil.
- MICHENERI Chillcott, 1961:164. Type-locality: Mexico, Tamaulipas, Antiguo Morelos. Distr. — Mexico.
- MOROSA (Wulp), 1896:339 (*Phorbia*). Type-locality: Mexico, Mexico City. Distr. — Mexico; North America. Ref. — Chillcott, 1961:208, figs. 141, 142, 142A, 214. N. COMB.
- laevis* Stein, 1898:174 (*Homalomyia*). Type-locality: Massachusetts. N. SYN.
- OBSCURINERVIS (Stein), 1900b:207 (*Homalomyia*). Type-locality: Bolivia, Songo. Distr. — Mexico, Venezuela, Guyana, Colombia, Peru, Bolivia, Brazil, Paraguay. Ref. — Albuquerque, 1946:1, 14 figs.
- OBSCURIPENNIS (Czerny), 1903:239 (*Homalomyia*). Type-locality: "Bolivia". Distr. — Colombia, Bolivia.
- PARASITICA Séguay, 1933:258, figs. 4-5. Type-locality: Ecuador, El Napo. Distr. — Ecuador.
- PENICILLARIS (Stein), 1900b:205 (*Homalomyia*). Type-locality: Bolivia, Songo. Distr. — Colombia, Peru, Bolivia, Brazil.
- PETROCCHIAE Shannon & Del Ponte, 1926:574. Type-locality: Argentina, Rio Bermejo. Distr. — Brazil, Argentina. Ref. — Albuquerque, 1945b:1, 12 figs.
- petrochiae*, error.
- solitaria* Shannon & Del Ponte, 1926:565. In key only.
- POLYCHAETA (Stein), 1895:108 (*Homalomyia*). Type-locality: West and Central Europe, many localities. Distr. — Bermuda Is.; Europe. Ref. — Hennig, 1955:71, pl. 3, fig. 52, pl. 4, fig. 71. Erroneous record: probably refers to *brevipalpis* Chillcott, 1961:133, figs. 64, 67, 182, 267-268.
- ? *lepidi* Wiedemann, 1817:82 (*Anthomyia*). Type-locality: Holstein.
- PSEUDOFALCINCTA Albuquerque, 1954i:75, figs. 9-15. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil.



- PUNCTIPENNIS Albuquerque, 1954f:319, figs. 6-17. Type-locality: Brazil, Minas Gerais, Camanducaia. Distr. — Brazil.
- PUNCTIVENTRIS Malloch, 1934:207, fig. 32b. Type-locality: Chile, Santiago. Distr. — Chile, Juan Fernández Is. Ref. — Seago, 1954:7, figs. 11-13.
- PUSILLA (Bigot), 1885:304 (*Ophyra*). Type-locality: Chile. Distr. — Chile.
- PUSIO (Wiedemann), 1830:437 (*Anthomyia*). Type-locality: "South America". Distr. — Bermuda Is., Mexico, Guatemala, El Salvador, Guadalupe Is., Galápagos Is., Puerto Rico, Cuba, Bahamas, Panama, Trinidad, Venezuela, Guyana, Chile; North America, Hawaii, Micronesia. Ref. — Chillcott, 1961:213, figs. 144, 144A, 209, 280.
- femorata* Loew, 1861:42 (*Homalomyia*). Type-locality: "Cuba".
- RUFITIBIA Stein, 1911:104. Type-locality: Peru, Paso de Lares. Distr. — Peru.
- SAEROSKYI Seago, 1954:5, figs. 7 and 16. Type-locality: Guyana, Kaieteur. Distr. — Guyana.
- SCALARIS (Fabricius), 1794:332 (*Musca*). Type-locality: "Hafniae" (= Copenhagen). Distr. — Chile, Argentina; cosmopolitan, most common in temperate areas. Ref. — Hennig, 1955:81, figs. 5, 11, 14, pl. 1, fig. 16, pl. 4, fig. 67. Chillcott, 1961:61, figs. 4, 4A, 222, 233, 273.
- SCHNUSEI Stein, 1911:106. Type-locality: Chile, Valparaíso, Santiago and Quillota. Distr. — Argentina, Chile. Ref. — Malloch, 1934:205.
- SETOSA (Bigot), 1885:303 (*Ophyra*). Type-locality: "Chile". Distr. — Chile, Argentina. Ref. — Malloch, 1934:203.
- armata* Bigot, 1883:clxxxviii (*Dasyphyma*; preocc. Meigen, 1826). Type-locality: "Chile". N. SYN.
- errata* Malloch, 1934:203 (nom. nov. for *armata* Bigot). Type-locality: "Chile". N. SYN.
- erratica*, error.
- steinella* Albuquerque, 1953b:496 (nom. nov. for *armata* Bigot). Type-locality: "Chile". N. SYN.
- TRICHOPODA (Blanchard), 1942:57 (*Mesazelia*). Type-locality: Argentina, Formosa and Tucumán. Distr. — Argentina. N. COMB.
- TRIMACULATA (Stein), 1898:176 (*Homalomyia*). Type-locality: North America and Jamaica. Distr. — British Honduras, Panama, Puerto Rico, Jamaica, Dominican Republic, Haiti, S. Domingo, Ecuador, Peru, Venezuela, Brazil, Uruguay, Argentina. Ref. — Albuquerque, 1945a:1, 21 figs.
- TUCUMANENSIS Albuquerque, 1957g:2, figs. 1-6. Type-locality: Argentina, Tucumán, Villa Pedro Monte. Distr. — Argentina.
- TUMIDIFEMUR Stein, 1911:104. Type-locality: Bolivia, Sorata. Distr. — Mexico, Bolivia. Ref. — Albuquerque, 1957g:12, figs. 14-19.
- VITTATA Malloch, 1912:1. Type-locality: Panama, Porto Bello, Alhajuelo and Araján. Distr. — Mexico, Panama.
- XANTHOCERA Albuquerque, 1954f:317, figs. 1-5. Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Distr. — Brazil.
- YENHEDI Albuquerque, 1957g:16, figs. 20-26. Type-locality: Brazil, São Paulo, Ilha Sêca. Distr. — Brazil.

Subfamily MUSCINAE

Tribe MUSCINI

Genus MUSCA Linnaeus

- MUSCA** Linnaeus, 1758:589. Type-species, *domestica* Linnaeus (I.C.Z.N., Opinion 82).
- Promusca** Townsend, 1915a:434. Type-species, *domestica* Linnaeus (orig. des.).
- ALBINA** Wiedemann, 1830:415. Type-locality: "East Indies". Distr. — Ecuador; Africa, Arabia, Indian subregion, a semi-desert species. Almost certainly an erroneous record.
- DOMESTICA** Linnaeus, 1758:596. Type-locality: Europe and America. Distr. — Guadalupe Is., Mexico, Guatemala, San Salvador, Nicaragua, Costa Rica, Panama, Bermuda Is., Bahamas, Nassau, Jamaica, Puerto Rico, Haiti, Dominica, Cuba, Barbados, Mona Is., St. Lucia, St. Vincent, Virgin Is., Grenada, Trinidad, Venezuela, Guyana, Surinam, Colombia, Peru, Ecuador, Brazil, Uruguay, Argentina, Chile, Juan Fernández Is., Galápagos Is.; cosmopolitan. Ref. — Hennig, 1964:997, figs. 371c, 383, 384c, 395-400.
- domestica major* Degeer, 1776:71, pl. 4, figs. 1-11. Type-locality: Sweden and Surinam.
- stomoxidea* Robineau-Desvoidy, 1830:396. Type-locality: "Brazil".
- stomoxides*, error.
- aurulans* Robineau-Desvoidy, 1830:397. Type-locality: "Brazil and Cayenne". N. SYN.
- fulvescens* Robineau-Desvoidy, 1830:397. Type-locality: French Guiana, Cayenne. N. SYN.
- basilaris* Macquart, 1843a:310; 1843b:153. Type-locality: "Brazil".
- basalaris*, error.
- analís* Macquart, 1843a:311, pl. 21, fig. 2; 1843b:154, pl. 21, fig. 2. Type-locality: "Chile".
- chilensis* Macquart, 1843a:311, pl. 20, fig. 6; 1843b:154, pl. 20, fig. 6. Type-locality: "Chile". Nec *chilensis* Walker, 1837.
- chiliensis*, error.
- rufiventris* Macquart, 1843a:312, pl. 20, fig. 9; 1843b:155, pl. 20, fig. 9. Type-locality: "America". Nec *rufiventris* Scopoli, 1763. N. SYN.
- rufiventris* Macquart, 1846a:328; 1846b:200. Type-locality: "Brazil". Nec *rufiventris* Scopoli, 1763. N. SYN.
- pusilla* Macquart, 1848a:219, pl. 6, fig. 13; 1848b:59, pl. 6, fig. 13. Type-locality: "Haiti". Nec *pusilla* Gmelin, 1790. N. SYN.
- consanguinea* Rondani, 1848:78. Type-locality: "Brazil".
- vicina* Macquart, 1851a:226; 1851b:253. Type-locality: "America".
- flavinervis* Thomson, 1869:547. Type-locality: Ross Islands.
- atrifrons* Bigot, 1888a:607; 1888b:clxxxi. Type-locality: "Cuba and Mexico".
- pampasiana* Bigot, 1888a:607. Type-locality: Argentina, Buenos Aires.
- XANTHOMELAS** Wiedemann, 1824a:49. Type-locality: Java. Distr. — Haiti; Ethiopian and Oriental regions. An erroneous record based on confusion between *Musca pusilla* Macquart, 1848 (Haiti = *domestica* Linnaeus) and *Musca pusilla* Macquart, 1851 (Egypt = *xanthomelas* Wiedemann).



Genus MORELLIA Robineau-Desvoidy

- MORELLIA Robineau-Desvoidy, 1830:405. Type-species, *agilis* Robineau-Desvoidy (Townsend, 1916:8) = *hortorum* (Fallén).
- Parapyrellia* Townsend, 1915b:97. Type-species, *Musca violacea* Fabricius (orig. des.) = *maculipennis* (Macquart).
- Sarcopromusca* Townsend, 1927:209. Type-species, *arcuata* Townsend (mon.) = *pruna* (Shannon and Del Ponte).
- Biopyrellia* Townsend, 1932:105. Type-species, *Musca bipuncta* Wiedemann (orig. des.).
- Chaetopyrellia* Townsend, 1932:106. Type-species, *Pyrellia ochrifacies* Rondani (orig. des.) = *flavicornis* (Macquart).
- Neopyrellia* Enderlein, 1935:236. Type-species, *Pyrellia violacea* Robineau-Desvoidy (orig. des.) = *flavicornis* (Macquart).
- AFFINIS Malloch, 1925:86. Type-locality: Panama Canal Zone. Distr. — Panama.
- BASALIS (Walker), 1852:347 (*Musca*). Type-locality: "Brazil". Distr. — Mexico, El Salvador, Costa Rica, Jamaica, Cuba, Haiti, Puerto Rico, Virgin Is., Venezuela; North America. Ref. — Albuquerque, 1956a:20, figs.
- surrepens* Walker, 1861:312 (*Lucilia*). Type-locality: "Mexico". N. SYN.
- scapulata* Bigot, 1878:35 (*Pyrellia*). Type-locality: "Mexico". N. SYN.
- flora* Bigot, 1878:36 (*Pyrellia*). Type-locality: "Haiti". N. SYN.
- BIPUNCTA (Wiedemann), 1830:417 (*Musca*). Type-locality: "Brazil". Distr. — Mexico, Costa Rica, Panama, Venezuela, Trinidad Is., Bolivia, Brazil, Paraguay, Argentina. Ref. — Albuquerque, 1956a:6, figs.
- bipunctata*, error.
- suspicax* Walker, 1861:312 (*Pyrellia*). Type-locality: "Mexico".
- CALLIMERA (Bigot), 1888a:615; 1888b:clxxxii (*Curtonevra*). Type-locality: "Mexico". Distr. — Mexico.
- callidimera*, error.
- DIVERSIPALPIS (Rondani), 1864:30 (*Pyrellia*; nom. nov. for *rufipalpis* Macquart, 1855). Type-locality: "South America". Distr. — South America.
- rufipalpis* Macquart, 1855a:134; 1855b:114 (*Pyrellia*; preocc. Macquart, 1851). Type-locality: "South America".
- FLAVICORNIS (Macquart), 1848a:218, pl. 6, fig. 11; 1848b:58, pl. 6, fig. 11 (*Pyrellia*). Type-locality: "Brazil". Distr. — Mexico, Costa Rica, Jamaica, Colombia, Ecuador, Venezuela, Peru, Brazil, Paraguay, Argentina. Ref. — Albuquerque, 1956a:24, figs.
- violacea* Robineau-Desvoidy, 1830:463 (*Pyrellia*). Type-locality: "Brazil". Nec Fabricius, 1805. N. SYN.
- ochrifacies* Rondani, 1850:177 (*Pyrellia*). Type-locality: Brazil, São Paulo, São Sebastião Island. N. SYN.
- facilis* Walker, 1858:214 (*Pyrellia*). Type-locality: "South America". N. SYN.
- brevis* Schiner, 1868:303 (*Cyrtoneura*). Type-locality: "South America". N. SYN.

- chloe* Bigot, 1878:36 (*Pyrellia*). Type-locality: Ecuador, Quito. N. SYN.
iris Bigot, 1878:36 (*Pyrellia*). Type-locality: "Mexico". N. SYN.
aconquija Shannon & Del Ponte, 1926:581. Type-locality: Argentina, Tucumán City, Parque Aconquija. N. SYN.
- HUMERALIS** (Stein), 1918:204 (*Pyrellia*). Type-locality: Brazil, Teresópolis, Esp. Santo; Peru; Bolivia. Distr. — Peru, Bolivia, Brazil. Ref. — Albuquerque, 1956a:16, figs.
- MACULIPENNIS** (Macquart), 1846a:327; 1846b:199 (*Pyrellia*). Type-locality: "Colombia and Brazil". Distr. — Mexico, Cuba, Puerto Rico, Jamaica, Dominica, Trinidad Is., Guadalupe Is., Colombia, Venezuela, Guyana, Peru, Bolivia, Brazil, Paraguay. Ref. — Albuquerque, 1956a:10.
- violacea* Fabricius, 1805:288 (*Musca*). Type-locality: "South America". Nec Scopoli, 1763. N. SYN.
- nitida* Wiedemann, 1830:410 (*Musca*). Proposed in synonymy with *violacea* Fabricius. N. SYN.
- maculipennata* Macquart, 1851a:225, pl. 23, fig. 7; 1851b:252, pl. 23, fig. 7 (*Pyrellia*). Type-locality: Brazil, Minas Gerais. N. SYN.
- specialis* Walker, 1861:313 (*Pyrellia*). Type-locality: "Mexico". N. SYN.
- centralis* Loew, 1869:37 (*Pyrellia*). Type-locality: "Cuba". N. SYN.
- MERIDENSIS** (Macquart), 1846a:327; 1846b:199 (*Lucilia*). Type-locality: Mexico, Merida. Distr. — Mexico. N. COMB.
- NEOTROPICA** Curran, 1934b:462. Type-locality: Guyana, Kartabo. Distr. — Guyana.
- NIGRICOSTA** Hough, 1900:216, figs. 7 and 8. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães). Distr. — Costa Rica, Guyana, Peru, Brazil, Paraguay. Ref. — Albuquerque, 1956a:29, figs.
- nitida* Stein, 1918:205 (*Pyrellia*). Type-locality: Peru, Vilcanota; Paraguay, S. Bernadino; and Costa Rica, Higuito, San Mateo.
- OCHRICORNIS** (Wiedemann), 1830:408 (*Musca*). Type-locality: "Brazil". Distr. — Mexico, Cuba, Puerto Rico, Brazil, Galápagos Is. Ref. — Albuquerque, 1956a:33, figs.
- rufipalpis* Macquart, 1851a:225; 1851b:252 (*Pyrellia*). Type-locality: Brazil, Minas Gerais.
- obscuripes* Bigot, 1888a:616; 1888b:clxxxix (*Pyrellia*). Type-locality: "Mexico". N. SYN.
- PRUNA** (Shannon & Del Ponte), 1926:581 (*Orthellia*). Type-locality: Argentina, Jujuy, Calilegua. Distr. — Guatemala, Nicaragua, Costa Rica, Brazil, Argentina. N. COMB.
- arcuata* Townsend, 1927:355 (*Sarcopromusca*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. N. SYN.
- SARCOPHAGINA** Wulp, 1896:302, pl. 7, fig. 20. Type-locality: Mexico, Atoyac in Vera Cruz and Teapa in Tabasco. Distr. — Mexico, Panama, Colombia, Brazil, Argentina.
- SEMIMARGINATA** (Stein), 1918:206 (*Pyrellia*). Type-locality: Peru, Callanga and Bolivia, S. Antonio. Distr. — Peru, Bolivia.

Genus ORTHELLIA Robineau-Desvoidy

- ORTHELLIA** Robineau-Desvoidy, 1863:837. Type-species, *rectinervis* Robineau-Desvoidy (Townsend, 1916:8) = *cornicina* (Fabricius).



CAESARION (Meigen), 1826:57 (*Musca*). Type-locality: Portugal. Distr. — Mexico, Costa Rica, Brazil; Nearctic and Palaearctic regions, north India. Ref. — Hennig, 1963:930.

argentifera Bigot, 1877:251 (*Somomyia*). Type-locality: "Mexico".

CORNICINA (Fabricius), 1781:438 (*Musca*). Type-locality: Italy. Distr. — Mexico; Palaearctic region. Ref. — Hennig, 1963:933, figs. 366b, 373c, 375, 377. Mexico and North America are erroneous records of this exclusively Palaearctic species.

Genus DASYMORELLIA Malloch

DASYMORELLIA Malloch, 1923d:527. Type-species, *trichops* Malloch (orig. des.).
Chlorellia Shannon & Del Ponte, 1926:580. Type-species, *aenula* Shannon & Del Ponte (orig. des.) = *trichops* Malloch.

FULVITIBIA Snyder, 1949a:22. Type-locality: Guatemala, Guatemala City. Distr. — Mexico, Guatemala.

NIGRITIBIA Snyder, 1949a:24. Type-locality: Colombia, Medellín. Distr. — Colombia.

SAPHIRINA (Séguy), 1935b:115 (*Trichomorellia*). Type-locality: Ecuador, Quito. Distr. — Ecuador.

SPINIFERA (Wulp), 1883:39 (*Dasyphora*). Type-locality: "Argentina". Distr. — Argentina. N. comb.

TRICHOPS Malloch, 1923d:527. Type-locality: Peru, Ollantaybamba and Paltaybamba. Distr. — Colombia, Ecuador, Peru, Bolivia, Brazil, Argentina.

tricops, error.

cyanea (Macquart) of Stein, 1919:109 (misidentification).

aenula Shannon & Del Ponte, 1926:580 (*Chlorellia*). Type-locality: Argentina, San Pedro de Jujuy.

boliviana Townsend, 1931b:314 (*Trichomorellia*). Type-locality: Bolivia, Lorenzopata.

Genus XENOMORELLIA Malloch

XENOMORELLIA Malloch, 1923d:524. Type-species, *holti* Malloch (orig. des.).

HOLTI Malloch, 1923d:526. Type-locality: Brazil, Rio de Janeiro, Alto do Itatiaia. Distr. — Brazil.

MONTANHESA Albuquerque, 1952b:2, figs. 1-13. Type-locality: Brazil, Rio de Janeiro, Itatiaia, Macieira. Distr. — Brazil.

Genus NEOPYRELLIA, nom. nov.

Neopyrellia Townsend, 1939:446 (preocc. Enderlein, 1935). Type-species, *neglecta* Townsend (orig. des.).

NEOPYRELLIA, nom. nov. for *Neopyrellia* Townsend, 1939. Type-species, *Neopyrellia neglecta* Townsend (aut.).

NEGLECTA (Townsend), 1939:447 (*Neopyrellia*). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil. N. comb.

Genus CYACYRTONEURA Townsend

CYACYRTONEURA Townsend, 1931b:479. Type-species, *cyanea* Townsend (orig. des.).

Cyrtoneuropsis Townsend, 1931a:68 (preocc. Malloch, 1925). Type-species, *cyanea* Macquart, sensu Townsend not Macquart (orig. des.).

CYANEA Townsend, 1931b:479. Type-locality: Peru. Distr. — Venezuela, Peru, Brazil.

Tribe HYDROTAEINI

Genus REINWARDTIA Brauer & Bergenstamm

REINWARDTIA Brauer & Bergenstamm, 1889:158. Type-species, *tachinina* Brauer & Bergenstamm (mon.).

TACHININA Brauer & Bergenstamm, 1889:158 and 171. Type-locality: "Venezuela". Distr. — Ecuador, Venezuela.

Genus MUSCINA Robineau-Desvoidy

MUSCINA Robineau-Desvoidy, 1830:406. Type-species, *Musca stabulans* Fallén (Coquillett, 1910:571).

DORSILINEA (Wulp), 1896:308 (*Clinopera*). Type-locality: Mexico, Guerrero, Amula. Distr. — Mexico; North America. Ref. — Snyder, 1955:448.

aurantiaca Hough, 1899:25, fig. 6. Type-locality: North America, Georgia, Tifton.

PRINCIPALIS (Schiner), 1868:301 (*Spilogaster*). Type-locality: "South America". Distr. — South America. N. comb.

STABULANS (Fallén), 1817:252 (*Musca*). Type-locality: Sweden. Distr. — Mexico, Venezuela, Brazil, Uruguay, Argentina, Chile; cosmopolitan. Ref. — Hennig, 1962:766, figs. 239, 244, 311, pl. 32, fig. 626, pl. 33, fig. 632.

tibialis Walker, 1837:355 (*Musca*). Type-locality: Brazil, Santa Catarina. N. SYN.

vicina Macquart, 1843a:314, pl. 21, fig. 7; 1843b:157, pl. 21, fig. 7 (*Curtonevra*). Type-locality: "Chile".

Genus SYNTHESIOMYIA Brauer & Bergenstamm

SYNTHESIOMYIA Brauer & Bergenstamm, 1893:96 and 110. Type-species, *brasiliانا* Brauer & Bergenstamm (orig. des.) = *nudiseta* (Wulp).

NUDISETA (Wulp), 1883:42 (*Cyrtoneura*). Type-locality: "Argentina". Distr. — Bermuda Is., Mexico, Nicaragua, Virgin Is., Jamaica, St. Domingo, Puerto Rico, Trinidad, Guyana, Venezuela, Ecuador, Bolivia, Brazil, Paraguay, Argentina, Chile, Galápagos Is.; tropicopolitan. Ref. — Hennig, 1963:771.

brasiliانا Brauer & Bergenstamm, 1893:96 and 110. Type-locality: "Brazil".



grisea Giglio-Tos, 1893:5; 1896:8, fig. 2 (*Hyadesimyia*). Type-locality: Mexico, Oaxaca.

Genus AZELIA Robineau-Desvoidy

- AZELIA Robineau-Desvoidy, 1830:592. Type-species, *duodecimpunctata* Robineau-Desvoidy (Hennig, 1956:103) = *triquetra* (Wiedemann).
NEOTROPICA Snyder, 1957c:466. Type-locality: Argentina, Tucumán, Queb. Carnzo. Distr. — Argentina.

Genus POGONOMYIA Rondani

- POGONOMYIA Rondani, 1871:336. Type-species, *alpicola* Rondani (orig. des.).
ATERRIMA Wulp, 1896:335, pl. 8, figs. 13 and 13a. Type-locality: Mexico, Durango Ciudad. Distr. — Mexico.

Genus HYDROTAEA Robineau-Desvoidy

- HYDROTAEA Robineau-Desvoidy, 1830:509. Type-species, *Musca meteorica* Linnaeus (Curtis, 1839:no. 768).
ACUTA Stein, 1898:167. Type-locality: North America, Georgia. Distr. — Chile; North America. Ref. — Hockett, 1954:329.
CYANEIVENTRIS Macquart, 1851a:236; 1851b:263. Type-locality: "Chile". Distr. — Colombia, Bolivia, Chile, Argentina, Juan Fernández Is. Ref. — Malloch, 1934:301.
cyaniventris, error.
stuebeli Röder, 1886:269. Type-locality: Colombia, Páramo.
fuscisquama Aldrich, 1928:6. Type-locality: Chile, Perales, Quilpue.
CYRTONEURINA (Zetterstedt), 1845:1486 (*Aricia*). Type-locality: Denmark, Copenhagen. Distr. — Chile; Europe, Asia, North India. Ref. — Hennig, 1962:717. An erroneous record, referring to *cyaneiventris* Macquart.
DENTIPES (Fabricius), 1805:303 (*Musca*). Type-locality: Denmark. Distr. — Mexico, Costa Rica, Peru, Bolivia, Chile; North America, Palaearctic region, North India. Ref. — Hennig, 1962:718.
FUSCOCALYPTRATA Macquart, 1855a:136, pl. 6, fig. 16; 1855b:119, pl. 6, fig. 16. Type-locality: Nouvelle-Hollande (= Australia). Type-locality in error: this is a South American species.
HOUGHII Malloch, 1916:110. Type-locality: U.S.A., Illinois, Homer. Distr. — Ecuador, Venezuela; North America, Hawaii. Ref. — Hockett, 1954:334.
LATA (Walker), 1849:963 (*Eriphia*). Type-locality: U.S.A., St. Martin's Falls. Distr. — North America. South American records refer to *acuta* Stein.
NICHOLSONI Curran, 1939:2. Type-locality: Peru, Arequipa. Distr. — Peru, Bolivia, Brazil, Chile, Argentina.
NUBILICOSTA Malloch, 1923b:668. Type-locality: Colombia, Villavicencio, Quitiquia River. Distr. — Colombia.
VILLOSA Stein, 1904:449. Type-locality: Colombia, Cordillera and Bogotá. Distr. — Colombia.

Genus OPHYRA Robineau-Desvoidy


- OPHYRA Robineau-Desvoidy, 1830:516. Type-species, *nitida* Robineau-Desvoidy (Hennig, 1962:688) = *leucostoma* (Wiedemann).
- AENESCENS (Wiedemann), 1830:435 (*Anthomyia*). Type-locality: "New Orleans and West Indies". Distr. — Mexico, Nicaragua, Bermuda Is., St. Vincent, Puerto Rico, Jamaica, Trinidad, Tobago, Guyana, Venezuela, Ecuador, Peru, Bolivia, Brazil, Uruguay, Argentina, Chile, Galápagos Is.; North America, south Europe, eastern Pacific. Ref. — Sabrosky, 1949:427.
- virscens* Macquart, 1843a:321; 1843b:164. Type-locality: Brazil, Paraná, Guaratuba. N. SYN.
- setia* Walker, 1849:956 (*Anthomyia*). Type-locality: Galápagos Is.
- argentina* Bigot, 1885:302. Type-locality: Argentina, Buenos Aires.
- carbonaria* Shannon & Del Ponte, 1926:576. Type-locality: Argentina, Tucumán.
- CAPENSIS (Wiedemann), 1818:46 (*Anthomyia*). Type-locality: South Africa, Cape of Good Hope. Distr. — Chile; North America, Palaearctic and Ethiopian regions, North India. Ref. — Hennig, 1962:690, figs. 283, 288-290, 293.
- villosa* Aldrich, 1928:6. Type-locality: Chile, Perales.
- CHALCOGASTER (Wiedemann), 1824:52 (*Anthomyia*). Type-locality: Java. Distr. — Bermuda Is., Chile; Ethiopian, Oriental and Australasian regions. Ref. — Hennig, 1962:692, figs. 284 and 286.
- metallica* Wiedemann, 1830:435 (*Anthomyia*). Type-locality: East Indies.
- LEUCOSTOMA (Wiedemann), 1817:82 (*Anthomyia*). Type-locality: Holstein. Distr. — Venezuela; North America, Palaearctic region, North India. Ref. — Hennig, 1962:693, figs. 227, 235, 260, 282, 292, pl. 32, fig. 623, pl. 33, fig. 636.
- SOLITARIA Albuquerque, 1958f:7, figs. 12-16. Type-locality: Brazil, Goiás, Anápolis. Distr. — Brazil.
- SP. NOV. Distr. — Brazil.
- cutilia* (Walker) of Albuquerque, 1958f:2, figs. 1-11. Misidentification. True *cutilia* belongs to the genus *Psilochaeta* Stein.

Genus DENDROPHAONIA Malloch

- DENDROPHAONIA Malloch, 1923a:237. Type-species, *Spilogaster hilariformis* Stein (orig. des.) = *scabra* (Giglio-Tos).
- SCABRA (Giglio-Tos), 1893:9; 1896:25 (*Spilogaster*). Type-locality: "Mexico". Distr. — Mexico; North America. Ref. — Snyder, 1955:451.

Genus COENOSOPSIA Malloch

- COENOSOPSIA Malloch, 1924:74. Type-species, *prima* Malloch (orig. des.).
- PRIMA Malloch, 1924:74. Type-locality: Costa Rica, Higuito, San Mateo. Distr. — Mexico, British Honduras, Costa Rica, Panama, Brazil; North America.



Genus PHAONANTHO Albuquerque

PHAONANTHO Albuquerque, 1957b:191. Type-species, *devia* Albuquerque (orig. des.).

DEVIA Albuquerque, 1957b:191, figs. 1-12. Type-locality: Brazil, Guanabara, Rio de Janeiro, Corcovado, Reprêsa do Cabeça. Distr. — Brazil.

Subfamily PHAONIINAE

Tribe PHAONIINI

Genus PHAONIA Robineau-Desvoidy

PHAONIA Robineau-Desvoidy, 1830:482. Type-species, *viarum* Robineau-Desvoidy (Coquillett, 1901:140).

ADVENA Snyder, 1957c:482. Type-locality: Argentina, Tucumán, Burruyacu, V. Padre Monti. Distr. — Argentina.

BRASILIENSIS Albuquerque, 1958e:14, figs. 19-23. Type-locality: Brazil, São Paulo, São José dos Campos. Distr. — Brazil.

CACHEUTA Snyder, 1957c:482. Type-locality: Argentina, Mendoza, Cacheuta. Distr. — Argentina.

CATAMACLA Snyder, 1957c:483. Type-locality: Argentina, Avolfín, Catamarca. Distr. — Argentina.

CILITIBIA Albuquerque, 1955e:375, figs. 8-13. Type-locality: Brazil, Mato Grosso, Salobra. Distr. — Brazil.

COMPRESSIPALPIS Stein, 1911:66. Type-locality: Peru, Urubamba River, Umuhuankiali and Meshaga. Distr. — Peru.

ELONGATA Albuquerque, 1958e:21, figs. 28-32. Type-locality: Brazil, Minas Gerais, Viçosa. Distr. — Brazil.

FEMORATA Stein, 1911:66. Type-locality: Chile, Concepción and Quillota. Distr. — Chile.

FUSCISQUAMA (Wulp), 1896:340 (*Phorbia*). Type-locality: Mexico, Guerrero, Omilteme. Distr. — Mexico.

GALLICOLA Albuquerque, 1958e:5, figs. 7-13. Type-locality: Brazil, Guanabara, Rio de Janeiro, Reprêsa dos Ciganos. Distr. — Brazil.

LATINERVIS (Stein), 1904:419 (*Aricia*). Type-locality: Peru, Callanga and Bolivia, Coroico. Distr. — Peru, Bolivia, Brazil.

LENTIGINOSA Snyder, 1957c:482. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

LIMBINERVIS Stein, 1918:208. Type-locality: U.S.A., California and Mexico, Oaxaca. Distr. — Mexico; North America.

MACHADOI Albuquerque, 1958e:10, figs. 14-18. Type-locality: Brazil, Guanabara, Rio de Janeiro, Corcovado. Distr. — Brazil.

?MACULOSA Stein, 1911:64. Type-locality: Peru, Puerto Bermúdez and Urubamba River and Bolivia, Sarampiuni. Distr. — Peru, Bolivia.

MARGINATA Stein, 1918:209. Type-locality: "Peru". Distr. — Peru.

MINUSCULA Albuquerque, 1955e:373, figs. 1-7. Type-locality: Brazil, Rio de Janeiro, Itatiaia, Lago Azul. Distr. — Brazil.


- MONOCHAETA Snyder, 1957c:481. Type-locality: Argentina, Tucumán, Queb. La Toma. Distr. — Argentina.
- NERVICINCTA Stein, 1918:210. Type-locality: "Paraguay". Distr. — Paraguay.
- NOCTILUCA Albuquerque, 1958e:28, figs. 38-42. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil.
- PERUVIANA Malloch, 1929b:325. Type-locality: Peru, Perales. Distr. — Peru.
- PTEROSPILA Stein, 1918:211. Type-locality: Ecuador and Peru, Ollachea. Distr. — Ecuador, Peru.
- pterostila*, error.
- PULVILLATA (Stein), 1904:422 (*Aricia*). Type-locality: U.S.A., Carolina. Distr. — North America. Erroneously recorded from South America.
- SANTOAMARENSIS Albuquerque, 1958e:25, figs. 33-37. Type-locality: Brazil, São Paulo, São Paulo, Sto. Amaro. Distr. — Brazil.
- SIMPLEX Albuquerque, 1958e:18, figs. 24-27. Type-locality: Brazil, Guanabara, Rio de Janeiro, Jardim Botânico. Distr. — Brazil.
- TACHINOIDES Albuquerque, 1958e:2, figs. 1-6. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil.
- TETRAGONA Gaminara, 1930:1256. Type-locality: Uruguay, Pando and Prado de Montevideo. Distr. — Uruguay.
- TEXENSIS Malloch, 1923a:271. Type-locality: U.S.A., Texas, Brownsville. Distr. — Mexico; North America.
- TRIGONA Shannon & Del Ponte, 1926:576. Type-locality: Argentina, San Isidro. Distr. — Uruguay, Argentina.
- TRIGONOTA (Wulp), 1896:320, pl. 8, figs. 4 and 4a (*Spilogaster*). Type-locality: Nicaragua, Chontales. Distr. — Nicaragua. N. COMB.

Unplaced Species of PHAONIA s. lat.

- CANALICULATA Robineau-Desvoidy, 1830:486. Type-locality: "Brazil". Distr. — Brazil.

Genus PSILOCHAETA Stein

- PSILOCHAETA Stein, 1911:72. Type-species, *Limnophora chlorogaster* Bigot (orig. des.) = *chlorogaster* (Wiedemann).
- Phyronota* Shannon & Del Ponte, 1926:577. Type-species, *portensis* Shannon & Del Ponte (Séguy, 1937:329) = *chalybea* (Wiedemann).
- CARNIFEX Stein, 1911:74. Type-locality: Chile, Palca. Distr. — Chile.
- CHALYBEA (Wiedemann), 1830:428 (*Anthomyia*). Type-locality: "Brazil". Distr. — Colombia, Brazil, Uruguay, Chile, Argentina, Juan Fernández Is. Ref. — Malloch, 1934:314.
- coerulea* Macquart, 1843a:322, pl. 22, fig. 5; 1843b:165, pl. 22, fig. 5 (*Ophyra*). Type-locality: "Chile".
- caerulea*, error.
- cutilla* Walker, 1849:954 (*Anthomyia*). Type-locality: Uruguay, Montevideo. N. SYN.
- hirtula* Bigot, 1885:303 (*Ophyra*). Type-locality: "Chile".



- portensis* Shannon & Del Ponte, 1926:577 (*Phyronota*). Type-locality: Argentina, Adrogué.
- cyanea* error, Malloch, 1934:338 (for *chalybea*).
- CHLOROGASTER (Wiedemann), 1830:427 (*Anthomyia*). Type-locality: Uruguay, Montevideo. Distr. — Peru, Bolivia, Brazil, Uruguay, Chile, Argentina. Ref. — Malloch, 1934:314.
- chlorogaster* Bigot, 1885:271 (*Limnophora*). Type-locality: Argentina, Buenos Aires. Nec *chlorogaster* (Wiedemann), 1830.
- platensis* Shannon & Del Ponte, 1926:578 (*Phyronota*). Type-locality: Argentina, San Isidro.
- FULVOLATERALIS Stein, 1911:73. Type-locality: Peru, Lago Titicaca. Distr. — Peru.
- PAMPEANA (Shannon & Del Ponte), 1926:577 (*Phyronota*). Type-locality: Argentina, Tucumán. Distr. — Argentina.
- pampiana*, error.

Genus BRACHYGASTERINA Macquart

- BRACHYGASTERINA Macquart, 1851a:232; 1851b:259. Type-species, *violaceiventris* Macquart (orig. des.).
- MAJOR Malloch, 1934:339, fig. 60b. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Chile, Argentina.
- VIOLACEIVENTRIS Macquart, 1851a:232, pl. 23, fig. 17; 1851b:259, pl. 23, fig. 17. Type-locality: "Chile". Distr. — Chile, Argentina, Tierra del Fuego. Ref. — Malloch, 1934:338, fig. 60a.

Genus EUPHAONIA Malloch


- EUPHAONIA Malloch, 1922:271. Type-species, *fulvohumeralis* Malloch (orig. des.).
- FULVOHUMERALIS Malloch, 1922:271. Type-locality: Tierra del Fuego, Port Famine. Distr. — Tierra del Fuego.

Genus CORRENTOSIA Malloch

- CORRENTOSIA Malloch, 1934:335. Type-species, *bicolor* Malloch (orig. des.).
- BICOLOR Malloch, 1934:335. Type-locality: Argentina, Rio Negro, Lago Correntoso. Distr. — Argentina.

Genus PALPIBRACUS Rondani

- Brachypalpus* Macquart, 1851a:233; 1851b:260 (preocc. Macquart, 1834). Type-species, *pilosus* Macquart (orig. des.).
- PALPIBRACUS Rondani, 1864:35 (nom. nov. for *Brachypalpus* Macquart). Type-species, *Brachypalpus pilosus* Macquart (aut.).
- Darwinomyia* Malloch, 1922:277. Type-species, *univittata* Malloch (orig. des.) = *confusus* (Malloch).



- Acrolasia* Enderlein, 1927:54. Type-species, *Yetodesia chilensis* Bigot (orig. des.).
- Anacanthosphys* Enderlein, 1935:243. Type-species, *Darwinomyia trivittata* Malloch (orig. des.). N. SYN.
- Acanthosphys* Enderlein, 1935:244. Type-species, *Darwinomyia reynoldsi* Malloch (orig. des.). N. SYN.
- Catantigymnus* Enderlein, 1935:244. Type-species, *Helomyza veneris* Bigot (orig. des.). N. SYN.
- ANGOLENSIS (Malloch), 1934:334, fig. 58c (*Darwinomyia*). Type-locality: Chile, Angol. Distr. — Chile, Argentina. N. COMB.
- APICALIS (Malloch), 1934:328 (*Darwinomyia*). Type-locality: Argentina, Puerto Blest. Distr. — Chile, Argentina. N. COMB.
- CHILENSIS (Bigot), 1885:294 (?*Yetodesia*). Type-locality: "Chile". Distr. — Chile. Ref. — Malloch, 1934:320, fig. 56a. N. COMB.
- CONFUSUS (Malloch), 1928b:314 (*Darwinomyia*; nom. nov. for *univittata* Malloch). Type-locality: Argentina, Chubut, Valle del Lago Blanco. Distr. — Chile, Argentina, Tierra del Fuego. N. COMB.
- univittata* Malloch, 1922:278 (*Darwinomyia*; preocc. Bigot, 1857). Type-locality: Argentina, Chubut, Valle del Lago Blanco.
- FASCICULATUS (Malloch), 1934:332, fig. 59b (*Darwinomyia*). Type-locality: Argentina, Lake Correntoso. Distr. — Argentina. N. COMB.
- LANCIFER (Malloch), 1934:326, figs. 57a-b (*Darwinomyia*). Type-locality: Argentina, Lake Correntoso. Distr. — Argentina. N. COMB.
- MALLOCHI (Curran), 1934b:468 (*Darwinomyia*). Type-locality: Guyana, Kartabo. Distr. — Panama, Guyana. N. COMB.
- NIGRIVENTRIS (Malloch), 1928b:314 (*Darwinomyia*). Type-locality: "Chile". Distr. — Chile, Argentina, Tierra del Fuego. N. COMB.
- PILOSUS (Macquart), 1851a:233, pl. 23, fig. 18; 1851b:260, pl. 23, fig. 18 (*Brachypalpus*). Type-locality: Chile, Magellan Straits (not Oceania, Triton Bay, as stated by Macquart). Distr. — Chile. Ref. — Albuquerque, 1951b:7, figs. 9-14.
- REYNOLDSI (Malloch), 1934:323 (*Darwinomyia*). Type-locality: Tierra del Fuego, Rio Grande. Distr. — Tierra del Fuego. N. COMB.
- SEPARATUS (Malloch), 1934:324 (*Darwinomyia*). Type-locality: Chile, Concepción. Distr. — Chile. N. COMB.
- SIMILIS (Malloch), 1934:321 (*Darwinomyia*). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Chile, Argentina. N. COMB.
- SPICATUS (Malloch), 1934:329, fig. 58a (*Darwinomyia*). Type-locality: Chile, Casa Pangué. Distr. — Chile, Argentina. N. COMB.
- SUTILIS Albuquerque, 1951b:9, figs. 15-25. Type-locality: Chile, Perales. Distr. — Chile.
- TRIVITTATUS (Malloch), 1934:333, fig. 58b (*Darwinomyia*). Type-locality: Argentina, Lake Correntoso. Distr. — Chile, Argentina. N. COMB.
- UNIVITTATUS (Bigot), 1857:303, pl. 7, figs. 2 and 2a (*Pegomyia*). Type-locality: "Chile". Distr. — Chile. Ref. — Snyder, 1957c:489. N. COMB.
- VENERIS (Bigot), 1888c:34 (*Helomyza*). Type-locality: Tierra del Fuego, Orange Bay. Distr. — Chile, Argentina, Tierra del Fuego. Ref. — Malloch, 1934:325, fig. 56b. N. COMB.

atricornis Malloch, 1934: fig. 56b (*Darwinomyia*). Published only in the legend of this figure. N. SYN.

Genus BIGOTOMYIA Malloch

- BIGOTOMYIA Malloch, 1921a:173. Type-species, *Spilogaster trispilus* Bigot (orig. des.).
- Pseudomyiospila* Vimmer, 1939:61. Type-species, *coquilletti* Vimmer (mon.). N. SYN.
- ABDITA (Giglio-Tos), 1893:8; 1896:23 (*Spilogaster*). Type-locality: "Mexico". Distr. — Mexico. N. COMB.
- ANNULATA Albuquerque, 1957c:1, figs. Type-locality: Brazil, Rio de Janeiro, Itatiaia, Lago Azul. Distr. — Brazil.
- BIGOTI Albuquerque, 1957c:5, figs. Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil.
- BIPUNCTATA (Schiner), 1868:299 (*Spilogaster*). Type-locality: "South America". Distr. — South America. N. COMB.
- COQUILLETII (emend.) (Vimmer), 1939:61 (*Pseudomyiospila*). Type-locality: Peru, Puno. Distr. — Peru. N. COMB.
- coquilletti*, error.
- cocquillet*, error.
- COSTIPENNIS Snyder, 1957c:486. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- CURVATA (Stein), 1904:423 (*Aricia*). Type-locality: Colombia, Cordillera and Bogotá. Distr. — Colombia, Peru, Bolivia. Ref. — Albuquerque, 1954j:219, figs. 1-9.
- GRAJAUENSIS Albuquerque, 1957c:9, figs. Type-locality: Brazil, Guanabara, Rio de Janeiro, Grajaú. Distr. — Brazil.
- MEXALA Snyder, 1957c:484 (erroneously printed here as *praedatoria*: correction in 1957, *Bull. Am. Mus. nat. Hist.*, 113:viii). Type-locality: Guatemala, Antigua. Distr. — Mexico, Guatemala.
- NIGRIVENTRIS Albuquerque, 1954j:223, figs. 10-17. Type-locality: Brazil, Paraná. Distr. — Brazil.
- NIGROCINCTA (Stein), 1918:210 (*Phaonia*). Type-locality: "Mexico, Jalapa and Venezuela". Distr. — Mexico, Venezuela. N. COMB.
- PRAEDATORIA Snyder, 1957c:485 (the *praedatoria* described on page 484 is *mezala* Snyder: see 1957, *Bull. Am. Mus. nat. Hist.*, 113:viii). Type-locality: Argentina, Tucumán, Queb. La Toma. Distr. — Argentina.
- PRAESUTURALIS (Stein), 1904:424 (*Aricia*). Type-locality: Peru, Callanga and Bolivia, Suapi. Distr. — Peru, Bolivia. N. COMB.
- presuturalis*, error.
- PUNCTINERVIS (Stein), 1911:63 (*Phaonia*). Type-locality: Bolivia, Lago Titicaca. Distr. — Bolivia, Argentina. Ref. — Snyder, 1957c:488.
- nigripuncta* Stein, 1911:63 (*Phaonia*). Type-locality: Bolivia, Lago Titicaca and Sorata.
- SIMILATA Albuquerque, 1957c:13, figs. Type-locality: Brazil, Rio de Janeiro, Jussara, Angra dos Reis. Distr. — Brazil.
- TRISPILA (Bigot), 1885:285 (*Spilogaster*). Type-locality: Argentina, Buenos Aires. Distr. — Venezuela, Brazil, Uruguay, Argentina. Ref. — Snyder, 1957c:487.



VENICURVA (Stein), 1904:417 (*Aricia*). Type-locality: Peru, Callanga and Bolivia, Sillutincara and Coroico. Distr. — Peru, Bolivia.

Genus HELINA Robineau-Desvoidy

HELINA Robineau-Desvoidy, 1830:493. Type-species, *euphemioidea* Robineau-Desvoidy (Coquillett, 1901:137) = *pertusa* (Meigen).

Ariciella Malloch, 1918a:66. Type-species, *flavicornis* Malloch (orig. des.) = *rubripalpis* (Wulp).

Tritonidis Séguy, 1937:461. Type-species, *Aricia nigrimanus* Macquart (orig. des.).

Trinodites, error.

ACHINIS Snyder, 1941:12. Type-locality: Peru, Chosica. Distr. — Ecuador, Peru.

ACROSTICHALIS Snyder, 1941:13. Type-locality: Ecuador, Quito. Distr. — Ecuador.

ACZELI Snyder, 1957c:477. Type-locality: Argentina, Salta, San Lorenzo. Distr. — Argentina.

ADELPHA (Schiner), 1868:300 (*Spilogaster*). Type-locality: "South America". Distr. — ?Mexico, Colombia, Peru. Ref. — Snyder, 1941:13.

ALBUQUERQUEI, nom. nov. for *discreta* Albuquerque. Type-locality: Brazil, Rio de Janeiro, Teresópolis, Parque Nacional da Serra dos Órgãos. Distr. — Brazil.

discreta Albuquerque, 1956j:11, figs. (preocc. Wulp, 1896). Type-locality: Brazil, Rio de Janeiro, Teresópolis, Parque Nacional da Serra dos Órgãos.

ANGUSTIPENNIS (Stein), 1911:79 (*Mydaea*). Type-locality: Peru, Urubamba River and Pachitea River. Distr. — Panama, Peru, Bolivia, Brazil. Ref. — Albuquerque, 1956j:1.

ANTHRACINA Albuquerque, 1956j:8, figs. Type-locality: Brazil, Rio de Janeiro, Itatiaia, Macieira. Distr. — Brazil.

ANUBES Snyder, 1941:18. Type-locality: Ecuador, Chimborazo, Urbino Cerro. Distr. — Ecuador.

ARGENTINA Snyder, 1957c:476. Type-locality: Argentina, Tucumán, V. Padre Monti, Burruyacu. Distr. — Argentina.

BIGOTI Malloch, 1934:312. Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.

BISETA (Stein), 1904:444 (*Spilogaster*). Type-locality: Peru, Callanga. Distr. — Peru. N. COMB.

BREVIVENA Snyder, 1941:19. Type-locality: Ecuador, Chimborazo, Cumbre Tililac. Distr. — Ecuador.

BROWN Snyder, 1941:17. Type-locality: Ecuador, Uyambicho. Distr. — Ecuador.

CANEO Snyder, 1941:10. Type-locality: Mexico, Mexico City. Distr. — Mexico.

CHILENSIS Malloch, 1934:312. Type-locality: Chile, Perales. Distr. — Chile.

CIRCULATRIX (Walker), 1861:316 (*Aricia*). Type-locality: "Mexico". Distr. — Mexico, Guatemala. Ref. — Snyder, 1940a:2.



- marginata* Snyder, 1940a:2. Type-locality: Guatemala, La Providencia, Obispo.
- CONFINIS (Wulp), 1896:319 (*Mydaea*). Type-locality: Mexico, Orizaba. Distr. — Mexico, Guatemala. Ref. — Snyder, 1940a:5.
- fulvapoda* Snyder, 1940a:5. Type-locality: Guatemala, La Providencia, Obispo.
- CONNEXA Malloch, 1934:309, figs. 55b-c. Type-locality: Argentina, Lake Correntoso. Distr. — Chile, Argentina.
- CONSANGUINEA (Stein), 1911:81 (*Mydaea*). Type-locality: Peru, Paso de Lares, Urubamba River and Chanchamayo. Distr. — Peru. N. comb.
- COPIOSA (Wulp), 1896:321, pl. 8, figs. 5 and 5a (*Spilogaster*). Type-locality: Mexico, Xucumanatlán, Omilteme and Sierra de las Aguas Escondidas. Distr. — Mexico. Ref. — Snyder, 1941:11.
- CREPEDOSETA Snyder, 1940a:3. Type-locality: Guatemala, La Providencia, Obispo. Distr. — Mexico, Guatemala, Costa Rica, Panama.
- CROCEA Snyder, 1940a:6. Type-locality: Brazil, Goiás, Campinas. Distr. — Brazil, Argentina.
- CYANICOLOR (Stein), 1911:65 (*Phaonia*). Type-locality: Bolivia, Cordillera. Distr. — Bolivia. N. comb.
- DASYOPHITHALMA Malloch, 1928a:473 (nom. nov. for *dasyops* Stein, 1911). Type-locality: Peru, Tarma, Cuzco, Lago Titicaca; Bolivia, Cordillera. Distr. — Peru, Bolivia.
- dasyops* Stein, 1911:82 (*Mydaea*; preocc. Macquart, 1843). Type-locality: Peru, Tarma, Cuzco and Lago Titicaca; Bolivia, Cordillera.
- DASYOPS (Macquart), 1843a:326, pl. 22, fig. 1; 1843b:169, pl. 22, fig. 1 (*Chortophila*). Type-locality: Falkland Islands. Distr. — Falkland Islands. Ref. — Malloch, 1928a:472.
- DISCOLOR (Stein), 1911:83 (*Mydaea*). Type-locality: Peru, Oroya and Cuzco. Distr. — Peru, Argentina. Ref. — Malloch, 1934:305.
- DISCRETA (Wulp), 1896:322 (*Spilogaster*). Type-locality: Mexico, Amula and Xucumanatlán. Distr. — Mexico, Jamaica. N. comb.
- DUBIA (Bigot), 1885:294 (*Yetodesia*). Type-locality: "Chile". Distr. — Chile. Ref. — Malloch, 1934:304.
- ECHINOASTER (Stein), 1911:78 (*Mydaea*). Type-locality: Peru, Cuzco. Distr. — Peru. N. comb.
- EQUATOR Snyder, 1941:18. Type-locality: Ecuador, Pichincha Prov. Distr. — Ecuador.
- ETESIA (Giglio-Tos), 1893:8; 1896:23 (*Spilogaster*). Type-locality: Mexico, Tuxpango. Distr. — Mexico. N. comb.
- EURYCEPHALA (Stein), 1911:80 (*Mydaea*). Type-locality: Peru, Urubamba River and Pachitea River. Distr. — Peru. N. comb.
- FALLAX (Stein), 1911:84 (*Mydaea*). Type-locality: Peru, Cuzco. Distr. — Peru. N. comb.
- FULVOCALYPTRATA Malloch, 1934:311. Type-locality: Argentina, Lake Correntoso. Distr. — Chile, Argentina.
- fulvisquama* Stein, 1904:447 (*Spilogaster*; preocc. Zetterstedt, 1845). Type-locality: Chile. N. syn.
- GIGANTEA Albuquerque, 1956j:15, figs. Type-locality: Brazil, Rio de Janeiro, Angra dos Reis. Distr. — Brazil.

- GRANDIS (Schiner), 1868:302 (*Spilogaster*). Type-locality: "South America". Distr. — South America. N. comb.
- INEPTA (Stein), 1911:81 (*Mydaea*). Type-locality: Peru, Calca and Cuzco. Distr. — Peru. Ref. — Snyder, 1941:20.
- LAETIFICA (Robineau-Desvoidy), 1830:500 (*Mydina*). Type-locality: France, Paris. Distr. — Mexico, Venezuela; North America, Europe, Asia, India. Ref. — Hennig, 1957:188, pl. 9, fig. 171.
- lucorum* Fallén, 1823:55 (*Musca*; preocc. Linnaeus, 1758). Type-locality: Sweden.
- abacta* Giglio-Tos, 1893:8; 1896:21 (*Hyetodesia*). Type-locality: "Mexico".
- abdicta* Giglio-Tos, 1893:8; 1896:21 (*Hyetodesia*). Type-locality: "Mexico". N. SYN.
- LASIOSTERNA Snyder, 1941:9. Type-locality: Mexico, Las Visayas, Chic. Distr. — Mexico; North America.
- LATIPENNIS (Stein), 1904:435 (*Spilogaster*). Type-locality: Colombia, Cordillera and Bolivia, San Antonio. Distr. — Colombia, Bolivia. N. comb.
- LEUCOCEPHALA (Wulp), 1896:318 (*Mydaea*). Type-locality: Mexico, Tabasco, Teapa. Distr. — Mexico, Panama. Ref. — Snyder, 1940a:4.
- compressifrons* Snyder, 1940a:4. Type-locality: Panama, Barro Colorado Island.
- LONGIPILA (Stein), 1918:217 (*Mydaea*). Type-locality: Peru, Ollachea. Distr. — Peru.
- LUTEOLA Albuquerque, 1956d:167, figs. 1-9. Type-locality: Brazil, Minas Gerais. Distr. — Brazil.
- LYNCHII (Wulp), 1883:43 (*Limnophora*). Type-locality: "Argentina". Distr. — Argentina. N. comb.
- MACULIPENNIS (Zetterstedt), 1845:1475 (*Aricia*). Type-locality: Sweden, Lule Lappmark, Kvikkjokk and Dalecarlia. Distr. — Brazil; North America, Europe, Asia. Ref. — Hennig, 1958:195, pl. 9, fig. 181. Almost certainly an erroneous record.
- poeciloptera* Malloch, 1918b:271 (*Aricia*; preocc. Schiner, 1868). Type-locality: U.S.A., New Mexico, Cloudcroft.
- MACULIPES (Stein), 1918:218 (*Mydaea*). Type-locality: Mexico, Jalapa. Distr. — Mexico. N. comb.
- MACULOSA (Walker), 1852:357 (*Anthomyia*). Type-locality: "Brazil". Distr. — Brazil, Paraguay. Ref. — Stein, 1901:201. N. comb.
- MARGINIPENNIS (Stein), 1904:440 (*Spilogaster*). Type-locality: Colombia, Cordillera. Distr. — Colombia, Peru, Bolivia. Ref. — Snyder, 1941:7.
- MERACA (Wulp), 1896:325 (*Limnophora*). Type-locality: Mexico, Omilteme. Distr. — Mexico. N. comb.
- MONACHA (Schiner), 1868:301 (*Spilogaster*). Type-locality: "South America". Distr. — South America. N. comb.
- MULCATA (Giglio-Tos), 1893:7; 1896:20, fig. 5 (*Hyetodesia*). Type-locality: "Mexico". Distr. — Mexico; ?North America.
- NEOSIMPLEX Snyder, 1957c:478. Type-locality: Argentina, Santa Cruz, Lago Argentina. Distr. — Argentina.



- NEOTROPICA (Snyder), 1951:7 (*Helinella*). Type-locality: Peru, Dept. of Junín, Chanchamayo. Distr. — Peru. N. comb.
- NIGRIMANA (Macquart), 1851a:231, pl. 23, fig. 15; 1851b:258, pl. 23, fig. 15 (*Aricia*). Type-locality: "South America" (not Oceania, Triton Bay, as stated by Macquart). Distr. — Chile, Argentina. Ref. — Malloch, 1934:307.
- bifimbriata* Malloch, 1934:307. Type-locality: Argentina, Lake Gutiérrez.
- SSP. BASALIS Malloch, 1934:308, fig. 55a. Type-locality: Argentina, Puerto Blest. Distr. — Chile, Argentina.
- SSP. GRISEA Malloch, 1934:308. Type-locality Argentina, Puerto Blest. Distr. — Argentina.
- NIGROMARGINATA (Stein), 1911:78 (*Mydaea*). Type-locality: Bolivia, Lorenzopata. Distr. — Bolivia. N. comb.
- NIVALOIDES Albuquerque, 1956j:5, figs. Type-locality: Brazil, Rio de Janeiro, Itatiaia, Macieira. Distr. — Brazil.
- NOBILIS Albuquerque, 1956d:171, figs. 10-18. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil.
- NOTHA Snyder, 1941:20. Type-locality: Ecuador, Miasa Ridge, Volcán Tungurahua. Distr. — Ecuador.
- PARSURA (Giglio-Tos), 1893:8; 1896:20 (*Hyetodesia*). Type-locality: "Mexico". Distr. — Mexico. N. comb.
- PARVULA (Wulp), 1896:321 (*Spilogaster*). Type-locality: Mexico, Yepetlapa, Amula and Omilteme. Distr. — Mexico. Ref. — Snyder, 1941:11.
- PILICEPS (Stein), 1911:84 (*Mydaea*). Type-locality: Bolivia, Sorata and Mapiri. Distr. — Bolivia.
- POECILOPTERA (Schiner), 1868:300 (*Spilogaster*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Snyder, 1941:15.
- PRAECIPUA (Walker), 1852:351 (*Anthomyia*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Snyder, 1940a:7.
- unicolor* Walker, 1852:353 (*Anthomyia*). Type-locality: "South America".
- ignava* Walker, 1858:217 (*Aricia*). Type-locality: "Brazil".
- pleuraseta* Snyder, 1940a:7. Type-locality: Brazil, São Paulo, São José dos Campos.
- PROCEDENS (Walker), 1861:315 (*Aricia*). Type-locality: "Mexico". Distr. — Mexico; North America. Ref. — Stein, 1901:205.
- PROLATIFRONS Snyder, 1940a:6. Type-locality: Panama, Barro Colorado Island. Distr. — Panama.
- REFUSA (Giglio-Tos), 1893:8; 1896:22 (*Spilogaster*). Type-locality: Mexico, Tuxpango. Distr. — Mexico, Costa Rica. Ref. — Snyder, 1941:14.
- decora* Snyder, 1941:14. Type-locality: Mexico, Huipulco.
- REGOBARROSI Albuquerque, 1958c:47, figs. 1-6. Type-locality: Brazil, Rio de Janeiro, Itatiaia, Lago Azul. Distr. — Brazil.
- RUBRIFALPIS (Wulp), 1896:320 (*Spilogaster*). Type-locality: Mexico, Dos Arroyos, Tepetlapa, Sierra de las Aguas Escondidas and Cuernavaca; Costa Rica, Caché. Distr. — Mexico, Costa Rica; North America. Ref. — Malloch, 1918a:66. N. comb.

- flavicornis* Malloch, 1918a:66 (*Ariciella*). Type-locality: U.S.A., Texas, Brownsville.
- RUFOAPICATA Malloch, 1934:310. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.
- SERA (Giglio-Tos), 1893:9; 1896:25 (*Spilogaster*). Type-locality: "Mexico". Distr. — Mexico. Ref. — Snyder, 1949a:6.
- arroya* Snyder, 1949a:6. Type-locality: Mexico, Mexico City.
- SIGNATIPENNIS (Wulp), 1896:322, pl. 8, figs. 6 and 6a (*Spilogaster*). Type-locality: Mexico, Chilpancingo, Xucumanatlán, Omilteme, Sierra de las Aguas Escondidas and Orizaba. Distr. — Mexico. Ref. — Snyder, 1941:16.
- SIMPLEX Malloch, 1934:309, figs. 55d-e. Type-locality: "Chile or Argentina". Distr. — Chile or Argentina.
- SOCIA (Wulp), 1896:326 (*Limnophora*). Type-locality: Mexico, Omilteme and Sierra de las Aguas Escondidas. Distr. — Mexico. Ref. — Snyder, 1941:11.
- TARSALIS (Stein), 1918:220 (*Mydaea*). Type-locality: Mexico, mountains near Cuernavaca. Distr. — Mexico. N. comb.
- TESSELATA (Stein), 1904:438 (*Spilogaster*). Type-locality: "Brazil". Distr. — Peru, Brazil. N. comb.
- TOWNSENDI Snyder, 1941:21. Type-locality: Peru, Huariaca. Distr. — Peru.
- TRICHOPS (Stein), 1918:222 (*Mydaea*). Type-locality: Peru, Matucana. Distr. — Peru. N. comb.
- TRICOLOR (Walker), 1852:358 (*Anthomyia*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Stein, 1901:214. N. comb.
- UMBROSA (Wulp), 1896:326 (*Limnophora*). Type-locality: Costa Rica, Rio Sucio and Volcán de Irazú. Distr. — Costa Rica, Venezuela.
- VENISETA Dodge, 1967:242. Type-locality: Chile, Santiago, Las Condes. Distr. — Chile.
- VIERECKI Snyder, 1941:8. Type-locality: Colombia, La Cumbee. Distr. — Colombia.
- VIOLA Malloch, 1934:303, fig. 54. Type-locality: Argentina, Lake Gutiérrez. Distr. — Argentina.
- VIOLESCENS Dodge, 1967:241. Type-locality: Chile, Santiago, Las Condes. Distr. — Chile.
- XENA Malloch, 1934:305. Type-locality: Chile, Casa Pangue. Distr. — Chile, Argentina.

Genus SOUZALOPESMYIA Albuquerque

- Peruvia* Malloch, 1929a:104 (proc. Scudder, 1890). Type-species, *Mydaea singularis* Stein (orig. des.). N. syn.
- SOUZALOPESMYIA Albuquerque, 1951a:53. Type-species, *carioca* Albuquerque (orig. des.).
- AMAZONICA Albuquerque, 1951a:56, figs. 11-12 and 16-18. Type-locality: Brazil, Amazonas, Manaus. Distr. — Brazil.
- CARIOCA Albuquerque, 1951a:53, figs. 1-10 and 13-15. Type-locality: Brazil, Guanabara, Rio de Janeiro, Grajaú. Distr. — Brazil.



SINGULARIS (Stein), 1911:91 (*Mydaea*). Type-locality: Bolivia, Sarampiuni.
Distr. — Peru, Bolivia. Ref. — Malloch, 1929a:105. N. comb.

Genus GYMNOIDIA Robineau-Desvoidy

- GYMNOIDIA Robineau-Desvoidy, 1863:635. Type-species, *pratensis* Robineau-Desvoidy (mon.) = *polystigma* (Meigen).
ARCUATA (Stein), 1898:201 (*Limnophora*). Type-locality: U.S.A., Georgia.
Distr. — Mexico, Puerto Rico, Peru, Bolivia, Brazil, Chile; North America, Hawaii.
AUCTILA (Stein), 1901:187 (*Limnophora*). Type-locality: "Brazil". Distr. — Brazil. N. comb.
DEBILIS (Williston), 1896:369 (*Limnophora*). Type-locality: St. Vincent Island. Distr. — Mexico, St. Vincent, Jamaica, Puerto Rico, Venezuela; U.S.A. Ref. — Hockett, 1932:54.
dorsovittata Malloch, 1920:146 (*Eulimnophora*). Type-locality: Jamaica, Kingston.
DELECTA (Wulp), 1896:325 (*Limnophora*). Type-locality: Mexico, Guerrero, Amula and Rincon. Distr. — Mexico. N. comb.
OVATIVENTRIS (Macquart), 1851a:240, pl. 24, fig. 12; 1851b:267, pl. 24, fig. 12 (*Anthomyia*). Type-locality: Uruguay, Montevideo. Distr. — Uruguay. Ref. — Albuquerque, 1950b:5, figs. 17-18.
SCATOPHAGA Curran, 1938:5. Type-locality: Haiti. Distr. — Haiti.

Genus PACHYCERAMYIA Albuquerque

- PACHYCERAMYIA Albuquerque, 1955d:253. Type-species, *enigmatica* Albuquerque (orig. des.).
ENIGMATICA Albuquerque, 1955d:253, figs. 1-6. Type-locality: Brazil, Guanabara, Rio de Janeiro, Manginhos. Distr. — Brazil.

Genus SCENETES Malloch

- SCENETES Malloch, 1936:9. Type-species, *cardini* Malloch (orig. des.).
CARDINI Malloch, 1936:10. Type-locality: Cuba, Santiago de las Vegas. Distr. — Cuba, Puerto Rico.

Tribe ATHERIGONINI

Genus ATHERIGONA Rondani

- ATHERIGONA Rondani, 1856:97. Type-species, *Anthomyia varia* Meigen (orig. des.).
Acritochaeta Grimshaw, 1901:41. Type-species, *pulvinata* Grimshaw (mon.) = *orientalis* Schiner.
ORIENTALIS Schiner, 1868:295. Type-locality: Nicobar Is., Tellnschong. Distr. — Mexico, El Salvador, Panama, Bermuda Is., Virgin Is., Puerto Rico, Barbados, Cuba, Colombia, Ecuador, Peru, Venezuela, Brazil, Galapagos Is.; cosmopolitan. Ref. — Hennig, 1961:500.

- excisa* Thomson, 1869:560 (*Coenosia*). Type-locality: Ross Island.
trilineata Stein, 1900a:157. Type-locality: New Guinea, Friedrich
 Wilhelmshafen (= Madang).
pulvinata Grimshaw, 1901:42, pl. 2, figs. 21-23 (*Acritochaeta*). Type-
 locality: Hawaii, Olaa.

Subfamily MYDAEINAE

Genus MYDAEA Robineau-Desvoidy

- MYDAEA Robineau-Desvoidy, 1830:479. Type-species, *scutellaris* Robineau-
 Desvoidy (Coquillett, 1901:139).
 CARAS Snyder, 1949a:5, figs. 1 and 3. Type-locality: Venezuela, Caracas.
 Distr. — Venezuela.
 CASTANEA Albuquerque, 1955e:377, figs. 14-20. Type-locality: Brazil, Gua-
 nabara, Rio de Janeiro, Tijuca. Distr. — Brazil.
 CENTRALIS Stein, 1911:85. Type-locality: Peru, Calca and Chile, Palca. Distr.
 — Peru, Chile.
 COMPRESSICORNIS Snyder, 1949a:4, figs. 2, 4 and 5. Type-locality: Colombia,
 Medellín. Distr. — Colombia.
 CRESA Snyder, 1949a:1. Type-locality: Mexico, Cuernavaca. Distr. — Mexico.
 FUMICOSTA Snyder, 1941:1. Type-locality: Ecuador, Prov. Bolívar, Hda.
 Talahua. Distr. — Ecuador.
 FUMIPENNIS (Wulp), 1896:324, pl. 8, fig. 7 (*Limnophora*). Type-locality:
 Mexico, Guerrero, Sierra de las Aguas Escondidas. Distr. —
 Mexico. N. comb.
 INSONS (Giglio-Tos), 1893:8; 1896:21 (*Hyetodesia*). Type-locality: "Mexico".
 Distr. — Mexico. N. comb.
 LATOMENSIS Snyder, 1957c:479. Type-locality: Argentina, Tucumán, Queb.
 La Toma. Distr. — Argentina.
 LONGISETA (Wulp), 1896:315 (*Hyetodesia*). Type-locality: Mexico, Guerrero,
 Sierra de las Aguas Escondidas. Distr. — Mexico. N. comb.
 MERIDA Snyder, 1941:3. Type-locality: Mexico, Coapa. Distr. — Mexico.
merida, error.
 NUBIVENA Snyder, 1941:2. Type-locality: Ecuador, Baños. Distr. — Vene-
 zuela, Ecuador, Bolivia.
 ORBA Stein, 1911:85. Type-locality: Bolivia, Lorenzopata. Distr. — Bolivia.
 PANSIA (Giglio-Tos), 1893:9; 1896:24, fig. 3 (*Spilogaster*). Type-locality:
 "Mexico". Distr. — Mexico.
 PAUCISETA Stein, 1911:77. Type-locality: Peru, Pichís. Distr. — Peru.
 PLAUMANNI Snyder, 1941:3. Type-locality: Brazil, Santa Catarina, Nova
 Teutônia. Distr. — Mexico, Brazil, Paraguay.
 SEXPUNCTATA (Wulp), 1883:43 (*Spilogaster*). Type-locality: "Argentina".
 Distr. — Argentina.
 TRISETA Curran, 1931:13. Type-locality: Brazil, Mato Grosso, Chapada (dos
 Guimarães). Distr. — Brazil.



Unplaced species of MYDAEA s. lat.

- ARCUATA (Wiedemann), 1830:425 (*Anthomyia*). Type-locality: "Brazil". Distr. — Brazil. Genus uncertain, probably Phaoniini.
- CONVEXA Stein, 1918:214. Type-locality: "Mexico". Distr. — Mexico. Genus uncertain, probably Cyrtoneurinae.
- DICHROMA (Wiedemann), 1830:425 (*Anthomyia*). Type-locality: "Brazil". Distr. — Brazil. Genus uncertain, probably Cyrtoneurinae or Phaoniini.
- INERMIS (Stein), 1904:433 (*Spilogaster*). Type-locality: Bolivia, S. Antonio and Coroico. Distr. — Bolivia. Genus uncertain, probably Cyrtoneurinae.
- LATIPENNIS (Hough), 1900:229, figs. 16 and 17 (*Muscina*). Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães). Distr. — Brazil. Genus uncertain, probably Cyrtoneurinae.
- LIMBATA (Wiedemann), 1830:425 (*Anthomyia*). Type-locality: "Brazil". Distr. — Brazil. Genus uncertain, probably Cyrtoneurinae or Phaoniini.
- MARGINALIS (Stein), 1904:445 (*Spilogaster*). Type-locality: Bolivia, Sillutincara. Distr. — Bolivia. Genus uncertain, possibly *Helina* Robineau-Desvoidy.
- NIGRINA (Wiedemann), 1830:424 (*Anthomyia*). Type-locality: "Brazil". Distr. — Peru, Brazil. Genus uncertain, probably Cyrtoneurinae or Phaoniini.
- PEDELLA (Wiedemann), 1830:423 (*Anthomyia*). Type-locality: "Brazil". Distr. — Peru, Brazil. Genus uncertain, probably Cyrtoneurinae or Phaoniini.
- calliphoroides* Jaenicke, 1867a:371; 1867b:63 (*Spilogaster*). Type-locality: "Brazil".

Genus MYOSPILA Rondani

- MYOSPILA Rondani, 1856:91. Type-species, *Musca mediatubunda* Fabricius (orig. des.).
- Myiospila*, error.
- Phasiophana* Brauer & Bergenstamm, 1891:390. Type-species, *obsoleta* Brauer & Bergenstamm (orig. des.).
- Trichomorellia* Stein, 1919:109. Type-species, *cyanea* (Macquart) (mon.).
- ABERRANS (Shannon & Del Ponte), 1926:579 (*Muscina*). Type-locality: Argentina, Tucumán. Distr. — Argentina.
- CYANEA (Macquart), 1843a:314, pl. 21, fig. 6; 1843b:157, pl. 21, fig. 6 (*Curtonevra*). Type-locality: Chile, Concepción. Distr. — Argentina, Chile. Ref. — Snyder, 1940b:6.
- semiclausula* Schiner, 1868:302 (*Aricia*). Type-locality: "Chile".
- semiclausula*, error.
- MEDIATUBUNDA (Fabricius), 1781:444 (*Musca*). Type-locality: Italy. Distr. — Mexico, Guatemala, Ecuador, Peru, Bolivia; North America; Palaearctic region. Ref. — Hennig, 1956:114, figs. 29 and 32.



- quadristigma* Thomson, 1869:551 (*Anthomyia*). Type-locality: Ecuador, Puna.
- quadrisetosa*, error.
- tetrastigma*, error.
- anthomydaea* Bigot, 1888a:614; 1888b:clxxxii (*Curtonevra*). Type-locality: U.S.A., Rocky Mountains.
- anthomyidaea*, error.
- nigriceps* Bigot, 1888a:615; 1888b:clxxxii (*Curtonevra*). Type-locality: U.S.A., Rocky Mountains.
- OBSCURA (Shannon & Del Ponte), 1926:580 (*Muscina*). Type-locality: Argentina, Buenos Aires. Distr. — Peru, Brazil, Argentina.
- OBSOLETA (Brauer & Bergenstamm), 1891:390 (*Phasiophana*). Type-locality: "Brazil". Distr. — Costa Rica, Panama, Puerto Rico, Haiti, Cuba, S. Domingo, Jamaica, Venezuela, Guyana, Peru, Brazil. Ref. — Snyder, 1940b:6.
- mendax* Stein, 1911:87 (*Mydaea*). Type-locality: Peru, Urubamba River, S. Carlos and Umuhuankiali.
- PALLIDICORNIS (Bigot), 1888a:614; 1888b:clxxxii (*Curtonevra*). Type-locality: "Mexico". Distr. — Mexico, Brazil. N. COMB.
- americana* Hough, 1900:222, figs. 12 and 13 (*Muscina*). Type-locality: Brazil, Rio de Janeiro, Mato Grosso and Chapada (dos Guimarães).

Genus GRAPHOMYA Robineau-Desvoidy

- GRAPHOMYA Robineau-Desvoidy, 1830:403. Type-species, *Musca maculata* Scopoli (Westwood, 1840:141).
- Graphomyia*, error.
- Protostomoxys* Enderlein, 1935:242. Type-species, *podexaureus* Enderlein (orig. des.). N. SYN.
- ANALIS (Macquart), 1851a:228, pl. 23, fig. 9; 1851b:255, pl. 23, fig. 9 (*Cyrtonevra*). Type-locality: "Chile" (not Tasmania as stated by Macquart). Distr. — Peru, Chile. Ref. — Albuquerque, 1951b:13, figs. 26-29.
- tropicalis* Malloch, 1934:345. Type-locality: Peru, Huascaray Ridge.
- AURICEPS Malloch, 1934:346. Type-locality: Argentina, Mendoza Province. Distr. — Argentina.
- CHILENSIS Bigot, 1888a:616. Type-locality: "Chile". Distr. — Peru, Argentina, Chile. Ref. — Malloch, 1934:345.
- CUBANA Dodge, 1965:394. Type-locality: Cuba, Paso Real. Distr. — Cuba.
- MACULATA (Scopoli), 1763:326 (*Musca*). Type-locality: Europe, Carniola. Distr. — Mexico, Puerto Rico, Venezuela, Uruguay, Argentina; North America; West Africa, Palaearctic and Oriental regions, Hawaii. Ref. — Hennig, 1959:233, figs. 48 and 49a.
- MALLOCHI Albuquerque, 1954b:79, figs. 1-12. Type-locality: Chile, Santiago. Distr. — Argentina, Chile.
- MENDOZANA (Enderlein), 1935:242 (*Protostomoxys*). Type-locality: Argentina, Mendoza. Distr. — Argentina. N. COMB.
- MERIDIONALIS Townsend, 1892:32 (nom. nov. for *americana* Schiner). Type-locality: "South America". Distr. — Brazil.

- americana* Schiner, 1868:304 (preocc. Robineau-Desvoidy, 1830). Type-locality: "South America".
- MEXICANA Giglio-Tos, 1893:5; 1896:9. Type-locality: Mexico, Tuxpango. Distr. — Mexico, Panama, Bolivia, Argentina.
- PANAMENSIS Dodge, 1965:395. Type-locality: Panama Canal Zone, Barro Colorado Island. Distr. — Panama.
- PODEXAUREUS (Enderlein), 1935:242 (*Protostomoxys*). Type-locality: Chile, Cordillera, Contulmo and Rengo. Distr. — Chile. N. comb.
- STIPATA (Walker), 1852:348 (*Musca*). Type-locality: Guyana, Demerara. Distr. — Puerto Rico, Venezuela, Guyana. Ref. — Curran, 1934:460.

Subfamily LIMNOPHORINAE

Genus LIMNOPHORA Robineau-Desvoidy

- LIMNOPHORA Robineau-Desvoidy, 1830:517. Type-species, *palustris* Robineau-Desvoidy (Coquillett, 1910:561).
- Leucomelina* Macquart, 1851a:234; 1851b:261. Type-species, *pica* Macquart (orig. des.).
- Microchylum* Macquart, 1851a:229; 1851b:256. Type-species, *vittatum* Macquart (orig. des.).
- Brachioophyra* Giglio-Tos, 1893:9. Type-species, *effrons* Giglio-Tos (orig. des.) = *normata* Bigot.
- Brachyophyra*, error.
- ALACRIS Stein, 1911:116. Type-locality: Bolivia, Sorata. Distr. — Bolivia.
- AURIFACIES Stein, 1911:130. Type-locality: Bolivia, La Paz and Sorata; Peru, Cuzco, Calca and Paso de Lares. Distr. — Peru, Bolivia, Argentina.
- BREVIHIRTIA Malloch, 1934:294. Type-locality: Peru, Chosica. Distr. — Peru, Argentina.
- CORVINA (Giglio-Tos), 1893:7; 1896:19 (*Leucomelina*). Type-locality: "Mexico". Distr. — Mexico, Puerto Rico, Peru, ?Brazil, Chile. Ref. — Stein, 1918:235.
- CUBANA Johnson, 1919:442. Type-locality: Cuba, Havana. Distr. — Cuba.
- DUBIA Bigot, 1885:264. Type-locality: "Mexico". Distr. — Mexico.
- ELEGANS Macquart, 1843a:322, pl. 22, fig. 6; 1843b:165, pl. 22, fig. 6. Type-locality: Guyana. Distr. — Guyana.
- EXUL Williston, 1896:370, pl. 12, figs. 122 and 122a. Type-locality: St. Vincent. Distr. — St. Vincent Is., Cuba, Jamaica.
- GARRULA (Giglio-Tos), 1893:7; 1896:18 (*Leucomelina*). Type-locality: Mexico, Tuxpango. Distr. — Mexico, Brazil.
- INIQUA (Stein), 1911:146 (*Calliophrys*). Type-locality: Peru, Arequipa and Sicuani; Chile, Valparaiso, Arica, Palca, Tacna. Distr. — Peru, Chile. N. comb.
- LAETA Stein, 1911:118. Type-locality: Chile, Quillota, Coquimbo. Distr. — Chile.
- LIMBATA Bigot, 1885:271. Type-locality: "Mexico, Chile". Distr. — Mexico, Chile. Ref. — Stein, 1907:278.
- MARGINATA Stein, 1904:461. Type-locality: Colombia, Cordillera. Distr. — Colombia, Argentina.



- MARGINIPENNIS Stein, 1911:115. Type-locality: Peru, Pichis. Distr. — Peru.
- MINUSCULA (Wulp), 1896:330 (*Leucomelina*). Type-locality: Mexico, Atoyac in Vera Cruz and Xucumanatlán in Guerrero. Distr. — Mexico.
- NARONA (Walker), 1849:945 (*Anthomyia*). Type-locality: U.S.A., Florida. Distr. — Mexico, Bermuda, Bahamas, Puerto Rico, Venezuela, Guyana, Peru, Brazil, Paraguay, Argentina, Chile; North America. Ref. — Stein, 1911:132.
- prominula* Thomson, 1869:550 (*Anthomyia*). Type-locality: Argentina, Buenos Aires.
- cyrtoneurina* Stein, 1898:203. Type-locality: U.S.A., various states.
- NORMATA Bigot, 1885:272. Type-locality: "Mexico". Distr. — Mexico, Venezuela, Colombia, Peru, Bolivia, Brazil, Paraguay. Ref. — Stein, 1911:136.
- effrons* Giglio-Tos, 1893:9 (*Brachiophyra*). Type-locality: Mexico, Tuxpango.
- PATAGONICA Malloch, 1934:295. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina, Chile, Juan Fernandez Is.
- PICA (Macquart), 1851a:235, pl. 24, fig. 3; 1851b:262, pl. 24, fig. 3 (*Leucomelina*). Type-locality: Brazil, Minas Gerais. Distr. — Jamaica, Brazil, Galápagos Is. Ref. — Albuquerque, 1950b:1, figs. 1-6.
- PILISETA Stein, 1919:136 (nom. nov. for *plumiseta* Stein, 1911). Type-locality: Peru, mouth of Pachitea River. Distr. — Peru, Guyana, Brazil, Argentina.
- plumiseta* Stein, 1911:117 (preocc. Stein, 1903). Type-locality: Peru, mouth of Pachitea River.
- PLATYSTOMA (Thomson), 1869:554 (*Anthomyia*). Type-locality: Ecuador, Puna. Distr. — Ecuador. Ref. — Stein, 1910:72.
- SAEVA (Wiedemann), 1830:430 (*Anthomyia*). Type-locality: "Brazil". Distr. — Mexico, ?Panama, Peru, Bolivia, ?Brazil, Argentina. Ref. — Stein, 1911:133.
- SPRETA Malloch, 1921d:422. Type-locality: Brazil, Teresópolis. Distr. — Brazil.
- STRIGATA (Giglio-Tos), 1893:7; 1896:19 (*Leucomelina*). Type-locality: Mexico, Tuxpango. Distr. — Mexico.
- VICARIA (Walker), 1852:361 (*Anthomyia*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Stein, 1901:216.
- VICINA Robineau-Desvoidy, 1830:520. Type-locality: "Brazil". Distr. — Brazil.
- VITTATA (Macquart), 1851a:230, pl. 23, fig. 11; 1851b:257, pl. 23, fig. 11 (*Microchylum*). Type-locality: Brazil, Bahia. Distr. — Brazil. Ref. — Albuquerque, 1950b:2, figs. 7-10.
- ZEBRINA Bigot, 1885:273. Type-locality: "Chile". Distr. — Chile. Ref. — Stein, 1907:292.

Genus SYLLIMNOPHORA Speiser

- Acanthoneura* Stein, 1919:140 (proc. Schiner, 1868). Type-species. *Limnophora aliena* Stein (Séguy, 1937:250). (*Acanthoneura* Schiner, 1868, is an unjustified emendation of *Acanthonevra* Macquart, 1843 (see Article 33 a ii of the Code) but is an available

name for the purposes of homonymy. *Acanthoneura* Stein is not preoccupied by *Acanthoneura* Macquart (see Article 56a.).

- SYLLIMNOPHORA Speiser, 1923:99 (nom. nov. for *Acanthoneura* Stein). Type-species, *Limnophora aliena* Stein (aut.).
- ALIENA (Stein), 1911:128 (*Limnophora*). Type-locality: Peru, Arequipa, Lago Titicaca, Sicuani, Calca, Tarma, Oroya; Bolivia, Tiahuanaco and Lago Titicaca; Chile, Palca. Distr. — Peru, Bolivia, Chile.
- ALTANEIRA (Albuquerque), 1954n:400, figs. 10-15 (*Heliographa*). Type-locality: Brazil, Minas Gerais, Serra do Cipó. Distr. — Brazil. N. comb.
- ANGUSTIFRONS Malloch, 1934:286. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.
- ATRA (Stein), 1904:469 (*Limnophora*). Type-locality: Colombia, Cordillera. Distr. — Colombia, Bolivia. N. comb.
- ATROVITTATA (Stein), 1904:463 (*Limnophora*). Type-locality: Peru, Callanga. Distr. — Colombia, Peru, Bolivia, Chile. N. comb.
- AURIFLUA (Stein), 1911:130 (*Limnophora*). Type-locality: not given, South America from the title of the paper. Distr. — Chile, Argentina.
- BARBITARSIS (Stein), 1911:115 (*Limnophora*). Type-locality: Peru, Tarma and Paso de Lares; Bolivia, Lorenzopata. Distr. — Peru, Bolivia. N. comb.
- BREVISETA (Stein), 1911:117 (*Limnophora*). Type-locality: Bolivia, Lorenzopata. Distr. — Bolivia. N. comb.
- BROWNI Snyder, 1957c:474. Type-locality: Ecuador, Chimborazo, Riobamba. Distr. — Ecuador.
- CANDIDIFRONS (Stein), 1911:128 (*Limnophora*). Type-locality: Peru, Sicuani and Cuzco; Bolivia, Island of Titicaca. Distr. — Peru, Bolivia.
- CLAVITIBIA (Stein), 1911:129 (*Limnophora*). Type-locality: Peru, Lago Titicaca, Juliaca and Sicuani; Bolivia, Lago Titicaca. Distr. — Peru, Bolivia.
- COMPRESSIFRONS (Stein), 1904:464 (*Limnophora*). Type-locality: Peru, Callanga. Distr. — Peru, Bolivia. N. comb.
- COMPRESSITARSIS (Stein), 1911:127 (*Limnophora*). Type-locality: Peru, Lago Titicaca and Bolivia, Tiahuanaco. Distr. — Peru, Bolivia.
- DELETA (Wulp), 1896:329 (*Leucomelina*). Type-locality: Mexico, Guerrero, Xucumanatlán, Omilteme and Sierra de las Aguas Escondidas. Distr. — Mexico, Peru, Brazil. N. comb.
- DUBITALIS Malloch, 1934:287. Type-locality: Chile, Chiloé Is., Castro. Distr. — Chiloé Is.
- FEMORATA (Stein), 1911:129 (*Limnophora*). Type-locality: Chile, Valparaiso, Quillota. Distr. — Argentina, Chile.
- FLAVIPALPIS Snyder, 1957c:471. Type-locality: Argentina, Neuquén, Zapala. Distr. — Argentina.
- GRACILITARSIS (Stein), 1911:116 (*Limnophora*). Type-locality: Peru, Pichis, and Bolivia, Lorenzopata. Distr. — Mexico, Peru, Bolivia. N. comb.
- INCONSPICUA Malloch, 1934:289. Type-locality: Chile, Osorno. Distr. — Chile, Chiloé Is.
- INTEGRA (Stein), 1911:119 (*Limnophora*). Type-locality: Bolivia, La Paz and Sorata. Distr. — Bolivia. N. comb.


- LATIFRONS Malloch, 1934:290. Type-locality: Tierra del Fuego, Estancia Viamonte. Distr. — Tierra del Fuego.
- LATIMANA Malloch, 1934:288, fig. 53. Type-locality: Argentina, Lake Nahuel Huapi. Distr. — Argentina.
- LATIMANODES Snyder, 1957c:473. Type-locality: Argentina, Tucumán, Tafi del Valle. Distr. — Argentina.
- LISPOMIMA Hennig, 1955:31. Type-locality: Juan Fernández Is., Masafuera Is., Quebrada de las Vacas. Distr. — Juan Fernández Is.
- NIGRARGENTATA (Albuquerque), 1954n:404, figs. 16-22 (*Heliographa*). Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil. N. COMB.
- NIGRIVENTRIS Malloch, 1934:287. Type-locality: Chile, Perales. Distr. — Chile.
- PARANAENSIS (Albuquerque), 1954n:396, figs. 1-9 (*Heliographa*). Type-locality: Brazil, Paraná, Iguazu. Distr. — Brazil. N. COMB.
- PAUCISETA (Stein), 1904:460 (*Limnophora*). Type-locality: Peru, Callanga. Distr. — Peru, Bolivia. N. COMB.
- PURA (Stein), 1911:119 (*Limnophora*). Type-locality: Peru, Paso de Lares and Tarma. Distr. — Peru. N. COMB.
- SETITIBIA (Stein), 1911:129 (*Limnophora*). Type-locality: Peru, Arequipa and Chile, Palca. Distr. — Peru, Chile.
- SPINULIGERA (Stein), 1911:130 (*Limnophora*). Type-locality: Peru, Oroya and Lago Titicaca; Bolivia, Cordillera and Lago Titicaca. Distr. — Peru, Bolivia.
- SUAVIS (Stein), 1911:130 (*Limnophora*). Type-locality: Peru, Paso de Lares. Distr. — Ecuador, Peru.
- TENUIPENNIS Malloch, 1934:284. Type-locality: Argentina, Bariloche. Distr. — Argentina.
- VARICEPS Malloch, 1934:291. Type-locality: Argentina, Buenos Aires. Distr. — Argentina.
- VITIVENTRIS Malloch, 1934:291. Type-locality: Peru, Huariaca. Distr. — Peru.

Genus TELEUTOCHAETA Stein

- TELEUTOCHAETA Stein, 1911:141. Type-species, *nigriceps* Stein (mon.).
- Mesochaeta* Stein, 1911:139 (preocc. Brauer & Bergenstamm, 1891). Type-species, *variegata* Stein (Séguy, 1937:273). N. SYN.
- INCAICA (Stein), 1911:140 (*Mesochaeta*). Type-locality: Bolivia, Titicaca Island. Distr. — Bolivia. N. COMB.
- NIGRICEPS Stein, 1911:141. Type-locality: Peru, Oroya. Distr. — Peru.
- VARIEGATA (Stein), 1911:139 (*Mesochaeta*). Type-locality: Peru, Lago Titicaca and Oroya and Bolivia, Lago Titicaca. Distr. — Peru, Bolivia. N. COMB.

Genus THAUMASIOCHAETA Stein

- THAUMASIOCHAETA Stein, 1911:137. Type-species, *pilitarsis* Stein (mon.).
- PILITARSIS Stein, 1911:138. Type-locality: Peru, Lago Titicaca and Bolivia, Tiahuanaco. Distr. — Peru, Bolivia.



Genus RHYNCHOLIMNOPHORA Stein

- RHYNCHOLIMNOPHORA Stein, 1911:142. Type-species, *compressitarsis* Stein (Séguy, 1937:273).
COMPRESSITARSIS Stein, 1911:142. Type-locality: Peru, Vally of Lares. Distr. — Peru.
HAUSTELLATA Stein, 1911:144. Type-locality: Peru, Cuzco. Distr. — Peru.
LONGIPALPIS Stein, 1911:143. Type-locality: Peru, Oroya. Distr. — Peru.

Genus BUCEPHALOMYIA Malloch


- BUCEPHALOMYIA Malloch, 1918b:273. Type-species, *Tetramerinx femorata* Malloch (orig. des.).
ACZELI Snyder, 1957c:468. Type-locality: Argentina, Tucumán, San Javier. Distr. — Argentina.

Genus AGENAMYIA Albuquerque

- AGENAMYIA Albuquerque, 1953c:519. Type-species, *fumipennis* Albuquerque (orig. des.).
FUMIPENNIS Albuquerque, 1953c:519, figs. 1-12. Type-locality: Brazil, Guanabara, Rio de Janeiro, Gávea. Distr. — Brazil.

Genus LISPOIDES Malloch

- LISPOIDES Malloch, 1920:146. Type-species, *Limnophora aequifrons* Stein (orig. des.).
ABNORMINERVIS (Stein), 1911:123 (*Limnophora*). Type-locality: Peru, Valley of Lares and Bolivia, Lago Titicaca. Distr. — Peru, Bolivia. N. COMB.
AEQUIFRONS (Stein), 1898:205 (*Limnophora*). Type-locality: U.S.A., Idaho and South Dakota. Distr. — Colombia; North America.
ARGENTICEPS Malloch, 1934:298. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina, ?Chile.
ARGENTINA Malloch, 1934:298. Type-locality: Argentina, Mendoza, Potrerillos. Distr. — Argentina.
ATRISQUAMA (Stein), 1904:466 (*Limnophora*). Type-locality: Peru, Callanga. Distr. — Peru.
DILUTA (Stein), 1911:126 (*Limnophora*). Type-locality: Chile, Tacna. Distr. — Chile. N. COMB.
ELEGANTULA, nom. nov. for *Limnophora elegans* Stein. Type-locality: Bolivia, Cordillera and Sorata. Distr. — Bolivia.
elegans Stein, 1911:120 (*Limnophora*; preocc. Macquart, 1843). Type-locality: Bolivia, Cordillera and Sorata.
GRACILIS (Stein), 1911:125 (*Limnophora*). Type-locality: Bolivia, Sorata and Chile, Palca. Distr. — Bolivia, Chile. N. COMB.
GUATEMALA Snyder, 1951:8. Type-locality: Guatemala, Guatemala City. Distr. — Guatemala.



- INAEQUIFRONS Malloch, 1934:296. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina, Chile.
- INSULARIS Hennig, 1957:409. Type-locality: Juan Fernández Is., Masafuera Is., La Correspondencia. Distr. — Juan Fernández Is.
- LAEVIS (Stein), 1911:125 (*Limnophora*). Type-locality: Peru, Chanchamayo. Distr. — Vcnezuela, Peru.
- LATIFRONS Snyder, 1957c:475. Type-locality: Argentina, Santa Cruz, Cristina, Lago Argentina. Distr. — Argentina.
- LOPESI Albuquerque, 1955c:153, figs. 1-9. Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil.
- NIGRIBASIS (Stein), 1911:122 (*Limnophora*). Type-locality: Peru, Sicuani, Cuzco, Tarma and Oroya; Bolivia, Lago Titicaca, Tiahuanaco and the Cordillera. Distr. — Peru, Bolivia. N. comb.
- PROPINQUA (Stein), 1911:124 (*Limnophora*). Type-locality: Peru, Valley of Lares. Distr. — Peru.
- TRIPLEX (Stein), 1911:123 (*Limnophora*). Type-locality: Peru, Cuzco; Bolivia, Lago Titicaca; Chile, Palca. Distr. — Peru, Bolivia, Chile.
- UNISETA Malloch, 1934:299. Type-locality: Peru, Lima, Verrugas Canyon. Distr. — Peru.

Genus SPILOGONA Schnabl

- Limnophora*, subg. SPILOGONA Schnabl, 1911:92. Type-species, *Aricia carbonella* Zetterstedt (mon.).
- ANTHRAX (Bigot), 1835:274 (*Limnophora*). Type-locality: "Mexico". Distr. — Mexico; North America. Ref. — Hockett, 1932:314.
- ARGENTIFRONTATA Snyder, 1957c:470. Type-locality: Argentina, Rio Negro, Isla Choelchoe. Distr. — Argentina.
- ATERRIMA (Stein), 1904:469 (*Limnophora*). Type-locality: Bolivia, San Antonio. Distr. — Colombia, Bolivia. N. comb.
- GOLBACHI Snyder, 1957c:469. Type-locality: Argentina, Tucumán, Queb. La Toina. Distr. — Argentina.
- HIRTICEPS (Stein), 1911:121 (*Limnophora*). Type-locality: Peru, Lago Titicaca and Sicuani; Bolivia, Lago Titicaca. Distr. — Peru, Bolivia. N. comb.
- IMMACULIVENTRIS (Malloch), 1923c:425 (*Melanochelia*). Type-locality: Gulf of California, San José Is. Distr. — San José Is., Mexico, San Francisco Is. (Gulf of Mexico). N. comb.
- PUBICEPS (Stein), 1911:122 (*Limnophora*). Type-locality: Peru, Lago Titicaca and Bolivia, Lago Titicaca. Distr. — Peru, Bolivia. N. comb.
- SEMIGINEREA (Stein), 1911:131 (*Limnophora*). Type-locality: Peru, Lago Titicaca and Bolivia, Lago Titicaca. Distr. — Peru, Bolivia. N. comb.
- TRICHOPS (Stein), 1911:120 (*Limnophora*). Type-locality: Peru, Valley of Lares, Chanchamayo, Tarma; Bolivia, Sorata. Distr. — Peru, Bolivia. N. comb.

Genus LISPE Latreille

- LISPE Latreille, 1796:169. Type-species, *Musca tentaculata* Degger (sub. mon., Latreille, 1802:462).
- Lispa*, error.

- ALBITARSIS Stein, 1898:277. Type-locality: U.S.A., various states. Distr. — Bermuda Is., Bahama Is.; North America. Ref. — Snyder, 1954b: 14, figs. 25, 40, 54, 68.
- BAHAMA Snyder, 1958b:1. Type-locality: South Caicos Is. Distr. — Bahama Is.
- COTIDIANA Snyder, 1954b:22, figs. 23, 42, 55, 69. Type-locality: Canada, Alberta, Suffield. Distr. — Mexico; North America, Central and East Asia. Ref. — Hennig, 1960:429.
- LATANA Snyder, 1949b:4. Type-locality: Brazil, Natal. Distr. — Brazil.
- LEVIS Stein, 1911:147. Type-locality: Chile, Arica. Distr. — Chile.
- LISARBA Snyder, 1949b:2. Type-locality: Brazil, Natal. Distr. — Brazil.
- NASONI Stein, 1898:280. Type-locality: U.S.A., South Dakota, Illinois, Georgia. Distr. — Bahama Is., Virgin Is., Cuba, Puerto Rico; North America. Ref. — Snyder, 1954b:16, figs. 6, 29, 37, 57, 70.
- RUFITIBIALIS Macquart, 1843a:325, pl. 22, fig. 7; 1843b:168, pl. 22, fig. 7. Type-locality: "Brazil or Chile". Distr. — Puerto Rico, Brazil or Chile.
- SEROTINA Wulp, 1896:342. Type-locality (restricted): Mexico, Veracruz, Atoyac. Distr. — Mexico, Puerto Rico, Guyana, Brazil, Paraguay. *edwardsi* Malloch, 1934:278. Type-locality: Paraguay, Encarnación.
- SETULIGERA Stein, 1911:146. Type-locality: Chile, Valparaiso and Coquimbo. Distr. — Argentina, Chile.
- TENTACULATA (Degeer), 1776:86 (*Musca*). Type-locality: Sweden. Distr. — Guadalupe Is., Guatemala, Peru; North America, Palaearctic region. Ref. — Snyder, 1954b:35, figs. 4, 9, 10; Hennig, 1960:456, figs. 78, 102, 111, 131, 159, 160a and b, pl. 18, fig. 369, pl. 19, fig. 370, pl. 20, fig. 407.
- ULIGINOSA Fallén, 1825:93. Type-locality: Sweden, Westrogothia and Scania. Distr. — Mexico, St. Vincent Is.; Palaearctic region. Ref. — Hennig, 1960:459, figs. 73, 86, 107, 120, 136, pl. 20, fig. 390, pl. 21, fig. 413. Erroneous records: certainly refer to *cotidiana* Snyder (see Snyder, 1954b:21 and 24).
- VILIS Stein, 1911:148. Type-locality: Bolivia, Sorata and Lorenzopata. Distr. — Bolivia.

Unplaced Species of LIMNOPHORA s. lat.

- PORTERI Brèthes, 1919:43 (*Limnophora*). Type-locality: Chile, Rio Blanco. Distr. — Chile. Probably not Limnophorinae.
- VIRGATA (Wiedemann), 1830:434 (*Anthomyia*). Type-locality: "Brazil". Distr. — Brazil.

Subfamily COENOSIINAE

Genus DREPANOCNEMIS Stein

- DREPANOCNEMIS Stein, 1911:178. Type-species, *dorae* Stein (Malloch, 1928a: 471).
- DORAE Stein, 1911:180. Type-locality: Peru, Tarma. Distr. — Peru.
- GRISOVIRENS Malloch, 1928a:470. Type-locality: Peru, Huariaca. Distr. — Peru.



HIRTICEPS Stein, 1911:179. Type-locality: Peru, Chanchamayo and Bolivia, Sorata and Mapiri. Distr. — Peru, Bolivia.

Genus OXYTONOCERA Stein

OXYTONOCERA Stein, 1919:139. Type-species, *Coenosia nigrohalterata* Stein (mon.).

Oxytonocera, error.

NIGROHALTERATA (Stein), 1904:484 (*Coenosia*). Type-locality: Peru, Callanga. Distr. — Peru, Bolivia.

Genus TETRAMERINX Berg

Tetrachaeta Stein, 1898:254 (preocc. Brauer & Bergenstamm, 1894). Type-species, *unica* Stein (mon.).

Phyllogaster Stein, 1898:256 (preocc. Gistel, 1834). Type-species, *cordyluroides* Stein (mon.).

TETRAMERINX Berg, 1898:17 (nom. nov. for *Tetrachaeta* Stein). Type-species, *Tetrachaeta unica* Stein (aut.).

CORDYLUROIDES (Stein), 1898:256 (*Phyllogaster*). Type-locality: U.S.A., Florida and Massachusetts. Distr. — Bermuda Is., Bahama Is.; North America. Ref. — Hockett, 1936:200. N. comb.

GENICULATA (Macquart), 1851a:244, pl. 25, fig. 2; 1851b:271, pl. 25, fig. 2 (*Cordylura*). Type-locality: "America". Distr. — America. N. comb.

LONGISPINA (Malloch), 1923c:426 (*Phyllogaster*). Type-locality: U.S.A., California, Gonzales Bay. Distr. — South California. Ref. — Hockett, 1936:198. N. comb.

NIGRIPES Stein, 1911:145. Type-locality: Chile, Arica. Distr. — Chile.

PANTHERINA (Wiedemann), 1824:53 (*Anthomyia*). Type-locality: "South America". Distr. — South America.

RUFITIBIA Stein, 1911:144. Type-locality: Peru, Mollendo and Chile, Arica and Tacna. Distr. — Bahama Is., Peru, Chile; North America. Ref. — Hockett, 1936:197.

Genus APSIL Malloch

APSIL Malloch, 1929a:103. Type-species, *maculiventris* Malloch (orig. des.).


APICATA Malloch, 1934:247, fig. 47b. Type-locality: Argentina, Lake Correntoso. Distr. — Chile, Argentina.

ATRIPES Malloch, 1934:245, figs. 46 and 47a. Type-locality: Chile, Concepción. Distr. — Chile.

BISETA Malloch, 1934:244. Type-locality: Argentina, Bariloche. Distr. — Argentina.

DILATA Malloch, 1934: fig. 47c. Type-locality: Chile, Chiloé Is., Puntra. Distr. — Chile, Chiloé Is.

MACULIPENNIS Malloch, 1934:243, figs. 44 and 45a-b. Type-locality: Argentina, Puerto Blest. Distr. — Argentina.



MACULIVENTRIS Malloch, 1929a:103. Type-locality: Chile, Perales. Distr. — Chile.

PENNATA Malloch, 1934:241, figs. 43a-c. Type-locality: Chile, Casa Pangué. Distr. — Chile, Argentina.

SPATULATA Malloch, 1934:250. Type-locality: Argentina, Bariloche. Distr. — Chile, Argentina.

Genus REYNOLDSIA Malloch

REYNOLDSIA Malloch, 1934:230. Type-species, *pectinata* Malloch (orig. des.).

AURIFERA (Bigot), 1885:273 (*Limnophora*). Type-locality: "Chile". Distr. — Chile. Ref. — Malloch, 1934:232.

BREVITARSIS Malloch, 1934:233. Type-locality: Argentina, Lake Gutiérrez.

COXATA Malloch, 1934:236. Type-locality: Argentina, Bariloche. Distr. — Argentina.

PECTINATA Malloch, 1934:235, fig. 41a. Type-locality: Tierra del Fuego, Río Grande, Estancia Viamonte. Distr. — Tierra del Fuego.

PTEROPLEURALIS Malloch, 1934:237, figs. 41b-c. Type-locality: Argentina, Lake Nahuel Huapi. Distr. — Chile, Argentina.
Distr. — Argentina.

?ROBUSTA (Stein), 1911:172 (*Coenosia*). Type-locality: Chile, Corral. Distr. — Chile. N. comp.

RUFOAPICATA Malloch, 1934:239, figs. 41d and 42. Type-locality: Argentina, Bariloche. Distr. — Argentina.

SCUTELLATA Malloch, 1934:234. Type-locality: Chile, Angol. Distr. — Chile.

TROCHANTERATA Malloch, 1934:233. Type-locality: Argentina, Puerto Blest. Distr. — Argentina.

Genus RAYMONDOMYIA Malloch

RAYMONDOMYIA Malloch, 1934:228. Type-species, *flavipalpis* Malloch (orig. des.).

FLAVIPALPIS Malloch, 1934:229, fig. 40. Type-locality: Argentina, Bariloche. Distr. — Argentina.

Genus BITHORACOAETA Stein

BITHORACOAETA Stein, 1911:177. Type-species, *Anthomyia despecta* Walker (orig. des.) = *leucoprocta* (Wiedemann).

ANNULATA Stein, 1911:178. Type-locality: Peru, Valley of Lares and mouth of Pachitea River. Distr. — Peru, Surinam, Brazil. Ref. — Albuquerque, 1956k:2.

ATRICORNIS Malloch, 1934:227, fig. 39d. Type-locality: Argentina, Viedma. Distr. — Brazil, Argentina. Ref. — Albuquerque, 1956k:6.

FLAVICOXA Malloch, 1934:228. Type-locality: Costa Rica, Farm la Caja, 8 km west of San José. Distr. — Costa Rica.

LEUCOPROCTA (Wiedemann), 1830:433 (*Anthomyia*). Type-locality: West Indies. Distr. — Mexico, Puerto Rico, Cuba, Jamaica, Venezuela,

Guyana, Peru, Brazil, Paraguay, Uruguay, Argentina; North America. Ref. — Malloch, 1934:226.

despecta Walker, 1852:364 (*Anthomyia*). Type-locality: "Brazil".

antica Walker, 1852:367 (*Coenosia*). Type-locality: "United States".

pipunculina Thomson, 1869:559 (*Coenosia*). Type-locality: Brazil, Guanabara, Rio de Janeiro.

rufipes Bigot, 1885:272 (*Limnophora*). Type-locality: "Mexico".

calopus Bigot, 1885:275 (?*Hydrophoria*). Type-locality: "Mexico".

uvens Giglio-Tos, 1894:11; 1896:32 (*Coenosia*). Type-locality: "Mexico".
N. SYN.

sevens, error.

femoralis Wulp, 1896:343 and 1897:345 (*Coenosia*). Type-locality: Mexico, Orizaba, Chilpancingo in Guerrero, Mexico City and Vera Cruz.

insignis Stein, 1898:257 (*Caricea*). Type-locality: U.S.A., Florida.

NIGRICORNIS Malloch, 1934:228. Type-locality: Costa Rica, Higuito. Distr. — Costa Rica.

PACIFERA (Giglio-Tos), 1893:11; 1896:31 (*Coenosia*). Type-locality: Mexico, Tuxpango. Distr. — Mexico. N. COMB.

PLUMATA Albuquerque, 1955b:169, figs. 1-6. Type-locality: Brazil, Minas Gerais, Salvaterra, Juiz de Fora. Distr. — Brazil.

SOCIABILIS Blanchard, 1937:41, figs. 4a-c. Type-locality: Argentina, Misiones, Loreto and Chaco. Distr. — Argentina.

VARICORNIS (Coquillett), 1900:256 (*Coenosia*). Type-locality: Puerto Rico, Bayamón, Aguadilla, Utuado, Vieques Is., Fajardo and Arroyo. Distr. — Puerto Rico.

Genus HAROLDOPSIS Albuquerque

HAROLDOPSIS Albuquerque, 1954c:119. Type-species, *cambuquirensis* Albuquerque (orig. des.).

CAMBUQUIRENSIS Albuquerque, 1954c:119, figs. 1-15. Type-locality: Brazil, Minas Gerais, Cambuquira. Distr. — Brazil.

Genus PILISPINA Albuquerque

PILISPINA Albuquerque, 1954d:180. Type-species, *pilitibia* Albuquerque (orig. des.).

PILITIBIA Albuquerque, 1954d:180, figs. 10-18. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil.

Genus LEVALLONIA Albuquerque

LEVALLONIA Albuquerque, 1958d:101. Type-species, *xanthogaster* Albuquerque (orig. des.).

XANTHOGASTER Albuquerque, 1958d:101, figs. 1-3. Type-locality: Brazil, Rio de Janeiro, Petrópolis, Le Vallon, Alto da Mosela. Distr. — Brazil.



Genus RHABDOPTERA Stein

RHABDOPTERA Stein, 1919:155. Type-species, *Schoenomyza striatipennis* Stein (mon.).

STRIATIPENNIS (Stein), 1911:185 (*Schoenomyza*). Type-locality: Peru, Tarma and Oroya. Distr. — Peru.

Genus SPATHIPHEROMYIA Bigot

SPATHIPHEROMYIA Bigot, 1884:lxxxix. Type-species, *stellata* Bigot (mon.). *Spathipheromya*, error.

ALBICEPS Malloch, 1934:255. Type-locality: Chile, Chiloé Island, Castro. Distr. — Chile, Chiloé Is.

ATRA Malloch, 1934:258. Type-locality: Argentina, Lake Nahuel Huapi. Distr. — Chile, Argentina.

AURICEPS (Stein), 1911:186 (*Schoenomyza*). Type-locality: Chile, Coquimbo and Magellan Straits. Distr. — Chile, N. comb.

CHILENSIS Malloch, 1934:254. Type-locality: Chile, Chiloé Island, Castro. Distr. — Chile, Chiloé Is.

FUSCIPALPIS (Stein), 1911:182 (*Schoenomyza*). Type-locality: Peru, Tarma and Oroya. Distr. — Peru, N. comb.

GUTTIPENNIS (Thomson), 1869:600 (*Ochthiphila*). Type-locality: Argentina, Buenos Aires. Distr. — Peru, Bolivia, Chile, Argentina. Ref. — Malloch, 1934:254.

guttipennis Stein, 1904:493 (*Schoenomyza*; preocc. Thomson, 1869). Type-locality: Argentina, Buenos Aires.

INSULARIS Malloch, 1934:257. Type-locality: Chile, Mechuque Island. Distr. — Chile, Mechuque Is.

MAGELLANI Malloch, 1934:256. Type-locality: Tierra del Fuego, Rio Grande. Distr. — Tierra del Fuego.

MINUTA (Stein), 1911:185 (*Schoenomyza*). Type-locality: Peru, Valley of Lares. Distr. — Peru, N. comb.

NIGRA (Stein), 1911:188 (*Schoenomyza*). Type-locality: Peru, Lago Titicaca. Distr. — Peru, N. comb.

PICTA (Stein), 1911:183 (*Schoenomyza*). Type-locality: Peru, Juliaca and Lago Titicaca. Distr. — Peru, N. comb.

STELLATA Bigot, 1884:lxxxix. Type-locality: Chile. Distr. — Chile. Ref. — Stein, 1907:287.

Genus NOTOSCHOENOMYZA Malloch

NOTOSCHOENOMYZA Malloch, 1934:259. Type-species, *chrysiceps* Malloch (orig. des.).

ANNULATA (Stein), 1911:181 (*Schoenomyza*). Type-locality: Peru, Chanchamayo and Pichis. Distr. — Peru, N. comb.

APICALIS (Stein), 1911:184 (*Schoenomyza*). Type-locality: Peru, Lago Titicaca, Cuzco and Valley of Lares; Bolivia, Lago Titicaca. Distr. — Peru, Bolivia.




- CHRYSEPS Malloch, 1934:259. Type-locality: Chile, Chiloé Island, Ancud. Distr. — Chile, Chiloé Is.
- COSTATA Snyder, 1957c:450. Type-locality: Argentina, Tucumán, Tafí del Valle. Distr. — Argentina.
- IMMACULATA (Walker), 1837:357 (*Anthomyia*). Type-locality: Chile, Port Famine. Distr. — Chile, Argentina. Ref. — Malloch, 1934:260.
- KUSCHELI Hennig, 1955:27. Type-locality: Juan Fernández, Masafuera Island, La Correspondencia. Distr. — Juan Fernández Is.
- SPINICOSTA (Stein), 1904:495 (*Schoenomyza*). Type-locality: Peru, Callanga. Distr. — Peru, Bolivia. N. COMB.
- SULFURICEPS Malloch, 1934:261. Type-locality: Uruguay, Montevideo. Distr. — Uruguay.

Genus SCHOENOMYZINA Malloch

- SCHOENOMYZINA Malloch, 1934:273. Type-species, *pallicornis* Malloch (orig. des.).
- BISETA (Stein), 1911:183 (*Schoenomyza*). Type-locality: Peru, Cuzco. Distr. — Peru. N. COMB.
- EMDENI Hennig, 1955:29. Type-locality: Juan Fernández, Masafuera Island, Inocentes Altos. Distr. — Juan Fernández Is.
- FUSCICOSTA Malloch, 1934:276. Type-locality: Argentina, Bariloche. Distr. — Argentina.
- PALLICORNIS Malloch, 1934:274, figs. 51 and 52a. Type-locality: Argentina, Lake Nahuel Huapi. Distr. — Argentina.
- TRIANGULARIS Malloch, 1934:275, figs. 52b-c. Type-locality: Argentina, Lake Gutiérrez. Distr. — Chile, Argentina.
- UNICOLOR (Stein), 1911:182 (*Coenosia*). Type-locality: Peru, Lago Titicaca. Distr. — Peru. N. COMB.

Genus SCHOENOMYZA Haliday

- SCHOENOMYZA Haliday, 1833:166. Type-species, *Sciomyza fasciata* Meigen (Westwood, 1840:143) = *litorella* (Fallén).
- ALBICEPS Malloch, 1934:268, fig. 50. Type-locality: Argentina, Bariloche. Distr. — Argentina.
- ALBOMEDIA Malloch, 1934:266. Type-locality: Chile, Chiloé Island, Castro. Distr. — Chile.
- ARGYRICEPS Malloch, 1934:273. Type-locality: Argentina, Buenos Aires. Distr. — Argentina.
- ARMIPES Malloch, 1934:266. Type-locality: Argentina, Lake Nahuel Huapi. Distr. — Argentina.
- BELLA Malloch, 1934:271. Type-locality: Peru, Lima. Distr. — Peru.
- BISERIATA Malloch, 1934:267. Type-locality: Argentina, Bariloche. Distr. — Argentina.
- BREVIVENTRIS Stein, 1911:187. Type-locality: Peru, Oroya and Tarma. Distr. — Peru.
- EVITTATA Malloch, 1934:272. Type-locality: Uruguay, Montevideo. Distr. — Uruguay.



- FUSCIFACIES Malloch, 1934:270. Type-locality: Chile, Mechuque Island. Distr. — Chile, Mechuque Is.
- LATIVITTATA Malloch, 1934:269. Type-locality: Chile, Peulla. Distr. — Chile.
- LITORELLA (Fallén), 1823a:10 (*Ochthipila*). Type-locality: Sweden. Distr. — Peru, Chile; Holarctic region, East Africa. Ref. — Hennig, 1961: 507, figs. 203, 209, pl. 22, fig. 447, pl. 23, fig. 458.
- MALLOCHII, nom. nov. for *apicalis* Malloch. Type-locality: Argentina, Potrerillos. Distr. — Argentina.
- apicalis* Malloch, 1934:271 (prcocc. Stein, 1911). Type-locality: Argentina, Potrerillos.
- NEOBISERIATA Snyder, 1957c:448. Type-locality: Argentina, Santa Cruz, Lago Argentina. Distr. — Argentina.
- NIGRITHORAX Stein, 1911:187. Type-locality: Peru, Lago Titicaca. Distr. — Peru.
- TARSALIS Malloch, 1934:265. Type-locality: Argentina, Puerto Blest. Distr. — Chile, Argentina.
- UNIVITTATA Malloch, 1934:264. Type-locality: Argentina, River Pichileufu. Distr. — Argentina.
- WILLINKI Snyder, 1957c:449. Type-locality: Argentina, Santa Cruz, La Cristina, Lago Argentina. Distr. — Argentina.

Unplaced species of *SCHOENOMYZA* s. lat.

- AUSTRALIS (Brèthes), 1924:111 (*Ochthipila*). Type-locality: Chile, Rio Blanco. Distr. — Chile.
- FENESTRATA Bigot, 1888c:30. Type-locality: Tierra del Fuego, Orange Bay. Distr. — Tierra del Fuego.

Genus *CHOLOMYIODES* Albuquerque

- CHOLOMYIODES Albuquerque, 1954e:225. Type-species, *fumipennis* Albuquerque (orig. des.).
- FUMIPENNIS Albuquerque, 1954e:225, figs. 1-6. Type-locality: Brazil, São Paulo, Salesópolis. Distr. — Brazil.

Genus *ANGOLINA* new name

- Angolia* Malloch, 1934:251 (prcocc. Obenberger, 1920). Type-species, *acuta* Malloch (orig. des.).
- ANGOLINA, nom. nov. for *Angolia* Malloch. Type-species, *Angolia acuta* Malloch (aut.).
- ACUTA (Malloch), 1934:251, fig. 48 (*Angolia*). Type-locality: Chile, Angol. Distr. — Chile. N. comb.
- ALBIBARBA (Snyder), 1957c:446 (*Angolia*). Type-locality: Argentina, Tucumán, Burruyacu, V. Padre Monte. Distr. — Argentina. N. comb.
- ARGENTINA (Snyder), 1957c:447 (*Angolia*). Type-locality: Argentina, Neuquén, San Martín de los Andes. Distr. — Argentina. N. comb.



Genus STOMOPOGON Malloch

STOMOPOGON Malloch, 1930:469. Type-species, *Coenosia albiseta* Stein (orig. des.).

ALBISETA (Stein), 1911:172 (*Coenosia*). Type-locality: Bolivia, Lago Titicaca; Peru, Calca and Valley of Lares.

Genus NOELIA Albuquerque

NOELIA Albuquerque, 1957e:359. Type-species, *benevenuta* Albuquerque (orig. des.).

BENEVENUTA Albuquerque, 1957e:360, figs. 1-9. Type-locality: Brazil, São Paulo, Serra da Bocaina, Fazenda Lajeado. Distr. — Brazil.

?CAPRIBARBA (Stein), 1911:160 (*Coenosia*). Type-locality: Bolivia, La Paz and Sorata and Argentina, Lules. Distr. — Bolivia, Argentina. N. comb.

?GARLEPPI (Stein), 1911:166 (*Coenosia*). Type-locality: Peru, mouth of Pachitea River; Bolivia, Sarampiuni and San Carlos. Distr. — Peru, Bolivia. N. comb.

Genus COENOSIA Meigen

COENOSIA Meigen, 1926:210. Type-species, *Musca tigrina* Fabricius (Westwood, 1840:143).

Caricea Robineau-Desvoidy, 1830:530. Type-species, *communis* Robineau-Desvoidy (Hennig, 1961:519) = *tigrina* (Fabricius).

Choetura Macquart, 1851a:244; 1851b:271. Type-species, *rufipes* Macquart (mon.) = *macquarti*, nom. nov. N. SYN.

Chaetura, error.

Austrocoenosia Malloch, 1934:217. Type-species, *nigerrima* Malloch (orig. des.). N. SYN.

Macquartiopsis Albuquerque, 1949b:439 (nom. nov. for *Choetura* Macquart, 1851, allegedly preoccupied by *Chaetura* Stephens, 1826). Type-species, *Choetura rufipes* Macquart (aut.) = *macquarti* nom. nov. N. SYN.

ALBISQUAMA (Albuquerque), 1959:9, figs. 15-19 (*Austrocoenosia*). Type-locality: Brazil, Rio de Janeiro, Teresópolis, Parque Nacional da Serra dos Órgãos. Distr. — Brazil. N. comb.


ARGENTIFRONS (Malloch), 1934:222, fig. 37a (*Austrocoenosia*). Type-locality: Argentina, Bariloche. Distr. — Chile, Argentina. N. comb.

ATRIFRONS Stein, 1911:164. Type-locality: Chile, Coronel, Valparaíso and Quillota. Distr. — Chile.

AURIFERA (Malloch), 1934:219, fig. 35a (*Austrocoenosia*). Type-locality: Argentina. Bariloche. Distr. — Argentina. N. comb.

BENOISTI Séguy, 1932:144, figs. 1-3. Type-locality: Ecuador, Coto Collao. Distr. — Ecuador.

CAMORINENSIS Albuquerque, 1956f:8, figs. 17-18. Type-locality: Brazil, Guanabara, Rio de Janeiro, Jacarepaguá, Represa do Camorim. Distr. — Brazil.



- CHAETOSA Malloch, 1934:212. Type-locality: Chile, Los Andes. Distr. — Chile.
- CHILENSIS, nom. nov. for *annulipes* Macquart. Type-locality: "Chile". Distr. — Chile. Ref. — Albuquerque, 1949b:442.
- annulipes* Macquart, 1843a:329, pl. 23, fig. 7; 1843b:172, pl. 23, fig. 7 (preocc. Macquart, 1835). Type-locality: "Chile".
- COTHURNATA (Bigot), 1885:298 (*Anthomyia*). Type-locality: "Chile". Distr. — Chile.
- CRASSICAUDA Stein, 1911:158. Type-locality: Peru, Lago Titicaca and Bolivia, Lago Titicaca. Distr. — Peru, Bolivia.
- CURVIVENTRIS Albuquerque, 1959:1, figs. 1-8. Type-locality: Brazil, Rio de Janeiro, Teresópolis, Parque Nacional da Serra dos Órgãos. Distr. — Brazil.
- DIAPHANA Stein, 1911:167. Type-locality: Peru, Tarma and Valley of Lares. Distr. — Peru.
- ENORMIS Albuquerque, 1956i:5, figs. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil.
- FEMORALIS (Robineau-Desvoidy), 1830:545 (*Palusia*). Type-locality: France, Saint-Sauveur. Distr. — Chile; Europe. Ref. — Hennig, 1961:550, pl. 27, fig. 511, pl. 31, fig. 607. Almost certainly an erroneous record.
- geniculata* Fallén, 1825:89 (*Musca*; preocc. Degger, 1776). Type-locality: Sweden.
- IGNOBILIS Stein, 1911:171. Type-locality: Chile, Quillota. Distr. — Chile, Argentina, Juan Fernández Is. Ref. — Malloch, 1934:221, fig. 35b.
- IMMACULATA Stein, 1911:158. Type-locality: Bolivia, Sorata and Peru, Valley of Lares. Distr. — Peru, Bolivia.
- INAEQUALIS Malloch, 1934:213. Type-locality: Chile, Concepción. Distr. — Chile.
- INCULTA Stein, 1911:173. Type-locality: Chile, Caldera and Arica. Distr. — Chile.
- INCURVA Stein, 1911:169. Type-locality: Bolivia, Sorata and Lorenzopata. Distr. — Brazil.
- INIQUA Stein, 1911:175. Type-locality: Chile, Coronel and Caldera. Distr. — Chile.
- INTERMEDIA (Fallén), 1825:87 (*Musca*). Type-locality: Sweden. Distr. — Uruguay; Palaearctic region. Ref. — Hennig, 1961:567, fig. 210a, pl. 29, fig. 538, pl. 30, fig. 571. Almost certainly an erroneous record.
- INUSITATA (Malloch), 1934:224, figs. 37c and 38a (*Austrocoenosia*). Type-locality: Chile, Angol. Distr. — Chile. N. comb.
- ITATIAIENSIS (Albuquerque), 1954d:184, figs. 27-30 (*Austrocoenosia*). Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil. N. comb.
- LATIMACULATA (Albuquerque), 1956c:74, figs. 5-9 (*Austrocoenosia*). Type-locality: Brazil, Rio de Janeiro, Juçara, Angra dos Reis. N. comb.
- LATITIBIA Albuquerque, 1957e:362, figs. 10-17. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil.
- LEUCOTRICHIA Albuquerque, 1956i:1, figs. Type-locality: Brazil, Rio de Janeiro, Itatiaia, Maromba. Distr. — Brazil.

- LONGIPEDE Albuquerque, 1956i:10, figs. Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil.
- LONGISETA (Gaminara), 1930:1250 (*Rhynchoconops*). Type-locality: "Uruguay". Distr. — Uruguay. N. comb.
- MACQUART, nom. nov. for *rufipes* Macquart. Type-locality: Uruguay, Maldonado. Distr. — Brazil, Uruguay. Ref. — Albuquerque, 1949b: 439, figs. 1-12.
- rufipes* Macquart, 1851a:245, pl. 25, fig. 3; 1851b:272, pl. 25, fig. 3 (*Choetura*; preocc. Macquart, 1835). Type-locality: Uruguay, Maldonado.
- MEDIOCRIS Stein, 1911:165. Type-locality: Chile, Quillota and Coquimbo. Distr. — Chile, Argentina. Ref. — Malloch, 1934:218.
- dubia* Bigot, 1885:298 (?*Anthomyia*; preocc. Macquart, 1835). Type-locality: Chile.
- MESOFULVATA (Albuquerque), 1959:13, figs. 20-26 (*Austrocoenosia*). Type-locality: Brazil, Rio de Janeiro, Teresópolis, Parque Nacional da Serra dos Órgãos. Distr. — Brazil. N. comb.
- MINUSCULARIS Albuquerque, 1956f:5, figs. 9-16. Type-locality: Brazil, Guanabara, Rio de Janeiro, Grajaú. Distr. — Brazil.
- NIGERRIMA (Malloch), 1934:219 (*Austrocoenosia*). Type-locality: Argentina, Bariloche. Distr. — Chile, Argentina. N. comb.
- NONIESMACULATA Gaminara, 1930:1249. Type-locality: "Uruguay". Distr. — Uruguay.
- PICTA Stein, 1911:162. Type-locality: Peru, Oroya. Distr. — Peru.
- PILITIBA Stein, 1911:167. Type-locality: Peru, Valley of Lares. Distr. — Peru.
- PLUMBEA Albuquerque, 1954d:182, figs. 19-26. Type-locality: Brazil, São Paulo, S. José dos Campos. Distr. — Brazil.
- PLUMBIFRONS (Albuquerque), 1956c:77, figs. 10-16 (*Austrocoenosia*). Type-locality: Brazil, Goiás, Goiânia. Distr. — Brazil. N. comb.
- PLUMISETA Stein, 1911:168. Type-locality: Bolivia, San Carlos and Sarampiuni; Peru, Chanchamayo. Distr. — Venezuela, Guyana, Peru, Bolivia, Brazil. Ref. — Albuquerque, 1956c:71, figs. 1-4.
- PROCERA Stein, 1911:166. Type-locality: Chile, Valparaíso and Quillota. Distr. — Chile.
- PROJECTA (Malloch), 1934:223, figs. 37a and 38b (*Austrocoenosia*). Type-locality: Argentina, River Pichileufu. Distr. — Argentina. N. comb.
- ROTUNDIVENTRIS Stein, 1911:174. Type-locality: Chile, Coquimbo. Distr. — Chile, Argentina.
- RUSTICA (Albuquerque), 1956i:7, figs. (*Austrocoenosia*). Type-locality: Brazil, Rio de Janeiro, Itatiaia, Lago Azul. Distr. — Brazil. N. comb.
- SETIVENTRIS Stein, 1911:164. Type-locality: Peru, Lago Titicaca and Cuzco; Bolivia, Lago Titicaca. Distr. — Peru, Bolivia.
- STRENUA Stein, 1911:175. Type-locality: Chile, Quillota. Distr. — Chile.
- TARSATA (Snyder), 1957c:465 (*Limosia*). Type-locality: Argentina, Santa Cruz, Lago Argentina. Distr. — Argentina. N. comb.
- TINCTIFACIES (Albuquerque), 1958a:265, figs. 1-6 (*Austrocoenosia*). Type-locality: Brazil, Rio de Janeiro, Petrópolis, Alto da Mosela. Distr. — Brazil. N. comb.

- TINCTIPENIS Albuquerque, 1956i:3, figs. Type-locality: Brazil, Rio de Janeiro, Teresópolis, Parque Nacional da Serra dos Órgãos. Distr. — Brazil.
- TRICHOPIHTHALMA Albuquerque, 1959:5, figs. 9-14. Type-locality: Brazil, Rio de Janeiro, Teresópolis, Parque Nacional da Serra dos Órgãos. Distr. — Brazil.
- TUMIDA Stein, 1911:163. Type-locality: Peru, Cuzco and Chile, Palca. Distr. — Peru, Chile.
- UNIFORMIS Malloch, 1934:214. Type-locality: Argentina, Bariloche. Distr. — ?Chile, Argentina.
- VITTITHORAX Stein, 1911:163. Type-locality: Peru, Oroya. Distr. — Peru.
- WULPI, nom. nov. for *pulicaria* Wulp. Type-locality: Mexico, Guerrero, Omilteme. Distr. — Mexico.
- pulicaria* Wulp, 1897:346, pl. 8, fig. 18 (*Schoenomyza*; preocc. Zetterstedt, 1845). Type-locality: Mexico, Guerrero, Omilteme.
- XANTHOPODA (Albuquerque), 1956f:1, figs. 1-8 (*Austrocoenosia*). Type-locality: Brazil, Guanabara, Rio de Janeiro, Represa do Camorim. Distr. — Brazil. N. comb.

Unplaced species of COENOSIA s. lat.

- ALBIPILA Stein, 1918:242. Type-locality: "South America or Africa". Holotype not located, so neither correct genus nor correct provenance could be ascertained. Cf. also under "Species erroneously described from South America".
- TENUIOR (Walker), 1852:365 (*Anthomyia*). Type-locality: "Brazil". Nec *tenuior* (Walker), 1849. Distr. — Brazil. Holotype not located.

Genus CORDILUROIDES Albuquerque

- CORDILUROIDES Albuquerque, 1954k:361. Type-species, *listrata* Albuquerque (orig. des.).
- BICOLOR (Bigot), 1885:302 (*Dialyta*). Type-locality: "Mexico". Distr. — Mexico. N. comb.
- INSULARIS (Williston), 1896:371, pl. 12, figs. 123 and 123a (September) (*Coenosia*). Type-locality: St. Vincent Island. Distr. — Mexico, St. Vincent Is., Colombia, Peru, Bolivia. N. comb.
- bistriata* Wulp, 1896:343 (December); 1897:345 (*Coenosia*). Type-locality: Mexico, Orizaba and Teapa in Tabasco. N. syn.
- vittifera* Stein, 1904:487 (*Coenosia*). Type-locality: Peru, Callanga. N. syn.
- LISTRATA Albuquerque, 1954k:364, figs. 8-15. Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil.
- MEGALOPYGA Albuquerque, 1954k:362, figs. 1-7. Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Distr. — Brazil.
- NEOTROPICA Albuquerque, 1954k:367, figs. 16-24. Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil.



Genus NEODEXIOPSIS Malloch

- NEODEXIOPSIS Malloch, 1920:162. Type-species, *Dexiopsis basalis* Stein (orig. des.).
- Xenocoenosia* Malloch, 1920:162. Type-species, *Coenosia calopyga* Loew (orig. des.).
- Plumispina* Albuquerque, 1954d:177. Type-species, *longipilis* Albuquerque (orig. des.). N. SYN.
- Paradexiopsis* Albuquerque, 1955g:391. Type-species, *vittiventris* Albuquerque (orig. des.). N. SYN.
- ARIZONA Snyder, 1958a:25, fig. 12. Type-locality: U.S.A., Arizona, Tucson. Distr. — Mexico; North America.
- BRASILIENSIS (Walker), 1852:366 (*Anthomyia*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Stein, 1901:190. N. COMB.
- BREVICORNIS (Malloch), 1934:221, fig. 36 (*Austrocoenosia*). Type-locality: Chile, Chiloé Island, Ancud. Distr. — Chile, Chiloé Is. N. COMB.
- CACUMINA Snyder, 1957c:457. Type-locality: Argentina, Tucumán, Lacavera, Tafi. Distr. — Argentina.
- CALOPYGA (Loew), 1872:94 (*Coenosia*). Type-locality: U.S.A., Pennsylvania. Distr. — Peru; North America. Ref. — Hockett, 1934:81, figs. 1, 16, 31.
- CAVALATA Snyder, 1957a:224. Type-locality: Puerto Rico, El Yunque. Distr. — Puerto Rico.
- CERA Snyder, 1958a:19, fig. 9. Type-locality: Brazil, Ceará. Distr. — Brazil.
- CIRRATIPILA Snyder, 1957c:463. Type-locality: Argentina, Tucumán, Burruyacu, V. Padre Monti. Distr. — Argentina.
- CLAVACULA Snyder, 1957c:461. Type-locality: Argentina, Tucumán, Burruyacu, V. Padre Monti. Distr. — Argentina.
- CRASSICRURUS Snyder, 1957a:218. Type-locality: Puerto Rico, El Yunque. Distr. — Puerto Rico.
- CRISPISETA Snyder, 1957a:215. Type-locality: Puerto Rico, El Yunque. Distr. — Puerto Rico.
- CROCEAFRONS Snyder, 1957c:460. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- DECLIVIS (Stein), 1904:486 (*Coenosia*). Type-locality: Bolivia, Sillutincara. Distr. — Bolivia. N. COMB.
- DEVIA (Curran), 1934a:164 (*Xenocoenosia*). Type-locality: Galápagos Islands, Chatham Island. Distr. — Galápagos Is. N. COMB.
- DISCOLORISEXUS Snyder, 1957a:212. Type-locality: Puerto Rico, Yauco-Lares Road, Kilometer 22. Distr. — Puerto Rico.
- DITIPORTUS Snyder, 1957a:221. Type-locality: Puerto Rico, Yauco-Lares Road, Kilometer 22. Distr. — Puerto Rico.
- EBENIFEMUR Snyder, 1957a:220. Type-locality: Puerto Rico, El Yunque. Distr. — Puerto Rico.
- EMMESA (Malloch), 1934:216, figs. 34e-f (*Coenosia*). Type-locality: Costa Rica, Farm la Caja 8 km west of San José. Distr. — Costa Rica, Panama, Venezuela.



- EQUATOR Snyder, 1958a:15, fig. 1. Type-locality: "Ecuador". Distr. — Ecuador.
- FLAVIPALPIS Albuquerque, 1956e:198, figs. 8-13. Type-locality: Brazil, Rio de Janeiro, Itatiaia, Lago Azul. Distr. — Brazil.
- FLAVIPES (Williston), 1896:370 (*Coenosia*). Type-locality: St. Vincent Island. Distr. — Jamaica, St. Vincent Is., Puerto Rico.
- GENICULATA (Bigot), 1885:289 (*Spilogaster*). Type-locality: Argentina, Buenos Aires. Distr. — Chile, Argentina. Ref. — Snyder, 1957c:457, *australis* Malloch, 1934:215, figs. 33a and 34a-b (*Coenosia*). Type-locality: Chile, Los Andes.
- GENUPUNCTA (Stein), 1904:491 (*Coenosia*). Type-locality: Peru, Callanga. Distr. — Peru. N. comb.
- HIRTITIBIA (Stein), 1911:160 (*Coenosia*). Type-locality: Bolivia, La Paz and Chile, Palca. Distr. — Bolivia, Chile. N. comb.
- HYDROTAEIFORMIS Snyder, 1958a:8, figs. 13-15. Type-locality: Costa Rica, Higuito, San Mateo. Distr. — Costa Rica, Panama.
- INTONICLUNIS Snyder, 1957c:454. Type-locality: Argentina, Tucumán, Queb. La Toma. Distr. — Argentina.
- LANIGERA (Stein), 1918:243 (*Coenosia*). Type-locality: Colombia, Sierra S. Lorenzo. Distr. — Colombia. N. comb.
- LATIFRONS (Thomson), 1869:559 (*Coenosia*). Type-locality: Ecuador, Puna. Distr. — Ecuador. Ref. — Stein, 1910:75. N. comb.
- LINEATA (Stein), 1904:492 (*Coenosia*). Type-locality: Colombia, Bogotá. Distr. — Colombia. N. comb.
- LONGIPILIS (Albuquerque), 1954d:177, figs. 1-9 (*Plumispina*). Type-locality: Brazil, Guanabara, Rio de Janeiro, Grajaú. Distr. — Brazil. N. comb.
- LUNATISIGNA Snyder, 1957c:454. Type-locality: Argentina, Tucumán, Queb. La Toma. Distr. — Argentina.
- MACROCERA (Wulp), 1896:344, pl. 8, figs. 17 and 17a (*Coenosia*). Type-locality: Mexico, Guerrero, Omilteme and Sierra de las Aguas Escondidas. Distr. — Mexico.
- MAGNICORNIS Snyder, 1958a:13, fig. 8. Type-locality: Panama, Boquete. Distr. — Panama.
- MALDONADOI Snyder, 1957a:227. Type-locality: Puerto Rico, El Yunque. Distr. — Puerto Rico.
- MEDINAI Snyder, 1957a:223. Type-locality: Puerto Rico, El Yunque. Distr. — Puerto Rico.
- MICANS Snyder, 1957a:213. Type-locality: Puerto Rico, El Yunque. Distr. — Puerto Rico.
- MICROCHAETA (Malloch), 1934:214 (*Coenosia*). Type-locality: Uruguay, Montevideo. Distr. — Uruguay. N. comb.
- NEOAUSTRALIS Snyder, 1957c:458. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil, Argentina.
- NEOFLAVIPES Snyder, 1957a:225. Type-locality: Puerto Rico, El Yunque. Distr. — Puerto Rico.
- NEOMACROCERA Snyder, 1957c:459. Type-locality: Argentina, Tucumán, Lacavera. Distr. — Argentina.

- OBTUSILOEA (Malloch), 1934:216, figs. 33c and 34g (*Coenosia*). Type-locality: Costa Rica, Farm la Caja, 8 km west of San José. Distr. — Costa Rica.
- OCULATA (Stein), 1911:169 (*Coenosia*). Type-locality: Bolivia, Lorenzopata. Distr. — Bolivia. N. comb.
- OSCILLANS (Wulp), 1896:344 (*Coenosia*). Type-locality: Mexico, Guerrero, Amula. Distr. — Mexico. N. comb.
- OVATA (Stein), 1898:262 (*Coenosia*). Type-locality: U.S.A., Georgia and Illinois. Distr. — Bahama Is., Peru, Bolivia, Chile, Argentina; North America. Ref. — Snyder, 1958a:21, figs. 2 and 5. Almost certainly erroneous records.
- PARVULA Albuquerque, 1958a:269, figs. 7-12. Type-locality: Brazil, Rio de Janeiro, Petrópolis, Alto da Mosela. Distr. — Brazil.
- PAULISTENSIS Albuquerque, 1956e:195, figs. 1-7. Type-locality: Brazil, São Paulo, Serra da Bocaina. Distr. — Brazil.
- PENINSULA Snyder, 1958a:23, fig. 3. Type-locality: U.S.A., Florida, Merritt Is. Distr. — Bahamas, Caicos Is.; North America.
- PERUVIANA Snyder, 1958a:18, fig. 10. Type-locality: "Peru". Distr. — Peru.
- PILOSA (Stein), 1904:483 (*Coenosia*). Type-locality: Colombia, Cordillera and Peru, Callanga. Distr. — Colombia, Peru. N. comb.
- PREACUTA Snyder, 1958a:16, fig. 11. Type-locality: Costa Rica, San José. Distr. — Costa Rica.
- PRISCIPAGUS Snyder, 1958a:20. Type-locality: ?Puerto Rico, Pueblo Viejo, San Juan. Distr. — ?Puerto Rico.
- PUNCTULATA (Wulp), 1896:343 (*Coenosia*). Type-locality: Mexico, Pátzcuaro, Xucumanatlán and Omilteme. Distr. — Mexico. Ref. — Snyder, 1958a:12.
- QUINTIVENA Snyder, 1957c:453. Type-locality: Argentina, Tucumán, Queb. Carnzo. Distr. — Argentina.
- RAVA Snyder, 1957c:462. Type-locality: Argentina, Tucumán, Tafí del Valle. Distr. — Argentina.
- RECEDENS (Stein), 1904:489 (*Coenosia*). Type-locality: Peru, Callanga. Distr. — Peru, Bolivia. N. comb.
- REX Curran, 1928a:88. Type-locality: Puerto Rico, Luquillo National Forest. Distr. — Puerto Rico. Ref. — Snyder, 1957a:210.
- RUFITIBIA (Stein), 1919:161 (*Coenosia*; nom. nov. for *tibialis* Stein). Type-locality: U.S.A., Georgia, Philadelphia, Illinois. Distr. — Panama, ?Jamaica, Ecuador, Peru, Brazil, Paraguay, Argentina; North America.
- tibialis* Stein, 1898:275 (*Coenosia*; preocc. Macquart, 1843). Type-locality: U.S.A., Georgia, Philadelphia, Illinois.
- SETIPUNCTA Snyder, 1957c:455. Type-locality: Argentina, Tucumán, Queb. La Toma. Distr. — Argentina.
- SIMA Snyder, 1957c:462. Type-locality: Argentina, Tucumán, San Javier. Distr. — Argentina.
- TENUICORNIS (Wulp), 1896:343 (*Coenosia*). Type-locality: Mexico, Vera Cruz, Atoyac. Distr. — Mexico, Jamaica. N. comb.
- TRUNCATA (Stein), 1911:170 (*Coenosia*). Type-locality: Peru, Valley of Lares and Cuzco. Distr. — Peru. N. comb.

- USPALLATA Snyder, 1957c:456. Type-locality: Argentina, Mendoza, Uspallata. Distr. — Argentina.
- VITILIS (Giglio-Tos), 1894:11; 1896:32 (*Coenostia*). Type-locality: Mexico, Tuxpango. Distr. — Mexico. N. comb.
- VITTIVENTRIS (Albuquerque), 1955g:391, figs. 1-8 (*Paradexiopsis*). Type-locality: Brazil, São Paulo, São Paulo, Ipiranga. Distr. — Brazil. N. comb.
- WILLISTONI Snyder, 1958a:9. Type-locality: “?Brazil”. Distr. — Brazil.

Subfamily CYRTONEURININAE

Genus MULFORDIA Malloch

- MULFORDIA Malloch, 1928a:466. Type-species, *ferruginea* Malloch (orig. des.).
- FERRUGINEA Malloch, 1928a:467. Type-locality: Bolivia, Tumupasa. Distr. — Bolivia.
- ?LONGIPES (Stein), 1911:80 (*Mydaea*). Type-locality: Bolivia, Lorenzopata. Distr. — Bolivia. N. comb.
- SECUNDA Snyder, 1951:2. Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.

Genus NEUROTRIXA Shannon & Del Ponte

- Steinella* Malloch, 1921a:171 (preocc. Cépède, 1910). Type-species, *prima* Malloch (orig. des.) = *felsina* (Walker).
- NEUROTRIXA Shannon & Del Ponte, 1926:575. Type-species, *mirata* Shannon & Del Ponte (orig. des.) = *felsina* (Walker).
- FELSINA (Walker), 1849:955 (*Anthomyia*). Type-locality: Uruguay, Montevideo. Distr. — Brazil, Uruguay, Argentina. Ref. — Albuquerque, 1950a:241, figs. 1-4; 1954a:156. N. comb.
- limbatinevris* Macquart, 1851a:234, pl. 24, fig. 2; 1851b:261, pl. 24, fig. 2 (*Spilogaster*). Type-locality: “Argentina” (not Tasmania, as stated by Macquart). N. SYN.
- limbatinervis*, error.
- fuscinevris* Macquart, 1851a:236, pl. 24, fig. 6; 1851b:263, pl. 24, fig. 6 (*Limnophora*). Type-locality: Argentina, Buenos Aires. N. SYN.
- fuscinervis*, error.
- prima* Malloch, 1921a:172 (*Steinella*). Type-locality: Uruguay, Montevideo. N. SYN.
- mirata* Shannon & Del Ponte, 1927:575. Type-locality: Argentina, San Isidro. N. SYN.

Genus XENOTHORACOAETA Malloch

- XENOTHORACOAETA Malloch, 1921a:170. Type-species, *prima* Malloch (orig. des.).
- Airalips* Snyder, 1949a:9. Type-species, *Spilogaster plumata* Stein (orig. des.).



- CAERULEA (Snyder), 1949a:12 (*Airalips*). Type-locality: Venezuela, Caracas. Distr. — Venezuela. N. comb.
- FUSCOMARGINATA (Snyder), 1949a:12 (*Airalips*). Type-locality: Colombia, Cali, western Cordillera. Distr. — Colombia. N. comb.
- GOLBACHI Snyder, 1957c:479. Type-locality: Argentina, Tucumán, Queb. la Toma. Distr. — Argentina.
- FLUMATA (Stein), 1904:446 (*Spilogaster*). Type-locality: Peru, Vilcanota. Distr. — Peru, Bolivia, Brazil, Paraguay, Argentina. Ref. — Snyder, 1949a:9. N. comb.
- PRIMA Malloch, 1921a:170. Type-locality: Colombia, Villavicencio, Quatiquia River. Distr. — Colombia, Peru.
- differentia* Snyder, 1949a:10 (*Airalips*). Type-locality: Peru, Rio Charape.
- RUFIGUTTATA (Macquart), 1851a:231, pl. 23, fig. 16; 1851b:258, pl. 23, fig. 16 (*Aricia*). Type-locality: Brazil, Bahia. Distr. — Brazil. Ref. — Albuquerque, 1951b:1, figs. 1-7. N. comb.

Genus PSEUDOPTILOLEPIS Snyder

- PSEUDOPTILOLEPIS Snyder, 1949a:14. Type-species, *fulvapoda* Snyder (orig. des.).
- CONFUSA Snyder, 1949a:19, fig. 11. Type-locality: Venezuela, Quebrada Secca. Distr. — Venezuela.
- FLUMINENSIS Albuquerque, 1954h:253, figs. 1-8. Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Distr. — Brazil.
- FULVAPODA Snyder, 1949a:16, figs. 6 and 8. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- LATIPALPIS (Stein), 1918:216 (*Mydaea*). Type-locality: Brazil, Teresópolis and São Paulo. Distr. — Brazil. N. comb.
- NIGRIPODA Snyder, 1949a:17, figs. 7 and 9. Type-locality: Panama, Balboa. Distr. — Nicaragua, Panama.
- NUDAPLEURA Snyder, 1949a:19, fig. 10. Type-locality: Paraguay, Villarica. Distr. — Paraguay.

Genus CHAETAGENIA Malloch

- CHAETAGENIA Malloch, 1928b:308. Type-species, *stigmatica* Malloch (orig. des.).
- STIGMATICA Malloch, 1928b:308. Type-locality: Brazil, São Vicente. Distr. — Brazil.

Genus HEMICHLORA Wulp

- HEMICHLORA Wulp, 1896:303. Type-species, *Curtonevra vittigera* Bigot (mon.) = *scordalus* (Walker).
- SCORDALUS (Walker), 1861:313 (*Pyrellia*). Type-locality: "Mexico". Distr. — Mexico, Guatemala, Costa Rica. Ref. — Snyder, 1954a:423.
- vittigera* Bigot, 1888a:613; 1888b:clxxxii (*Curtonevra*). Type-locality: "Mexico".

Genus NEOMUSCINA Townsend

- NEOMUSCINA Townsend, 1919:541. Type-species, *cavicola* Townsend (orig. des.) = *tripunctata* (Wulp).
- Spilopteromyia* Malloch, 1921d:422. Type-species, *Spilogaster apicata* Stein (orig. des.).
- Scutellomusca* Townsend, 1931b:313. Type-species, *Musca scutellaris* Fabricius (orig. des.). N. SYN.
- APICATA (Stein), 1904:442 (*Spilogaster*). Type-locality: Peru, Vilcanota. Distr. — Colombia, Peru. Ref. — Snyder, 1949c:37; 1954a:428.
- ATINCTA Snyder, 1949c:27. Type-locality: Panama, Barro Colorado Is. Distr. — Costa Rica, Panama.
- ATINCTICOSTA Snyder, 1949c:37. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- CAPALTA Snyder, 1949c:17. Type-locality: Trinidad, Port of Spain. Distr. — Barro Colorado Is., Trinidad.
- CURRANI Snyder, 1949c:35. Type-locality: Panama, Barro Colorado Is. Distr. — Barro Colorado Is.
- DORSIPUNCTA (Stein), 1918:216 (*Mydaea*). Type-locality: Mexico, Tapachula and Oaxaca. Distr. — Mexico, Costa Rica, Panama, Venezuela, Brazil. N. COMB.
- davida* Snyder, 1949c:14. Type-locality: Panama, David. N. SYN.
- INFLEXA (Stein), 1918:224 (*Cyrtoneurina*). Type-locality: Brazil, São Paulo. Distr. — Brazil. Ref. — Snyder, 1949c:18.
- INSTABILIS Snyder, 1949c:30. Type-locality: Panama, David. Distr. — Panama, Trinidad, Brazil.
- MACROPS Snyder, 1949c:16. Type-locality: Venezuela, Caracas. Distr. — Venezuela.
- MEDIANA Snyder, 1949c:19. Type-locality: Brazil, Guanabara, Rio de Janeiro, Botanical Garden. Distr. — Brazil.
- MEXICANA (Macquart), 1843a:315, pl. 21, fig. 9; 1843b:158, pl. 21, fig. 9 (*Curtonevra*). Type-locality: "Mexico". Distr. — Mexico.
- NEOSIMILIS Snyder, 1949c:33. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil, Paraguay.
- NIGRICOSTA Snyder, 1949c:22. Type-locality: Trinidad, Port of Spain. Distr. — Trinidad, Guyana.
- NUDISTIGMA Snyder, 1949c:25. Type-locality: Venezuela, Caracas. Distr. — Venezuela.
- PARILIS (Giglio-Tos), 1893:6; 1896:14 (*Cyrtoneurina*). Type-locality: Mexico, Oaxaca. Distr. — Mexico. Ref. — Snyder, 1954a:426.
- paralis*, error.
- PICITIPPENNIS (Bigot), 1878:39 (*Curtonevra*). Type-locality: "Brazil". Distr. — Venezuela, Brazil, Paraguay. Ref. — Snyder, 1949c:29.
- pictipennis* Bigot, 1875:lxv (*Curtonevra*), *nomen nudum*.
- fulva* Snyder, 1949c:29 (as a ssp. of *nudinervis* Stein). Type-locality: Paraguay, Villarica.
- ssp. *NUDINERVIS* (Stein), 1918:227 (*Cyrtoneurina*, as a species). Type-locality: Mexico, Cordoba. Distr. — Mexico, Guatemala, El Salvador, Costa Rica, Panama, Venezuela, Guyana, Peru. Ref. — Snyder, 1949c:27.



- PRAETASETA Snyder, 1954a:428. Type-locality: Guatemala, La Providencia. Distr. — Mexico, Guatemala, ?Brazil.
- RUFOSCUTELLA Dodge, 1955:150. Type-locality: U.S.A., Florida, Keys. Distr. — Cuba; U.S.A., Florida.
- ssp. FARRI Dodge, 1955:151. Type-locality: Jamaica, Clarendon, 1 mile W of Jacob's Hut. Distr. — Jamaica.
- SANESPRA Snyder, 1949c:12. Type-locality: Brazil, Espírito Santo. Distr. — Brazil.
- SCHADEI Snyder, 1949c:32. Type-locality: Paraguay, Villarica. Distr. — Paraguay.
- SCUTELLARIS (Fabricius), 1805:293 (*Musca*). Type-locality: "South America". Distr. — South America.
- SIMILATA Snyder, 1949c:33. Type-locality: Trinidad, Port of Spain. Distr. — Trinidad, Venezuela, Guyana.
- SPARSIPLUMATA (Stein), 1918:219 (*Mydaea*). Type-locality: Brazil, Rio Grande do Sul. Distr. — Brazil. N. comb.
- STABILIS (Stein), 1911:86 (*Mydaea*). Type-locality: Peru, mouth of Pachitea River. Distr. — Peru.
- TINCTINERVIS (Stein), 1918:221 (*Mydaea*). Type-locality: Brazil, São Paulo and S. Catarina. Distr. — Brazil.
- TRANSPORTA Snyder, 1949c:23. Type-locality: Africa, Ghana, Accra, presumed to be imported from South America. Distr. — South America. (Letter from Sabrosky to Emden, 9.3.1951: "The lot from which it came consisted of 'Insects collected on desinsected aircraft, Sta. Accra'.")
- TRIPUNCTATA (Wulp), 1896:305 (*Muscina*). Type-locality (restricted): Mexico, Guerrero, Amula. Distr. — Mexico, Puerto Rico, Venezuela; southern U.S.A. Ref. — Snyder, 1949c:6; 1954a:425.
- texana* Hough, 1899:25, fig. 7 (*Muscina*). Type-locality: U.S.A., Texas.
- cavicola* Townsend, 1919:541. Type-locality: U.S.A., Arizona, East Verde River.
- TRISETA Snyder, 1949c:10. Type-locality: Guatemala, Guatemala City. Distr. — Mexico, Guatemala.
- VECTA (Giglio-Tos), 1893:6; 1896:14, fig. 4 (*Cyrtoneurina*). Type-locality: Mexico, Tuxpango. Distr. — Mexico. Ref. — Snyder, 1954a:427.
- linea* Wulp, 1896:304, pl. 7, fig. 22 (*Muscina*). Type-locality (restricted): Mexico, Guerrero, Xucumanatlán.
- ZOSTERIS (Shannon & Del Ponte), 1926:579 (*Muscina*). Type-locality: Argentina, Tucumán. Distr. — Brazil, Argentina. Ref. — Snyder, 1949c:8; 1954a:427.
- fulvifrons* Snyder, 1949c:20. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus CYRTONEURINA Giglio-Tos

- CYRTONEURINA Giglio-Tos, 1893:5. Type-species, *uber* Giglio-Tos (Coquillett, 1910:530).
- Clinopera* Wulp, 1896:305. Type-species, *hieroglyphica* Wulp (Coquillett, 1910:535) = *uber* Giglio-Tos.

- Cyrtoneuropsis* Malloch, 1925:91. Type-species, *Spilogaster veniseta* Stein (orig. des.).
- Mallocharia* Curran, 1934b:462. Type-species, *beebei* Curran (orig. des.).
- Dyadimyia* Séguy, 1937:238 and 340. Type-species, *aurea* Séguy (orig. des.) = *conspersa* (Stein). N. SYN.
- ARMIPES (Stein), 1911:88 (*Mydaea*). Type-locality: Peru, Urubamba River, mouth of Pachitea River and Pto. Bermúdez. Distr. — Panama, Trinidad, Guyana, Surinam, Brazil. Ref. — Snyder, 1954a:455.
- BEEBET (Curran), 1934b:462 (*Mallocharia*). Type-locality: Guyana, Kartabo. Distr. — Trinidad, Guyana, Surinam, Brazil, Paraguay, Argentina. Ref. — Snyder, 1954a:445.
- BISETA Snyder, 1954a:436. Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil.
- BRUNNEA (Hough), 1900:220, fig. 10 (*Muscina*). Type-locality: Brazil, Mato Grosso, Corumbá. Distr. — Brazil. N. COMB.
- CONFUSA Snyder, 1954a:437. Type-locality: Panama, Partilla Point. Distr. — Costa Rica, Panama, Brazil.
- CONSPERSA (Stein), 1911:89 (*Mydaea*). Type-locality: Peru, Pto. Bermúdez. Distr. — Peru, Guyana, French Guiana, Brazil. Ref. — Snyder, 1954a:457.
- aurea* Séguy, 1937:341 (*Dyadimyia*). Type-locality: French Guyana, Charvein. N. SYN.
- CONTINENS Snyder, 1954a:459. Type-locality: Brazil, São Paulo. Distr. — Brazil, Argentina.
- COSTALIS (Walker), 1852:408 (?*Notiphila*). Type-locality: "Brazil" Distr. — Brazil. N. COMB.
- nearmipes* Snyder, 1954a:437. Type-locality: Brazil, São Paulo. N. SYN.
- CRISPASETA Snyder, 1954a:438, figs. 24, 35, 50 and 51. Type-locality: Brazil, Natal. Distr. — Peru, Brazil.
- CYLINDRICA (Macquart), 1846a:328; 1846b:200 (*Cyrtonevra*). Type-locality: "Brazil". Distr. — Brazil. N. COMB.
- DUBIA Snyder, 1954a:453, figs. 19, 31, 42 and 58. Type-locality: Peru, Junín, Chanchamayo. Distr. — Guatemala, Honduras, Costa Rica, Panama, Peru, Guyana, Brazil.
- FUSCICOSTA (Curran), 1934b:466 (*Cyrtoneuropsis*). Type-locality: Guyana, Kartabo. Distr. — Guyana, Brazil. Ref. — Snyder, 1954a:457.
- FUSCISQUAMA Snyder, 1954a:450. Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil.
- GEMINA (Wiedemann), 1830:436 (*Anthomyia*). Type-locality: "Brazil". Distr. — Guatemala, Honduras, Nicaragua, El Salvador, Costa Rica, Panama, Grenada, Trinidad, Colombia, Peru, Venezuela, Guyana, Brazil. Ref. — Snyder, 1954a:454.
- immunda* Stein, 1911:90 (*Mydaea*). Type-locality: Peru, mouth of Pachitea River and Urubamba River.
- GEMINATA (Stein), 1904:439 (*Spilogaster*). Type-locality: La Pita (Guatemala or Panama?). Distr. — Mexico, Guatemala, Honduras, El Salvador, Costa Rica, Panama, Colombia, Peru, Venezuela, Brazil, Paraguay. Ref. — Snyder, 1954a:436.

- GLUTA Giglio-Tos, 1893:6; 1896:15. Type-locality: Mexico, Tuxpango. Distr. — Mexico, Guatemala, Costa Rica, Panama, Venezuela, Guyana. Ref. — Snyder, 1954a:458.
- Umbisquama* Stein, 1918:225. Type-locality: Mexico, Vera Cruz, Dona Maria and Santa Lucrezia.
- GRISEA (Fabricius), 1805:293 (*Musca*; preocc. Villers, 1789). Type-locality: "South America". Distr. — South America.
- INCOGNITA Snyder, 1954a:452, fig. 17. Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil.
- INUBER Giglio-Tos, 1893:6; 1896:15. Type-locality: Mexico, Solco. Distr. — Mexico, Guatemala, El Salvador, Costa Rica, Panama, Guyana, ?Argentina. Ref. — Snyder, 1954a:440.
- pellez* Giglio-Tos, 1893:6; 1896:16. Type-locality: Mexico, Tuxpango.
- digramma* Wulp, 1896:308 (*Clinopera*). Type-locality (restricted): Mexico, Guerrero, Chilpancingo.
- diagramma*, error.
- MACULIPENNIS (Macquart), 1843a:320, pl. 22, fig. 3; 1843b:163, pl. 22, fig. 3 (*Spilogaster*). Type-locality: "Guiana". Distr. — French Guyana. N. COMB.
- MELLINA Stein, 1918:226. Type-locality: Brazil, Santa Catarina. Distr. — Brazil, Paraguay. Ref. — Snyder, 1954a:440.
- MIMICA Snyder, 1954a:448. Type-locality: Panama, Quipo. Distr. — Panama.
- MONSTRATA (Wulp), 1896:309 (*Clinopera*). Type-locality: Mexico, Guerrero, Xucumanatlán. Distr. — Mexico.
- MULTOMACULATA (Stein), 1904:434 (*Spilogaster*). Type-locality: Colombia, Cordillera. Distr. — Costa Rica, Colombia, Peru, Guyana, Brazil, Bolivia. Ref. — Snyder, 1954a:443.
- multimaculata*, error.
- NEOTRITA Snyder, 1954a:445. Type-locality: Peru, Iquitos. Distr. — Peru.
- PALLIPES Stein, 1918:228. Type-locality: "Paraguay". Distr. — Paraguay.
- POLYSTIGMA (Wulp), 1896:309, pl. 7, fig. 25 (*Clinopera*). Type-locality (restricted): Mexico, Medellín near Vera Cruz. Distr. — Mexico, Guatemala, El Salvador, Costa Rica, Panama, Brazil, Paraguay. Ref. — Snyder, 1954a:451.
- PRAENUBILA Snyder, 1954a:444, figs. 8, 9, 26, 37 and 53. Type-locality: Trinidad, Port of Spain. Distr. — Panama, Grenada, Trinidad, Peru, Brazil.
- PROTOSSETOSA Snyder, 1954a:446. Type-locality: Brazil, Natal. Distr. — Brazil.
- RESCITA (Walker), 1861:315 (*Aricia*). Type-locality: Mexico. Distr. — Mexico, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Cuba, St. Vincent Is., Trinidad, Colombia, Venezuela, Guyana, Peru, Bolivia, Brazil, Paraguay. Ref. — Snyder, 1954a:448.
- setinervis* Thomson, 1869:549 (*Anthomyia*). Type-locality: Ecuador, Puna.
- frontina* Wulp, 1896:306 (*Clinopera*). Type-locality: Mexico, Tabasco, Teapa.
- pterostigma* Wulp, 1896:309 (*Clinopera*). Type-locality: Mexico, Tabasco, Teapa and Frontera.
- maculipennis* Williston, 1896:368, pl. 12, figs. 121 and 121a (*Cyrtoneura*; preocc. Macquart, 1843). Type-locality: St. Vincent Is.

- SCUTELLATA (Johnson), 1919:440 (*Clinopera*). Type-locality: Jamaica, Liguanea Plain near Kingston. Distr. — Jamaica.
- SERIATA (Stein), 1911:91 (*Mydaea*). Type-locality: Peru, Valley of Lares and Urubamba River. Distr. — Colombia, Peru, Brazil. Ref. — Snyder, 1954a:457.
- SPILOPTERA (Wiedemann), 1830:433 (*Anthomyia*). Type-locality: "Brazil". Distr. — Honduras, Panama, Ecuador, Surinam, Peru, Bolivia, Brazil. Ref. — Snyder, 1954a:447.
- STEINI Snyder, 1954a:456. Type-locality: Mexico, Tapachula. Distr. — Mexico, Guatemala, El Salvador, Honduras, Panama, Colombia.
- TRITA (Stein), 1911:87 (*Mydaea*). Type-locality: Bolivia, San Carlos. Distr. — Costa Rica, Panama, Bolivia, ?Brazil. Ref. — Snyder, 1954a:452.
- UBER Giglio-Tos, 1893:7; 1896:17. Type-locality: Mexico, Tuxpango. Distr. — Mexico, Guatemala, El Salvador, Honduras, Nicaragua, Panama, Grenada, Trinidad, Colombia, Peru, Venezuela, Guyana. Ref. — Snyder, 1954a:434.
- hieroglyphica* Wulp, 1896:307, pl. 7, fig. 23 (*Clinopera*). Type-locality: Mexico, Guerrero, Tierra Colorada; Tabasco, Teapa and Frontera.
- perspicua* Stein, 1911:86 (*Mydaea*). Type-locality: Peru, mouth of Pachitea River.
- VARICOLOR (Hough), 1900:226, figs. 14 and 15 (*Muscina*). Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães). Distr. — Brazil. Ref. — Albuquerque, 1955f:8, figs. 11-15.
- VENISETA (Stein), 1904:443 (*Spilogaster*). Type-locality: Brazil, Pará. Distr. — Mexico, El Salvador, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Trinidad, Guyana, Peru, Bolivia, Brazil, Argentina. Ref. — Snyder, 1954a:439.
- WALKERI, nom. nov. for *maculipennis* Walker. Type-locality: "Brazil". Distr. — Brazil. Ref. — Snyder, 1954a:458.
- maculipennis* Walker, 1852:357 (*Anthomyia*; preocc. Macquart, 1843). Type-locality: "Brazil".
- WÜLPF Snyder, 1954a:449, figs. 14 and 15. Type-locality: Panama, Balboa. Distr. — Mexico, Nicaragua, Costa Rica, Panama, Trinidad.

Genus POLIETINA Schnabl & Dziedzicki


- POLIETINA Schnabl & Dziedzicki, 1911:218. Type-species, *Aricia pruinosa* Macquart (as *pruinus* Bigot) (orig. des.).
- Poecilophaonia* Malloch, 1921a:171. Type-species, *Aricia flavithorax* Stein (orig. des.).
- Smithomyia* Malloch, 1921b:42. Type-species, *Mydaea concinna* Wulp (orig. des.) = *pruinosa* (Macquart).
- Chaetypopleura* Enderlein, 1927:52. Type-species, *steini* Enderlein (orig. des.).
- Lasiomala* Enderlein, 1927:52. Type-species, *Aricia flavithorax* Stein (orig. des.).
- BASICINCTA (Stein), 1904:421 (*Aricia*). Type-locality: Peru, Callanga Distr. — Peru, Bolivia.



- BICOLOR Albuquerque, 1956h:10, figs. Type-locality: Brazil, Rio de Janeiro, Tinguá. Distr. — Brazil.
- FLAVIDICINCTA (Stein), 1904:420 (*Aricia*). Type-locality: Bolivia, Songo and Colombia, Cordillera. Distr. — Venezuela, Colombia, Bolivia.
- FLAVITHORAX (Stein), 1904:416 (*Aricia*). Type-locality: Peru, Callanga. Distr. — Venezuela, Peru, Bolivia, Brazil. Ref. — Albuquerque, 1956h:13, figs.
- MAJOR Albuquerque, 1956h:17, figs. Type-locality: Brazil, Mato Grosso, Salobra. Distr. — Brazil.
- MELINA (Stein), 1904:426 (*Aricia*). Type-locality: Bolivia, San Antonio. Distr. — Bolivia. N. comb.
- MINOR Albuquerque, 1956h:7, figs. Type-locality: Brazil, Rio de Janeiro, Angra dos Reis, Fazenda Jaquiba. Distr. — Brazil.
- ORBITALIS (Stein), 1904:427 (*Aricia*). Type-locality: Peru, Vilcanota. Distr. — Peru, Bolivia, Brazil, Argentina. Ref. — Albuquerque, 1956h:3, figs.
- PRUINOSA (Macquart), 1846a:329; 1846b:201 (*Aricia*). Type-locality: U.S.A., Texas, Galveston. Distr. — Mexico; North America.
- concinna* Wulp, 1896:317 (*Mydaea*). Type-locality: Mexico, Guerrero, Xucumanatlán.
- RUBELLA (Wulp), 1896:314, pl. 8, figs. 1 and 1a (*Hyetodesia*). Type-locality: Mexico, Guerrero, Omilteme. Distr. — Mexico, Venezuela.
- STEINI (Enderlein), 1927:53 (*Chaetypopleura*). Type-locality: Brazil, Santa Catarina, Corupá (as Colonie Hansa). Distr. — Brazil.

Genus NEOMUSCA Malloch

- Philornis* Meinert, 1890:315 (preocc. Selys, 1839). Type-species, *molesta* Meinert (mon.). N. SYN.
- NEOMUSCA Malloch, 1921b:41. Type-species, *Mydaea obscura* Wulp (orig. des.).
- Mesembrinellopsis* Townsend, 1927:208 and 328. Type-species, *mima* Townsend (orig. des.). N. SYN.
- AITKENI (Dodge), 1963:241, figs. 1-6 (*Philornis*). Type-locality: Trinidad, Fort Read. Distr. — Trinidad. N. comb.
- ANGUSTIFRONS (Loew), 1861:41 (*Hylemyia*). Type-locality: Cuba. Distr. — Costa Rica, Cuba, Trinidad, Guyana, Brazil. Ref. — Dodge & Aitken, 1968:144. N. comb.
- anomala* Jaennicke, 1867a:377, pl. 2, fig. 4 (*Mesembrina*). Type-locality: "Cuba".
- BLANCHARDI (Garcia), 1952:289 (*Philornis*). Type-locality: Argentina, Corrientes. Distr. — Argentina. N. comb.
- CARINATA (Dodge), 1968:160, fig. 1c (*Philornis*). Type-locality: Costa Rica, Higuito, San Mateo. Distr. — Costa Rica. N. comb.
- CINNAMOMINA (Stein), 1918:214 (*Mydaea*). Type-locality: "Paraguay". Distr. — Brazil, Paraguay.
- DECEPTIVA (Dodge & Aitken), 1968:149 (*Philornis*). Type-locality: Trinidad, Fishing Pond. Distr. — Trinidad. N. comb.



- DOWNSI (Dodge & Aitken), 1968:138, figs. 1-5 (*Philornis*). Type-locality: Trinidad, Cunaripo R. C. School. Distr. — Trinidad Is. N. COMB.
- FALSIFICA (Dodge & Aitken), 1968:141, figs. 11-13 (*Philornis*). Type-locality: Trinidad, Bonaire Road, Cunaripo. Distr. — Costa Rica, Panama, Trinidad, Ecuador. N. COMB.
- FASCIVENTRIS (Wulp), 1896:318 (*Mydaea*). Type-locality: Mexico, Tabasco, Teapa. Distr. — Mexico. Ref. — Dodge, 1963:243, fig. 9. N. COMB.
- FUMICOSTA (Dodge), 1968:163 (*Philornis*). Type-locality: Brazil, Goiás, Campinas. Distr. — Brazil. N. COMB.
- GLAUCINIS (Dodge & Aitken), 1968:150 (*Philornis*). Type-locality: Trinidad, St. Patrick's Estate, Arima Valley. Distr. — Trinidad. N. COMB.
- MIMA (Townsend), 1927:208 and 328 (*Mesembrinellopsis*). Type-locality: Brazil, São Paulo, Alto da Serra (= Paranapiacaba). Distr. — Brazil. N. COMB.
- MIMICOLA (Dodge), 1968:159, fig. 1b (*Philornis*). Type-locality: U.S.A., Texas, Uvalde. Distr. — Peru; U.S.A., Texas. N. COMB.
- MOLESTA (Meinert), 1890:304-317, pl. 6, figs. 1-8 (*Philornis*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Dodge, 1968:155. N. COMB.
- NIELSENI (Dodge), 1968:157 (*Philornis*). Type-locality: "Argentina". Distr. — Argentina. N. COMB.
- NIGER (Dodge & Aitken), 1968:152 (*Philornis*). Type-locality: Trinidad, St. Patrick's Estate, Arima Valley. Distr. — Trinidad. N. COMB.
- OBSCURA (Wulp), 1896:317, pl. 8, figs. 3 and 3a (*Mydaea*). Type-locality: Mexico, Northern Yucatán. Distr. — Mexico, Puerto Rico, Ecuador, Peru; U.S.A. Ref. — Dodge, 1968:158, fig. 1a.
- PICI (Macquart), 1854:659, pl. 20, II figs. 1-4 (*Aricia*). Type-locality: "S. Domingo". Distr. — Costa Rica, Panama, Cuba, S. Domingo, Puerto Rico, Trinidad, Brazil, Argentina.
- QUERULA (Dodge & Aitken), 1968:152 (*Philornis*). Type-locality: Trinidad, Arima Textile Mills, Arima. Distr. — Trinidad. N. COMB.
- RETTEMEYER (Dodge), 1963:245, fig. 10 (*Philornis*). Type-locality: Panama, Barro Colorado Is. Distr. — Panama. N. COMB.
- SABROSKYI (Albuquerque), 1957f:11 (*Philornis*; nom. nov. for *blanchardi* Albuquerque). Type-locality: Brazil, Guanabara, Rio de Janeiro, Jardim Botânico. Distr. — French Guyana, Brazil.
- blanchardi* Albuquerque, 1953a:141, figs. 1-8 (*Philornis*; preocc. Garcia, 1952). Type-locality: Brazil, Guanabara, Rio de Janeiro, Jardim Botânico.
- SANGUINIS (Dodge & Aitken), 1968:148 (*Philornis*). Type-locality: Trinidad, Cunapo Southern Road, 3/4 mile post, Sangre Grande. Distr. — Trinidad. N. COMB.
- SCHILDI (Dodge), 1963:242, figs. 7-8 (*Philornis*). Type-locality: Costa Rica, Higuito, San Mateo. Distr. — Costa Rica. N. COMB.
- SEGUYI (Garcia), 1952:286 (*Philornis*). Type-locality: Argentina, Misiones. Distr. — Argentina. N. COMB.
- SETINERVIS (Dodge), 1963:246 (*Philornis*). Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil. N. COMB.
- SPERMOPHILA (Townsend), 1895a:79; 1895b:174 (*Mydaea*). Type-locality: Jamaica, Kingston. Distr. — Jamaica. N. COMB.

- STEINI, nom. nov. for *angustifrons* Stein. Type-locality: Colombia, Cordillera. Distr. — Colombia.
- angustifrons* Stein, 1904:448 (*Spilogaster*; preocc. Loew, 1861). Type-locality: Colombia, Cordillera.
- TORQUANS (Nielsen), 1913:252, fig. 3 (*Mydaea*). Type-locality: Chile, Concepcion. Distr. — Chile. Ref. — Dodge, 1968:156. N. comb.
- TRINITENSIS (Dodge & Aitken), 1968:146, figs. 14-17 (*Philornis*). Type-locality: Trinidad, Vega de Oropouche. Distr. — Trinidad. N. comb.
- UMANANI (Garcia), 1952:284 (*Philornis*). Type-locality: Argentina, Corrientes. Distr. — Argentina. N. comb.
- UNIVITTATA (Dodge), 1968:161 (*Philornis*). Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil. N. comb.
- VESPIDICOLA (Dodge), 1968:161 (*Philornis*). Type-locality: Peru, Oxapampa. Distr. — Peru. N. comb.
- ZETEKI (Dodge), 1963:246, fig. 11 (*Philornis*). Type-locality: Panama, El Cermeno. Distr. — Panama. N. comb.

Genus CHARADRELLA Wulp

- CHARADRELLA Wulp, 1896:341. Type-species, *macrosoma* Wulp (mon.).
- MACROSOMA Wulp, 1896:341, pl. 8, figs. 16, 16a and 16b. Type-locality: Mexico, Northern Yucatán. Distr. — Mexico, Ecuador, Venezuela, Brazil.
- MALACOPHAGA Souza Lopes, 1938a:927, figs. 1-3. Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil.

Genus ARTHURELLA Albuquerque

- ARTHURELLA Albuquerque, 1954a:151. Type-species, *nudiseta* Albuquerque (orig. des.).
- NUDISETA Albuquerque, 1954a:151, figs. 1-8. Type-locality: Chile, El Canelo. Distr. — Chile.

Genus HEMIXANTHOMYIA Albuquerque


- HEMIXANTHOMYIA Albuquerque, 1954g:323. Type-species, *marginata* Albuquerque (orig. des.).
- MARGINATA Albuquerque, 1954g:323, figs. 1-5. Type-locality: Brazil, Minas Gerais, Rio Doce, Calado. Distr. — Brazil.

Genus CHORTINUS Aldrich

- CHORTINUS Aldrich, 1932:23. Type-species, *bequaerti* Aldrich (orig. des.).
- BEQUAERTI Aldrich, 1932:23. Type-locality: Guatemala, Santa Emilia, Pochuta and La Providencia, Obispo. Distr. — Guatemala.

Genus CARIOCAMYIA Snyder

- CARIOCAMYIA Snyder, 1951:1. Type-species, *maculosa* Snyder (orig. des.).



- MACULOSA Snyder, 1951:1. Type-locality: Brazil, Guanabara, Rio de Janeiro, Grajaú. Distr. — Colombia, Brazil.
- VIBRISSATA (Stein), 1918:222 (*Mydaea*). Type-locality: Brazil, São Paulo, Iguape. Distr. — Brazil. N. comb.

Unplaced Species of Cyrtoneurinae

- BREVIPECTINATA Stein, 1918:215 (*Mydaea*). Type-locality: Brazil, São Paulo. Distr. — Brazil.
- FULVIPES Bigot, 1888a:613; 1888b:clxxxii (*Curtonevra*). Type-locality: Mexico. Distr. — Mexico.

Subfamily STOMOXYINAE

Genus STOMOXYS Geoffroy

- STOMOXYS Geoffroy, 1762:449 and 538. Type-species, *Conops calcitrans* Linnaeus (Opinion 441).
- CALCITRANS (Linnaeus), 1758:604 (*Conops*). Type-locality: Sweden. Distr. — Mexico, Guatemala, Nicaragua, Costa Rica, Panama, Bermuda, Jamaica, Bahamas, Grenada, Barbados, Puerto Rico, St. Vincent, Haiti, Dominica, Trinidad, Venezuela, Guyana, Colombia, Ecuador, Bolivia, Brazil, Uruguay, Chile, Argentina, Galápagos Is.; cosmopolitan. Ref. — Hennig, 1964:1039, figs. 407, 410-414, 417.
- nebulosa* Fabricius, 1805:282. Type-locality: "In Americae meridionalis insulis".
- sugillatrix* Robineau-Desvoidy, 1830:386. Type-locality: "Brazil".
- geniculata* Macquart, 1846a:320; 1846b:192. Type-locality: "Brazil".
- NIGRA Macquart, 1851a:212, pl. 22, figs. 5 and 5a; 1851b:239, pl. 22, figs. 5 and 5a. Type-locality: "Île Bourbon" (= Reunion). Distr. — Venezuela; Ethiopian and Malagasy regions. Almost certainly an erroneous record.

Genus HAEMATOBIA Le Peletier & Serville

- HAEMATOBIA Le Peletier & Serville in Latreille et al., 1828:499. Type-species, *Conops irritans* Linnaeus (Westwood, 1840:140).
- Siphona* Meigen, 1803:281, of authors not of Meigen. (True *Siphona* belongs to the family Tachinidae).
- Lyperosia* Rondani, 1856:93. Type-species, "*Stomoxis irritans* Mg." (orig. des.).
- IRRITANS (Linnaeus), 1758:604 (*Conops*). Type-locality: Sweden. Distr. — Mexico, Guatemala, El Salvador, Nicaragua, Costa Rica, Panama, Puerto Rico, Jamaica, Virgin Is., Haiti, S. Domingo, Colombia, Venezuela, Guyana, Brazil, Chile; North America, Europe, Asia, north Africa, Hawaii. Ref. — Hennig, 1964:1054, figs. 420, 424, 425, 428.
- serrata* Robineau-Desvoidy, 1830:389. Type-locality: South France.

MINUTA (Bezzi), 1892:192 (*Lyperosia*). Type-locality: Somalia, Obbia. Distr. — Venezuela; Mediterranean subregion, Africa, Indian subregion. Almost certainly an erroneous record.

Genus NEIVAMYIA Pinto & Fonseca

NEIVAMYIA Pinto & Fonseca, 1930a:243. Type-species, *lutzi* Pinto & Fonseca (orig. des.) = *flavicornis* (Malloch).

ANTUNESI Souza Lopes, 1955:415, figs. 2-8. Type-locality: Colombia, Sardinata, Meta. Distr. — Colombia.

FLAVICORNIS (Malloch), 1928b:318 (*Bdellolarynx*). Type-locality: Surinam, Upper Para. Distr. — Trinidad, Guyana, Surinam, Brazil.

lutzi Pinto & Fonseca, 1930a:244, plates 1-3, fig. 1. Type-locality: "Brazil".

LATIFRONS Malloch, 1932:431. Type-locality: Peru, Iquitos. Distr. — Peru, Brazil.

TRAVASSOSI Souza Lopes & Mangabeira, 1938:288, pl. 1, figs. 5-6, pl. 2, figs. 1-2, pl. 3, figs. 3-4, pl. 4, figs. 1-2. Type-locality: Brazil, Amazonas, Manacapuru. Distr. — Brazil.

SUMMARY OF ERRONEOUS OR DUBIOUS RECORDS FROM SOUTH AMERICA

Full reference is made to these species and records in the main body of the Catalogue.

FANNIA POLYCHAETA Stein
MUSCA ALBINA Wiedemann
MUSCA XANTHOMELAS Wiedemann
ORTHELLIA CORNICINA Fabricius
HYDROTAEA CYRTONEURINA Zetterstedt
HYDROTAEA LATA Walker
PHAONIA PULVILLATA Stein
HELINA MACULIPENNIS Zetterstedt
LISPE ULIGINOSA Fallén
COENOSIA FEMORALIS Robineau-Desvoidy
COENOSIA INTERMEDIA Fallén
NEODEXIOPSIS OVATA Stein
STOMOXYS NIGRA Macquart
HAEMATOBIA MINUTA Bezzi

SPECIES ERRONEOUSLY PLACED IN THE MUSCIDAE

ACROMION Wiedemann, 1824:47 (*Musca*). Tachinidae.
AENEIVENTRIS Wiedemann, 1830:376 (*Dexia*). Calliphoridae.
AETHIOPS Walker, 1852:365 (*Anthomyia*). Agromyzidae.
ANDINA Bigot, 1885:304 (*Ophyra*). Lonchaeidae.
CLUVIA Walker, 1849:885 (*Musca*). Calliphoridae.
COSTALIS Walker, 1852:344 (*Musca*). Tachinidae.



- ELEODIVORA Walton, 1918:25, pl. 5, figs. 5-8 (*Biomyia*). Tachinidae, North America.
- FLAVIPENNIS Macquart, 1843a:291, pl. 17, fig. 3; 1843b:134, pl. 17, fig. 3 (*Ochromyia*). Calliphoridae.
- GIBBA Fabricius, 1805:297 (*Musca*). Curtonotidae.
- PROBATA Walker, 1861:318 (*Hylemyia*). Tachinidae.
- PROTRITA Walker, 1861:317 (*Anthomyia*). Sarcophagidae.
- SENSIFERA Walker, 1861:314 (*Musca*). Tachinidae.

DOUBTFUL SPECIES WHICH MAY NOT BE MUSCIDAE

- EQUESTRIS Fabricius, 1775:782 (*Musca*). Type-locality: "Brazil".
- MACTANS Fabricius, 1787:344 (*Musca*). Type-locality: Cayenne.
- MORIO Fabricius, 1775:797 (*Stomoxys*). Type-locality: "Brazil".
- PINOCHETI Brèthes, 1924:107 (*Gimnomera?*). Type-locality: Chile, Bio-Bio. Possibly Coenosiainae.

SPECIES ERRONEOUSLY DESCRIBED FROM SOUTH AMERICA

- albipila* Stein, 1918:242 (*Coenosia*). Type-locality: "South America or Africa". Holotype not located, so neither correct genus nor correct provenance could be ascertained. cf. also under "Unplaced Species of *Coenosia* s.lat."
- bicolor* Macquart, 1855a:137, pl. 6, fig. 14; 1855b:117, pl. 6, fig. 14 (*Aricia*). Type-locality: "?Brazil". An African species of the genus *Dichatomyia* Malloch, subgenus *Panaga* Curran. N. comb.
- concolor* Walker, 1852:333 (*Musca*). Type-locality: Australia (not South America, as stated by Walker). An Australian species of the genus *Helina* Robineau-Desvoidy. N. comb.
- flavicornis* Macquart, 1855a:137, pl. 6, fig. 12; 1855b:117, pl. 6, fig. 12 (*Aricia*). Type-locality: Fernando Po Island (not South America, Saint-Fernando, as stated by Macquart). An African species of *Alluaudinella* Giglio-Tos, the holotype of which is labelled Fernando Po.
- plumifera* Bigot, 1885:288 (*Spilogaster*). Type-locality: "Cuba". An Oriental species of the genus *Pygophora* Schiner. N. comb.
- supera* Walker, 1853:131 (*Anthomyia*). Type-locality: England. Described by Stein (1901:212) as *supera* Walker MS from Brazil, but in fact already described by Walker and identical with the Palearctic *Helina quadrum* (Fabricius), 1805.


LITERATURE CITED

- AITKEN, T. H. G., DOWNS, W. G., & ANDERSON, C. R.
1958. Parasitic *Philornis* Flies as Possible Sources of Arbor Virus Infections (Diptera, Anthomyiidae). *Proc. Soc. exp. Biol. Med.* 99:635-637.



- 1945a. Sobre *Fannia trimaculata* (Stein, 1897) Malloch, 1913 (Diptera, Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 34:11 pp., 21 figs.
- 1945b. Descrição do Alótipo Macho de *Fannia petrochiae* Shannon & Del Ponte, 1926 e notas sobre a fêmea (Diptera, Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 41:4 pp., 12 figs.
1946. Sobre *Fannia obscurinervis* (Stein, 1900) Stein, 1911 (Diptera, Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)* 57:9 pp., 14 figs.
1947. Contribuição ao Conhecimento de *Charadrella malacophaga* Lopes, 1938 (Muscidae, Diptera). *Rev. Ent., Rio de J.*, 18: 101-112, 35 figs.
- 1949a. Sobre um novo gênero neotrópico de «Anthomyiinae» (Diptera, Muscidae). *Rev. bras. Biol.* 9:419-423, 11 figs.
- 1949b. Sobre alguns tipos de Macquart existentes no Museu Nacional de História Natural de Paris e dados sobre «*Coenosia benoisti*» Séguy. 3ª nota. (Muscidae, Coenosiinae). *Rev. bras. Biol.* 9:439-442, 12 figs.
- 1950a. Sobre Alguns Tipos de Macquart (Diptera, Muscidae), existentes na Coleção do Museu Nacional de História Natural de Paris. (3a Nota). *Rev. Ent., Rio de J.*, 21:241-246, 10 figs.
- 1950b. Primeiras notas sobre alguns Muscidae de Macquart, da coleção do Museu de Paris. *Bol. Mus. nac. R. de J. (Zoologia)*, 98:7 pp., 18 figs.
- 1951a. Sobre um gênero e duas espécies novas de «Phaoniinae» neotropical (Diptera, Muscidae). *Rev. bras. Biol.* 11:53-57, 18 figs.
- 1951b. Quinta nota sobre tipos de Macquart (Diptera — Muscidae), existentes no Museu Nacional de História Natural de Paris e Descrição de uma espécie nova, proveniente do Chile. *Bol. Mus. nac. R. de J. (Zoologia)*, 105:17 pp., 29 figs.
- 1952a. Sobre «*Hylemyioides*» Albuquerque, 1949, com descrição de espécies novas (Diptera, Muscidae, Anthomyiinae). *Rev. bras. Biol.* 12:25-31, 19 figs.
- 1952b. Sobre alguns gêneros metálicos de Phaoniinae Neotropicais e descrição de uma *Xenomorellia* nova (Diptera: Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 116:21 pp., 24 figs.
- 1953a. Fauna do Distrito Federal. IV. Nova espécie baseada em «*Philornis pici*» (Blanchard, 1896) nec Macquart (Diptera, Muscidae). *Rev. bras. Biol.* 13:141-144, 8 figs.
- 1953b. Sobre a Identidade de «*Fannia erythropsis*» (Bigot) e «*Fannia armata*» (Bigot) nec Meigen (Diptera — Muscidae). *Anais Acad. bras. Cienc.* 25:495-498, 55 figs.
- 1953c. Fauna do Distrito Federal. VII. Sobre um Novo Gênero e Espécie Nova de «Limnophorini», com Descrição do Alótipo Fêmea de «*Coenosopsia prima*» Mall., capturada no Rio de Janeiro (Diptera — Muscidae). *Anais Acad. bras. Cienc.* 25:519-525, 18 figs.
- 1954a. Sobre um gênero novo de Phaoniinae do Chile e notas sobre *Steinella fuscinervis* (Macquart) (Diptera, Muscidae). *Rev. bras. Ent.* 1:151-158, 17 figs.

- 1954b. «*Graphomyia analis*» (Malloch, 1934) *nec* Macquart, 1850, uma espécie nova (Diptera, Muscidae). *Rev. bras. Biol.* 14:79-82, 12 figs.
- 1954c. Sobre um gênero e uma espécie nova de Coenosiinae do Brasil (Diptera, Muscidae). *Rev. bras. Biol.* 14:119-122, 15 figs.
- 1954d. Fauna do Distrito Federal. XV: Descrição de dois gêneros e quatro espécies novas de Coenosiinae (Diptera, Muscidae). *Rev. bras. Biol.* 14:177-186, 30 figs.
- 1954e. «*Cholomyioides fumipennis*» gen. n. e sp.n. de Coenosiinae do Brasil (Diptera, Muscidae). *Rev. bras. Biol.* 14:225-228, 6 figs.
- 1954f. Sobre duas Novas Espécies de *Fannia* R. D. do Brasil (Diptera-Muscidae). *Anais Acad. bras. Cienc.* 26: 317-322, 17 figs.
- 1954g. Um Gênero e uma espécie Nova de Phaoninae do Brasil (Diptera — Muscidae). *Anais Acad. bras. Cienc.* 26:323-326, 5 figs.
- 1954h. Contribuição ao conhecimento do gênero «*Pseudoptilolepis*» Snyder, 1949 (Diptera, Muscidae). *Rev. bras. Biol.* 14:253-256, 8 figs.
- 1954i. Descrição de nova espécie do gênero *Fannia* e redescritção de *F. flavicincta* (Stein, 1904) (Diptera, Muscidae). *Rev. bras. Ent.* 2: 71-80, 15 figs.
- 1954j. Sobre *Bigotomyia* Malloch, 1921, com descrição de uma espécie nova e catálogo das espécies (Diptera, Muscidae). *Rev. bras. Ent.*, 2:219-228, 17 figs.
- 1954k. Fauna do Distrito Federal. XXI: Um gênero e três espécies novas de «Coenosiinae» (Diptera, Muscidae). *Rev. bras. Biol.* 14:361-369, 24 figs.
- 1954l. Um novo gênero e uma espécie nova de «Anthomyiinae» proveniente da Argentina (Dipt., Muscidae). *Rev. bras. Biol.* 14: 493-496, 10 figs.
- 1954m. Descrição de três espécies novas de *Fannia* R.-D. brasileiras, com palpos e antenas amarelas (Diptera-Muscidae). *Anais Acad. bras. Cienc.* 26:385-394, 25 figs.
- 1954n. Fauna do Distrito Federal. XVIII — Sobre três espécies de *Heliographa* Malloch, 1921 (Diptera — Muscidae). *Anais Acad. bras. Cienc.* 26:395-409, 22 figs.
- 1955a. Fauna do Distrito Federal. XVII — Descrição do alotipo ♀ de *Cariocamyia* Snyder, 1951 e notas sobre o ♂ (Diptera-Muscidae). *Dusenía* 6:41-46, 9 figs.
- 1955b. Nova espécie de *Bithoracochaeta* Stein (Diptera, Muscidae). *Rev. bras. Ent.* 3:169-172, 6 figs.
- 1955c. Uma nova espécie de «*Lispoides*» Malloch, 1920 (Diptera, Muscidae). *Rev. bras. Biol.* 15:153-156, 9 figs.
- 1955d. Fauna do Distrito Federal. XXXII. Um gênero e uma espécie nova de Phaoninae (Diptera, Muscidae). *Rev. bras. Biol.* 15:253-256, 6 figs.
- 1955e. Fauna do Distrito Federal: XXXV. Três novas espécies de Phaoninae (Diptera-Muscidae). *Anais Acad. bras. Cienc.* 27:373-380, 20 figs.



- 1955f. Fauna do Distrito Federal. VI. Sobre alguns Phaoninae pouco conhecidos (Dipt. Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 130:11 pp., 15 figs.
- 1955g. Um gênero e uma espécie nova de Coenosiinae (Diptera, Muscidae). *Rev. bras. Biol.* 15: 391-394, 8 figs.
- 1956a. Fauna do Distrito Federal. XII. Sobre o gênero *Morellia* R.-D. (Diptera-Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 131:45 pp., 69 figs.
- 1956b. Uma nova espécie de «*Fannia*» R.-D. (Diptera, Muscidae). *Rev. bras. Biol.* 16:33-35, 4 figs.
- 1956c. Redescrição de *Caricea plumiseta* (Stein, 1911) e descrição de duas espécies novas de *Austrocoenosia* Mall., 1934 (Diptera-Muscidae). *Rev. bras. Ent.* 5: 71-80, 16 figs.
- 1956d. Fauna do Distrito Federal: XXXIX. Duas novas espécies de *Helina* R.-D. (Diptera-Muscidae). *Rev. bras. Ent.* 5:167-176, 18 figs.
- 1956e. Duas espécies novas de «*Neodexiopsis*» Mall., 1920 (Diptera, Muscidae). *Rev. bras. Biol.* 16:195-200, 13 figs.
- 1956f. Fauna do Distrito Federal. XXX. Três espécies novas de Coenosiinae (Diptera, Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 133:11 pp., 18 figs.
- 1956g. Fauna do Distrito Federal. XXXI. Contribuição ao conhecimento de *Euryomma* Stein, 1899, com descrição de uma espécie nova (Diptera-Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 136:6pp., 7 figs.
- 1956h. Fauna do Distrito Federal. XIII. Sobre o gênero *Polietina* Schnabl & Dziedzicki, 1911, com descrições de espécies novas (Diptera — Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 139: 31 pp., 54 figs.
- 1956i. Novas espécies de Coenosiinae (Diptera — Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 146:23 pp., 27 figs.
- 1956j. Fauna do Distrito Federal. XLI. Redescrição de *Helina angustipennis* (Stein, 1911) e descrição de quatro espécies novas (Diptera-Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 148:33 pp., 38 figs.
- 1956k. Contribuição ao conhecimento de *Bithoracochaeta* Stein, 1911 (Diptera-Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 149:17 pp., 14 figs.
- 1957a. Uma espécie nova de «*Pegomyza*» Schnable & Dziedzicki, 1911 (Diptera, Muscidae). *Rev. bras. Biol.*, 17: 119-122 8 figs.
- 1957b. Fauna do Distrito Federal: XLII. Um novo gênero e uma espécie nova de Anthomyinae (Diptera, Muscidae). *Rev. bras. Biol.* 17:191-196, 12 figs.
- 1957c. Fauna do Distrito Federal. XXXVI. Novas espécies de *Bigotomyia* Malloch, 1921 (Diptera — Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 159:17 pp., 25 figs.
- 1957d. Contribuição ao conhecimento de *Hydrotaea* R.-D., 1830 na América do Sul (Diptera — Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 160:18 pp., 18 figs.



- 1957e. Um gênero e duas novas espécies de «Coenosiinae» do Brasil (Diptera, Muscidae). *Rev. bras. Biol.* 17:359-365, 17 figs.
- 1957f. Fauna do Distrito Federal. XLI. Notas sobre alguns Muscideos, com descrição de dois alótipos (Diptera — Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 164:13 pp., 17 figs.
- 1957g. Sobre espécies de *Fannia* R.-D., 1830 novas ou pouco conhecidas (Diptera — Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 172: 31 pp., 39 figs.
- 1958a. Duas Espécies Novas de Coenosiinae (Dipt. Muscidae). *Studia ent. (N. S.)* 1:265-272, 12 figs.
- 1958b. Sobre uma nova espécie de *Fannia* R.-D., 1830 (Diptera Muscidae). *Rev. bras. Ent.* 8:21-24, 7 figs.
- 1958c. Sobre uma nova espécie de *Helina* R.-D., 1830 e notas sobre *Helina pleuraseta* Snyder, 1940 (Diptera, Muscidae). *Rev. bras. Ent.* 8:47-54, 12 figs.
- 1958d. Descrição de um gênero e uma espécie nova de «Coenosiinae» (Diptera, Muscidae). *Rev. bras. Biol.* 18:101-104, 3 figs.
- 1958e. Fauna do Distrito Federal. V. Contribuição ao conhecimento de *Phaonia* R.-D., 1830, com descrição de novas espécies (Diptera — Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 179:38 pp., 42 figs.
- 1958f. Sobre *Ophyra* R.-D., 1830 na América do Sul, com descrição de uma espécie nova (Diptera — Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 181:13 pp., 21 figs.
1959. Novas espécies de Coenosiinae (Diptera — Muscidae). *Bol. Mus. nac. R. de J. (Zoologia)*, 193: 17 pp., 26 figs.
- ALDRICH, J. M.
1921. The Anthomyiid genus *Atherigona* in America. *Insecutor Inscit. menstr.* 9:93-98.
1923. The Genus *Philornis* — A Bird-Infesting Group of Anthomyiidae. *Ann. ent. Soc. Am.* 16 (4):304-309, 1 fig.
1928. New Diptera or two-winged flies from South America. *Proc. U.S. nat. Mus.* 74 (art. 1): 25 pp.
1932. A new genus and two new species of Muscoid flies from Guatemala. *Proc. ent. Soc. Wash.* 34:23-25.
- ANON.
1938. Report of the Puerto Rico Experiment Station, 1937. U.S.D.A., Washington, 115 pp., 28 figs., 39 tables.
1968. Catálogo de insectos en cultivos de importancia económica en Colombia. *Publn. Asoc. latinoam. Ent.*, 1: 156 pp.
- D'ARAUJO E SILVA, A. G., ET AL.
1968. Quarto catálogo dos insetos que vivem nas plantas do Brasil, seus parasitos e predadores. Parte II, 1.º Tomo. Min. Agric., Rio de Janeiro, xvi + 622 pp.
- ARNAUD, P. J.
1959. Records of Diptera from Guadalupe Island, Mexico. *Ent. News* 70:182-185.



1963. Records of Diptera from Guadalupe Island, Mexico. Second paper. *Ent. News* 74:117-129, 7 figs.
- BALLOU, C. H.
1945. Notas sobre insectos dañinos observados en Venezuela 1938-1943. *Cuad. verde com. organ. 3a Conf. Interamer. Agric.* 34:151 pp., 6 plates.
- BALLOU, H. A.
1912. Insect Pests of the Lesser Antilles. *Pamph. Ser. Imp. Dept. Agric. W. Indies* 71:210 pp., 185 figs.
- BARTLETT, K. A.
1939. The Introduction into Puerto Rico of Beneficial Insects to aid in the Control of the Horn Fly of Cattle. *Agric. Notes Porto Rico agric. Exp. Stn* 88:6 pp.
- BEATTY, H. A.
1944. The Insects of St. Croix, V. I. *J. Agric. Univ. P. Rico* 28: 114-172.
- BEQUAERT, J. C.
1926. Part II. Medical and Economic Entomology. In Medical Report of the Hamilton Rice Seventh Expedition to the Amazon, in Conjunction with the Department of Tropical Medicine of Harvard University, 1924-5. *Contr. Harv. Inst. trop. Biol. Med.* 4:xvi + 313 pp., 70 plates, 16 figs.
1938. Chapter 16. Notes on the Arthropoda of Medical Importance in Guatemala. In Shattuck G. C., A Medical Survey of the Republic of Guatemala. *Publ. Carnegie Inst.* 499: xi + 253 pp., 76 tables, 5 graphs, 2 plates, 1 fig.
- BERG, C.
1898. Substitución de nombres genéricos. *Comun. Mus. nac. B. Aires* 1 (1):16-19.
1899. Apuntes dipterológicos. *Comun. Mus. nac. B. Aires* 1 (4):124-130.
- BERRY, P. A., & VAQUERO, M. S.
1957. Lista de Insectos Clasificados de El Salvador. *Bol. téc. Min. Agric. Ganad. El Salvador* 21:134 pp.
- BEZZI, M.
1892. Di alcuni Ditteri raccolti nel paese dei Somali dall'ingegnere L. Bricchetti-Robecchi. *Annali Mus. civ. Stor. nat. Giacomo Doria* (2) 12:181-196.
1911. Eine seltene Fliege von Weltverbreitung (Dipt.). *Societas ent.* 26:65-67.
1922. On the Dipterous Genera *Passeromyia* and *Ornithomusca*, with Notes and Bibliography on the Non-Pupiparous Myioidaria parasitic on Birds. *Parasitology* 14: 29-46.
- BIGOT, J.-M.-F.
1857a. Diptères nouveaux provenant du Chili. *Ann. Soc. ent. Fr.* (3) 5:277-308, plates 6-7.

- 1857b. Diptera. In Sagra, R. de la, *Histoire Physique, Politique et Naturelle de l'Île de Cuba*. [Volume 7] Animaux Articulés à Pieds Articulés, par F. E. Guérin-Méneville. Bertrand, Paris, lxxxvii + 868 pp., 20 plates.
1875. (Notice of a forthcoming paper, in Séance of 14 April 1875.) *Ann. Soc. ent. Fr.* (5) 5: lxxv.
1877. Diptères nouveaux ou peu connus. 8e partie. X. Genre *Somomya* (Rondani) *Calliphora*, *Melinda*, *Mufetia*, *Lucilia*, *Chrysomya* (alias *Microchrysa* Rond.) Robineau-Desvoidy. (Suite.) XI. Notes et mélanges diptérologiques. *Ann. Soc. ent. Fr.* (5) 7: 243-262.
1878. Diptères nouveaux ou peu connus. 9e partie. XII. Genus *Phumosia*, *Pyrellia*, *Cosmina*, *Ochromyia* et *Curtonevra*. XIII. Genes *Ocyptera* (Latr.), *Ocypterula*, *Exogaster* (Rond.). XIV. Notes et mélanges. *Ann. Soc. ent. Fr.* (5) 8: 31-48.
1883. (Note, in Séance of 13 December 1882.) *Ann. Soc. ent. Fr.* (6) 2 [1882]: clxxxviii.
1884. (Note, in Séance of 9 July 1884.) *Ann. Soc. ent. Fr.* (6) 4: lxxxix.
1885. Diptères nouveaux ou peu connus. 25e partie. XXXIII. Anthomyzides nouvelles. *Ann. Soc. ent. Fr.* (6) 4 [1884]: 263-304.
- 1888a. Diptères nouveaux ou peu connus. Muscidi (J.B.). *Bull. Soc. zool. Fr.* 12 [1887]: 581-617.
- 1888b. (Note, in Séance of 9 November 1887.) *Ann. Soc. ent. Fr.* (6) 7 [1887]: clxxx-clxxxii.
- 1888c. Diptères. In *Mission Scientifique du Cap Horn, 1882-1883*. Tome VI, Zoologie, 2. Paris, 45 pp., 4 plates.

BLANCHARD, E. E.

1933. Parasitos animales de la langosta. *Bol. Minist. Agric., B. Aires*, 34: 247-266, 5 plates.
1937. Dipteros Argentinos Nuevos o Poco Conocidos. *Rev. Soc. ent. argent.* 9: 35-58, 8 figs.
1942. Parasitos de *Alabama argillacea* Hbn. en la Republica Argentina. Estudio preliminar. *An. Soc. cient. argent.* 134: 54-63 and 94-128, 17 figs.
1963. Dipteros parásitos de Noctuidae argentinos. *Rev. Invest. agric., B. Aires* 17: 129-254, 38 figs.

BLANCHARD, R.

1895. (Note, in Séance of 14 May 1895). *Bull. Soc. zool. Fr.* 20: 118.
1897. Contributions à l'étude des Diptères parasites. X. Sur l'*Aricia pici* Macquart. *Ann. Soc. ent. Fr.* 65(4) [1896]: 652-654, plate 17, figs. 5-9.

BODKIN, G. E.

1917. «Cowfly Tigers.» An account of the Hymenopterous family *Bembecidae* in British Guiana. *J. Bd. Agric. Br. Guiana* 10: 119-125.



BORGMEIER, T.

1931. Nota sobre *Neivamyia lutzi* Pinto et Fonseca (Dipt., Muscidae). *Rev. Ent., Rio de J.* 1:224-228, 1 fig.

ROUCHÉ, P. F.

1834. *Naturgeschichte der Insekten, besonders in Hinsicht ihrer ersten Zustände als Larven und Puppen*. Erste Lieferung. Berlin, v + 216 pp., 10 plates.

BRAUER, F., & BERGENSTAMM, J. VON.

1889. Die Zweiflügler des Kaiserlichen Museums zu Wien. IV. Vorarbeiten zu einer Monographie der Muscaria Schizometopa (exclusive Anthomyidae). *Denkschr. Kais. Akad. Wiss. Wien (Math.-Nat. Cl.)* 56:69-180, 11 plates.
1891. Die Zweiflügler des Kaiserlichen Museums zu Wien. V. Vorarbeiten zu einer Monographie der Muscaria Schizometopa (exclusive Anthomyidae), Pars II. *Denkschr. Kais. Akad. Wiss. Wien (Math.-Nat. Cl.)* 58:305-446.
1893. Die Zweiflügler des Kaiserlichen Museums zu Wien. VI. Vorarbeiten zu einer Monographie der Muscaria Schizometopa (exclusive Anthomyidae), Pars III. *Denkschr. Kais. Akad. Wiss. Wien (Math.-Nat. Cl.)* 60:89-240.

BRÉTHES, J.

1908. Catálogo de los Dípteros de las Repúblicas del Plata. *An. Mus. nac. B. Aires (III)* 9:277-305.
1915. Sur la *Prospalangia platensis* (n. gen., n. sp.) (Hymén.) et sa Biologie. *An. Soc. cient. argent.* 79:314-320, 8 figs.
- 1918a. Quelques Díptères du Chili. *Rev. chil. Hist. nat.* 22:49-50.
- 1918b. Sur quelques insectes du Pérou. *Rev. chil. Hist. nat.* 22:122-125, fig. 7.
1919. Cueillette d'Insectes au Rio Blanco. III. Díptères. *Rev. chil. Hist. nat.* 23:40-44.
1924. Sur quelques Díptères chiliens. *Rev. chil. Hist. nat.* 28:104-111.

RUSCK, A.

1906. (Exhibit at meeting of 2 November 1905.) *Proc. ent. Soc. Wash.* 8:2-3.

CALHOUN, E. L., DODGE, H. R., & FAY, R. W.

1956. Description and rearing of various stages of *Dendrephaonia scabra* (Giglio-Tos) (Diptera, Muscidae). *Ann. ent. soc. Am.* 49:49-54, 3 tables, 14 figs.

CAMPOS, R. F.

1960. Las moscas (Brachycera) del Ecuador. *Rev. ecuat. Hig. Med. trop.* 17(1):1-66.

CASSAMAGNAGHI, A.

1946. Miasis primitiva, a forma cutanca-forunculosa en las aves del Uruguay. (Comunicación Previa.) *Bol. mens. Dir. Ganad., Montev.* 29(2):517-530, 5 photographs.



CHILLCOTT J. G.

1958. Two new species of *Euryomma* Stein (Muscidae: Diptera) from Panama. *Can. Ent.* 90:725-731, 21 figs.
1961. A Revision of the Nearctic Species of Fanniinac (Diptera: Muscidae). *Can. Ent.* 92, Suppl. 14 [1960]: 295 pp., 289 figs., 1 table, 61 maps.
1965. New Species and Stages of Nearctic *Fannia* R.-D. (Diptera: Muscidae) Associated with Nests of Hymenoptera. *Can. Ent.* 97:640-647, 18 figs.

COCKERELL, T. D. A.

1892. Additions to the Museum. *J. Inst. Jamaica* 1(2):54-56.
1894. A Supplementary Note to Mr. Johnson's List of Jamaican Diptera. *Proc. Acad. nat. Sci. Philad.* 46:419-420.

COLLIER, W. A.

1928. Die Derrengadera und ihre Bekämpfung. *Dt. tierärztl. Wschr. tierärztl. Rdsch.* 36. (45):749-752.

COQUILLETT, D. W.

1900. Report on a collection of Dipterous insects from Puerto Rico. *Proc. U.S. natn. Mus.* 22, (1198): 249-270.
- 1901a. Types of Anthomyid Genera. *Jl N. Y. ent. Soc.* 9:134-146.
- 1901b. Papers from the Hopkins Stanford Galapagos Expedition, 1898-1899. II. Entomological Results (2): Diptera [sic]. *Proc. Wash. Acad. Sci.* 3:371-379.
1910. The Type-Species of the North American Genera of Diptera. *Proc. U.S. natn. Mus.* 37: 499-647.

CURRAN, C. H.

- 1928a. Insects of Porto Rico and the Virgin Islands. Diptera or Two-winged Flies. In Scientific Survey of Porto Rico and the Virgin Islands, Vol. XI, part. 1:118 pp., 39 figs. New York.
- 1928b. Records and Descriptions of Diptera, mostly from Jamaica. In Gowdey, G. C., Catalogus Insectorum Jamaicaensis. *Ent. Bull. Dep. Agric. Jamaica* 4(3):45 + iii pp.
1931. Twelve New Diptera. *Am. Mus. Novit.* 492:13 pp., 7 figs.
1932. The Norwegian Zoological Expedition to the Galapagos Islands 1925, conducted by Alf Wollebaek. IV. Diptera (excl. of Tipulidae and Culicidae). *Nyt Mag. Naturvid.* 71:347-366, 1 fig.
- 1934a. The Templeton Crocker Expedition of the California Academy of Sciences, 1932. No. 13, Diptera. *Proc. Calif. Acad. Sci.* 21:147-172, 4 figs.
- 1934b. The Diptera of Kartabo, Bartica District, British Guiana, with descriptions of new species from other British Guiana localities. *Bull. Am. Mus. nat. Hist.* 66:287-532, 1 map, 54 figs.
1938. New American Diptera. *Am. Mus. Novit.* 975:7 pp.
1939. Two new American Diptera with notes on *Ascosyrphus* Bigot. *Am. Mus. Novit.* 1031: 3 pp.

CURTIS, J.

1839. *British Entomology; being illustrations and descriptions of the genera of insects found in Great Britain and Ireland*. Volume 16, plates 722-769. London.

CZERNY, L.

1903. Zwei neue *Homalomyia*-Arten aus Bolivien. *Z. syst. Hymenopt. Dipterol.* 3:239-240.

DEGEER, C.

1776. *Mémoires pour servir à l'Histoire des Insectes*. Volume 6, 523 pp., 30 plates. P. Hesselberg, Stockholm.

DODGE, H. R.

1955. New Muscid Flies from Florida and the West Indies (Diptera: Muscidae). *Fla. Ent.* 38:147-151.
1963. A new *Philornis* with coprophagous larva, and some related species (Diptera: Muscidae). *J. Kans. ent. Soc.* 36:239-247, 11 figs.
1965. Two new neotropical *Graphomya* (Diptera: Muscidae). *J. Kans. ent. Soc.* 38:394-397.
1967. Two new metallic species of *Helina* from Chile (Diptera: Muscidae). *Proc. ent. Soc. Wash.* 69:241-243.
1968. Some new and little-known species of *Philornis* (Diptera: Muscidae). *J. Kans. ent. Soc.* 41: 155-164, 1 fig.

DODGE, H. R., AND AITKEN, T. H. G.

1968. *Philornis* flies from Trinidad. *J. Kans. ent. Soc.* 41:134-154, 22 figs.

DUNN, L. H.

1929. Notes on some insects and other arthropods affecting man and animals in Colombia. *Am. J. trop. Med.* 9:493-508.
1934. Entomological Investigations in the Chiriqui Region of Panama. *Psyche, Camb.* 4:166-183.

ELDRIDGE, B. F., & JAMES, M. T.

1957. The Typical Muscid Flies of California. *Bull. Calif. Insect Surv.*, 6 (1): 17 pp., 4 maps, 3 plates.

ENDERLEIN, G.

1912. Die Insekten des Antarkto-Archiplata-Gebietes (Feuerland, Falklands Inseln, Süd Georgien). 20. Beitrag zur Kenntnis der antarktischen Fauna. *K. svenska Vetenskapskad. Handl.* 48(3): 170 pp., 4 plates, 35 figs.
1927. Dipterologische Studien. XVIII. *Konowia* 6:50-56.
1934. Dipterologica, I. *Sber. Ges. naturf. Freunde Berl.* (1933):416-429.
1935. Dipterologica, III. *Sber. Ges. naturf. Freunde Berl.* (1935): 235-246.
1940. Die Dipterenfauna der Juan-Fernandez-Inseln und der Oster-Insel. *Nat. Hist. Juan Fernandez and Easter Island*, Uppsala, III, Zool., part 5: 643-680, 49 figs.



ENGEL, E. O.

1931. Die Ausbeute der deutschen Chaco-Expedition 1925/26. Diptera. XXVI. Anthomyidae, XXVII. Muscidae und XXVIII. Sarcophagidae. *Konowia* 10:133-154, plates 1-6.

EVANS, J. W.

1952. *The Injurious Insects of the British Commonwealth (except the British Isles, India and Pakistan). With a Section on the Control of Weeds by Insects.* C.I.E., London, vii + 242 pp., 1 plate.

FABRICIUS, J. C.

1775. *Systema entomologiae, sistens insectorum classes, ordines, genera, species, adiectis synonymis, locis, descriptionibus, observationibus.* Flensbvirgi et Lipsiae, 32 + 382 pp.
1781. *Species insectorum exhibentes eorum differentias specificas, synonyma, auctorum, loca natalia, metamorphosin adiectis observationibus, descriptionibus.* Volume 2. Hambvrgi et Kilonii, 517 pp.
1787. *Mantissa insectorum sistens species nuper detectas, adiectis observationibus, descriptionibus, emendationibus.* Volume 2. Hafniae, 382 pp.
1794. *Entomologia systematica emendata et aucta. Secundum classes, ordines, genera, species adiectis synonymis, locis, observationibus, descriptionibus.* Volume 4. Hafniae, 472 pp.
1805. *Systema anthiatorum secundum ordines, genera, species adiecta synonymis, locis, observationibus, descriptionibus.* Brunsvigae, i-xiv + 15-372 + 1 + 30 pp.

FAIRCHILD, G. B.

1943. An Annotated List of the Bloodsucking Insects, Ticks and Mites Known from Panama. *Am. J. trop. Med.* 23: 569-591.

FALLÉN, C. F.

1817. Beskrifning Ofver de i Sverige funna Fluge Arter, som kunna foras till Slagtet Musca. Forsta Afdelningen. *K. svenska Vetensk. Akad. Handl.* 1816 (2):226-254.
1819. *Scatomyzides Sueciae.* Lundae, 10 pp.
- 1823a. *Phytomyzides et Ochtidae Sueciae.* Lundae, 10 pp.
- 1823b. *Monographia Muscidum Sueciae.* Part V, pages 49-56. Lundae.
1825. *Monographia Muscidum Sueciae.* Part VIII, pages 73-80. Part IX, pages 81-94. Lundae.

FENNAH, R. G.

1947. The Insects Pests of Food-crops in the Lesser Antilles. *Dept. of Agric., Grenada and Antigua*, ii + 207 pp., 139 figs.

FIASSON, R.

1943. Notes sur les parasites animaux du Haut-Apure (Venezuela). *Revue Sci. méd. pharm. vét. Afr. fr. libre* 2:125-151, 1 map, 5 col. plates.



FRAUENFELD, G. VON

1867. Zoologische Miscellen. XI. *Verh. zool.-bot. Ges. Wien* 17:425-502.

GALLEGO, F. L.

1967. Lista preliminar de insectos de importancia economica y secundarios, que afectan les principales cultivos, animales domesticos y al hombre, en Colombia. *Rev. Fac. nac. Agron., Medellin*, 26(65):32-66.

GAMINARA, A.

1930. Clasificación de algunos Muscoideos Uruguayos (Muscidae y Calliphoridae). *An. Fac. Med. Univ. Montevideo*, 14 [1929]: 1235-1282, 19 figs.

1931. Clasificación de algunos Muscoideos Uruguayos (Muscidae y Calliphoridae). *Arch. Soc. Biol. Montev., Suppl.* 5: 1237-1280, 19 figs.

GARCIA, M.

1952. Las especies argentinas del género *Philornis* Mein., con descripción de especies nuevas (Dipt. Anthom.). *Rev. Soc. ent. argent.* 15:277-293, 6 figs.

GARCIA, P. C.

1964. *Moscas de Venezuela*. 17 unnumbered pp., 30 plates, Caracas.

GAY, C.

1852. *Historia fisica y politica de Chile*. Zoologia, volume 7. *Insectos*. 471 pp., Paris and Santiago.

GEOFFROY, E. L.

1762. *Histoire abrégée des Insectes qui se trouvent aux environs de Paris*. Volume 2. 690 pp., plates XI-XXI, Paris.

GIGLIO-TOS E.

1893. Diagnosi di nuovi generi e di nuove specie di Ditteri. VIII. *Bol. Musei Zool. Anat. comp. R. Univ. Torino*, VIII, No. 147:11 pp.

1896. Ditteri del Messico. Parte Quarta. Muscidae calypteratae: Muscinae, Anthomyiinae. Muscidae acalypteratae: Scatophaginae, Helomyzinae, Tetanocerinae, Ortalinae, Ulidinae, Sapromyzinae, Trypetinae, Sepsinae, Tanypezinae, Psilinae, Chloropinae, Ephydrinae, Drosophilinae. *Memorie Accad. Sci. Torino* (2)45:1-74, plate 1.

GONZALES, R. H.

1968. La «mosca de los cuernos» *Haematobia irritans* (L.) en Chile. *Rev. chil. Ent.* 6:142.

GOWDEY, G. C.

1926. Catalogus Insectorum Jamaicensis. *Ent. Bull. Dep. Agric. Jamaica* 4:114 +xiv and 10 + ii pp.

1928. Catalogus Insectorum Jamaicensis. *Ent. Bull. Dep. Agric. Jamaica* 4:45 + iii pp.

GREENBERG, B., VARELA, G., BORNSTEIN, A., & HERNANDEZ, H.

1963. Salmonellae from flies in a Mexican slaughterhouse. *Amer. J. Hyg.* 77:177-183, 4 tables.

GREENBERG, B., & BORNSTEIN, A. A.

1964. Fly dispersion from a rural Mexican slaughterhouse. *Am. J. trop. Med. Hyg.* 13:881-886, 2 figs., 2 tables.

GRIMSHAW, P. H.

1901. Part I, Diptera. In Sharp, D., Fauna Hawaiiensis, Volume 3: 1-77, plates 1-3.

GUNDLACH, J.

1887. Apuntes para la Fauna Puerto-Riqueña. VIII. Insectos. *An. Soc. esp. Hist. nat.* 16: 137-199.

HALIDAY, A. H.

1833. Catalogue of Diptera occurring about Holywood in Downshire. *Ent. Mag.* 1:147-180.
1838. New British Insects indicated in Mr. Curtis's Guide. *Ann. nat. Hist.* 2:112-121 and 183-190.

HENNIG, W.

1955. Los Insectos de las Islas Juan Fernandez. 16. Phryneidae, Helomyzidae, Lonchacidae, Piophilidae, Anthomyzidae und Muscidae (Diptera). *Rev. chil. Ent.* 4:21-34.
- 1955-1964. Muscidae. In Lindner, E., *Fliegen palaearkt. Reg.* 63b: 1110 pp., 429 text-figs., 33 plates. Stuttgart.
1957. Los Insectos de las Islas Juan Fernandez. 34. Therevidae, Helomyzidae, Piophilidae, Milichiidae und Muscidae (Diptera). (Nachtrag). *Rev. chil. Ent.* 5:409-412.
1965. Vorarbeiten zu einem phylogenetischen System der Muscidae (Diptera: Cyclorrhapha). *Stuttg. Beitr. Naturk.* 141:100 pp., 53 figs.

HOUGH, G. DE N.

1899. Some Muscinae of North America. *Biol. Bull. mar. biol. Lab., Woods Hole* 1: 19-33, 19 figs.
1900. South American Muscidae in the Collection of S. W. Williston. *Kans. Univ. Q.* 9:203-232, plates 44-45.

HOWARD, L. O.

1900. A Contribution to the Study of the Insect Fauna of Human Excrement. *Proc. Wash Acad. Sci.* 2:541-604, figs. 17-38, plates 30-31.

HUCKETT, H. C.

1932. The North American Species of the Genus *Limnophora* Robineau-Desvoidy, with Descriptions of New Species (Muscidae, Diptera). *Jl. N. Y. ent. Soc.* 40:25-76, 105-158, and 280-339, plates 7-13.
1934. A Revision of the North American Species belonging to the Genus *Coenosia* (Diptera: Muscidae). I. The Subgenera *Neodeixiopsis*, *Coenosia*, *Hoplogaster* and related Genera *Allognota*, *Bithoracochaeta* and *Schoenomyza*. *Trans. Am. ent. Soc.* 60: 57-119, plates 4-7.



1936. A Revision of Connectant Forms between Coenosian and Limnophorine Genera occurring in North America (Diptera, Muscidae). *Jl. N. Y. ent. Soc.*, 44 (3): 187-223, pls. IV-V.
1954. A Review of the North American Species belonging to the genus *Hydrotaea* Robineau-Desvoidy (Diptera, Muscidae). *Ann. ent. Soc. Am.* 47: 316-342.
- IRIARTE, D. R.
1943. Contribución a la Fauna Venezolana. *Bol. Lab. Clin. Luis Razetti* 4 (11): 189-203.
- JACOBS, J.-CH., BECKER, T., & RUEBSAAMEN, E. H.
1906. Diptères. In *Expédition Antarctique Belge. Résultats du Voyage du S. Y. Belgica en 1897-1898-1899 sous le commandement de A. de Gerlache de Gomery. Volume 8, Zoologie, Insectes*, 92 pp., 5 plates. Anvers.
- JAENNICKE, F.
1867a. Neue Exotische Dipteren. *Abh. senckenb. naturforsch. Ges.* 6:311-408, plates 43 and 44.
1867b. *Ibidem*, reprinted with pagination 3-99, 2 plates.
- JOHNSON, C. W.
1894. List of the Diptera of Jamaica with descriptions of new species. *Proc. Acad. nat. Sci. Philad.* 46: 271-281.
1904. A revised list of the Diptera of Bermuda. *Psyche, Camb.* 11: 76-80.
1908. The Diptera of the Bahamas, with notes and description of one new species. *Psyche, Camb.* 15:69-80.
1913. The Dipteran Fauna of Bermuda. *Ann. ent. Soc. Am.* 6:443-452, 2 figs.
1919. A Revised List of the Diptera of Jamaica. Phoridae by Charles T. Brues. *Bull. Am. Mus. nat. Hist.* 41, art. 8:421-449.
1924. Diptera of the Williams Galapagos Expedition. *Zoologica, N. Y.* 5:85-92, 2 figs.
- JOHNSON, W. T., & VENARD, C. E.
1957. Observation son the Biology and Morphology of *Ophyra aenes-cens* (Diptera: Muscidae). *Ohio J. Sci.* 57:21-26, 11 figs., 1 table.
- KARL, O.
1935. Aussereuropäische Musciden (Anthomyiden) aus dem Deutschen Entomologischen Institut. *Arb. morph. taxon. Ent. Berl.* 2: 29-49
- LAAKE, E. W.
1949. Livestock Parasite Control Investigations and Demonstrations in Brazil. *J. econ. Ent.*, 42:276-280, 1 table.
- LAHILLE, F.
1907. La langosta y sus Moscas Parasitarias. *An. Minist. Agric. (Zool.)* 3, (4) 1-136, 29 figs., 6 plates.



LAINSON, R., & SOUTHGATE, B. E.

1965. Mechanical Transmission of *Leishmania mexicana* by *Stomoxys calcitrans*. *Trans. R. Soc. trop. Med. Hyg.* 59:716.

LATREILLE, P. A.

1796. *Précis des caractères génériques des Insectes, disposés dans un ordre naturel*. xiv + 208 pp., Paris.

1802. *Histoire naturelle, générale et particulière, des Crustacés et des Insectes*. Volume 3. xii + 468 pp., Paris.

LATREILLE, P. A., LEPELETIER, A. L. M., SERVILLE, J. G. A., & GUÉRIN-MÉNEVILLE, F. E.

1828. *Entomologie, ou histoire naturelle des Crustacés, des Arachnides et des Insectes*. *Encyclopédie méthodique, Histoire naturelle*, Volume 10. 833 pp., Paris.

LEFÈVRE, E.

1887. (Note, in Séance of 27 April 1887.) *Ann. Soc. ent. Fr.* (6)7: lxxiv-lxxx.

LEGNER, E. F., BAY, E. C., & MCCOY, C. W.

1965. Parasitic natural regulatory agents attacking *Musca domestica* L. in Puerto Rico. *J. Agric. Univ. P. Rico* 49 (3):368-376, 1 fig., 2 tables.

LEGNER, E. F., BAY, E. C., & WHITE, E. B.

1967. Activity of Parasites from Diptera: *Musca domestica*, *Stomoxys calcitrans*, *Fannia canicularis*, and *F. femoralis*, at Sites in the Western Hemisphere. *Ann. ent. Soc. Am.* 60:462-468, 1 fig., 1 table.

LE PELLEY, R. H.

1968. *Pests of Coffee*. Longmans, London, xii + 590 pp., 21 figs., plates XIII + 15.

LINNAEUS, C.

1758. *Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis*. 10th Edition. Volume 1. 824 pp., Holmiae.

1761. *Fauna Svecica Sistens Animalia Sveciae Regni*. 2nd Edition. 578 pp., 2 plates, Stockholmiae.

LINS DE ALMEIDA, J.

1933. Nouveaux agents de transmission de la Berne [*Dermatobia hominis* (L. junior, 1781)] au Brésil. *C. r. Séanc. Sec. Biol.* 113, (27):1274-1275.

LINSLEY, E. G., & USINGER, R. L.

1966. Insects of the Galápagos Islands. *Proc. Calif. Acad. Sci.* 33 (7):113-196, 1 table.

LIOY, P.

1864. I Ditteri distribuiti secondo un nuovo metodo di classificazione naturale. [Part.] *Atti R. Ist. veneto Sci.* (3) 9:989-1027.



LOEW, H.

1861. Diptera aliquot in insula Cuba collecta. *Wien. ent. Monatschr.* 5:33-43.
1869. Diptera Americae septentrionalis indigena. Centuria octava. *Berl. ent. Z.* 13:1-52.
1872. Diptera Americae septentrionalis indigena. Centuria decima. *Berl. ent. Z.* 13:1-52.

LUNDBECK, W.

1901. Diptera groenlandica. *Vidensk. Meddr dansk. naturh. Foren.* (6) 2 [1900]: 281-316, 5 figs.

LUTZ, A., & NEIVA, A.

1912. Notas Dipterológicas. A proposito da *Mydaea pici* Macquart. *Mem. Inst. Oswaldo Cruz* 4:130-135.

MACDONALD, A.

1917. Netes on Blood-sucking Flies in Grenada. *Bull. ent. Res.* 7 (3):259-264, plates III and IV.

MACQUART, J.

1835. *Histoire Naturelle des Insectes. Diptères*, Tome deuxième. 703 pp., 12 plates, Paris.
1938. Notice sur trois Diptères nouveaux des genres *Medeterus* et *Scatophaga*. *Ann. Soc. ent. Fr.* 7:421-424, plate 11 figs. 1-3.
1843a. Diptères Exotiques nouveaux ou peu connus. Tome deuxième. 3^e partie. *Mém. Soc. Sci. Agric. Lille* 1843:162-460, 36 plates.
1843b. *Ibidem*, reprinted with pagination 5-304.
1846a. Diptères Exotiques nouveaux ou peu connus. Supplément. *Mém. Soc. Sci. Agric. Lille* 1844:133-366, 20 plates.
1846b. *Ibidem*, reprinted with pagination 5-238.
1848a. Diptères Exotiques nouveaux ou peu connus. Suite du 2^{me} Supplément. *Mém. Soc. Sci. Agric. Lille* 1847(2): 161-237, 7 plates.
1848b. *Ibidem*, reprinted with pagination 1-77.
1851a. Diptères Exotiques nouveaux ou peu connus. 4^e Supplément (part). *Mém. Soc. Sci. Agric. Lille* 1850:134-294, plates 15-28.
1851b. *Ibidem*, reprinted with pagination 161-336.
1854. Notice sur une nouvelle espèce d'Aricie, Diptère de la tribu des Anthomyzides. *Ann. Soc. ent. Fr.* (3) 1 [1853]: 657-660, plate 20, II, figs. 1-4.
1855a. Diptères Exotiques nouveaux ou peu connus. 5^e Supplément. *Mém. Soc. Sci. Agric. Lille* 1854:25-156, 7 plates.
1855b. *Ibidem*, reprinted with pagination 5-136.

MAGARINOS TORRES, C.-B. DE, FONSECA, O. DA, ARÊA LEÃO, A.-E. DE


1923. Sur la «Esponja» (Habronémose cutanée des Equidés). Du Parasitisme des Mouches par l'*Habronema muscae* (Carter). *C. r. Séanc. Soc. Biol.*, 89 (27):767-768.

MALLOCH, J. R.

1912. New Diptera from Panama. *Smithson. misc. Collns* 59 (17): 8 pp.



1913. Notes on some American Diptera of the genus *Fannia*, with descriptions of new species. *Proc. U.S. natn. Mus.* 44:621-631, plate 7.
1916. A key to the males of the Anthomyid genus *Hydrotaea* recorded from North America. *Bull. Brooklyn ent. Soc.* 11:108-111.
- 1917a. A new genus and species of Anthomyiidae (Diptera). *Bull. Brooklyn ent. Soc.* 12:37-38.
- 1917b. A new genus of Anthomyiidae (Diptera). *Bull. Brooklyn ent. Soc.* 12:113-115.
- 1918a. Notes and descriptions of some Anthomyiid genera. *Proc. biol. Soc. Wash.* 31:65-68.
- 1918h. Diptera from the Southwestern United States. Paper IV. Anthomyiidae. *Trans. Am. ent. Soc.* 44: 263-319, plate 17.
1920. Descriptions of New North American Anthomyiidae (Diptera). *Trans. Am. ent. Soc.* 46 (2):133-196, plates 4-6.
- 1921a. Exotic Muscaridae (Diptera), I. *Ann. Mag. nat. Hist.* (9)7: 161-173.
- 1921b. Notes on some of van der Wulp's Species of North American Anthomyiidae (Diptera). *Ent. News* 32:40-45.
- 1921c. Exotic Muscaridae (Diptera), II. *Ann. Mag. nat. Hist.* (9)7: 420-431.
- 1921d. Exotic Muscaridae (Diptera), IV. *Ann. Mag. nat. Hist.* (9)8: 414-425.
1922. Exotic Muscaridae (Diptera), V. *Ann. Mag. nat. Hist.* (9)9: 271-280.
- 1923a. Flies of the Anthomyiid Genus *Phaonia* and Related Genera. *Trans. Am. ent. Soc.* 48 [1922]:227-282, plates 11-13.
- 1923b. Exotic Muscaridae (Diptera), IX. *Ann. Mag. nat. Hist.* (9) 11: 664-675.
- 1923c. Expedition of the California Academy of Sciences to the Gulf of California in 1921. Anthomyidae and Lonchaeidae (Kelp Flies and their allies). *Proc. Calif. Acad. Sci.* (4)12:425-428.
- 1923d. Exotic Muscaridae (Diptera), XI. *Ann. Mag. nat. Hist.* (9)12: 505-528.
1924. A new Genus and Species of Muscidae (Diptera). *Proc. ent. Soc. Wash.* 26:74.
1925. Exotic Muscaridae (Diptera), XVI. *Ann. Mag. nat. Hist.* (9)16:81-100.
- 1928a. Exotic Muscaridae (Diptera), XXI. *Ann. Mag. nat. Hist.* (10) 1: 465-494.
- 1928b. Exotic Muscaridae (Diptera), XXII. *Ann. Mag. nat. Hist.* (10)2:307-319.
- 1929a. Exotic Muscaridae (Diptera), XXVI. *Ann. Mag. nat. Hist.* (10)4:97-120.
- 1929b. Exotic Muscaridae (Diptera), XXVIII. *Ann. Mag. nat. Hist.* (10)4:322-341.
1930. Exotic Muscaridae (Diptera), XXIX. *Ann. Mag. nat. Hist.* (10) 5:465-484.



1932. Exotic Muscaridae (Diptera), XXXVI. *Ann. Mag. nat. Hist.* (10)9:377-405, 421-447 and 501-518.
1934. Muscidae. In *Diptera of Patagonia and South Chile*, VII (2): 171-346, figs. 22-60. London.
1936. A New Genus and Species of Muscidae from Puerto Rico. *Proc. ent Soc. Wash.* 38:9-10.
- MARCHIONATTO, J. B.
1945. Nota sobre algunos hongos entomogenos. *Publ. Inst. Sanid. reg., B. Aires, ser. A*, 1(8): 11 pp., 1 col. pl., 3 figs.
- MARTINEZ, M.
1928. La mosca domestica en la ciudad de Mexico. *Mem. Soc. «Ant. Alzate»*, 48:391-418, 8 plates. [Not seen.]
- MARTORELL, L. F.
1939. Insects observed in the State of Aragua, Venezuela, South America. *J. Agric. Univ. P. Rico*, 23(4):177-232.
- MAZZA, S., & JOERG, M. E.
1939. *Cochliomyia hominivorax* = *americana* C. y P., Estudio de sus Larvas y Consideraciones sobre Miasis. In *Investigaciones sobre Dipteros Argentinos. I. Miasis. Publ. Misión Estud. Patol. reg. argent. Jujuy* 41:1-46, 38 figs., 1 plate.
- MAZZA, S., & ORIBE, H. R.
1939. Miasis urinaria por *Fannia fusconotata* Rondania, en Formosa. In *Investigaciones sobre Dipteros Argentinos. I. Miasis. Publ. Misión Estud. Patol. Reg. argent. Jujuy* 41:66-69, 2 figs.
- MCLINTOCK, J., & DEPNER, K. R.
1954. A Review of the Life-History and Habits of the Horn Fly, *Siphona irritans* (L.) (Diptera: Muscidae). *Can. Ent.* 86:20-33, 1 table.
- MEIFERT, D. W., LABRECQUE, G. C., SMITH, C. N., & MORGAN, P. B.
1967. Control of House Flies on Some West Indian Islands with Metepa, Apholate, and Trichlorfon Baits. *J. econ. Ent.* 60:480-485.
- MEIGEN, J. W.
1803. Versuch einer neuen Gattungs Eintheilung der europäischen zweiflügeligen Insekten. *Illig. Mag. Ins.* 2:259-281.
1826. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten*. Volume 5. xii + 412 pp., plates 42-54, Hamm.
1830. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten*. Volume 6. xi + 401 pp., plates 55-56, Hamm.
1838. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten*. Volume 7. xii + 434 pp., plates 67-74, Hamm.
- MEINERT, F.
1890. *Philornis molesta*, en paa Fugle snyltende Tachinarie. *Vidensk. Meddr. dansk naturh. Foren. (V)* 1 [1889]: 304-317, plate 6.



MELLO, M. J., & CUOCOLO, R.

- 1943a. Técnica para o xenodiagnóstico da habronemose gástrica dos equídeos. *Arch. Inst. Biol. S. Paulo* 14:217-226, 1 fig., plate 20.
1943b. Alguns aspetos das relações do «Habronema muscae (Carter, 1861)» com a mosca doméstica. *Arch. Inst. Biol. S. Paulo* 14: 227-234, 2 tables, plata 21.

MELLO, M. J. DE, & PEREIRA, C.

1946. Determinismo de evasão das larvas de "Habronema" sp. da tromba da mosca doméstica. *Arch. Inst. Biol. S. Paulo* 17:259-266, 1 fig., 1 table, 1 graph.

MYERS, J. G.

1934. Field Observations on some Guiana Insects of Medical and Veterinary Interest. *Trop. Agric., Trin.* 11 (11):279-283.
1938. Report on the Natural Enemies in Haiti of the Horn-fly (*Lyperosia irritans*) and the Green Tomato Bug (*Nezara viridula*). *J. Coun. scient. ind. Res. Aust.* 11(1):35-46, 1 map.

NEIVA, A., & PENNA, B.

1916. Viagem científica pelo Norte da Bahia, sudoeste de Pernambuco, sul do Piauí e de norte a sul de Goiaz. *Mem. Inst. Oswaldo da Cruz* 8 (3): 74-224, 27 plates.

NIELSEN, J. C.

1911. *Mydaea anomala* Jaenn., a parasite of South-American birds. *Vidensk. Meddr. dansk naturh. Foren.* 63:195-208, 14 figs.
1913. On some South-American species of the genus *Mydaea*, parasitic on birds. *Vidensk. Meddr. dansk naturh. Foren.* 65:251-256, 4 figs.

NUÑEZ TOVAR, M.

1924. Índice Dipterológico de Venezuela con la distribución geográfica por Estados. *Trab. Contrib.* 4º Congreso Venez. Med.: 14 pp.

OGILVIE, L.

1928. The Insects of Bermuda. *Dept. of Agriculture, Bermuda*, 52 pp.

OLIVEIRA, S. J. DE

1941. Sobre *Ophyra aenescens* (Wiedemann, 1830) (Diptera: Anthomyiidae). *Arch. Zool. S. Paulo* 2(15):341-355, 3 plates.

OLMEDO, I. H.

1957. Investigaciones sobre el Minador de la hoja del Huauzontle. *Revista Chapingo* 64:117-131, 8 figs.

OPINION 82

1925. Opinions rendered by the International Commission on Zoological Nomenclature. Opinion 82. Suspension of Rules for *Musca* Linnaeus, 1758A, Type *M. domestica*. *Smithson. misc. Collns* 73(3): 1-7.

OPINION 441

1957. Validation under the Plenary Powers of the names for five genera in the Order Diptera (Class Insecta) published in 1762 by

Geoffroy (E. L.) in the work entitled *Histoire abrégée des Insectes qui se trouvent aux Environs de Paris*. (Opinion supplementary to Opinion 228). *Opinions and Declarations rendered by the International Commission on Zoological Nomenclature* 15(6): 83-120.

OPINION 884

1969. *Musca lateralis* Linnaeus, 1758 (Insecta, Diptera): suppressed under the plenary powers. *Bull. zool. Nom.* 26:123-124.

ORTIZ, C. I.

1944. Contribución al Estudio de la Entomología Médica del Estado Falcón. *Bol. Lab. Clín. Luis Razetti* 4(14):247-251.

ORTIZ, C. STUARDO

1946. *Catalogo de los Dipteros de Chile*. 250 pp., Ministry of Agriculture, Santiago.

OSTEN-SACKEN, C. R.

1877. Note on some Diptera from the Island Guadalupe (Pacific Ocean), collected by Mr. E. Palmer. *Proc. Boston Soc. nat. Hist.* 18:133-134.

PATÍÑO-CAMARGO, L.

1940. Artrópodos Hematófagos de la Fauna colombiana. *Rev. Acad. colomb. Cienc. exact. fis. nat.* 3(11):337-344.

PEREIRA, C., & DE CASTRO, M. P.

1945. Contribuição para o conhecimento da espécie tipo de "Macrocheles Latr." ("Acarina"): "*M. muscaedomesticae* (Scopoli, 1772)" emend. *Arch. Inst. Biol. S. Paulo* 16:153-186, 22 figs.

PIMENTEL, D.

1955. Relationship of ants to fly control in Puerto Rico. *J. econ. Ent.* 48:28-30, 2 tables.

PINTO, C.

- 1931a. Hypopygio ou terminalia das moscas hematophagas Stomoxydinae (Diptera, Muscidae). *Rev. med-cirúrg. Braz.* 39(7):248-254, 7 figs.

- 1931b. A proposito da "Nota sobre Nelvamyia lutzi Pinto et Fonseca por T. Borgmeier". *Bol. biol. Lab. Parasit. Fac. Med. S. Paulo* 17:19-20.

PINTO, C., & FONSECA, F. DA

- 1930a. Novo genero e nova especie de mosca hematophaga da sub-familia Stomoxydinae, hospedadora intermediaria da *Dermatobia hominis*. *Rev. med-cirúrg. Braz.* 38(7):241-247, 1 fig., 3 plates.

- 1930b. Novos hospedadores intermediarios da *Dermatobia hominis* (L. Junior, 1781) "Diptera. Oestridae". *Rev. med-cirúrg. Braz.* 38(7):248-250, 1 plate.

1931. Chave practica para a classificação dos generos de moscas hematophagas "Stomoxydinae" (Diptera, Muscidae). *Rev. med-cirúrg. Braz.* 39(8):261-268, 12 figs.

PINTO, C., & SOUZA LOPES, H. DE

1933. Anatomia, biologia e papel patogenico da "Neivamyia lutzi", mosca sugadora de sangue dos equideos do Brasil. *Arch. Esc. sup. Agric. Med. vet.* 10(1):77-88, plates 6-21.

PONT, A. C.

1966. The identity of two Neotropical species of *Fannia* Desvoidy, *anthracina* Walker and *albibasis* Malloch (Diptera: Muscidae). *Ann. Mag. nat. Hist.* (13)8[1965]:427-433, 12 figs.

1967. Notes on some Australasian Muscidae (Diptera) described by J. Macquart. *Ann. Soc. ent. Fr. (N. S.)* 3(1):181-190.

RABINOVICH, J. E.

1970. Vital Statistics of *Synthesiomyia nudisetia* (Diptera: Muscidae). *Ann. ent. Soc. Am.*, 63:749-752, 6 figs., 2 tables.

RAMOS, J. A.

1946. The Insects of Mona Island (West Indies). *J. Agric. Univ. P. Rico* 20:1-74, plates 1-2.

REED, E. C.

1888. Catalogo de los Insectos Dipteros de Chile. *An. Univ. Chile* 73:v + 46 pp.

ROBERTS, R. A.

1934. Some Insects collected in Mexico, mostly in association with Man and Animals or animal Products. *Jl N. Y. ent. Soc.* 42(3): 249-262, 5 tables.

ROBINEAU-DESVOIDY, A. J. B.

1830. Essai sur les Myodaires. *Mém. prés. div. Sav. Acad. Sci. Inst. Fr.* 2:813 pp.

1863. *Histoire Naturelle des Diptères des Environs de Paris*. Volume 2. 920 pp., Paris, Leipzig, London.

ROEDER, V. VON

1885. Dipteren von der Insel Portorico, erhalten durch Herrn Consul Krug in Berlin. *Stettin. ent. Ztg* 46:337-349.

1886. Dipteren von den Cordilleren in Columbien. Gesammelt durch Herrn Alphons Stübel. *Stettin. ent. Ztg* 47:257-270.

1892. *Dipteren gesammelt in den Jahren 1868-1877 auf eine Reise durch Süd-Amerika von Alphons Stübel*. (Vermehrter und verbesserter Abdruck aus der Stettiner Entomologischen Zeitung 47. Jahrgang 1886 pag. 257-270.) 16 pp., 1 plate, Berlin.

RONDANI, C.

1848. Esame di varie specie d'Insetti Ditteri Brasilliani. *Studi Entomologici* 1:63-112, plate III bis.

1850. Osservazioni sopra alquante specie di Esapodi Ditteri del Museo Torinese. *Nuovi Annali Sci. Nat. (III)* 2:165-197, plate 4.

1856. *Dipterologiae Italicae Prodrromus*. Volume 1. 228 pp., Parmae.

1864. Dipteriorum species et genera aliqua exotica revisa et annotata novis nonnullis descriptis. *Arch. Zool. Anat. Fis., Modena* 3 [1863]: 1-99, plate 5.

1866. Anthomyinae Italicac collectae distinctae et in ordinem dispositae. *Atti Soc. ital. Sci. nat.* 9:68-217.
1868. Diptera aliqua in America meridionali lecta a Prof. P. Strobel annis 1866-67. *Annular. Soc. Nat. Modena* 3:24-40, plate 4.
1871. Diptera Italica non vel minus cognita descripta aut annotata. Fasc. IV. Addenda Anthomyinis Prodr. Vol. VI. *Boll. Soc. ent. ital.* 2 [1870]:317-338.
1877. *Dipterologiae Italicæ Prodrum*. Volume 6. 304 pp., Parmae.
- ROSSI, P.
1790. *Fauna Etrusca. Sistens insecta quae in provinciis Florentina et Pisana praesertim collegit*. Volume 2. 348 pp., 10 plates, Liburni.
- RUFFINELLI, A., & CARBONELL, C. S.
1954. Segunda lista de insectos y otros artrópodos de importancia económica en el Uruguay. *Rev. Asoc. Ing. agron., Montev.* 94:33-82.
- RUSSO, G.
1927. Informe de entomologia agricola dominicana. *Infme. Estac. agron. Moca* 2:33-244, 30 plates, 2 maps.
- SABROSKY, C. W.
1949. The Muscid Genus *Ophyra* in the Pacific Region (Diptera). *Proc. Hawaii ent. Soc.* 13(3):423-432, 12 figs.
- SCHINER, J. R.
1868. Diptera. In *Reise der österreichischen Fregatte Novara*, Zoologischer Theil, Bd.II. vi + 388 pp., 4 plates, Wien.
- SCHNABL, J.
1902. Eine neue *Coelomyia* (Diptera, Muscidae) vom Cap Horn. *Russk. ent. Obozr.* 2(3):160-162.
1911. Anthomyidae. In Becker, T., Kuntze, A., Schnabl, J., and Villeneuve, J., *Dipterologische Sammelreise nach Korsika* (Dipt.). *Dt. ent. Z.* 1911:62-100.
- SCHNABL, J., & DZIEDZICKI, H.
1911. Die Anthomyiden. *Nova Acta Acad. Caesar. Leop.-Carol.* 95: 55-358, plates 3-37.
- SCOPOLI, J. A.
1763. *Entomologia carniolica exhibens insecta carniolae indigine et distributa in ordines, genera, species, varietates methodo Linnaeana*. 421 pp., Vindobonae.
- SEAGO, J. M.
1954. The *pusio* Group of the Genus *Fannia* Robineau-Desvoidy, with Descriptions of New Species (Diptera, Muscidae). *Am. Mus. Novit.* 1699:13 pp., 20 figs.
- SÉGUY, E.
- 1932a. Étude sur les Diptères parasites ou prédateurs des sauterelles. *Encycl. ent. B II Dipt.* 6:11-40.
- 1932b. Études sur les Anthomyides. 8e note. Un *Coenosia* nouveau de l'Amérique Equatoriale. *Encycl. ent. B II Dipt.* 6:144, figs. 1-3.

1933. Diptères Néotropiques Nouveaux. *Rev. chil. Hist. nat.* 37:256-260, plate 8.
1934. Étude sur Quelques Muscides de l'Amérique Latine. *Rev. Soc. ent. argent.* 6:9-16, 3 figs.
- 1935a. Étude sur les Stomoxydines et particulièrement des mouches charbonneuses du genre *Stomoxys*. *Encycl. ent. B II Dipt.* 8: 15-58.
- 1935b. Études sur les Anthomyides. 9e, 10e et 11e notes. 9. Trois Phaoniines nouveaux de Madagascar. 10. Étude sur le genre *Morellia* R.-D. 11. Un nouveau *Trichomorellia* de l'Equateur. *Encycl. ent. B II Dipt.* 8:97-116.
1937. Diptera, Family Muscidae. In Wytsman, P., Genera Insect. Fasc. 205: 604 pp., 9 plates. Bruxelles.
- 1938a. Notes sur les Anthomyides (Muscidae), 12e note. *Encycl. ent. B II Dipt.* 9:109-120.
- 1938b. Précisions sur deux Muscides du Chili. *Rev. chil. Hist. nat.* 41 [1937]:39-41.
1941. Une nouvelle espèce d'*Euryomma* Stein (Dipt., Muscidae) et synopsis des espèces du genre. *Bull. Soc. ent. Fr.* 46:54-55, 3 figs.
1946. Notes biologiques et taxonomiques sur les Diptères. *Encycl. ent. B II Dipt.* 10:105-150.
1952. Diptera, Family Scatophagidae. In Wytsman, P., Genera Insect. Fasc. 209: 107 pp., 44 figs. Bruxelles.
- SHANNON, R. C., & DEL PONTE, E.
1926. Sinopsis parcial de los Muscoideos Argentinos. *Rev. Inst. bact., B. Aires* 4(5):549-590, 4 plates.
1928. Sinopsis parcial de los Muscoideos Argentinos (Addenda et Corrigenda). *Rev. Inst. bact., B. Aires* 5(1):141-147.
- SMYTH, E. G.
1919. Dominio de la Plaga de la Mosca del Ganado en Puerto Rico. *Rev. Agric. P. Rico* 3(5):10-24, 4 plates, and 3(6):17-28.
- SNYDER, F. M.
- 1940a. New and little known tropical *Helina* Desvoidy, allied to *Ariciella* Malloch (Diptera: Muscidae). *Am. Mus. Novit.* 1060:9 pp.
- 1940b. A Review of the Genus *Myospila* Rondani with Descriptions of New Species (Diptera: Muscidae). *Am. Mus. Novit.* 1087:10 pp., 2 figs.
1941. Contribution to a Revision of Neotropical Mydaeini (Diptera: Muscidae). *Am. Mus. Novit.* 1134:22 pp.
- 1949a. Notes and Descriptions of some Phaoniinae (Diptera: Muscidae). *Am. Mus. Novit.* 1402:25 pp., 14 figs.
- 1949b. New Genera and Species of Lispinae (Diptera, Muscidae). *Am. Mus. Novit.* 1403:9 pp., 5 figs.
- 1949c. Revision of *Neomuscina* Townsend. *Am. Mus. Novit.* 1404: 39 pp.
1951. New Neotropical Muscidae (Diptera). *Am. Mus. Novit.* 1494: 11 pp., 1 fig.



- 1954a. A Revision of *Cyrtoneurina* Giglio-Tos, with Notes on Related Genera (Diptera, Muscidae). *Bull. Am. Mus. nat. Hist.* 103(6): 417-464, 63 figs., 1 table.
- 1954b. A Review of Nearctic *Lispe* Latreille (Diptera, Muscidae). *Am. Mus. Novit.* 1675:40 pp., 77 figs.
1955. Notes and Descriptions of *Muscina* and *Dendrophæonia* (Diptera: Muscidae). *Ann. ent. Soc. Am.* 48:445-452, 14 figs.
- 1957a. Puerto Rican *Neodexiopsis* (Diptera: Muscidae: Coenosiniinae). *J. Agric. Univ. P. Rico* 41(3):207-229.
- 1957b. A New *Pegomya* Leaf Miner in *Chenopodium* from Mexico (Diptera: Muscidae). In Olmedo, I. H., Investigaciones sobre el Minador de la Hoja del Huauzontle. *Revista Chapingo* 64:121-131.
- 1957c. Notes and Descriptions of some Neotropical Muscidae (Diptera). *Bull. Am. Mus. nat. Hist.* 113(6):437-490.
- 1958a. A Review of New World *Neodexiopsis* (Diptera, Muscidae). The *ovata* group. *Am. Mus. Novit.* 1892:27 pp., 16 figs.
- 1958b. Muscidae from the Bahama Islands (Diptera). *Am. Mus. Novit.* 1893:4 pp.

SOBÉRON Y PARRA, G.

1946. Algo acerca de la transmisión del Mal del Pinto. *Kuba* 2(2): 40-43.

SOUZA LOPES, H. DE

- 1938a. Sur une espèce du genre *Charadrella* Wulp (Dipt.-Anthomyiidae) trouvée au Brésil et vivant aux dépens de *Bulinus taumayi* Fer. (Mollusca-Pulmonata). *C. r. Séanc. Soc. Biol.* 128: 926-928, 3 figs.
- 1938b. Sur quelques Diptères porteurs d'oeufs de la *Dermatobia hominis* L. J. au Brésil (Dipt.-Oestridae). *C. r. Séanc. Soc. Biol.* 129:427-428.
1943. Contribuição ao conhecimento das larvas dos Sarcophagidae com especial referência ao esqueleto cefálico (Diptera). *Mem. Inst. Oswaldo Cruz* 38:127-163, 90 figs.
1955. Sobre uma nova espécie de mosca hematófaga do gênero "Neivamyia" P. & F., da Colômbia (Diptera, Muscidae). *Rev. bras. Biol.* 15:415-418, 8 figs.

SOUZA LOPES, H. DE, & MANGABEIRA, O.


1938. Contribuição ao conhecimento do genero *Neivamyia* Pinto et Fonseca, 1930 e descrição de uma nova especie (Diptera: Muscidae). *Livro jubilar Prof. Travassos, R. de J.*: 287-290, 4 plates.

SPEISER, P.

1923. Aethiopische Dipteren. *Wien. ent. Ztg* 40:81-99.

STEIN, P.

1895. Die Anthomyiden-gruppe *Homalomyia* nebst ihren Gattungen und Arten. *Berl. ent. Z.* 40:1-141.
1898. Nordamerikanische Anthomyiden. Beitrag zur Dipterenfauna der Vereinigten Staaten. *Berl. ent. Z.* 42[1897]:161-288.
1899. *Euryomma*, eine neue Gattung der Anthomyiden-gruppe *Homalomyia*. *Ent. Nachr.* 25:19-22.



- 1900a. Anthomyiden aus Neu-Guinea, gesammelt von Herrn L. Biró. *Természetr. Füzet.* 23:129-159.
- 1900b. Vier neue aus Bolivia stammende *Homalomyia*-Arten des Ungarischen National-Museums. *Természetr. Füzet.* 23:205-212.
1901. Die Walker'schen aussereuropäischen Anthomyiden in der Sammlung des British Museum zu London (Dipt.). *Z. syst. Hymenopt. Dipterol.* 1:185-221.
1902. Einige Wiedemann'sche u. Jacnicke'sche aussereuropäische Anthomyiden. (Dipt.) *Z. syst. Hymenopt. Dipterol.* 2:129-137.
1904. Die Amerikanischen Anthomyiden des Königlichen Museums für Naturkunde zu Berlin und des Ungarischen National-Museums zu Budapest. *Ann. hist.-nat. Mus. natn. hung.* 2:414-495.
1907. Revision der Bigot'schen und einiger von Macquart beschriebenen aussereuropäischen Anthomyiden (Dipt.). *Z. syst. Hymenopt. Dipterol.* 7:209-217 and 273-293.
1911. Die von Schunze in Südamerika gefangenen Anthomyiden. *Arch. Naturgesch.* 77(1):61-189.
1918. Zur weiteren Kenntnis aussereuropäischer Anthomyiden. *Ann. hist.-nat. Mus. natn. hung.* 16:147-244.
1919. Die Anthomyidengattungen der Welt, analytisch bearbeitet, nebst einem kritisch-systematischen Verzeichnis aller aussereuropäischen Arten. *Arch. Naturgesch.* 83 A 1 [1917]:85-178, 1 fig.
1920. Nordamerikanische Anthomyiden. 2. Beitrag. *Arch. Naturgesch.* 84 A 9 [1918]:1-106.
- STONE, A., SABROSKY, C. W., WIRTH, W. W., FOOTE, R. H., & COULSON, J. R.
1965. A Catalog of the Diptera of America North of Mexico. *U. S. Dep. Agric., Agric. Handbook* No. 276:1696 pp.
- THOMSON, C. G.
1869. Diptera. In Kongliga svenska fregatten Eugenie's Resa omkring jorden, part 2, Zoologie, 1, Insekter: 443-614, plate 9. Stockholm.
- TOWNSEND, C. H. T.
1892. Catalogue of the described South American Species of Calyptrate Muscidae. *Ann. N. Y. Acad. Sci.* 7:1-44.
1893. A Bot-like Anthomyiid Parasite of a Bird. [Published anonymously.] *J. Inst. Jamaica* 1(8):381-382.
1895a. Contributions to the Dipterology of North America. II. — Tabanidae, Conopidae, Tachinidae, etc. *Trans. Am. ent. Soc.* 22: 55-80.
1895b. The Grass-quit Bot, an Anthomyiid parasite of nestling birds in Jamaica. *J. Inst. Jamaica* 2(2):173-174.
1915a. Correction of the misuse of the generic name *Musca*, with description of two new genera. *J. Wash. Acad. Sci.* 5:433-436.
1915b. Nine new tropical American genera of Muscoidea. *Insecutor Inscit. menstr.* 3:91-97.
1916. Designations of Muscoid Genotypes, with New Genera and Species. *Insecutor Inscit. menstr.* 4:4-12.
1919. New Genera and Species of Muscoid Flies. *Proc. U. S. natn. Mus.* 56:541-592.



1927. Synopse dos generos muscoideos da região humida tropical da America, com generos e especies novas. *Rev. Mus. Paulista* 15: 203-385, 7 figs.
- 1931a. Notes on American Oestromuscoid flies. *Rev. Ent., Rio de J.* 1:65-104 and 157-183.
- 1931b. New Genera and Species of American Oestromuscoid Flies. *Rev. Ent., Rio de J.* 1:313-354 and 437-479.
- 1931c. On *Neivamyia* and its Allies. *Rev. Ent., Rio de J.* 1:479-482.
1932. Five new Brazilian Oestromuscoid genera. *Rev. Ent., Rio de J.* 2:105-107.
1935. Notes on Hough's 1900 species of South American Calliphorinac and Muscinae Musciformes. *Rev. Ent., Rio de J.* 5:366.
1939. Seven new genera of Brazilian Oestromuscarian flies. *Rev. Ent., Rio de J.* 10:446-452.
- TUCKER, R. W. E.
1952. The Insects of Barbados. *J. Agric. Univ. P. Rico* 36:330-363.
1953. Insects of Barbados. *J. Barbados Mus.* 20(4):155-181.
- VIMMER, A.
1939. *Pseudomyiospila*, novy rod Calyptrat (Dipt.). *Cas. esl. Spol. ent.* 36:61.
- VOGELSANG, E. G., & ARMAS, J. C. DE
1940. La Mosquilla del Ganado, *Lyperosia irritans* [L. 1761] en Venezuela. *Rev. Med. vet. Parasit., Caracas* 2:95-98, 2 plates.
- VOLKENBERG, H. L. VAN
1932. Report of the Parasitologist. *Rep. P. Rico fed. agric. Exp. Stn.* 1931: 24-27, fig. 8.
1934. Parasites and Parasitic Diseases of Cattle in Puerto Rico. *Bull. Porto Rico agric. Exp. Stn fed. Stn Mayaguez n.* 36:26 pp., 4 figs.
1935. Parasites and Parasitic Diseases of Horses in Puerto Rico. *Bull. Porto Rico agric. Exp. Stn fed. Stn Mayaguez n.* 37:19 pp., 3 figs.
- WALKER, F.
1837. Descriptions, & c. of the Insects collected by Captain P. P. King, R. N., F. R. S., in the Survey of the Straits of Magellan. Diptera. *Trans. Linn. Soc. Lond.* 17:331-359.
1849. *List of the Specimens of Dipterous Insects in the collection of the British Muscum.* Part 4: 689-1172. London.
1852. *Insecta Saundersiana: or characters of undescribed insects in the collection of William Wilson Saunders, Esq., F. R. S., F. L. S., & c.* Volume I, Diptera, part 4: 253-414. London.
1853. *Insecta Britannica, Diptera.* Volume 2. vi + 297 pp. London, Rceve and Benham.
1857. Characters of undescribed Diptera in the Collection of W. W. Saunders, Esq., F. R. S., & c. [Part.] *Trans. ent. Soc. Lond.* (2)4:119-158.



1858. Characters of undescribed Diptera in the Collection of W. W. Saunders, Esq., F. R. S., & c. [Concl.] *Trans. ent. Soc. Lond.* (2)4:190-235.
1860. Characters of undescribed Diptera in the Collection of W. W. Saunders, Esq., F. R. S., & c. [Part.] *Trans. ent. Soc. Lond.* (2)5:268-296.
1861. Characters of undescribed Diptera in the Collection of W. W. Saunders, Esq., F. R. S., & c. [Concl.] *Trans. ent. Soc. Lond.* (2)5:297-334.
- WALTON, W. R.
1918. Three New Tachinid Parasites of Eleodes. *Proc. ent. Soc. Wash.* 19 [1917]:22-26, plate 5.
- WEST, L. S.
1951. *The Housefly. Its Natural History, Medical Importance, and Control.* xi + 584 pp., 176 figs., 16 tables, 1 photograph, New York.
- WESTWOOD, J. O.
1840. *An Introduction to the Modern Classification of Insects. Synopsis of the Genera of British Insects.* 158 pp., London.
- WIEDEMANN, C. R. W.
1817. Neue Zweiflügler (Diptera Linn.) aus der Gegend um Kiel. *Zoologisches Magazin* 1(1):61-86.
1818. Neue Insecten vom Vorgebirge der guten Hoffnung. *Zoologisches Magazin* 1(2):40-48.
1819. Brasilianische Zweiflügler. *Zoologisches Magazin* 1(3):40-56.
1824. *Analecta entomologica ex Museo Regio Havniensi.* 60 pp., 12 figs., Killiae.
1830. *Aussereuropäische zweiflügelige Insekten.* Part II. xii + 684 pp., Hamm.
- WILLIAMS, R. W.
1958. A study of the summer filth fly population of the Bermuda Islands. *J. Parasit.* 44:339-342, 1 table.
1959. The reported biting and filth frequenting arthropods of the Bermuda Islands exclusive of the Ixodidae and Araneida. *Proc. ent. Soc. Wash.* 61:234-238.
- WILLISTON, S. W.
1896. On the Diptera of St. Vincent (West Indies). *Trans. ent. Soc. Lond.* 1896(3):253-449, plates 8-14.
- WOKE, P. A.
1947. Arthropods of sanitary importance in the Republic of Nicaragua, Central America. *Am. J. trop. Med.* 27(3):357-375, 1 map, 1 table.
- WOLCOTT, G. N.
1924. "Insectae Portoricensis". A preliminary annotated check-list of the insects of Porto Rico, with descriptions of some new species. *J. Dcp. Agric. P. Rico* 7(1):1-313, 2 plates.



1927. *Entomologie d'Haiti*. 440 pp., 133 figs., Agric. Dept. Haiti.
1936. "Insectae Borinquenses". A Revised Annotated Check-List of the Insects of Puerto Rico. *J. Agric. Univ. P. Rico* 20(1):627 + 3 pp., illustr.
1941. Supplement to "Insectae Borinquenses". *J. Agric. Univ. P. Rico* 25(2):33-158.
1948. The Insects of Puerto Rico. Diptera. *J. Agric. Univ. P. Rico* 32(3):417-532, figs.
1955. Entomología económica puertorriqueña. *Bull. Porto Rico agric. Exp. Stn insular Stn Rio Pedras* no. 125:208 pp., 1 plate, 133 figs.
- WULP, F. M. VAN DER
1883. Amerikaansche Diptera (III). *Tijdschr. Ent.* 26:1-60, plates 1-2.
1888. Nieuwe Argentijnsche Diptera van wijlen Prof. H. Weyenbergh Jr. *Tijdschr. Ent.* 31:359-376.
- 1888-1903. Insecta, Diptera, In Godman, F. D., and Salvin, O., *Biology Centrali-Americana*. Volume 2: x + 489 pp., 11 figs., 13 plates. London.
- ZELEDÓN, R.
1957. Algunas observaciones sobre la biología de la *Dermatobia hominis* (L. Jr.) y el problema del tórsalo en Costa Rica. *Rev. Biol. Trop.* 5(1):63-74, 1 plate, 1 graph, 2 tables.
- ZETTERSTEDT, J. W.
- 1838-1840. *Insecta Lapponica*. vi + 1140 pp., Lipsiae.
1845. *Diptera Scandinaviae. Disposita et descripta*. Volume 4:1281-1738. Lundae.
- ZUMPT, F.
1938. 1. Vorstudie zu einer monographischen Bearbeitung der Stomoxydinae. Taxonomische Ergebnisse des Studiums einiger Sammlungen, besonders der von Bezzi und Enderlein. *Z. angew. Ent.* 25:337-353.
1939. Das System der Stomoxydinae. Ein Beitrag zu schwebenden Fragen der Taxonomie. 2. Vorstudie zu einer monographischen Bearbeitung der Stomoxydinae. VII. *Int. Congr. Ent., Berlin 1938*: 1723-1733.
1940. 4. Vorstudie zu einer monographischen Bearbeitung der Stomoxydinae. Ueber einige afrikanische und orientalische *Stomoxys*-Arten. *Z. angew. Ent.* 27:126-141.

Index


<i>abacta</i> (Giglio-Tos), 1893 (Hyetodesia), <i>Helina</i>	97.21
<i>abdicta</i> (Giglio-Tos), 1893 (Hycotodesia), <i>Helina</i>	97.21
<i>abdicta</i> (Giglio-Tos), 1893 (Spilogaster), <i>Bigotomyia</i>	97.18
<i>aberrans</i> (Shannon & Del Ponte), 1926 (Muscina), <i>Myospila</i>	97.26
<i>abnorminervis</i> (Stein), 1911, (Limnophora), <i>Lispoides</i>	97.32
<i>abnormis</i> (Stein), 1900 (Homalomyia), <i>Fannia</i>	97.3
<i>Acanthoneura</i> Stein, 1919	97.30
<i>Acanthosphys</i> Enderlein, 1935	97.17

97.87

<i>aconquija</i> Shannon & Del Ponte, 1926, Morellia	97.9
<i>acrinis</i> Snyder, 1941, Helina	97.19
<i>Acritochaeta</i> Grimshaw, 1901	97.24
<i>Acrolasia</i> Enderlein, 1927	97.17
<i>acromion</i> Wiedemann, 1824 (Musca), Tachinidae	97.59
<i>acrostichalis</i> Snyder, 1941, Helina	97.19
<i>acuta</i> (Malloch), 1934 (Angolia), Angolina	97.40
<i>acuta</i> Stein, 1898, Hydrotaea	97.12
<i>aczeli</i> Snyder, 1957, Bucephalomya	97.32
<i>aczeli</i> Snyder, 1957, Helina	97.19
<i>adelpha</i> (Schiner), 1868 (Spilogaster) Helina	97.19
<i>admirabilis</i> Albuquerque, 1958, Fannia	97.3
<i>advena</i> Snyder, 1957, Phaonia	97.14
<i>aeneiventris</i> Wiedemann, 1830 (Dexia), Calliphoridae	97.59
<i>aenescens</i> (Wiedemann), 1830 (Anthomyia), Ophyra	97.13
<i>aenula</i> (Shannon & Del Ponte), 1926 (Chlorellia), Dasymorellia ..	97.10
<i>aequifrons</i> (Stein), 1898 (Limnophora), Lispoides	97.32
<i>aethiops</i> Walker, 1852 (Anthomyia), Agromyzidae	97.59
<i>affinis</i> Malloch, 1952, Morellia	97.8
<i>Agenamyia</i> Albuquerque, 1953	97.32
<i>Airalips</i> Snyder, 1949	97.48
<i>aitkeni</i> (Dodge), 1963 (Philornis), Neomusca	97.55
<i>alacris</i> Stein, 1911, Limnophora	97.28
<i>alibiarba</i> (Snyder), 1957 (Angolia), Angolina	97.40
<i>albibasis</i> Malloch, 1934, Fannia	97.3
<i>albiceps</i> Malloch, 1934, Schoenomyza	97.39
<i>albiceps</i> Malloch, 1934, Spathipheromyia	97.38
<i>albina</i> Wiedemann, 1830, Musca	97.59
<i>alpipila</i> Stein, 1918, Coenosia	97.44,60
<i>albiseta</i> (Stein), 1911 (Coenosia), Stomopogon	97.41
<i>albisquama</i> (Albuquerque), 1959 (Austrocoenosia), Coenosia	97.41
<i>albitarsis</i> Stein, 1911, Fannia	97.3
<i>albitarsis</i> Stein, 1898, Lispe	97.34
<i>albomedia</i> Malloch, 1934, Schoenomyza	97.39
<i>albuquerquei</i> new name, Helina	97.19
<i>aliena</i> (Stein), 1911 (Limnophora), Syllimnophora	97.30
<i>altaneira</i> (Albuquerque), 1954 (Heliographa), Syllimnophora	97.30
<i>amazonica</i> Albuquerque, 1951, Souzalopesmyia	97.23
<i>americana</i> (Hough), 1900 (Muscina), Myospila	97.27
<i>americana</i> Schiner, 1868, Graphomya	97.28
<i>Anacanthosphys</i> Enderlein, 1935	97.17
<i>analís</i> (Macquart), 1851 (Cyrtonevra), Graphomya	97.27
<i>analís</i> Macquart, 1843, Musca	97.7
<i>andina</i> Bigot, 1885 (Ophyra), Lonchaeidae	97.59
<i>angolensis</i> (Malloch), 1934 (Darwinomyia), Palpibracus	97.17
<i>Angolia</i> Malloch, 1934	97.40
<i>Angolina</i> new name	97.40
<i>angustifrons</i> (Loew), 1861 (Hylemyia), Neomusca	97.56
<i>angustifrons</i> Malloch, 1934, Syllimnophora	97.30
<i>angustifrons</i> (Stein), 1904 (Spilogaster), Neomusca	97.57
<i>angustipennis</i> (Stein), 1911 (Mydaea), Helina	97.19
<i>annosa</i> Chillcott, 1961, Fannia	97.3
<i>annulata</i> Albuquerque, 1957, Bigotomyia	97.18
<i>annulata</i> Stein, 1911, Bithoracochaeta	97.36



<i>annulata</i> (Stein), 1911 (<i>Schoenomyza</i>), <i>Notoschoenomyza</i>	97.38
<i>annulipes</i> Macquart, 1843, <i>Coenosia</i>	97.42
<i>anomala</i> (Jaennicke), 1867 (<i>Mesembrina</i>), <i>Neomusca</i>	97.55
<i>antennata</i> Stein, 1911, <i>Fannia</i>	97.3
<i>anthomydea</i> (Bigot), 1888 (<i>Curtonevra</i>), <i>Myospila</i>	97.27
<i>anthomyidaea</i> , error.	97.27
<i>anthracina</i> Albuquerque, 1956, <i>Helina</i>	97.19
<i>anthracina</i> (Walker), 1837 (<i>Anthomyia</i>), <i>Fannia</i>	97.3
<i>anthrax</i> (Bigot), 1885, (<i>Limnophora</i>), <i>Spilogona</i>	97.33
<i>antica</i> (Walker), 1852 (<i>Coenosia</i>), <i>Bithoracochaeta</i>	97.37
<i>antunesi</i> Souza Lopes, 1955, <i>Neivamyia</i>	97.59
<i>anubes</i> Snyder, 1941, <i>Helina</i>	97.19
<i>apicalis</i> (Malloch), 1934 (<i>Darwinomyia</i>), <i>Palpibracus</i>	97.17
<i>apicalis</i> Malloch, 1934, <i>Schoenomyza</i>	97.40
<i>apicalis</i> (Stein), 1911 (<i>Schoenomyza</i>), <i>Notoschoenomyza</i>	97.38
<i>apicata</i> Malloch, 1934, <i>Apsil</i>	97.35
<i>apicata</i> (Stein), 1904 (<i>Spilogaster</i>), <i>Neomuscina</i>	97.50
<i>Apsil</i> Malloch, 1929	97.35
<i>arcuata</i> Chillcott, 1961, <i>Fannia</i>	97.3
<i>arcuata</i> (Stein), 1898 (<i>Limnophora</i>), <i>Gymnodia</i>	97.24
<i>arcuata</i> (Townsend), 1927 (<i>Sarcopromusca</i>), <i>Morellia</i>	97.9
<i>arcuata</i> (Wiedemann), 1830 (<i>Anthomyia</i>), <i>Mydaea</i>	97.26
<i>argenticeps</i> Malloch, 1934, <i>Lispoides</i>	97.32
<i>argentifera</i> (Bigot), 1877 (<i>Somomyia</i>), <i>Orthellia</i>	97.10
<i>argentifrons</i> (Malloch), 1934 (<i>Austrocoenosia</i>), <i>Coenosia</i>	97.41
<i>argentifrontata</i> Snyder, 1957, <i>Spilogona</i>	97.33
<i>argentina</i> Bigot, 1885, <i>Ophyra</i>	97.13
<i>argentina</i> Malloch, 1934, <i>Lispoides</i>	97.32
<i>argentina</i> (Snyder), 1957 (<i>Angola</i>), <i>Angolina</i>	97.40
<i>argentina</i> Snyder, 1957, <i>Helina</i>	97.19
<i>argyriceps</i> Malloch, 1934, <i>Schoenomyza</i>	97.39
<i>Ariciella</i> Malloch, 1918	97.19
<i>arizona</i> Snyder, 1958, <i>Neodexiopsis</i>	97.45
<i>armata</i> (Bigot), 1883 (<i>Dasyphyma</i>), <i>Fannia</i>	97.6
<i>armipes</i> Malloch, 1934, <i>Schoenomyza</i>	97.39
<i>armipes</i> (Stein), 1911 (<i>Mydaea</i>), <i>Cyrtoneurina</i>	97.52
<i>arroya</i> Snyder, 1949, <i>Helina</i>	97.23
<i>Arthurella</i> Albuquerque, 1954	97.57
<i>aterrima</i> (Stein), 1904 (<i>Limnophora</i>), <i>Spilogona</i>	97.33
<i>aterrima</i> Wulp, 1896, <i>Pogonomyia</i>	97.12
<i>atincta</i> Snyder, 1949, <i>Neomuscina</i>	97.50
<i>atinctiocosta</i> Snyder, 1949, <i>Neomuscina</i>	97.50
<i>Atherigona</i> Rondani, 1856	97.24
<i>Atherigonini</i>	97.24
<i>atra</i> Malloch, 1934, <i>Spathipheromya</i>	97.38
<i>atra</i> (Stein), 1904 (<i>Limnophora</i>), <i>Syllimnophora</i>	97.30
<i>atricornis</i> Malloch, 1934, <i>Bithoracochaeta</i>	97.36
<i>atricornis</i> (Malloch), 1934 (<i>Darwinomyia</i>), <i>Palpibracus</i>	97.18
<i>atrifrons</i> Bigot, 1888, <i>Musca</i>	97.7
<i>atrifrons</i> Stein, 1911, <i>Coenosia</i>	97.41
<i>atripes</i> Malloch, 1934, <i>Apsil</i>	97.35
<i>atrisquama</i> (Stein), 1904 (<i>Limnophora</i>), <i>Lispoides</i>	97.32
<i>atrovittata</i> (Stein), 1904 (<i>Limnophora</i>), <i>Syllimnophora</i>	97.30
<i>auctila</i> (Stein), 1901 (<i>Limnophora</i>), <i>Gymnodia</i>	97.24



<i>aurantiaca</i> Hough, 1899, <i>Muscina</i>	97.11
<i>aurea</i> (Séguy), 1937 (<i>Dyadimya</i>), <i>Cyrtoneurina</i>	97.52
<i>auriceps</i> Malloch, 1934, <i>Graphomya</i>	97.27
<i>auriceps</i> (Stein), 1911 (<i>Schoenomyza</i>), <i>Spathipheromyia</i>	97.38
<i>aurifacies</i> Stein, 1911, <i>Limnophora</i>	97.28
<i>aurifera</i> (Bigot), 1885 (<i>Limnophora</i>), <i>Reynoldsia</i>	97.36
<i>aurifera</i> (Malloch), 1934 (<i>Austrocoenosia</i>), <i>Coenosia</i>	97.41
<i>auriflua</i> (Stein), 1911 (<i>Limnophora</i>), <i>Syllimnophora</i>	97.30
<i>aurulans</i> Robineau-Diesvoidy, 1830, <i>Musca</i>	97.7
<i>australis</i> (Brèthes), 1924 (<i>Ochthiphila</i>), <i>Schoenomyza</i>	97.40
<i>australis</i> (Malloch), 1934, (<i>Coenosia</i>), <i>Neodexiopsis</i>	97.46
<i>Austrocoenosia</i> Malloch, 1934	97.41
<i>Azella</i> Robineau-Desvoidy, 1830	97.12
<i>bahama</i> Snyder, 1958, <i>Lispe</i>	97.34
<i>bahiensis</i> Albuquerque, 1954, <i>Fannia</i>	97.3
<i>barbitarsis</i> (Stein), 1911 (<i>Limnophora</i>), <i>Syllimnophora</i>	97.30
<i>basalaris</i> , error	97.7
<i>basalis</i> Malloch, 1934, <i>Helina</i>	97.22
<i>basalis</i> (Walker), 1852 (<i>Musca</i>), <i>Morellia</i>	97.8
<i>basicincta</i> (Stein), 1904 (<i>Aricia</i>), <i>Polietina</i>	97.54
<i>basilaris</i> Macquart, 1843, <i>Musca</i>	97.7
<i>beebei</i> (Curran), 1934 (<i>Mallocharia</i>), <i>Cyrtoneurina</i>	97.52
<i>bela</i> , error	97.3
<i>bella</i> Albuquerque, 1957 <i>Fannia</i>	97.3
<i>bella</i> Malloch, 1934, <i>Schoenomyza</i>	97.39
<i>benevenuta</i> Albuquerque, 1957, <i>Noelia</i>	97.41
<i>benjamini</i> Malloch, 1913, <i>Fannia</i>	97.3
<i>benoisti</i> Séguy, 1932, <i>Coenosia</i>	97.41
<i>bequaerti</i> Aldrich, 1932, <i>Chortinus</i>	97.57
<i>bicolor</i> Albuquerque, 1956, <i>Polietina</i>	97.55
<i>bicolor</i> (Bigot), 1885 (<i>Dialyta</i>), <i>Cordiluroides</i>	97.44
<i>bicolor</i> (Macquart), 1855 (<i>Aricia</i>), <i>Diachatomyia</i>	97.60
<i>bicolor</i> Malloch, 1934, <i>Correntosia</i>	97.16
<i>bifimbriata</i> Malloch, 1934, <i>Helina</i>	97.22
<i>bifimbriata</i> <i>vasalis</i> Malloch, 1934, <i>Helina</i>	97.22
<i>bifimbriata</i> <i>grisea</i> Malloch, 1934, <i>Helina</i>	97.22
<i>bigoti</i> Albuquerque, 1957, <i>Bigotomyia</i>	97.18
<i>bigoti</i> Malloch, 1934, <i>Helina</i>	97.19
<i>bigoti</i> (Stein), 1907 (<i>Homalomyia</i>), <i>Fannia</i>	97.3
<i>Bigotomyia</i> Malloch, 1921	97.18
<i>binotata</i> Chillcott, 1961, <i>Fannia</i>	97.3
<i>Biopyrellia</i> Townsend, 1932	97.8
<i>bipuncta</i> (Wiedemann), 1830 (<i>Musca</i>), <i>Morellia</i> :.....	97.8
<i>bipunctata</i> (Schiner), 1868 (<i>Spilogaster</i>), <i>Bigotomyia</i>	97.18
<i>bipunctata</i> , error	97.8
<i>biseriata</i> Malloch, 1934, <i>Schoenomyza</i>	97.39
<i>biseta</i> Malloch, 1934, <i>Apsil</i>	97.35
<i>biseta</i> Snyder, 1954, <i>Cyrtoneurina</i>	97.52
<i>biseta</i> (Stein), 1904 (<i>Spilogaster</i>), <i>Helina</i>	97.19
<i>biseta</i> (Stein), 1911 (<i>Schoenomyza</i>), <i>Schoenomyzina</i>	97.39
<i>bistriata</i> (Wulp), 1896 (<i>Coenosia</i>), <i>Cordiluroides</i>	97.44
<i>Bithoracochaeta</i> Stein, 1911	97.36



<i>blanchardi</i> (Albuquerque), 1953 (Philornis), Neomusca	97.56
<i>blanchardi</i> (García), 1952 (Philornis), Neomusca	97.55
<i>boliviana</i> (Townsend), 1931 (Trichomorellia), Dasymorellia	97.10
<i>Brachiophyra</i> Giglio-Tos, 1893	97.28
<i>Brachygasterina</i> Macquart, 1851	97.16
<i>Brachyophyra</i> , error	97.28
<i>Brachypalpus</i> Macquart, 1851	97.16
<i>brasiliana</i> Brauer & Bergenstamm, 1893, Synthesiomyia	97.11
<i>brasiliensis</i> Albuquerque, 1958, Phaonia	97.14
<i>brasiliensis</i> (Walker), 1852 (Anthomyia), Neodexiopsis	97.45
<i>brevicornis</i> (Malloch), 1934 (Austrocoenosia), Neodexiopsis	97.45
<i>brevihirta</i> Malloch, 1934, Limnophora	97.28
<i>brevipalpis</i> (Thomson), 1869 (Anthomyia), Euryomma	97.2
<i>brevipectinata</i> (Stein), 1918 (Mydaea), Cyrtoneuriniinae	97.58
<i>brevis</i> (Schiner), 1868 (Cyrtoneura), Morellia	97.8
<i>breviseta</i> (Stein), 1911 (Limnophora), Syllimnophora	97.30
<i>brevitarsis</i> Malloch, 1934, Reynoldsia	97.36
<i>brevivena</i> Snyder, 1941, Helina	97.19
<i>breviventris</i> Stein, 1911, Schoenomyza	97.39
<i>browni</i> Snyder, 1941, Helina	97.19
<i>browni</i> Snyder, 1957, Syllimnophora	97.30
<i>brunnea</i> (Hough), 1900 (Muscina), Cyrtoneurina	97.52
<i>Bucephalomyia</i> Malloch, 1918	97.32
<i>cacheuta</i> Snyder, 1957, Phaonia	97.14
<i>cacumina</i> Snyder, 1957, Neodexiopsis	97.45
<i>cacrulea</i> (Snyder), 1949 (Airalips), Xenothoracochaeta	97.49
<i>caerulea</i> , error	97.15
<i>caesarion</i> (Meigen), 1826 (Musca), Orthellia	97.10
<i>calcitrans</i> (Linnaeus), 1758 (Conops), Stomoxys	97.58
<i>callidimera</i> , error	97.8
<i>callimera</i> (Bigot), 1888 (Curtonevra), Morellia	97.8
<i>calliphoroides</i> (Jaenicke), 1867 (Spilogaster), Mydaea	97.26
<i>calopus</i> (Bigot), 1885 (?Hydrophoria), Bithoracochaeta	97.37
<i>calopyga</i> (Loew), 1872 (Coenosia), Neodexiopsis	97.45
<i>cambuquirensis</i> Albuquerque, 1954, Haroldopsis	97.37
<i>camorinensis</i> Albuquerque, 1956, Coenosia	97.41
<i>canaliculata</i> Robineau-Desvoidy, 1830, Phaonia	97.15
<i>candidifrons</i> (Stein), 1911 (Limnophora), Syllimnophora	97.30
<i>caneo</i> Snyder, 1941, Helina	97.14
<i>canicularis</i> (Linnaeus), 1758 (Musca), Fannia	97.3
<i>capalta</i> Snyder, 1949, Neomuscina	97.50
<i>capensis</i> (Wiedemann), 1818 (Anthomyia), Ophyra	97.13
<i>capribarba</i> (Stein), 1911 (Coenosia), ?Noelia	97.41
<i>caras</i> Snyder, 1949, Mydaea	97.25
<i>carbonaria</i> Shannon & Del Ponte, 1926, Ophyra	97.13
<i>cardini</i> Malloch, 1936, Scenetes	97.24
<i>Caricea</i> Robineau-Desvoidy, 1830	97.41
<i>carinata</i> (Dodge), 1968, (Philornis), Neomusca	97.55
<i>carioca</i> Albuquerque, 1956, Euryomma	97.2
<i>carioca</i> Albuquerque, 1951, Souzalopesmyia	97.23
<i>Cariocamyia</i> Snyder, 1951	97.57
<i>carnifex</i> Stein, 1911, Psilochaeta	97.15



<i>castanea</i> Albuquerque, 1955, Mydaea	97.25
<i>catamacla</i> Snyder, 1957, Phaonia	97.14
<i>Catantigymnus</i> Enderlein, 1935	97.17
<i>cavalata</i> Snyder, 1957, Neodexiopsis	97.45
<i>cavicola</i> Townsend, 1919, Neomuscina	97.51
<i>centralis</i> (Loew), 1869 (Pyrellia), Morellia	97.9
<i>centralis</i> Stein, 1911, Mydaea	97.25
<i>cera</i> Snyder, 1958, Neodexiopsis	97.45
<i>Chaetagenia</i> Malloch, 1928	97.49
<i>Chaetopyrellia</i> Townsend, 1932	97.8
<i>chaetosa</i> Malloch, 1934, Coenosia	97.42
<i>Chaetura</i> error	97.41
<i>Chaetypopleura</i> Enderlein, 1927	97.54
<i>chalcogaster</i> (Wiedemann), 1824 (Anthomya), Ophyra	97.13
<i>chalybea</i> (Wiedemann), 1830 (Anthomyia), Psilochaeta	97.15
<i>Charadrella</i> Wulp, 1896	97.57
<i>chilensis</i> Bigot, 1888, Graphomya	97.27
<i>chilensis</i> (Bigot), 1885 (?Yctodesia), Palpibracus	97.17
<i>chilensis</i> (Macquart), 1843 (Anthomyia), Fannia	97.4
<i>chilensis</i> Macquart, 1843, Musca	97.7
<i>chilensis</i> Malloch, 1934, Helina	97.19
<i>chilensis</i> Malloch, 1934, Spathipheromyia	97.38
<i>chilensis</i> new name, Coenosia	97.42
<i>chiliensis</i> , error	97.7
<i>chloe</i> (Bigot), 1878 (Pyrellia), Morellia	97.7
<i>Chlorellia</i> Shannon & Del Ponte, 1926	97.10
<i>chlorogaster</i> (Bigot), 1885 (Limnophora), Psilochaeta	97.16
<i>chlorogaster</i> (Wiedemann), 1830 (Anthomyia), Psilochaeta	97.16
<i>Choetura</i> Macquart, 1851	97.41
<i>Cholomyioides</i> Albuquerque, 1954	97.40
<i>Chortinus</i> Aldrich, 1932	97.57
<i>chrysiceps</i> Malloch, 1934, Notoschoenomyza	97.39
<i>cilitibia</i> Albuquerque, 1955, Phaonia	97.4
<i>cinnamomina</i> (Stein), 1918 (Mydaea), Neomusca	97.55
<i>circulatrix</i> (Walker), 1861 (Aricia), Helina	97.19
<i>cirratipila</i> Snyder, 1957, Neodexiopsis	97.45
<i>clavacula</i> Snyder, 1957, Neodexiopsis	97.45
<i>clavata</i> Chillcott, 1961, Fannia	97.4
<i>clavitibia</i> (Stein), 1911 (Limnophora), Syllimnophora	97.30
<i>Clinopera</i> Wulp, 1896	97.51
<i>cluvia</i> Walker, 1849 (Musca), Calliphoridae	97.59
<i>cocquilletti</i> , error	97.18
<i>Coenosia</i> Meigen, 1826	97.47
<i>Coenosinae</i>	97.34
<i>Coenosopsia</i> Malloch, 1924	97.13
<i>coerulea</i> (Macquart), 1843 (Ophyra), Psilochaeta	97.15
<i>compressicornis</i> Snyder, 1949, Mydaea	97.25
<i>compressifrons</i> Snyder, 1940, Helina	97.21
<i>compressifrons fulvapoda</i> Snyder, 1940, Helina	97.20
<i>compressifrons</i> (Stein), 1904 (Limnophora), Syllimnophora	97.30
<i>compressipalpis</i> Stein, 1911, Phaonia	97.14
<i>compressitarsis</i> (Stein), 1911 (Limnophora), Syllimnophora	97.30
<i>compressitarsis</i> Stein, 1911, Rhyncholimnophora	97.31



<i>concinna</i> (Wulp), 1896 (Mydaea), Polletina	97.55
<i>eoneolor</i> (Walker), 1852 (Musea), Helina	97.60
<i>confinis</i> ((Wulp), 1896 (Mydaea), Helina	97.20
<i>confusa</i> Snyder, 1954, Cyrtoneurina	97.52
<i>econfusa</i> Snyder, 1949, Pseudoptilolepis	97.49
<i>econfusus</i> (Malloch), 1928 (Darwinomyia), Palpibraeus	97.17
<i>connexa</i> Malloch, 1934, Helina	97.20
<i>consanguinea</i> Rondani, 1848, Musea	97.7
<i>consanguinea</i> (Stein), 1911 (Mydaea), Helina	97.20
<i>econspersa</i> (Stein), 1911 (Mydaea), Cyrtoneurina	97.52
<i>eontinens</i> Snyder, 1954, Cyrtoneurina	97.52
<i>econvexa</i> Stein, 1918, Mydaea	97.26
<i>eopiosa</i> (Wulp), 1896 (Spilogaster), Helina	97.20
<i>coquilleti</i> , error	97.18
<i>eoquilletti</i> (Vimmer), 1939 (Pseudomyospila), Bigotomyia	97.18
<i>Cordiluroides</i> Albuquerque, 1954	97.44
<i>eordyduroides</i> (Stein), 1898 (Phyllogaster), Tetramerinx	97.35
<i>cornicina</i> (Fabrieius), 1781 (Musca), Orthellia	97.10,59
<i>Correntosia</i> Malloch, 1934	97.16
<i>corvina</i> (Giglio-Tos), 1893 (Leucomelina), Limnophora	97.28
<i>eostalis</i> Walker, 1852 (Musca), Taehinidae	97.59
<i>costalis</i> (Walker), 1852 (?Notiphila), Cyrtoneurina	97.52
<i>eostata</i> Snyder, 1957, Notosehoenomyza	97.39
<i>eostipennis</i> Snyder, 1957, Bigotomyia	97.18
<i>eothurnata</i> (Bigot), 1885 (?Anthomyia), Coenosia	97.42
<i>cotidiana</i> Snyder, 1954, Lisse	97.34
<i>eoxata</i> Malloch, 1934, Reynoldsia	97.36
<i>eoxata</i> Shannon & Del Ponte, 1926, Fannla	97.4
<i>erassieauda</i> Stein, 1911, Coenosia	97.42
<i>erassierurus</i> Snyder, 1957, Neodexiopsis	97.45
<i>erepedoseta</i> Snyder, 1940, Helina	97.20
<i>eresa</i> Snyder, 1949, Mydaea	97.25
<i>erispaseta</i> Snyder, 1954, Cyrtoneurina	97.52
<i>erlspiseta</i> Snyder, 1957, Neodexiopsis	97.45
<i>crocea</i> Snyder, 1940, Helina	97.20
<i>eroceafrons</i> Snyder, 1957, Ncodexiopsis	97.45
<i>eubana</i> Dodge, 1965, Graphomya	97.27
<i>cubana</i> Johnson, 1919, Limnophora	97.28
<i>eurrani</i> Snyder, 1949, Neomuseina	97.50
<i>eurvata</i> (Stein), 1904 (Arieia), Bigotomyia	97.18
<i>eurviventris</i> Albuquerque, 1959, Coenosia	97.42
<i>cutilia</i> (Walker), 1849 (Anthomyia), Psilochaeta	97.15
<i>Cyaeyrtoneura</i> Townsend, 1931	97.11
<i>eyanea</i> (Macquart), 1843 (Curtonevra), Myospila	97.26
<i>cyanea</i> (Macquart) of Stein, 1919, Dasymorellia	97.10
<i>eyanea</i> Townsend, 1931, Cyaeirtoneura	97.11
<i>cyanea</i> , error	97.16
<i>cyaneiventris</i> Macquart, 1851, Hydrotaea	97.12
<i>eyanieolor</i> (Stein), 1911 (Phaonia), Helina	97.20
<i>cyaniventris</i> , error	97.12
<i>eyllndria</i> (Macquart), 1846 (Cyrtonevra), Cyrtoneurina	97.52
<i>Cyrtoneurina</i> Giglio-Tos, 1893	97.51
<i>cyrtoneurina</i> Stein, 1898, Limnophora	97.29

<i>cyrtoneurina</i> (Zetterstedt), 1845 (<i>Aricia</i>), <i>Hydrotaea</i>	97.12,59
<i>Cyrtoneurininae</i>	97.48
<i>Cyrtoneuropsis</i> Malloch, 1925	97.52
<i>Cyrtoneuropsis</i> Townsend, 1931	97.11
<i>Darwinomyia</i> Malloch, 1922	97.16
<i>Dasymorellia</i> Malloch, 1923	97.10
<i>dasyophthalma</i> Malloch, 1928, <i>Helina</i>	97.20
<i>dasyops</i> (Macquart), 1843 (<i>Chortophila</i>), <i>Helina</i>	97.20
<i>dasyops</i> (Stein), 1900 (<i>Homalomyia</i>), <i>Fannia</i>	97.4
<i>dasyops</i> (Stein), 1911 (<i>Mydaea</i>), <i>Helina</i>	97.20
<i>Dasyphyra</i> Bigot, 1883	97.3
<i>davida</i> Snyder, 1949, <i>Neomuscina</i>	97.50
<i>debilis</i> (Williston), 1896 (<i>Limnophora</i>), <i>Gymnodia</i>	97.24
<i>deceptiva</i> (Dodge & Aitken), 1968 (<i>Philornis</i>), <i>Neomusca</i>	97.55
<i>declivis</i> (Stein), 1904 (<i>Coenosia</i>), <i>Neodexiopsis</i>	97.45
<i>decora</i> Snyder, 1941, <i>Helina</i>	97.22
<i>delecta</i> (Wulp), 1896 (<i>Limnophora</i>), <i>Gymnodia</i>	97.24
<i>deleta</i> (Wulp), 1896 (<i>Leucomelina</i>), <i>Syllimnophora</i>	97.30
<i>Dendrophaonia</i> Malloch, 1923	97.13
<i>dentipes</i> (Fabricius), 1805 (<i>Musca</i>), <i>Hydrotaca</i>	97.12
<i>despecta</i> (Walker), 1852 (<i>Anthomyia</i>), <i>Bithoracochaeta</i>	97.37
<i>devia</i> Albuquerque, 1957, <i>Phaonantho</i>	97.14
<i>devia</i> (Curran), 1934 (<i>Xenocoenosia</i>), <i>Neodexiopsis</i>	97.45
<i>diagramma</i> , error	97.53
<i>diaphana</i> Stein, 1911, <i>Coenosia</i>	97.42
<i>dichroma</i> (Wiedemann), 1830 (<i>Anthomyia</i>), <i>Mydaea</i>	97.26
<i>differentia</i> (Snyder), 1949 (<i>Airalips</i>), <i>Xenothoracochaeta</i>	97.49
<i>digramma</i> (Wulp), 1896 (<i>Clinopera</i>), <i>Cyrtoneurina</i>	97.53
<i>dilata</i> Malloch, 1934, <i>Apsil</i>	97.35
<i>diluta</i> (Stein), 1911 (<i>Limnophora</i>), <i>Lispoides</i>	97.32
<i>discolor</i> (Stein), 1911 (<i>Mydaea</i>), <i>Helina</i>	97.20
<i>discolorisexus</i> Snyder, 1957, <i>Neodexiopsis</i>	97.45
<i>discreta</i> Albuquerque, 1956, <i>Helina</i>	97.19
<i>discreta</i> (Wulp), 1896 (<i>Spilogaster</i>), <i>Helina</i>	97.20
<i>ditiportus</i> Snyder, 1957, <i>Neodexiopsis</i>	97.45
<i>diversipalpis</i> (Rondani), 1864 (<i>Pyrellia</i>), <i>Morellia</i>	97.8
<i>dodgei</i> Seago, 1954, <i>Fannia</i>	97.4
<i>domestica</i> Linnaeus, 1758, <i>Musca</i>	97.7
<i>domestica major</i> Degeer, 1776, <i>Musca</i>	97.7
<i>dorae</i> Stein, 1911, <i>Drepanocnemis</i>	97.34
<i>dorsilinea</i> (Wulp), 1896 (<i>Clinopera</i>), <i>Muscina</i>	97.11
<i>dorsipuncta</i> (Stein), 1918 (<i>Mydaea</i>), <i>Neomuscina</i>	97.50
<i>dorsovittata</i> (Malloch), 1920 (<i>Eulimnophora</i>), <i>Gymnodia</i>	97.24
<i>downsi</i> (Dodge & Aitken), 1968 (<i>Philornis</i>), <i>Neomusca</i>	97.56
<i>Drepanocnemis</i> Stein, 1911	97.34
<i>dubia</i> (Bigot), 1885 (? <i>Anthomyia</i>), <i>Coenosia</i>	97.43
<i>dubia</i> Bigot, 1885, <i>Limnophora</i>	97.28
<i>dubia</i> (Bigot), 1885 (<i>Yetodesia</i>), <i>Helina</i>	97.20
<i>dubia</i> Snyder, 1954, <i>Cyrtoneurina</i>	97.52
<i>dubitalis</i> Malloch, 1934, <i>Syllimnophora</i>	97.30
<i>Dyadimyia</i> Séguy, 1937	97.52



<i>ebenifemur</i> Snyder, 1957, <i>Neodexiopsis</i>	97.45
<i>echinogaster</i> (Stein), 1911 (<i>Mydaea</i>), <i>Helina</i>	97.20
<i>edwardsi</i> Malloch, 1934, <i>Lispe</i>	97.34
<i>effrons</i> (Giglio-Tos), 1893 (<i>Brachiophyra</i>), <i>Limnophora</i>	97.29
<i>elegans</i> Macquart, 1843, <i>Limnophora</i>	97.28
<i>elegans</i> (Stein), 1911 (<i>Limnophora</i>), <i>Lispoides</i>	97.32
<i>elegantula</i> new name, <i>Lispoides</i>	97.32
<i>eleodivora</i> Walton, 1918 (<i>Biomya</i>), <i>Tachinidae</i>	97.60
<i>elongata</i> Albuquerque, 1958, <i>Phaonia</i>	97.14
<i>emdeni</i> Hennig, 1955, <i>Schoenomyzina</i>	97.39
<i>emmesa</i> (Malloch), 1934 (<i>Coenosia</i>), <i>Neodexiopsis</i>	97.45
<i>enigmatica</i> Albuquerque, 1955, <i>Pachyceramyia</i>	97.24
<i>enormis</i> Albuquerque, 1956, <i>Coenosia</i>	97.42
<i>equator</i> Snyder, 1941, <i>Helina</i>	97.20
<i>equator</i> Snyder, 1958, <i>Neodexiopsis</i>	97.46
<i>equestris</i> Fabricius, 1775, <i>Musca</i>	97.60
<i>errata</i> Malloch, 1934, <i>Fannia</i>	97.6
<i>erratica</i> , error	97.6
<i>erythrogaster</i> Séguy, 1941, <i>Euryomma</i>	97.2
<i>erythroptis</i> (Bigot), 1888 (<i>Homalomyia</i>), <i>Fannia</i>	97.4
<i>erythroptis</i> (Schnabl), 1902 (<i>Coelomyia</i>), <i>Fannia</i>	97.4
<i>etesia</i> (Giglio-Tos), 1893 (<i>Spilogaster</i>), <i>Helina</i>	97.20
<i>Euphaonia</i> Malloch, 1922	97.16
<i>eurycephala</i> (Stein), 1911 (<i>Mydaea</i>), <i>Helina</i>	97.20
<i>Euryomma</i> Stein, 1899	97.2
<i>evittata</i> Malloch, 1934, <i>Schoenomyza</i>	97.39
<i>excisus</i> (Thomson), 1869 (<i>Coenosia</i>), <i>Atherigona</i>	97.25
<i>exilis</i> (Williston), 1896 (<i>Limnophora</i>), <i>Fannia</i>	97.4
<i>exul</i> Williston, 1896, <i>Limnophora</i>	97.28
<i>facilis</i> (Walker), 1858 (<i>Pyrellia</i>), <i>Morellia</i>	97.8
<i>fallax</i> (Stein), 1911 (<i>Mydaea</i>), <i>Helina</i>	97.20
<i>falsifica</i> (Dodge & Aitken), 1968 (<i>Philornis</i>), <i>Neomusca</i>	97.56
<i>Fannia</i> Robineau-Desvoidy, 1830	97.3
<i>Fanniinae</i>	97.2
<i>farri</i> Dodge, 1955, <i>Neomuscina</i>	97.51
<i>fasciculatus</i> (Malloch), 1934 (<i>Darwinomyia</i>), <i>Palpibracus</i>	97.17
<i>fasciventris</i> (Wulp), 1896 (<i>Mydaea</i>), <i>Neomusca</i>	97.56
<i>felsina</i> (Walker), 1849 (<i>Anthomyia</i>), <i>Neurotrixa</i>	97.48
<i>femoralis</i> (Robineau-Desvoidy), 1830 (<i>Palusia</i>), <i>Coenosia</i>	97.42,59
<i>femoralis</i> (Stein), 1898 (<i>Homalomyia</i>), <i>Fannia</i>	97.4
<i>femorals</i> (Wulp), 1896 (<i>Coenosia</i>), <i>Bithoracochaeta</i>	97.37
<i>femorata</i> (Loew), 1861 (<i>Homalomyia</i>), <i>Fannia</i>	97.6
<i>femorata</i> Stein, 1911, <i>Phaonia</i>	97.14
<i>femorata</i> (Stein), 1911 (<i>Limnophora</i>), <i>Syllimnophora</i>	97.30
<i>fenestrata</i> Bigot, 1888, <i>Schoenomyza</i>	97.40
<i>ferruginea</i> Malloch, 1928, <i>Mulfordia</i>	97.48
<i>flavicincta</i> (Stein), 1904 (<i>Homalomyia</i>), <i>Fannia</i>	97.4
<i>flavicornis</i> (Macquart), 1855 (<i>Aricla</i>), <i>Alluaudinella</i>	97.60
<i>flavicornis</i> (Macquart), 1848 (<i>Pyrellia</i>), <i>Morellia</i>	97.8
<i>flavicornis</i> (Malloch), 1918 (<i>Ariclella</i>), <i>Helina</i>	97.23
<i>flavicornis</i> (Malloch), 1928 (<i>Bdellolarynx</i>), <i>Neivamyia</i>	97.59
<i>flavicornis</i> Stein, 1911, <i>Fannia</i>	97.4

<i>flavicoxa</i> Malloch, 1934, <i>Bithoracochaeta</i>	97.36
<i>flavidicincta</i> (Stein), 1904 (<i>Aricia</i>), <i>Polietina</i>	97.55
<i>flavinervis</i> Thomson, 1869, <i>Musca</i>	97.7
<i>flavipalpis</i> Albuquerque, 1956, <i>Neodexiopsis</i>	97.46
<i>flavipalpis</i> Malloch, 1934, <i>Raymondomyia</i>	97.36
<i>flavipalpis</i> Snyder, 1957, <i>Syllimnophora</i>	97.30
<i>flavipalpis</i> Stein, 1911, <i>Fannia</i>	97.4
<i>flavipennis</i> Macquart, 1843 (<i>Ochromyia</i>), <i>Calliphoridae</i>	97.60
<i>flavipes</i> Stein, 1918, <i>Fannia</i>	97.4
<i>flavipes</i> (Williston), 1896 (<i>Coenosia</i>), <i>Neodexiopsis</i>	97.46
<i>flavithorax</i> (Stein), 1904 (<i>Aricia</i>), <i>Polietina</i>	97.55
<i>flora</i> (Bigot), 1878 (<i>Pyrellia</i>), <i>Morellia</i>	97.8
<i>fluminensis</i> Albuquerque, 1954, <i>Pseudoptilolepis</i>	97.49
<i>frontina</i> (Wulp), 1896 (<i>Clinopora</i>), <i>Cyrtoneurina</i>	97.53
<i>fulva</i> Snyder, 1949, <i>Neomuscina</i>	97.50
<i>fulvapoda</i> Snyder, 1940, <i>Helina</i>	97.20
<i>fulvapoda</i> Snyder, 1949, <i>Pseudoptilolepis</i>	97.49
<i>fulvescens</i> Robineau-Desvoidy, 1830, <i>Musca</i>	97.7
<i>fulvifrons</i> Snyder, 1949, <i>Neomuscina</i>	97.51
<i>fulvipes</i> (Bigot), 1888 (<i>Curtonevra</i>), <i>Cyrtoneurinae</i>	97.53
<i>fulvisquama</i> (Stein), 1904 (<i>Spilogaster</i>), <i>Helina</i>	97.20
<i>fulvitibia</i> Snyder, 1949, <i>Dasymorellia</i>	97.10
<i>fulvocalyptrata</i> Malloch, 1934, <i>Helina</i>	97.20
<i>fulvohumeralis</i> Malloch, 1922, <i>Euphaonia</i>	97.16
<i>fulvolateralis</i> Stein, 1911, <i>Psilocheaeta</i>	97.16
<i>fumicosta</i> (Dodge), 1968 (<i>Philornis</i>), <i>Neomusca</i>	97.56
<i>fumicosta</i> Snyder, 1941, <i>Mydaea</i>	97.25
<i>fumipennis</i> Albuquerque, 1953, <i>Agenamyia</i>	97.32
<i>fumipennis</i> Albuquerque, 1954, <i>Cholomyioides</i>	97.40
<i>fumipennis</i> (Wulp), 1896 (<i>Limnophora</i>), <i>Mydaea</i>	97.25
<i>fuscicosta</i> (Curran), 1934 (<i>Cyrtoneuropsis</i>), <i>Cyrtoneurina</i>	97.52
<i>fuscicosta</i> Malloch, 1934, <i>Schoenomyzina</i>	97.39
<i>fuscifacies</i> Malloch, 1934, <i>Schoenomyza</i>	97.40
<i>fuscinervis</i> , error	97.48
<i>fuscinevis</i> (Macquart), 1851 (<i>Limnophora</i>), <i>Neurotrixa</i>	97.48
<i>fuscipalpis</i> (Stein), 1911 (<i>Schoenomyza</i>), <i>Spathipheromyia</i>	97.38
<i>fuscisquama</i> Aldrich, 1928, <i>Hydrotaea</i>	97.12
<i>fuscisquama</i> Snyder, 1954, <i>Cyrtoneurina</i>	97.52
<i>fuscisquama</i> (Wulp), 1896 (<i>Phorbia</i>), <i>Phaonia</i>	97.14
<i>fuseocalyptrata</i> Macquart, 1855, <i>Hydrotaea</i>	97.12
<i>fuscomarginata</i> (Snyder), 1949 (<i>Airalips</i>), <i>Xenothoracochaeta</i> ...	97.49
<i>fusconotata</i> (Rondani), 1868 (<i>Myantha</i>), <i>Fannia</i>	97.4
<i>gallicola</i> Albuquerque, 1958, <i>Phaonia</i>	97.14
<i>garleppi</i> (Stein), 1911 (<i>Coenosia</i>), ? <i>Noelia</i>	97.41
<i>garrula</i> (Giglio-Tos), 1893 (<i>Leucomelina</i>), <i>Limnophora</i>	97.28
<i>gemina</i> (Wiedemann), 1830 (<i>Anthomyia</i>), <i>Cyrtoneurina</i>	97.52
<i>geminata</i> (Stein), 1904 (<i>Spilogaster</i>), <i>Cyrtoneurina</i>	97.52
<i>geniculata</i> (Bigot), 1885 (<i>Spilogaster</i>), <i>Neodexiopsis</i>	97.46
<i>geniculata</i> (Fallén), 1825 (<i>Musca</i>), <i>Coenosia</i>	97.42
<i>genupuncta</i> (Stein), 1904 (<i>Coenosia</i>), <i>Neodexiopsis</i>	97.46
<i>geniculata</i> Macquart, 1846, <i>Stomoxys</i>	97.58
<i>geniculata</i> (Macquart), 1851 (<i>Cordylura</i>), <i>Tetramerinx</i>	97.35




<i>gibba</i> Fabricius, 1805 (Musca), Curtonotidae	97.60
<i>gigantea</i> Albuquerque, 1956, Helina	97.20
<i>gilvitaris</i> Chillcott, 1961, Fannia	97.4
<i>glaucinis</i> (Dodge & Aitken), 1968 (Philornis), Neomusca	97.56
<i>gluta</i> Giglio-Tos, 1893, Cyrtoneurina	97.53
<i>golbachii</i> Snyder, 1957, Spiliogona	97.33
<i>golbachii</i> Snyder, 1957, Xenothoracochaeta	97.49
<i>gracilis</i> (Stein), 1911 (Limnophora), Lispoides	97.32
<i>gracilitarsis</i> (Stein), 1911 (Limnophora), Syllimnophora	97.30
<i>grajauensis</i> Albuquerque, 1957, Bigotomyia	97.18
<i>grandis</i> Malloch, 1912, Fannia	97.4
<i>grandis</i> (Schiner), 1868 (Spilogaster), Helina	97.21
<i>Graphomya</i> Robineau-Desvoidy, 1830	97.27
<i>Graphomya</i> , error	97.27
<i>grisea</i> (Fabricius), 1805 (Musca), Cyrtoneurina	97.53
<i>grisea</i> (Giglio-Tos), 1893 (Hyadesimyia), Synthesiomysia	97.12
<i>grisea</i> Malloch, 1934, Helina	97.22
<i>griseovirens</i> Malloch, 1928, Drepanocnemis	97.34
<i>guatemala</i> Snyder, 1951, Lispoides	97.32
<i>guttipennis</i> (Stein), 1904 (Schoenomyza), Spathipheromyia	97.38
<i>guttipennis</i> (Thomson), 1869 (Ochthiphia), Spathipheromyia ..	97.38
<i>Gymnodia</i> Robineau-Desvoidy, 1863	97.24
<i>Haematobia</i> Le Peletier & Serville, 1828	97.58
<i>Haroldopsis</i> Albuquerque, 1954	97.37
<i>haustellata</i> Stein, 1911, Rhyncholimnophora	97.32
<i>heydeni</i> , error	97.4
<i>Helina</i> Robineau-Desvoidy, 1830	97.19
<i>Hemichlora</i> Wulp, 1896	97.49
<i>Hemixanthomyia</i> Albuquerque, 1954	97.57
<i>heydeni</i> , error	97.4
<i>heydenii</i> (Wiedemann), 1830 (Anthomyia), Fannia	97.4
<i>hieroglyphica</i> (Wulp), 1896 (Clinopera), Cyrtoneurina	97.54
<i>hirticeps</i> Stein, 1911, Drepanocnemis	97.35
<i>hirticeps</i> (Stein), 1911 (Limnophora), Spilogona	97.33
<i>hirtifemur</i> (Stein), 1904 (Homalomyia), Fannia	97.5
<i>hirtitibia</i> (Stein), 1911 (Coenosia), Neodexiopsis	97.46
<i>hirtula</i> (Bigot), 1885 (Ophyra), Psilochaeta	97.15
<i>hispaniense</i> Stein, 1899, Euryomma	97.2
<i>holii</i> Malloch, 1923, Xenomorellia	97.10
<i>Homalomyia</i> Bouché, 1835	97.3
<i>houghi</i> Malloch, 1916, Hydrotaea	97.12
<i>humeralis</i> (Stein), 1918 (Pyrellia), Morellia	97.9
<i>hyalinipennis</i> (Czerny), 1903 (Homalomyia), Fannia	97.5
<i>Hydrotaea</i> Robineau-Desvoidy, 1830	97.12
<i>hydrotaeiformis</i> Snyder, 1958, Neodexiopsis	97.46
<i>Hydrotaeini</i>	97.11
<i>ignava</i> (Walker), 1858 (Aricia), Helina	97.22
<i>ignobilis</i> Stein, 1911, Coenosia	97.42
<i>immaculata</i> Stein, 1911, Coenosia	97.42
<i>immaculata</i> (Walker), 1837 (Anthomyia), Notoschoenomyza	97.39

97.97

<i>immaculiventris</i> (Malloch), 1923 (Melanochelia), <i>Spilogona</i>	97.33
<i>immunda</i> (Stein), 1911 (Mydaea), <i>Cyrtoneurina</i>	97.52
<i>inaequalis</i> Malloch, 1934, <i>Coenosia</i>	97.42
<i>inaequifrons</i> Malloch, 1934, <i>Lispoides</i>	97.33
<i>incaica</i> (Stein), 1911 (Mesochaeta), <i>Teleutochaeta</i>	97.31
<i>incisurata</i> (Zetterstedt), 1838 (Anthomyza), <i>Fannia</i>	97.5
<i>incognita</i> Snyder, 1954, <i>Cyrtoneurina</i>	97.53
<i>inconspicua</i> Malloch, 1934, <i>Syllimnophora</i>	97.30
<i>inculta</i> Stein, 1911, <i>Coenosia</i>	97.42
<i>incurva</i> Stein, 1911, <i>Coenosia</i>	97.42
<i>inducta</i> (Walker), 1852 (Anthomyia), <i>Fannia</i>	97.5
<i>inepta</i> (Stein), 1911 (Mydaea), <i>Helina</i>	97.21
<i>inermipennis</i> Albuquerque, 1954, <i>Fannia</i>	97.5
<i>inermis</i> (Stein), 1904 (Spilogaster), <i>Mydaea</i>	97.26
<i>inflexa</i> (Stein), 1918 (Cyrtoneurina), <i>Neomuscina</i>	97.50
<i>iniqua</i> Stein, 1911, <i>Coenosia</i>	97.42
<i>iniqua</i> (Stein), 1911 (Calliophrys), <i>Limnophora</i>	97.28
<i>insignis</i> (Stein), 1898 (Caricea), <i>Bithoracochaeta</i>	97.37
<i>insons</i> (Giglio-Tos), 1893 (Hyetodesia), <i>Mydaea</i>	97.25
<i>instabilis</i> Snyder, 1949, <i>Neomuscina</i>	97.50
<i>insularis</i> Hennig, 1957, <i>Lispoides</i>	97.33
<i>insularis</i> Malloch, 1934, <i>Spathipheromyia</i>	97.38
<i>insularis</i> (Williston), 1896 (Coenosia), <i>Cordiluroides</i>	97.44
<i>integra</i> (Stein), 1911 (Limnophora), <i>Syllimnophora</i>	97.30
<i>intensica</i> Curran, 1928, <i>Fannia</i>	97.5
<i>intermedia</i> (Fallén), 1825 (Musca), <i>Cocnosia</i>	97.42,59
<i>intoniclunis</i> Snyder, 1957, <i>Neodexiopsis</i>	97.46
<i>introducta</i> (Walker), 1852 (Anthomyia), <i>Fannia</i>	97.4
<i>inuber</i> Giglio-Tos, 1893, <i>Cyrtoneurina</i>	97.53
<i>inusitata</i> (Malloch), 1934 (Austrocoenosia), <i>Coenosia</i>	97.42
<i>iris</i> (Bigot), 1878 (Pyrellia), <i>Morellia</i>	97.9
<i>irritans</i> (Linnaeus), 1758 (Conops), <i>Haematobia</i>	97.58
<i>itaiensis</i> (Albuquerque), 1954 (Austrocoenosia), <i>Coenosia</i>	97.42
<i>itaiensis</i> Albuquerque, 1956, <i>Fannia</i>	97.5
 <i>kuscheli</i> Hennig, 1955, <i>Notoschoenomyza</i>	 97.39
 <i>laeta</i> Stein, 1911, <i>Limnophora</i>	 97.28
<i>laetifica</i> (Robineau-Desvoidy), 1830 (Mydina), <i>Helina</i>	97.21
<i>laevis</i> (Stein), 1898 (Homalomyia), <i>Fannia</i>	97.5
<i>laevis</i> (Stein), 1911 (Limnophora), <i>Lispoides</i>	97.33
<i>lancifer</i> (Malloch), 1934 (Darwinomyia), <i>Palpibracus</i>	97.17
<i>lanigera</i> (Stein), 1918 (Coenosia), <i>Neodexiopsis</i>	97.46
<i>Lasiomala</i> Enderlein, 1927	97.54
<i>laslosterna</i> Snyder, 1941, <i>Helina</i>	97.21
<i>lata</i> (Walker), 1849 (Eriphia), <i>Hydrotaea</i>	97.12,59
<i>latana</i> Snyder, 1949, <i>Lispe</i>	97.34
<i>lateralis</i> (Linnaeus), 1758 (Musca), <i>Fannia</i>	97.4
<i>latifrons</i> Malloch, 1932, <i>Neivamyia</i>	97.59
<i>latifrons</i> Malloch, 1934, <i>Syllimnophora</i>	97.31
<i>latifrons</i> Snyder, 1957, <i>Lispoides</i>	97.33
<i>latifrons</i> (Thomson), 1869 (Coenosia), <i>Neodexiopsis</i>	97.46




<i>latimaculata</i> (Albuquerque), 1956 (Austrocoenosia), Coenosia ...	97.42
<i>latimana</i> Malloch, 1934, Syllimnophora	97.31
<i>latimanoides</i> Snyder, 1957, Syllimnophora	97.31
<i>latinervis</i> (Stein), 1904, (Aricia), Phaonia	97.14
<i>latipalpis</i> (Stein), 1918 (Mydaea), Pseudoptilolepis	97.49
<i>latipennis</i> (Hough), 1900 (Muscina), Mydaea	97.26
<i>latipennis</i> (Stein), 1904 (Spilogaster), Helina	97.21
<i>latitibia</i> Albuquerque, 1957, Coenosia	97.42
<i>lativittata</i> Malloch, 1934, Schoenomyza	97.40
<i>latomensis</i> Snyder, 1957, Mydaea	97.25
<i>lentiginosa</i> Snyder, 1957, Phaonia	97.14
<i>lepida</i> (Wiedemann), 1817 (Anthomyia), Fannia	97.5
<i>leucocephala</i> (Wulp), 1896 (Mydaea), Helina	97.21
<i>Leucomelina</i> Macquart, 1851	97.28
<i>leucoprocta</i> (Wiedemann), 1830 (Anthomyia), Bithoracochaeta	97.36
<i>leucostoma</i> (Wiedemann), 1817 (Anthomyia), Ophyra	97.13
<i>leucotrichia</i> Albuquerque, 1956; Coenosia	97.42
<i>Levallonia</i> Albuquerque, 1958	97.37
<i>levis</i> Stein, 1911, Lispe	97.34
<i>limbata</i> Bigot, 1885, Limnophora	97.28
<i>limbata</i> (Wiedemann), 1830 (Anthomyia), Mydaea	97.26
<i>limbatinervis</i> , error	97.48
<i>limbatinevris</i> (Macquart), 1851 (Spilogaster), Neurotrixa	97.48
<i>limbinervis</i> Stein, 1918, Phaonia	97.14
<i>limbisquama</i> Stein, 1918, Cyrtoneurina	97.53
<i>Limnophora</i> Robineau-Desvoidy, 1830	97.28
<i>Limnophorinae</i>	97.28
<i>linea</i> (Wulp), 1896 (Muscina), Neomuscina	97.51
<i>lineata</i> (Stein), 1904 (Coenosia), Neodexiopsis	97.46
<i>lisarba</i> Snyder, 1949, Lispe	97.34
<i>Lispa</i> , error	97.33
<i>Lispe</i> Latreille, 1796	97.33
<i>Lispoides</i> Malloch, 1920	97.32
<i>lispomima</i> Hennig, 1955, Syllimnophora	97.31
<i>listrata</i> Albuquerque, 1954, Cordiluroides	97.44
<i>litorella</i> Fallén, 1823 (Ochtiphila), Schoenomyza	97.40
<i>longicorne</i> Stein, 1911, Euryomma	97.2
<i>longipalpis</i> Stein, 1911, Rhyncholimnophora	97.32
<i>longipede</i> Albuquerque, 1956, Coenosia	97.43
<i>longipes</i> (Stein), 1911 (Mydaea), ?Mulfordia	97.48
<i>longipila</i> Albuquerque, 1954, Fannia	97.5
<i>longipila</i> (Stein), 1918 (Mydaea), Helina	97.21
<i>longipilis</i> (Albuquerque), 1954 (Plumispina), Neodexiopsis	97.46
<i>longiseta</i> (Gaminara), 1930 (Rhynchocoenops), Coenosia	97.43
<i>longiseta</i> (Wulp), 1896 (Hyetodesia), Mydaca	97.25
<i>longispina</i> (Malloch), 1923, (Phyllogaster), Tetramerinx	97.35
<i>lopesi</i> Albuquerque, 1955, Lispoides	97.33
<i>lucorum</i> (Fallén), 1823 (Musca), Helina	97.21
<i>lunatisigna</i> Snyder, 1957, Neodexiopsis	97.46
<i>luteola</i> Albuquerque, 1956, Helina	97.21
<i>lutzi</i> Pinto & Fonseca, 1930, Neivamyia	97.59
<i>lynchii</i> (Wulp), 1883 (Limnophora), Helina	97.21
<i>Lyperosia</i> Rondani, 1856	97.58



<i>machadoi</i> Albuquerque, 1958, <i>Phaonia</i>	97.14
<i>macquarti</i> new name, <i>Coenosia</i>	97.43
<i>Macquartiopsis</i> Albuquerque, 1949	97.41
<i>macrocera</i> (Wulp), 1896 (<i>Coenosia</i>), <i>Neodexiopsis</i>	97.46
<i>macrops</i> Snyder, 1949, <i>Neomuscina</i>	97.50
<i>macrosoma</i> Wulp, 1896, <i>Charadrella</i>	97.57
<i>mactans</i> Fabricius, 1787, <i>Musca</i>	97.60
<i>maculata</i> (Scopoli), 1763 (<i>Musea</i>), <i>Graphomya</i>	97.27
<i>maculipennata</i> (Macquart), 1851 (<i>Pyrellia</i>), <i>Morellia</i>	97.9
<i>maculipennis</i> (Macquart), 1843 (<i>Spilogaster</i>), <i>Cyrtoneurina</i>	97.53
<i>maculipennis</i> (Macquart), 1846 (<i>Pyrellia</i>), <i>Morellia</i>	97.9
<i>maculipennis</i> Malloch, 1934, <i>Apsil</i>	97.35
<i>maculipennis</i> (Walker), 1852 (<i>Anthomyia</i>), <i>Cyrtoneurina</i>	97.54
<i>maculipennis</i> (Williston), 1896 (<i>Cyrtoneura</i>), <i>Cyrtoneurina</i>	97.53
<i>maculipennis</i> (Zetterstedt), 1845 (<i>Aricia</i>), <i>Helina</i>	97.21,59
<i>maculipes</i> (Stein), 1918 (<i>Mydaea</i>), <i>Helina</i>	97.21
<i>maculiventris</i> Malloch, 1929, <i>Apsil</i>	97.36
<i>maculosa</i> Snyder, 1951, <i>Cariocamyia</i>	97.58
<i>maculosa</i> Stein, 1911, ? <i>Phaonia</i>	97.14
<i>maculosa</i> (Walker), 1852 (<i>Anthomyia</i>), <i>Helina</i>	97.21
<i>magellani</i> Malloch, 1934, <i>Spathipheromyia</i>	97.38
<i>magnicornis</i> Snyder, 1958, <i>Neodexiopsis</i>	97.46
<i>major</i> Albuquerque, 1956, <i>Polietina</i>	97.55
<i>major</i> Degeer, 1776, <i>Musca</i>	97.7
<i>major</i> Malloch, 1934, <i>Brachygasterina</i>	97.16
<i>malacophaga</i> Souza Lopes, 1938, <i>Charadrella</i>	97.57
<i>maldonadoi</i> Snyder, 1957, <i>Neodexiopsis</i>	97.46
<i>Mallocharia</i> Curran, 1934	97.52
<i>mallochi</i> Albuquerque, 1954, <i>Graphomya</i>	97.27
<i>mallochi</i> (Curran), 1934 (<i>Darwinomyia</i>), <i>Palpibracus</i>	97.17
<i>mallochi</i> , new name, <i>Schoenomyza</i>	97.40
<i>marginalis</i> (Stein), 1904 (<i>Spilogaster</i>), <i>Mydaea</i>	97.26
<i>marginata</i> Albuquerque, 1954, <i>Hemixanthomyia</i>	97.57
<i>marginata</i> Snyder, 1940, <i>Helina</i>	97.20
<i>marginata</i> Stein, 1904, <i>Limnophora</i>	97.28
<i>marginata</i> Stein, 1918, <i>Phaonia</i>	97.14
<i>marginipennis</i> (Stein), 1904 (<i>Spilogaster</i>), <i>Helina</i>	97.21
<i>marginipennis</i> Stein, 1911, <i>Limnophora</i>	97.29
<i>mediana</i> Snyder, 1949, <i>Neomuscina</i>	97.50
<i>medinai</i> Snyder, 1957, <i>Neodexiopsis</i>	97.46
<i>mediocris</i> Stein, 1911, <i>Coenosia</i>	97.43
<i>meditabunda</i> (Fabricius), 1781 (<i>Musca</i>), <i>Myospila</i>	97.26
<i>megalopyga</i> Albuquerque, 1954, <i>Cordiluroides</i>	97.44
<i>mellina</i> (Stein), 1904 (<i>Aricia</i>), <i>Polietina</i>	97.55
<i>mellina</i> Stein, 1918, <i>Cyrtoneurina</i>	97.53
<i>mendax</i> (Stein), 1911 (<i>Mydaea</i>), <i>Myospila</i>	97.27
<i>mendozana</i> (Enderlein), 1935 (<i>Protostomoxys</i>), <i>Graphomya</i>	97.27
<i>meraca</i> (Wulp), 1896 (<i>Limnophora</i>), <i>Helina</i>	97.21
<i>merida</i> , error	97.25
<i>meridensis</i> (Macquart), 1846 (<i>Lucilia</i>), <i>Morellia</i>	97.9
<i>meridia</i> Snyder, 1941, <i>Mydaea</i>	97.25
<i>meridionalis</i> Townsend, 1892, <i>Graphomya</i>	97.27
<i>Mesazelia</i> Blanchard, 1942	97.3



<i>Mesembrinellopsis</i> Townsend, 1927	97.55
<i>Mesochaeta</i> Stein, 1911	97.31
<i>mesofulvata</i> (Albuquerque), 1959 (Austrocoenosia), Coenosia ...	97.43
<i>metallica</i> (Wiedemann), 1830 (Anthomyia), Ophyra	97.17
<i>mexala</i> Snyder, 1957, Bigotomyia	97.18
<i>mexicana</i> (Bigot), 1885 (Homalomyia), Fannia	97.4
<i>mexicana</i> Giglio-Tos, 1893, Graphomya	97.28
<i>mexicana</i> (Macquart), 1843 (Curtonevra), Neomuscina	97.50
<i>micans</i> Snyder, 1957, Neodexiopsis	97.46
<i>micheneri</i> Chillcott, 1961, Fannia	97.5
<i>microchaeta</i> (Malloch), 1934 (Coenosia), Neodexiopsis	97.46
<i>Microchylum</i> Macquart, 1851	97.28
<i>mima</i> (Townsend), 1927 (Mesembrinellopsis), Neomusca	97.56
<i>mimica</i> Snyder, 1954, Cyrtoneurina	97.53
<i>mimicola</i> (Dodge), 1968 (Philornis), Neomusca	97.56
<i>minor</i> Albuquerque, 1956, Polietina	97.55
<i>minuscula</i> Albuquerque, 1955, Phaonia	97.14
<i>minuscula</i> (Wulp), 1896 (Leucomelina), Limnophora	97.29
<i>minuscularis</i> Albuquerque, 1956, Coenosia	97.43
<i>minuta</i> (Bezzi), 1892 (Lyperosia), Haematobia	97.59
<i>minuta</i> (Stein), 1911 (Schoenomyza), Spathippheromyia	97.38
<i>mirata</i> Shannon & Del Ponte, 1927, Neurotrixa	97.48
<i>molesta</i> (Meinert), 1890 (Philornis), Neomusca	97.56
<i>monacha</i> (Schiner), 1868 (Spilogaster), Helina	97.21
<i>monochaeta</i> Snyder, 1957, Phaonia	97.15
<i>monstrata</i> (Wulp), 1896 (Clinopera), Cyrtoneurina	97.53
<i>montanhesa</i> Albuquerque, 1952, Xenomorellia	97.10
<i>Morellia</i> Robineau-Desvoidy, 1830	97.8
<i>morio</i> Fabricius, 1775, Stomoxys	97.60
<i>morosa</i> (Wulp), 1896 (Phorbia), Fannia	97.5
<i>mulcata</i> (Giglio-Tos), 1893 (Hyetodesia), Helina	97.21
<i>Mulfordia</i> Malloch, 1928	97.48
<i>multimaculata</i> , error	97.53
<i>multomaculata</i> (Stein), 1904 (Spilogaster), Cyrtoneurina	97.53
<i>Musca</i> Linnaeus, 1758	97.7
<i>Muscidae</i>	97.1
<i>Muscina</i> Robineau-Desvoidy, 1830	97.11
<i>Muscinae</i>	97.7
<i>Muscini</i>	97.7
<i>Mydaea</i> Robineau-Desvoidy, 1830	97.25
<i>Mydaeinae</i>	97.25
<i>Myiospila</i> , error	97.26
<i>Myospila</i> Rondani, 1856	97.26
<i>narona</i> (Walker), 1849 (Anthomyia), Limnophora	97.29
<i>nasoni</i> Stein, 1893, Lispe	97.34
<i>nearmipes</i> Snyder, 1954, Cyrtoneurina	97.52
<i>nebulosa</i> Fabricius, 1805, Stomoxys	97.58
<i>neglecta</i> (Townsend), 1939 (Neopyrellia), Neorypellia	97.10
<i>Neivamyia</i> Pinto & Fonseca, 1930	97.59
<i>neoaustralis</i> Snyder, 1957, Neodexiopsis	97.46
<i>neobiseriata</i> Snyder, 1957, Schoenomyza	97.40
<i>Neodexiopsis</i> Malloch, 1920	97.45



neoflavipes Snyder, 1957, Neodexiopsis	97.46
neomacrocera Snyder, 1957, Neodexiopsis	97.46
Ncomusca Malloch, 1921	97.55
Neomuscina Townsend, 1919	97.50
Neopyrellia Enderlein, 1935	97.8
Neopyrellia Townsend, 1939	97.10
Neorypellia, new name	97.10
neosimilis Snyder, 1949, Neomuscina	97.50
neosimplex Snyder, 1957, Helina	97.21
neotrita Snyder, 1954, Cyrtoneurina	97.53
neotropica Albuquerque, 1954, Cordiluroides	97.44
neotropica Curran, 1934, Morellia	97.9
neotropica Snyder, 1957, Azelia	97.12
neotropica (Snyder), 1951 (Helinella), Helina	97.22
nervicincta Stein, 1918, Phaonia	97.15
Neurotrixa Shannon & Del Ponte, 1926	97.48
nicholsoni Curran, 1939, Hydrotaea	97.12
nielsenii (Dodge), 1968 (Philornis), Neomusca	97.56
niger (Dodge & Aitken), 1968 (Philornis), Neomusca	97.56
nigerrima (Malloch), 1934 (Austrocoenosia), Coenosia	97.43
nigra Macquart, 1851, Stomoxys	97.58,59
nigra (Stein), 1911 (Schoenomyza), Spathipheromyia	97.38
nigrargentina (Albuquerque), 1954 (Heliographa), Syllimnophora	97.31
nigribasis (Stein), 1911 (Limnophora), Lispoldes	97.33
nigriceps (Bigot), 1888 (Curtonevra), Myospila	97.27
nigriceps Stein, 1911, Teleutochaeta	97.31
nigricornis Malloch, 1934, Bithoracochaeta	97.37
nigricosta Hough, 1900, Morellia	97.9
nigricosta Snyder, 1949, Neomuscina	97.50
nigrifemur Stein, 1911, Euryomma	97.2
nigrimana (Macquart), 1851 (Aricia), Helina	97.22
nigrimana basalis Malloch, 1934, Helina	97.22
nigrimana grisea Malloch, 1934, Helina	97.22
nigrina (Wiedemann), 1830 (Anthomyia), Mydaea	97.26
nigripes Stein, 1911, Tetramerinx	97.35
nigripoda Snyder, 1949, Pseudoptilolepis	97.49
nigripuncta (Stein), 1911 (Phaonia), Bigotomyia	97.18
nigrithorax Stein, 1911, Schoenomyza	97.40
nigritibia Snyder, 1949, Dasymorellia	97.10
nigriventris Albuquerque, 1954, Bigotomyia	97.18
nigriventris (Malloch), 1928 (Darwinomyia), Palpibracus	97.17
nigriventris Malloch, 1934, Syllimnophora	97.31
nigrocincta (Stein), 1918 (Phaonia), Bigotomyia	97.18
nigrohalterata (Stein), 1904 (Coenosia), Oxytonocera	97.35
nigromarginata (Stein), 1911 (Mydaea), Helina	97.22
nitida (Stein), 1918 (Pyrellia), Morellia	97.9
nitida (Wiedemann), 1830 (Musca), Morellia	97.9
nivaloides Albuquerque, 1956, Helina	97.22
nobilis Albuquerque, 1956, Helina	97.22
noctiluca Albuquerque, 1958, Phaonia	97.15
Noelia Albuquerque, 1957	97.41
noniesmaculata Gaminara, 1930, Coenosia	97.43
normata Bigot, 1885, Limnophora	97.29



notha Snyder, 1941, Helina	97.22
Notoschoenomyza Malloch, 1934	97.38
nubilicosta Malloch, 1923, Hydrotaea	97.12
nubivena Snyder, 1941, Mydaea	97.25
nudaplcura Snyder, 1949, Pseudoptilolepis	97.49
nudinervis (Stein), 1918 (Cyrtoneurina), Neomuscina	97.50
nudinervis fulva Snyder, 1949, Neomuscina	97.50
nudiseta Albuquerque, 1954, Arthurella	97.57
nudiseta (Wulp), 1883 (Cyrtoneura), Synthesiomyia	97.11
nudistigma Snyder, 1949, Ncomuscina	97.50
obscura (Shannon & Del Ponte), 1926 (Muscina), Myospila	97.27
obscura (Wulp), 1896 (Mydaea), Neomusca	97.56
obscurinervis (Stein), 1900 (Homalomyia), Fannia	97.5
obscuripennis (Czerny), 1903 (Homalomyia), Fannia	97.5
obscuripes (Bigot), 1888 (Pyrellia), Morellia	97.9
obsoleta (Brauer & Bergenstamm), 1891 (Phasiophana), Myospila.	97.27
obtusiloba (Malloch), 1934 (Coenosia), Neodexiopsis	97.47
ochricornis (Wiedemann), 1830 (Musca), Morellia	97.9
ochrifacies (Rondani), 1850 (Pyrellia), Morellia	97.8
oculata (Stein), 1911 (Coenosia), Neodexiopsis	97.47
Ophyra Robineau-Desvoidy, 1830	97.13
orba Stein, 1911, Mydaea	97.25
orbitalis (Stein), 1904 (Aricia), Polietina	97.55
orientalis Schiner, 1868, Atherigona	97.24
Orthellia Robineau-Desvoidy, 1863	97.9
oscillans (Wulp), 1896 (Coenosia), Neodexiopsis	97.47
ovata (Stein), 1898 (Coenosia), Neodexiopsis	97.47,59
ovativentris (Macquart), 1851 (Anthomyia), Gymnodia	97.24
Oxytonocera, error	97.35
Oxytonocera Stein, 1919	97.35
Pachyccramyia Albuquerque, 1955	97.24
pacifera (Giglio-Tos), 1893 (Coenosia), Bithoracochaeta	97.37
pallicornis Malloch, 1934, Schoenomyzina	97.39
pallidicornis (Bigot), 1888 (Curtonevra), Myospila	97.27
pallipes Stein, 1918, Cyrtoneurina	97.53
Palpibracus Rondani, 1864	97.16
pampasiana Bigot, 1888, Musca	97.7
pampeana (Shannon & Del Ponte), 1926 (Phyronota), Psilo- chaeta	97.16
pampiana, error	97.16
panamensis Chillcott, 1958, Euryomma	97.2
panamensis Dodge, 1965, Graphomya	97.28
pansa (Giglio-Tos), 1893 (Spilogaster), Mydaea	97.25
pantherina (Wiedemann), 1824 (Anthomyia), Tetramerinx	97.35
Paradexiopsis Albuquerque, 1955	97.45
paralis, error	97.50
paranaensis (Albuquerque), 1954 (Heliographa), Syllimnophora.	97.31
Parapyrellia Townsend, 1915	97.8
parasitica Séguy, 1933, Fannia	97.5
parilis (Giglio-Tos), 1893 (Cyrtoneurina), Neomuscina	97.50


<i>parsura</i> (Giglio-Tos), 1893 (<i>Hyetodesia</i>), <i>Helina</i>	97.22
<i>parvula</i> Albuquerque, 1958, <i>Neodexiopsis</i>	97.47
<i>parvula</i> (Wulp), 1896 (<i>Spilogaster</i>), <i>Helina</i>	97.22
<i>patagonica</i> Malloch, 1934, <i>Limnophora</i>	97.29
<i>pauciseta</i> (Stein), 1904 (<i>Limnophora</i>), <i>Syllimnophora</i>	97.31
<i>pauciseta</i> Stein, 1911, <i>Mydaea</i>	97.25
<i>paulistensis</i> Albuquerque, 1956, <i>Neodexiopsis</i>	97.47
<i>pectinata</i> Malloch, 1934, <i>Reynoldsia</i>	97.36
<i>pedella</i> (Wiedemann), 1830 (<i>Anthomya</i>), <i>Mydaea</i>	97.26
<i>pellea</i> Giglio-Tos, 1893, <i>Cyrtoneurina</i>	97.53
<i>penicillaris</i> (Stein), 1900 (<i>Homalomya</i>), <i>Fannia</i>	97.5
<i>peninsula</i> Snyder, 1958, <i>Neodexiopsis</i>	97.47
<i>pennata</i> Malloch, 1934, <i>Apsil</i>	97.36
<i>peregrinum</i> (Meigen), 1826 (<i>Anthomyia</i>), <i>Euryomma</i>	97.2
<i>perspicua</i> (Stein), 1911 (<i>Mydaea</i>), <i>Cyrtoneurina</i>	97.54
<i>Peruvia</i> Malloch, 1929	97.23
<i>peruviana</i> Malloch, 1929, <i>Phaonia</i>	97.15
<i>peruviana</i> Snyder, 1958, <i>Neodexiopsis</i>	97.47
<i>petrocchia</i> Shannon & Del Ponte, 1926, <i>Fannia</i>	97.5
<i>petrocchia</i> , error	97.5
<i>Phaonaltho</i> Albuquerque, 1957	97.14
<i>Phaonia</i> Robineau-Desvoidy, 1830	97.14
<i>Phaoniinac</i>	97.14
<i>Phaoniini</i>	97.14
<i>Phasiophana</i> Brauer & Bergenstamm, 1891	97.26
<i>Philornis</i> Meinert, 1890	97.55
<i>Phyllogaster</i> Stein, 1898	97.35
<i>Phyronota</i> Shannon & Del Ponte, 1926	97.15
<i>pica</i> (Macquart), 1851 (<i>Leucomelina</i>), <i>Limnophora</i>	97.29
<i>pici</i> (Macquart), 1854 (<i>Aricia</i>), <i>Neomusca</i>	97.56
<i>picta</i> Stein, 1911, <i>Coenosia</i>	97.43
<i>picta</i> (Stein), 1911 (<i>Schoenomyza</i>), <i>Spathipheromyia</i>	97.38
<i>pictipennis</i> (Bigot), 1875 (<i>Curtonevra</i>), <i>Neomuscina</i>	97.50
<i>pictipennis</i> (Bigot), 1878 (<i>Curtonevra</i>), <i>Neomuscina</i>	97.50
<i>pictipennis nudicrvis</i> (Stein), 1918 (<i>Cyrtoneurina</i>), <i>Neomuscina</i> ..	97.50
<i>piliceps</i> (Stein), 1911 (<i>Mydaea</i>), <i>Helina</i>	97.22
<i>piliseta</i> Stein, 1919, <i>Limnophora</i>	97.29
<i>Pilispina</i> Albuquerque, 1954	97.37
<i>pilitarsis</i> Stein, 1911, <i>Thaumasiochaeta</i>	97.31
<i>pilitibia</i> Albuquerque, 1954, <i>Pilispina</i>	97.37
<i>pilitibia</i> Stein, 1911, <i>Coenosia</i>	97.43
<i>pilosa</i> (Stein), 1904 (<i>Coenosia</i>), <i>Neodexiopsis</i>	97.47
<i>pilosus</i> (Macquart), 1851 (<i>Brachypalpus</i>), <i>Palpibracus</i>	97.17
<i>pinocheti</i> Brèthes, 1924, ? <i>Gimnomera</i>	97.60
<i>pipunculina</i> (Thomson), 1869 (<i>Coenosia</i>), <i>Bithoracochaeta</i>	97.37
<i>platense</i> , error	97.4
<i>platensis</i> (Brèthes), 1908 (<i>Homalomyia</i>), <i>Fannia</i>	97.4
<i>platensis</i> (Shannon & Del Ponte), 1926 (<i>Phyronota</i>), <i>Psilochaeta</i> ..	97.16
<i>platystoma</i> (Thomson), 1869 (<i>Anthomyia</i>), <i>Limnophora</i>	97.29
<i>plaumanni</i> Snyder, 1941, <i>Mydaea</i>	97.25
<i>plenruseta</i> Snyder, 1940, <i>Helina</i>	97.22
<i>plumata</i> Albuquerque, 1955, <i>Bithoracochaeta</i>	97.37



plumata (Stein), 1904 (Spilogaster), Xenothoracochaeta	97.49
plumbea Albuquerque, 1954, Coenosia	97.43
plumbifrons (Albuquerque), 1956 (Austrocoenosia), Coenosia	97.43
plumifera (Bigot), 1885 (Spilogaster), Pygophora	97.60
plumiseta Stein, 1911, Coenosia	97.43
plumiseta Stein, 1911, Limnophora	97.29
Plumispina Albuquerque, 1954	97.45
podexaureus (Enderlein), 1935 (Protostomoxys), Graphomya ..	97.28
Poecilophaonia Malloch, 1921	97.54
poeciloptera (Malloch), 1918 (Aricia), Helina	97.21
poeciloptera (Schiner), 1868 (Spilogaster), Helina	97.22
Pogonomyia Rondani, 1871	97.12
Polietina Schnabl & Dziedzicki, 1911	97.54
polychaeta (Stein), 1895 (Homalomyia), Fannia	97.5, 59
portensis (Shannon & Del Ponte), 1926 (Phyronota), Psilochaeta.	97.16
porteri Brèthes, 1919, Limnophora	97.34
praecipua (Walker), 1852 (Anthomyia), Helina	97.22
praedatoria Snyder, 1957, Bigotomyia	97.18
praenubila Snyder, 1954, Cyrtoneurina	97.53
praesuturalis (Stein), 1904 (Aricia), Bigotomyia	97.18
praetaseta Snyder, 1954, Neomuscina	97.51
precuta Snyder, 1958, Neodexiopsis	97.47
presuturalis, error	97.18
prima Malloch, 1924, Coenosopsia	97.13
prima (Malloch), 1921 (Steinella), Neurotrixa	97.48
prima Malloch, 1921, Xenothoracochaeta	97.49
principalis (Schiner), 1868 (Spilogaster), Muscina	97.11
priscipagus Snyder, 1958, Neodexiopsis	97.47
probata Walker, 1861 (Hylemyia), Tachinidae	97.60
procedens (Walker), 1861 (Aricia), Helina	97.22
procera Stein, 1911, Coenosia	97.43
Profannia Séguy, 1937	97.3
projecta (Malloch), 1934 (Austrocoenosia), Coenosia	97.43
prolatifrons Snyder, 1940, Helina	97.22
prominula (Thomson), 1869 (Anthomyia), Limnophora	97.29
Promusca Townsend, 1915	97.7
propinqua (Stein), 1911 (Limnophora), Lispoides	97.33
prostrata (Scopoli), 1763 (Musca), Fannia	97.5
protosetosa Snyder, 1954, Cyrtoneurina	97.53
Protostomoxys Enderlein, 1935	97.27
protrita Walker, 1861 (Anthomyia), Sarcophagidae	97.60
pruinosa (Macquart), 1846 (Aricia), Polletina	97.55
pruinosa Shannon & Del Ponte, 1926, Fannia	97.5
pruna (Shannon & Del Ponte), 1926 (Orthellia), Morellia	97.9
pseudoflavicincta Albuquerque, 1954, Fannia	97.5
Pseudomyiospila Vimmer, 1939	97.18
Pseudoptilolepis Snyder, 1949	97.49
Psilochaeta Stein, 1911	97.15
pteropteurialis Malloch, 1934, Reynoldsia	97.36
pterospila Stein, 1918, Phaonia	97.15
ptero stigma (Wulp), 1896 (Clinopera), Cyrtoneurina	97.53
ptero stila, error	97.15



<i>pubiceps</i> (Stein), 1911 (Limnophora), <i>Spilogona</i>	97.33
<i>pulicaria</i> (Wulp), 1897 (Schoenomyza), <i>Coenosia</i>	97.44
<i>pulvillata</i> (Stein), 1904 (Aricia), <i>Phaonia</i>	97.15,59
<i>pulvinata</i> (Grimshaw), 1901 (Acritochaeta), <i>Atherigona</i>	97.25
<i>punctinervis</i> (Stein), 1911 (Phaonia), <i>Bigotomyia</i>	97.18
<i>punctipennis</i> Albuquerque, 1954, <i>Fannia</i>	97.6
<i>punctiventris</i> Malloch, 193, <i>Fannia</i>	97.6
<i>punctulata</i> (Wulp), 1896 (Coenosia), <i>Neodexiopsis</i>	97.47
<i>pura</i> (Stein), 1911 (Limnophora), <i>Syllimnophora</i>	97.31
<i>pusilla</i> (Bigot), 1885 (Ophyra), <i>Fannia</i>	97.6
<i>pusilla</i> Macquart, 1848, <i>Musca</i>	97.7
<i>pusio</i> (Wiedemann), 1830 (Anthomyia), <i>Fannia</i>	97.6
<i>quadrisetosa</i> , error	97.27
<i>quadristigma</i> (Thomson), 1869 (Anthomyia), <i>Myospila</i>	97.27
<i>querula</i> (Dodge & Aitken), 1968 (Philornis), <i>Neomusca</i>	97.56
<i>quintivena</i> Snyder, 1957, <i>Neodexiopsis</i>	97.47
<i>rava</i> Snyder, 1957, <i>Neodexiopsis</i> ,	97.47
<i>Raymondomyia</i> Malloch, 1934	97.36
<i>recedens</i> (Stein), 1904 (Coenosia), <i>Neodexiopsis</i>	97.47
<i>refusa</i> (Giglio-Tos), 1893 (Spilogaster), <i>Helina</i>	97.22
<i>regobarrosi</i> Albuquerque, 1958, <i>Helina</i>	97.22
<i>Reinwardtia</i> Brauer & Bergenstamm, 1889	97.11
<i>rescita</i> (Walker), 1861 (Aricia), <i>Cyrtoneurina</i>	97.53
<i>rettenmeyeri</i> Chillcott, 1958, <i>Euryomma</i>	97.2
<i>rettenmeyeri</i> (Dodge), 1963 (Philornis), <i>Neomusca</i>	97.56
<i>rex</i> Curran, 1928, <i>Neodexiopsis</i>	97.47
<i>reynoldsi</i> (Malloch), 1934 (Darwinomyia), <i>Palpibracus</i>	97.17
<i>Reynoldsia</i> Mallocha, 1934	97.36
<i>Rhabdoptera</i> Stein, 1919	97.38
<i>Rhyncholimnophora</i> Stein, 1911	97.32
<i>robusta</i> (Stein), 1911 (Coenosia), ? <i>Reynoldsia</i>	97.36
<i>rotundiventris</i> Stein, 1911, <i>Coenosia</i>	97.43
<i>rubella</i> (Wulp), 1896 (Hyetodesia), <i>Polietina</i>	97.55
<i>rubripalpis</i> (Wulp), 1896 (Spilogaster), <i>Helina</i>	97.22
<i>rufifrons</i> Stein, 1911, <i>Euryomma</i>	97.2
<i>rufiguttata</i> (Macquart), 1851 (Aricia), <i>Xenothoracochaeta</i>	97.49
<i>rufipalpis</i> (Macquart), 1851 (Pyrellia), <i>Morellia</i>	97.9
<i>rufipalpis</i> (Macquart), 1855 (Pyrellia), <i>Morellia</i>	97.9
<i>rufipes</i> (Bigot), 1885 (Limnophora), <i>Bithoracochaeta</i>	97.37
<i>rufipes</i> (Macquart), 1851 (Choetura), <i>Coenosia</i>	97.43
<i>rufitibia</i> Stein, 1911, <i>Fannia</i>	97.6
<i>rufitibia</i> (Stein), 1919 (Coenosia), <i>Neodexiopsis</i>	97.47
<i>rufitibia</i> Stein, 1911, <i>Tetramerinx</i>	97.35
<i>rufitibialis</i> Macquart, 1843, <i>Lispe</i>	97.34
<i>rufiventris</i> Macquart, 1843, <i>Musca</i>	97.7
<i>rufiventris</i> Macquart, 1846, <i>Musca</i>	97.7
<i>rufoapicata</i> Malloch, 1934, <i>Helina</i>	97.23
<i>rufoapicata</i> Malloch, 1934, <i>Reynoldsia</i>	97.36
<i>rufoscutella</i> Dodge, 1955, <i>Neomuscina</i>	97.51




rufoscutella farri Dodge, 1955, Neomuscina	91.51
sabroskyi (Albuquerque), 1957 (Philornis), Neomusca	97.56
rustica (Albuquerque), 1956 (Austrocoenosia), Coenosia	97.43
sabroskyi Seago, 1954, Fannia	97.6
saeva (Wiedemann), 1830 (Anthomyia), Limnophora	97.29
sanespra Snyder, 1949, Neomuscina	97.51
sanguinis (Dodge & Aitken), 1968 (Philornis), Neomusca	97.56
santoamarensis Albuquerque, 1958, Phaonia	97.15
saphirina (Séguy), 1935 (Trichomorellia), Dasymorellia	97.10
sarcophagina Wulp, 1896, Morellia	97.9
Sarcopromusca Townsend, 1927	97.8
scabra (Giglio-Tos), 1893 (Spilogaster), Dendrophaonia	97.13
scalaris (Fabricius), 1794 (Musca), Fannia	97.6
scapulata (Bigot), 1878 (Pyrellia), Morellia	97.8
scatophaga Curran, 1938, Gymnodia	97.24
Scenetes Malloch, 1936	97.24
schadei Snyder, 1949, Neomuscina	97.51
schildi (Dodge), 1963 (Philornis), Neomusca	97.56
schnusei Stein, 1911, Fannia	97.6
Schoenomyza Haliday, 1833	97.39
Schoenomyzina Malloch, 1934	97.39
scordalus (Walker), 1861 (Pyrellia), Hemichlora	97.49
scutellaris (Fabricius), 1805 (Musca), Neomuscina	97.51
scutellata (Johnson), 1919 (Clinopera), Cyrtoneurina	97.54
scutellata Malloch, 1934, Reynoldsia	97.36
Scutellomusca Townsend, 1931	97.50
secunda Snyder, 1951, Mulfordia	97.48
seguyi (Garcia), 1952 (Philornis), Neomusca	97.56
semicinerea (Stein), 1911 (Limnophora), Spilogona	97.33
semiclausula (Schiner), 1868 (Aricia), Myospila	97.26
semiclausula, error	97.26
semimarginata (Stein), 1918 (Pyrellia), Morellia	97.9
sensifera Walker, 1861 (Musca), Tachinidae	97.60
separatus (Malloch), 1934 (Darwinomyia), Palpibracus	97.17
sera (Giglio-Tos), 1893 (Spilogaster), Helina	97.23
seriata (Stein), 1911 (Mydaea), Cyrtoneurina	97.54
serotina Wulp, 1896, Lispe	97.34
serrata Robineau-Desvoidy, 1830, Haematobia	97.58
setia (Walker), 1849 (Anthomyia), Ophyra	97.13
setinervis (Dodge), 1963 (Philornis), Neomusca	97.56
setinervis (Thomson), 1869 (Anthomyia), Cyrtoneurina	97.53
setipuncta Snyder, 1957, Neodexiopsis	97.47
setitibia (Stein), 1911 (Limnophora), Syllimnophora	97.31
setiventris Stein, 1911, Coenosia	97.43
setosa (Bigot), 1885 (Ophyra), Fannia	97.6
setuligera Stein, 1911, Lispe	97.34
sevens, error	97.37
sexpunctata (Wulp), 1883 (Spilogaster), Mydaea	97.25
signatipennis (Wulp), 1896 (Spilogaster), Helina	97.23
sima Snyder, 1957, Neodexiopsis	97.47
similata Albuquerque, 1957, Bigotomyia	97.18
similata Snyder, 1949, Neomuscina	97.51



<i>similis</i> (Malloch), 1934 (Darwinomyia), <i>Palpibracus</i>	97.17
<i>simplex</i> Albuquerque, 1958, <i>Phaonia</i>	97.15
<i>simplex</i> Malloch, 1934, <i>Helina</i>	97.23
<i>singularis</i> (Stein), 1911 (Mydaea), <i>Souzalopesmyia</i>	97.24
<i>Siphona</i> Meigen, 1803, of authors	97.58
<i>Smithomyia</i> Malloch, 1921	97.54
<i>socia</i> (Wulp), 1896 (Limnophora), <i>Helina</i>	97.23
<i>sociabilis</i> Blanchard, 1937, <i>Bithoracochaeta</i>	97.37
<i>solitaria</i> Albuquerque, 1958, <i>Ophyra</i>	97.13
<i>solitaria</i> Shannon & Del Ponte, 1926, <i>Fannia</i>	97.5
<i>Souzalopesmyia</i> Albuquerque, 1951	97.23
<i>sp.</i> , <i>Ophyra</i>	97.13
<i>sparsiplumata</i> (Stein), 1918 (Mydaea), <i>Neomuscina</i>	97.51
<i>Spathipheromyia</i> , error	97.38
<i>Spathipheromyia</i> Bigot, 1884	97.38
<i>spatulata</i> Malloch, 1934, <i>Apsil</i>	97.36
<i>specialis</i> (Walker), 1861 (Pyrellia), <i>Morellia</i>	97.9
<i>spermophila</i> (Townsend), 1895 (Mydaea), <i>Neomusca</i>	97.56
<i>spicatus</i> (Malloch), 1934 (Darwinomyia), <i>Palpibracus</i>	97.17
<i>Spilogona</i> Schnabl, 1911	97.33
<i>spiloptera</i> (Wiedemann), 1830 (Anthomyia), <i>Cyrtoneurina</i>	97.54
<i>Spilopteromyia</i> Malloch, 1921	97.50
<i>spiniocosta</i> (Sten), 1904 (Schoenomyza), <i>Notoschoenomyza</i>	97.39
<i>spinifera</i> (Wulp), 1883 (Dasyphora), <i>Dasymorellia</i>	97.10
<i>spinuligera</i> (Stein), 1911 (Limnophora), <i>Syllimnophora</i>	97.31
<i>spretata</i> Malloch, 1921, <i>Limnophora</i>	97.29
<i>stabilis</i> (Stein), 1911 (Mydaea), <i>Neomuscina</i>	97.51
<i>stabulans</i> (Fallén), 1817 (Musca), <i>Muscina</i>	97.11
<i>steinella</i> Albuquerque, 1953, <i>Fannia</i>	97.6
<i>Steinella</i> Malloch, 1921	97.48
<i>steini</i> (Enderlein), 1927 (Chaetypoplcura), <i>Polietina</i>	97.55
<i>steini</i> Snyder, 1954, <i>Cyrtoneurina</i>	97.54
<i>stcini</i> , new name, <i>Neomusca</i>	97.57
<i>stellata</i> Bigot, 1884, <i>Spathipheromyia</i>	97.38
<i>stigmatica</i> Malloch, 1928, <i>Chaetagenia</i>	97.49
<i>stipata</i> (Walker), 1852 (Musca), <i>Graphomya</i>	97.28
<i>Stomopogon</i> Malloch, 1930	97.41
<i>stomoxidea</i> Robineau-Desvoidy, 1830, <i>Musca</i>	97.7
<i>stomoxides</i> , error	97.7
<i>Stomoxys</i> Geoffroy, 1762	97.58
<i>strenua</i> Stein, 1911, <i>Coenosia</i>	97.43
<i>striatipennis</i> (Stein), 1911 (Schoenomyza), <i>Rhabdoptera</i>	97.38
<i>strigata</i> (Giglio-Tos), 1893 (Leucomelina), <i>Limnophora</i>	97.29
<i>stuebeli</i> Röder, 1886, <i>Hydrotaca</i>	97.12
<i>sauvis</i> (Stein), 1911 (Limnophora), <i>Syllimnophora</i>	97.31
<i>subtilis</i> Albuquerque, 1951, <i>Palpibracus</i>	97.17
<i>sugillatrix</i> Robineau-Desvoidy, 1830, <i>Stomoxys</i>	97.58
<i>sulfuriceps</i> Malloch, 1934, <i>Notoschoenomyza</i>	97.39
<i>supera</i> (Walker), 1853 (Anthomyia), <i>Helina</i>	97.60
<i>surrepens</i> (Walker), 1861 (Lucilia), <i>Morellia</i>	97.8



<i>susficax</i> (Walker), 1861 (Pyrellia), Morellia	97.8
Syllimnophora Speiser, 1923	97.30
Synthesiomya Brauer & Bergenstamm, 1893	97.11
tachinina Brauer & Bergenstamm, 1889, Reinwardtia	97.11
tachinoides Albuquerque, 1958, Phaonia	97.15
tarsalis Malloch, 1934, Schoenomyza	97.40
tarsalis (Stein), 1918 (Mydaea), Helina	97.23
tarsata (Snyder), 1957 (Limosia), Coenosia	97.43
Teleutochaeta Stein, 1911	97.31
<i>tentaculata</i> (Degeer), 1776 (Musca), Lispe	97.34
tenuicornis (Wulp), 1896 (Coenosia), Neodexiopsis	97.47
tenuior (Walker), 1852 (Anthomyia), Coenosia	97.44
tenuipennis Malloch, 1934, Syllimnophora	97.31
tesselata (Stein), 1904 (Spilogaster), Helina	97.23
<i>Tetrachaeta</i> Stein, 1898	97.35
tetragona Gaminara, 1930, Phaonia	97.15
Tetramerinx Berg, 1898	97.35
<i>tetrastigma</i> , error	97.27
<i>texana</i> (Hough), 1899 (Muscina), Neomuscina	97.51
texensis Malloch, 1923, Phaonia	97.15
Thaumasiochaeta Stein, 1911	97.31
<i>tibialis</i> (Stein), 1898 (Coenosia), Neodexiopsis	97.47
tibialis (Walker), 1837 (Musca), Muscina	97.11
tinctifacies (Albuquerque), 1958 (Austrocoenosia), Coenosia	97.43
tinctinervis (Stein), 1918 (Mydaea), Neomuscina	97.51
tinctipenis Albuquerque, 1956, Coenosia	97.44
<i>torquans</i> (Nielsen), 1913 (Mydaea), Neomusca	97.57
townsendi Snyder, 1941, Helina	97.23
transporta Snyder, 1949, Neomuscina	97.51
travassosi Souza Lopes & Mangabeira, 1938, Neivamyia	97.59
triangularis Malloch, 1934, Schoenomyzina	97.39
<i>Trichomorellia</i> Stein, 1919	97.26
trichophthalma Albuquerque, 1959, Coenosia	97.44
trichopoda (Blanchard), 1942 (Mesazelia), Fannia	97.6
trichops Malloch, 1923, Dasymorellia	97.10
trichops (Stein), 1918 (Mydaea), Helina	97.23
trichops (Stein), 1911 (Limnophora), Spilogona	97.33
tricolor (Walker), 1852 (Anthomyia), Helina	97.23
<i>tricops</i> , error	97.10
trigona Shannon & Del Ponte, 1926, Phaonia	97.15
trigonota (Wulp), 1896 (Spilogaster), Phaonia	97.15
<i>trilineata</i> Stein, 1990, Atherigona	97.25
trimaculata (Stein), 1898 (Homalomyia), Fannia	97.6
trinitensis (Dodge & Aitken), 1968 (Philornis), Neomusca	97.57
<i>Trinodites</i> , error	97.19
triplex (Stein), 1911 (Limnophora), Lispoides	97.33
tripunctata (Wulp), 1896 (Muscina), Neomuscina	97.51
triseta Curran, 1931, Mydaea	97.25
triseta Snyder, 1949, Neomuscina	97.51
trispila (Bigot), 1885 (Spilogaster), Bigotomyia	97.18
trita (Stein), 1911 (Mydaea), Cyrtoneurina	97.54




<i>Tritonidis</i> Séguy, 1937	97.19
<i>trivittatus</i> (Malloch), 1934 (Darwinomyia), Palpibracus	97.17
<i>trochanterata</i> Malloch, 1934, Reynoldsia	97.36
<i>tropicalis</i> Malloch, 1934, Graphomya	97.27
<i>truncata</i> (Stein), 1911 (Coenosia), Neodexiopsis	97.47
<i>tucumanensis</i> Albuquerque, 1957, Fannia	97.6
<i>tumida</i> Stein, 1911, Coenosia	97.44
<i>tumidifemur</i> Stein, 1911, Fannia	97.6
<i>uber</i> Giglio-Tos, 1893, Cyrtoneurina	97.54
<i>uliginosa</i> Fallén, 1825, Lispe	97.34,59
<i>umanani</i> (García), 1952 (Philornis), Neomusca	97.57
<i>umbrosa</i> (Wulp), 1896 (Limnophora), Helina	97.23
<i>unicolor</i> (Stein), 1911 (Schoenomyza), Schoenomyzina	97.39
<i>unicolor</i> (Walker), 1852 (Anthomyia), Helina	97.22
<i>uniformis</i> Malloch, 1934, Coenosia	97.44
<i>uniseta</i> Malloch, 1934, LISPoides	97.33
<i>univittata</i> (Dodge), 1968 (Philornis), Neomusca	97.57
<i>univittata</i> Malloch, 1934, Schoenomyza	97.40
<i>univittatus</i> (Bigot), 1857 (Pegomyia), Palpibracus	97.17
<i>univittatus</i> (Malloch), 1922 (Darwinomyia), Palpibracus	97.17
<i>uspallata</i> Snyder, 1957, Neodexiopsis	97.48
<i>uens</i> (Giglio-Tos), 1894 (Coenosia), Bithoracochaeta	97.37
<i>variceps</i> Malloch, 1934, Syllimnophora	97.31
<i>varicolor</i> (Hough), 1900 (Muscina), Cyrtoneurina	97.54
<i>varicornis</i> (Coquillett), 1900 (Coenosia), Bithoracochaeta	97.37
<i>variegata</i> (Stein), 1911 (Mesochaeta), Teleutochaeta	97.31
<i>vecta</i> (Giglio-Tos), 1893 (Cyrtoneurina), Neomuscina	97.51
<i>veneris</i> (Bigot), 1888 (Helomyza), Palpibracus	97.17
<i>venicurva</i> (Stein), 1904 (Aricia), Bigotomyia	97.19
<i>veniseta</i> Dodge, 1967, Helina	97.23
<i>veniseta</i> (Stein), 1904 (Spilogaster), Cyrtoneurina	97.54
<i>vespidicola</i> (Dodge), 1968 (Philornis), Neomusca	97.57
<i>vibrissata</i> (Stein), 1918 (Mydaea), Cariocamylia	97.58
<i>vicaria</i> (Walker), 1852 (Anthomyia), Limnophora	97.29
<i>vicina</i> Macquart, 1851, Musca	97.7
<i>vicina</i> (Macquart), 1843 (Curtonevra), Muscina	97.11
<i>vicina</i> Robineau-Desvoidy, 1830, Limnophora	97.29
<i>vierecki</i> Snyder, 1911, Helina	97.23
<i>vilis</i> Stein, 1911, Lispe	97.34
<i>villosa</i> Aldrich, 1928, Ophyra	97.13
<i>villosa</i> Stein, 1904, Hydrotaea	97.12
<i>viola</i> Malloch, 1934, Helina	97.23
<i>violacea</i> (Fabricius), 1805 (Musca), Morellia	97.9
<i>violacea</i> (Robineau-Desvoidy), 1830 (Pyrellia), Morellia	97.8
<i>violaceiventris</i> Macquart, 1851, Brachygasterina	97.16
<i>violescens</i> Dodge, 1967, Helina	97.23
<i>virescens</i> Macquart, 1843, Ophyra	97.13
<i>virgata</i> (Wiedemann), 1830 (Anthomyia), Limnophora	97.34
<i>vitilis</i> (Giglio-Tos), 1894 (Coenosia), Neodexiopsis	97.48
<i>vittata</i> (Macquart), 1851 (Microchylum), Limnophora	97.29



<i>vittata</i> Malloch, 1912, <i>Fannia</i>	97.6
<i>vittifera</i> (Stein), 1904 (<i>Coenosia</i>), <i>Cordiluroides</i>	97.44
<i>vittigera</i> (Bigot), 1888 (<i>Curtonevra</i>), <i>Hemichlora</i>	97.44
<i>vittithorax</i> Stein, 1911, <i>Coenosia</i>	97.44
<i>vittiventris</i> (Albuquerque), 1955 (<i>Paradexiopsis</i>), <i>Neodexiopsis</i> ..	97.48
<i>vittiventris</i> Malloch, 1934, <i>Syllimnophora</i>	97.31
<i>willinki</i> Snyder, 1957, <i>Schoenomyza</i>	97.40
<i>willistoni</i> Snyder, 1958, <i>Neodexiopsis</i>	97.48
<i>wulpi</i> Snyder, 1954, <i>Cyrtoneurina</i>	97.54
<i>wulpi</i> , new name, <i>Coenosia</i>	97.44
<i>xanthocera</i> Albuquerque, 1954, <i>Fannia</i>	97.6
<i>xanthogaster</i> Albuquerque, 1958, <i>Levallonia</i>	97.37
<i>xanthomelas</i> Wiedemann, 1824, <i>Musca</i>	97.7,59
<i>xanthopoda</i> (Albuquerque), 1956 (<i>Austrocoenosia</i>), <i>Coenosia</i>	97.44
<i>xena</i> Malloch, 1934, <i>Helina</i>	97.23
<i>Xenocoenosia</i> Malloch, 1920	97.45
<i>Xenomorellia</i> Malloch, 1923	97.10
<i>Xenothoracochaeta</i> Malloch, 1921	97.48
<i>yenhedii</i> Albuquerque, 1957, <i>Fannia</i>	97.6
<i>zebrina</i> Bigot, 1885, <i>Limnophora</i>	97.29
<i>zeteki</i> (Dodge), 1963 (<i>Philornis</i>), <i>Neomusca</i>	97.57
<i>zosteris</i> (Shannon & Del Ponte), 1926 (<i>Muscina</i>), <i>Neomuscina</i>	97.51





SciELO₉

DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

98

FAMILY GASTEROPHILIDAE

J. H. GUIMARÃES

*Departamento de Zoologia, Secretaria da Agricultura
São Paulo, Brasil*

The Gasterophilidae, commonly called horse bot-flies, are myasis-producing Diptera. Their larvae develop in the digestive tract of equids, rhinoceroses and elephants. In the New World only the genus *Gasterophilus* occurs. The occurrence and distribution of the Neotropical species of that genus are not well known. The papers of Lahille (1911), Ihering (1929), Silva Junior & Proença (1934) and Vogelsang (1956) must be consulted on this subject. Keys to adults and larvae were provided by James (1948). For biology, anatomy and physiology, see Dinulescu, (1932) and for taxonomy Zumpt (1957) and Zumpt & Paterson (1953). The genitalia of the most common species were illustrated by Patton (1937).

Subfamily GASTEROPHILINAE

Genus GASTEROPHILUS Leach

GASTEROPHILUS Leach, 1817:2 (Leach in Brewster, 1817:162). Type-species, *Oestrus equi* Clark (Curtis, 1826: pl. 146) = *intestinalis* (De Geer).

Gastrophilus, emend.

Gastrus, emend.

Enteromyza Rondani, 1857:20 (unjustified change of name for *Gasterophilus* Leach). Type-species, *Oestrus equi* Clark (aut.) = *intestinalis* (De Geer).

Rhinogastrophilus Townsend, 1918:152. Type-species, *Oestrus nasalis* Linnaeus (orig. des.).

Haemorrhoeostrus Townsend, 1934:406. Type-species, *Oestrus haemorrhoidalis* Linnaeus (orig. des.).

Enteromyia Enderlein, 1934:425. Type-species, *Oestrus haemorrhoidalis* Linnaeus (orig. des.).

HAEMORRHOIDALIS (Linnaeus), 1758:584 (*Oestrus*). Type-locality: "Europe". Distr. — Almost cosmopolitan. In the Neotropical Region was recorded from Mexico, Venezuela, and Argentina.

98.1

DEPARTAMENTO DE ZOOLOGIA
BIBLIOTECA

- INTESTINALIS (De Geer), 1776:292 (*Oestrus*). Type-locality: "Europe".
Distr. — Almost cosmopolitan. In the Neotropical Region was
recorded from Jamaica, Venezuela, Brazil, and Argentina.
equi Clark, 1797:326 (*Oestrus*). Type-locality: "Europe".
- NASALIS (Linnaeus), 1758:584. (*Oestrus*). Type-locality: "Europe". Distr.
— Cosmopolitan. This is the commonest species in the Neotro-
pical Region.
veterinus Clark, 1797:328 (*Oestrus*). Type-locality: "Europe".
subjacens Walker, 1849:687 (*Gastrus*). Type-locality: Canada, Nova
Scotia.

LITERATURE CITED

CLARK, B.

1797. Observations on the genus *Oestrus*. *Trans. Linn. Soc. London*
3:289-329, 1 pl.

CURTIS, J.

1826. *British Entomology. Being illustrations and descriptions of
the genera of insects found in Great Britain and Ireland* 3:
pls. 99-146.

DE GEER, C.

1776. *Mémoires pour servir à l'histoire des Insectes* 6:523 pp., 30
pls. Stockholm.

DINULESCU, G.

1932. Recherches sur la biologie des Gastrophiles. Anatomie, phy-
siologie, cycle évolutif. *Ann. Sci. Nat.* 15 (1): 1-183, 8 pls.

ENDERLEIN, G.

1934. *Dipterologica I. Sitzber. Ges. Naturf. Freunde*, Berlin (1933):
416-429.

IHERING, R. VON

1929. Os Oestrideos importados, seu papel como parasitas e em
particular os *Gasterophilus* no Brasil. *Bol. Agric., São Paulo*
30:863-883.

JAMES, M. T.

- (1948) The flies that cause myiasis in man. *U.S. Dept. Agric. Misc.
Publ.* (1947) 631:1-157, 98 figs.

LAHILLE, F.

1911. Los Gastrófilos en la Republica Argentina. *Bol. Minist
Agric. Rep. Argentina* 13(12):836-856, 8 figs., col. pl.

LEACH, W. E.

1817. *On the genera and species of eproboscideous insects*, 20 pp.,
3 pls. Edinburgh.
1817. In D. Brewster, ed.: *The Edinburgh Encyclopaedia* 12:1-376,
pls. Edinburgh.



LINNAEUS, C.

1758. *Systema naturae per regna tria naturae*. Ed. 10. Vol. 1:824 pp. Holmiae.

PATTON, W. S.

1937. Studies on the higher Diptera of medical and veterinary importance. Illustrations of the terminalia of some species of *Gasterophilus*. *Ann. Trop. Med. Parasit.* 31(1):351-359, figs. 1-7.

RONDANI C.

1857. *Dipterologiae Italicae Prodrumus 2: Species Italicae ordinis dipteriorum in genera characteribus definita, ordinatim collectae, methodo analitica, distinctae, et novis vel minus cognitis descriptis*. Pars prima: Oestridae, Syrphidae [sic], Conopidae, 264 pp., 1 fig. Parmae.

SILVA JUNIOR, J. O. & M. C. PROENÇA

1934. Contribuição ao estudo da Gasterophilose e da espécie *Gasterophilus nasalis* (L. 1758) no Brasil. (Com a descrição da larva de 2ª fase e primeira observação no Brasil do inseto adulto). *Bol. Vet. do Exército*, Rio de Janeiro 1(1):163-185.

TOWNSEND, C. H. T.

1918. New muscoid genera, species and synonymy (Diptera). *Ins. Insc. Menstr.* 6:151-156.

1934. The Neotropical oestromuscoid flies. *Rev. Entom.*, Rio de Janeiro 4:201-212, 390-406.

VOGELSANG, E. G.

1956. Los *Gasterophilus* de los equinos en Venezuela. *Rev. Med. Vet. Parasit.*, Caracas 15(1-4):7-11, 1 pl.

WALKER, F.

1849. *List of the specimens of dipterous insects in the collection of the British Museum* 3:485-687, London.

ZUMPT, F.


1957. Some remarks on the classification of the Oestridae s. lat. (Diptera). *Jour. Ent. Soc. South Africa* 20:154-161, 1 pl.

ZUMPT, F. & H. E. PATERSON

1953. Studies on the family Gasterophilidae, with keys to the adult and maggots. *Jour. Ent. Soc. South Africa* 16:59-72, 17 figs.

INDEX

<i>Enteromyia</i> Enderlein, 1934	98.1
<i>Enteromyza</i> Rondani, 1857	98.1
<i>equi</i> (Clark), 1797 (Oestrus), <i>Gasterophilus</i>	98.2
<i>Gasterophilinae</i>	98.1
<i>Gasterophilus</i> Leach, 1817	98.1
<i>Gastrophilus</i> , emend.	98.1



<i>Gastrus</i> , emend.	98.1
<i>Haemorrhoestrus</i> Townsend, 1934	98.1
<i>haemorrhoidalis</i> (Linnaeus), 1758 (<i>Oestrus</i>), <i>Gasterophilus</i>	98.1
<i>intestinalis</i> (De Geer), 1776 (<i>Oestrus</i>), <i>Gasterophilus</i>	98.2
<i>nasalis</i> (Linnaeus), 1758 (<i>Oestrus</i>), <i>Gasterophilus</i>	98.2
<i>Rhinogasterophilus</i> Townsend, 1918	98.1
<i>subjacens</i> (Walker), 1849 (<i>Gastrus</i>), <i>Gasterophilus</i>	98.2
<i>veterinus</i> (Clark), 1797 (<i>Oestrus</i>), <i>Gasterophilus</i>	98.2



DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

99

FAMILY HIPPOBOSCIDAE

LINDOLPHO R. GUIMARÃES

*Departamento de Zoologia, Secretaria da Agricultura
São Paulo, Brazil*

These are blood-sucking flies, living as ectoparasites on birds and certain groups of mammals. Practically little is known about their biology. All species are larviparous, the larva pupating immediately after it is laid. Schhurmans-Stekhoven *et al.* (1957) published an account of the biology of *Pseudolynchia canariensis* (Macquart). Bequaert (1954-1957), revised the entire family, and gave important data on morphology, anatomy, hosts, and distribution. Maa (1963) gives a checklist of the family for the entire world, and his paper is especially relevant in what he examined several types, clearing the situation of many genera and species.

Subfamily ORNITHOICINAE

Genus ORNITHOICA Rondani

- ORNITHOICA Rondani, 1878:159. Type-species, *beccariina* Rondani (mon.) = *confluenta* (Say).
- CONFLUENTA (Say), 1823:103 (1859:88) (*Ornithomyia*). Type-locality: "U.S.A." Distr. — U.S.A., Bahama Islands, West Indies, Venezuela, Brazil; Old World.
- confluens* Wiedemann, 1830:611 (*Ornithomyia*), emend.
- beccariina* Rondani, 1878:160, unnumbered text-figs. Type-locality: "Amboina".
- VICINA (Walker), 1849:1144 (*Ornithomyia*). Type-locality: "Jamaica". Distr. — widespread in the Americas; established in the Hawaiian Islands.
- peroneura* Speiser, 1902c:91 (as *confluenta* var.). Type-locality: Hawaiian Islands.
- promiscua* Ferris & Cole, 1922:203, figs. 19, 20a-b. Type-locality: U.S.A., California, Lake Co., Castle Hot Springs.
- melaleuca* Bau, 1922:276, 278. Type-locality: Cuba, Santa Clara, Estación Meyer.
- confluens* authors, not Say.
- confluenta* authors, not Say.

Subfamily ORNITHOMYIINAE

Genus ORNITHOMYIA Latreille

ORNITHOMYIA Latreille, 1802:466 (as *Ornithomya*). Type-species, *Hippobosca avicularia* Linnaeus (mon.).

Subgenus ORNITHOMYIA Latreille

FRINGILLINA Curtis, 1836: pl. 585, colour-fig., and figs. 2-4, 8. Type-locality: "England". Distr. — Holarctic region; Mexico.

pallida Say, 1823:103 (1859:87) (preocc. Latreille, 1812). Type-locality: "U.S.A."

variegata Bigot, 1885:245. Type-locality: "New Zealand".

chloropus Bergroth, 1901:146. Type-locality: "Finland".

anchincuria Speiser, 1905:348 (nom. nov. for *pallida* Say). Type-locality: "U.S.A."

avicularia American authors, not Linnacus.

PARVA Macquart, 1843:436 (sep., p. 279). Type-locality: "Brazil or Chile".

Distr. — Chile, Argentina, Tristan da Cunha Island, Gough Island.

remota Walker, 1849:1144. Type-locality: Tristan da Cunha Island.

?synallaxidis Lynch Arribálzaga, 1881:90. Type-locality: Argentina, Rio Negro.

chilensis Reed, 1904:152 (preocc. Guérin-Ménéville, 1835). Type-locality: Chile, Concepción.

paricella Speiser, 1905:349 (nom. nov. for *chilensis* Reed). Type-locality: Chile, Concepción.

anchineuria Speiser of Donoso Barros, 1949:404, misident.

Subgenus PSEUDORNITHOMYIA Lutz, Neiva & Lima

PSEUDORNITHOMYIA Lutz, Neiva & Lima, 1915:176, 185 (as genus). Type-species, *ambigua* Lutz, Neiva & Lima (mon.).

AMBIGUA (Lutz, Neiva & Lima), 1915:185, pl. 27, fig. 5, pl. 28, fig. 7 (*Pseudornithomyia*). Type-locality: Brazil, Minas Gerais and Santa Catarina. Distr. — Venezuela, Colombia, Peru, Brazil.

HOFFMANNÆ Bequaert, 1954:158, figs. 31f-i, 32. Type-locality: Mexico, Chiapas, Cacahoatán. Distr. — Mexico.

Genus CRATERINA Ôlfers

CRATERINA Ôlfers, 1816:101. Type-species, *lonchoptera* Ôlfers (mon.) = *pallida* (Latreille).

Oxypterum Leach, 1817a:3. Type-species, *kyrbyanum* Leach (Westwood, 1840:154) = *pallida* (Latreille).



Anapera Meigen, 1830:234 (nom. nov. for *Oxypterum* Leach, on the wrong assumption that it was preocc. by *Ocyptera* Latreille, 1804). Type-species, *Oxypterum kyrbyanum* Leach (aut.).

Chelidomyia Rondani, 1879:15. Type-species, *Hippobosca hirundinis* Linnaeus (orig. des.).

SEGUYI Falcoz, 1930:53, fig. 8. Type-locality: Peru, Arequipa. Distr. — Colombia, Venezuela, Peru, Chile, Argentina.

Genus MYIOPHTHIRIA Rondani

MYIOPHTHIRIA Rondani, 1875:464 (as *Myophthiria*). Type-species, *reduvioides* Rondani (mon.).

Myocoryza Rondani, 1878:155. Type-species, *Myiophthiria lygaeoides* Rondani (mon.).

Subgenus BRACHYPTEROMYIA Williston

BRACHYPTEROMYIA Williston, 1896:184 (as genus). Type-species, *femorata* Williston (mon.) = *fimbriata* (Waterhouse).

NEOTROPICA (Bequaert), 1943b:113, fig. 1 (*Brachypteromyia*). Type-locality: Venezuela, D.F., Galipán, near Pico de Ávila. Distr. — Venezuela.

Genus ORNITHOCTONA Speiser

ORNITHOCTONA Speiser, 1902a:328. Type-species, *Ornithomyia erythrocephala* Leach (orig. des.).

Subgenus ORNITHOCTONA Speiser

ERYTHROCEPHALA (Leach), 1817a:13, pl. 27, figs. 4-6 (1818:559) (*Ornithomyia*). Type-locality: "Brazil". Distr. — widespread in the New World.

nebulosa Say, 1823:102 (1859:87) (*Ornithomyia*). Type-locality: "U.S.A."

gemina Thomson, 1869:611 (*Ornithomyia*). Type-locality: Peru, Callao.

bellardiana Rondani, 1878:157 (*Ornithomyia*). Type-locality: "Mexico".

haitiensis Bigot, 1885:242 (*Ornithomyia*). Type-locality: "Haiti".

robusta Wulp, 1903:431, pl. 13, figs. 5, 5a-c (*Ornithomyia*). Type-locality: Costa Rica, Caché (lectotype selected by Austen, 1903:261).

- pyrrhocephala* Speiser, 1904a:344 (*Ornithomyia*). Type-locality: "Mexico".
- buteonis* Swenk, 1916:133 (*Ornithomyia*). Type-locality: U.S.A., Nebraska, Neligh.
- costaricensis* Swenk, 1916:134 (*Ornithomyia*). Type-locality: Costa Rica, Juan Viñas.
- albiventris* Bau, 1922:274, 276. Type-locality: Cuba, Santa Clara, Estación Meyer.
- FUSCIVENTRIS (Wiedemann), 1830:611 (*Ornithomyia*). Type-locality: U.S.A., Kentucky. Distr. — widespread in the New World.
- chilensis* Guérin-Ménéville, 1835: pl. 104, fig. 5 (*Ornithomyia*). Type-locality: "Chile".
- chilensis* Macquart, 1843:437 (sep., p. 280) (*Ornithomyia*; preocc. Guérin-Ménéville, 1835). Type-locality: "Chile".
- testacea* Macquart, 1846:346 (sep., p. 218) (*Ornithomyia*). Type-locality: "Colombia".
- varipes* Walker, 1849:1146 (*Ornithomyia*). Type-locality: "Colombia".
- parva* Macquart of Speiser, 1902b:180, misident.
- pilosula* Wulp, 1903:432, pl. 13, figs. 6, 6a (*Ornithomyia*). Type-locality: Costa Rica, Volcán de Irazú.
- pirangae* Swenk, 1916:135 (*Ornithomyia*). Type-locality: Costa Rica, Juan Viñas.
- strigilecula* Ferris, 1923:57, figs. 3, 14a-c (*Ornithomyia*). Type-locality: Peru, Lima.
- ORIZABAE Bequaert, 1954:208, figs. 39a, 43g-h, 44b. Type-locality: Mexico, Veracruz, Orizaba. Distr. — Mexico.
- OXYCERA Falcoz, 1930:36, figs. 1-3. Type-locality: Colombia, Bogotá. Distr. — Colombia, Venezuela.

Subgenus ORNITHOPERTHA Speiser

- ORNITHOPERTHA Speiser, 1902b:167 (as genus). Type-species, *Ornithomyia nitens* Bigot (mon.).
- NITENS (Bigot), 1885:241 (*Ornithomyia*). Type-locality: "Panama". Distr. — Mexico, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela.
- geniculata* Bigot, 1892:49 (*Ornithomyia*). Type-locality: "Costa Rica".
- anthracina* Speiser, 1904b:393 (*Ornithopertha*). Type-locality: "Honduras".

Genus STILBOMETOPA Coquillett

- STILBOMETOPA Coquillett, 1899:336. Type-species, *Ornithomyia fulvifrons* Walker (mon.).
- FULVIFRONS (Walker), 1849:1145 (*Ornithomyia*). Type-locality: "Jamaica". Distr. — U.S.A., Costa Rica, Bahama Islands, Cuba, Isle of Pines, Jamaica.



- IMPRESSA (Bigot), 1885:237 (*Olfersia*). Type-locality: U.S.A., California. Distr. — U.S.A., Mexico.
- LEGTERSI Bequaert, 1955:256, figs. 46e, 48d, 49d, 59j, n. Type-locality: Mexico, Yucatán, Xocempich near Chichen Itzá. Distr. — Mexico, Brazil, Argentina.
- PODOPOSTYLA Speiser, 1904b:394. Type-locality: Brazil, Rio Grande do Sul (lectotype selected by Maa, 1963:59). Distr. — U.S.A. and continental Neotropical region.
columbarum Lutz, Neiva & Lima, 1915:178 (name proposed tentatively for specimens from Brazil referred to *podopostyla*).
- RAMPHASTONIS Ferris, 1930:63, figs. 1, 2a-d, f. Type-locality: Panama, Chiriquí, Camp Pitál. Distr. — Guatemala, Costa Rica, Panama, Ecuador, Brazil, Cuba, Hispaniola.

Genus LYNCHIA Weyenbergh

- LYNCHIA Weyenbergh, 1881:195. Type-species, *penelopes* Weyenbergh (mon.).
- Icosta* Speiser, 1905:358. Type-species, *Olfersia dioxyrhina* Speiser (mon.).
- Ornithoponus* Aldrich, 1923:73, 77. Type-species, *Feronia americana* Leach (orig. des.).
- ALBIPENNIS (Say), 1823:101 (1859:87) (*Olfersia*). Type-locality: "U.S.A." Distr. — New World, from Canada to Argentina, Bahama Islands, Cuba, Isle of Pines, Jamaica, Puerto Rico, Galápagos; Old World.
ardeae Macquart, 1835:640, pl. 24, fig. 10 (*Olfersia*). Type-locality: "Sicily".
propinqua Walker, 1849:1141 (*Olfersia*). Type-locality: "Jamaica".
botauri Rondani, 1879:22 (*Olfersia*). Type-locality: "Italy".
coriacea Wulp, 1903:430 (part: paratype of the species, from Mexico, Presidio) (*Olfersia*).
pahustris Lutz, Neiva & Lima, 1915:183, pl. 28, fig. 4 (*Olfersia*). Type-locality: Brazil, Piauí; Rio São Francisco; Minas Gerais, Lassance.
botaurinorum Swenk, 1916:128 (*Olfersia*). Type-locality: U.S.A., Nebraska, Omaha.
scutellaris Swenk, 1916:129 (*Olfersia*). Type-locality: U.S.A., Michigan, Watkin's Station near Manchester.
- AMERICANA (Leach), 1817a:11, pl. 27, figs. 1-3 (1818:557) (*Feronia*). Type-locality: U.S.A., Georgia. Distr. — continental New World.
falconis Harris, 1833:594 (*Ornithomyia*). *Nomen nudum*.
fusa Macquart, 1846:346 (sep., p. 218) (*Olfersia*). Type-locality: "Colombia" (as "Nouvelle Grenade").
bubonis Packard, 1869:417, fig. 340 (*Hippobosca*). Type-locality: U.S.A., Massachusetts.
rufiventris Bigot, 1885:243 (*Ornithomyia*). Type-locality: Brazil, Rio Grande do Sul, Pôrto Alegre.
- ANGUSTIFRONS (Wulp), 1903:430, pl. 13, figs. 3, 3a (*Olfersia*). Type-locality: Mexico, Tabasco, Teapa; Costa Rica, Río Sucio. Distr. — conti-

mental Neotropical region, from southern Canada (Ontario) to southern Brazil (Santa Catarina).

fusca Macquart of Lutz, Neiva & Lima, 1915:182 (*Olfersia*), misident.

HOLOPTERA (Lutz, Neiva & Lima), 1915:184 (*Olfersia*). Type-locality: Brazil, Rio de Janeiro. Distr. — U.S.A., Colombia, Brazil.

LATIFACIES Bequaert, 1955:302, figs. 56a-e. Type-locality: Brazil, São Paulo, Tabatinguera. Distr. — Brazil, Paraguay.

NIGRA (Perty), 1833:190, pl. 37, fig. 15 (*Hippobosca*). Type-locality: Brazil, Piauí. Distr. — New World, from Canada to Argentina, Galápagos; Hawaiian Islands.

fuscipennis Macquart, 1835:640 (*Olfersia*). Type-locality: "Brazil".

mexicana Macquart, 1843:435 (sep., p. 278) (*Olfersia*). Type-locality: "Mexico".

intertropica Walker, 1849:1144 (*Ornithomyia*). Type-locality: Galápagos Islands.

macquartii Rondani, 1878:160 (*Olfersia*). Type-locality: "Colombia" (as "Nouvelle Grenade").

pallidilabris Rondani, 1878:161 (*Olfersia*). Type-locality: "Mexico".

obliquinervis Rondani, 1878:162 (*Olfersia*). Type-locality: "Mexico".

villadae Dugès, 1887:20, pl. 3, figs. 3a-h (*Ornithomyia*). Type-locality: "Mexico".

acarta Speiser, 1902b:149 (*Olfersia*). Type-locality: Hawaiian Islands, Molokai Island.

raptatorum Lutz, Neiva & Lima, 1915:181, pl. 28, fig. 3 (*Olfersia*). Type-locality: "Brazil".

PENELOPES Weyenbergh, 1881:199. Type-locality: Argentina, Tucumán. Distr. — Argentina.

PLAUMANNI Bequaert, 1943a:132, figs. 1a-e. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Mexico, Guyana, Brazil, Bolivia.

WOLCOTTI (Swenk), 1916:132 (*Olfersia*). Type-locality: U.S.A., Michigan, Ann Arbor. Distr. — U.S.A., Mexico, Costa Rica, Panama, Venezuela, Brazil, Argentina.

Genus MICROLYNCHIA Lutz, Neiva & Lima

MICROLYNCHIA Lutz, Neiva & Lima, 1915:185. Type-species, *Lynchia pusilla* Speiser (mon.).

CRYPTURELLI Bequaert, 1938:346, figs. 1-2. Type-locality: Panama, Chiriquí, La Vaca. Distr. — Panama, Brazil.

FURTIVA Bequaert, 1955:382, figs. 75a-c. Type-locality: British Honduras, El Cayo. Distr. — British Honduras.

GALAPAGOENSIS Bequaert, 1955:384, figs. 76a-d. Type-locality: Galápagos Islands, Indefatigable Island, Academy Bay. Distr. — Galápagos Islands.



PUSILLA (Speiser), 1902b:157 (*Lynchia*). Type-locality: "Cuba". Distr. — Neotropical region, extending to the U.S.A., Cuba, St. Croix, St. Thomas, Granada.

Genus PSEUDOLYNCHIA Bequaert

PSEUDOLYNCHIA Bequaert, 1926:271. Type-species, *Olfersia maura* Bigot (orig. des.) = *canariensis* (Macquart).

BRUNNEA (Latreille) in Olivier, 1812:544 (*Ornithomyia*). Type-locality: "Carolina" (probably U.S.A., South Carolina). Distr. — Canada, U.S.A., Mexico, Panama, Cuba, Brazil, Argentina.

CANARIENSIS (Macquart), 1840:119 (*Olfersia*). Type-locality: Canary Islands. Distr. — Old World tropics and subtropics, introduced and established in the New World.

testacea Macquart, 1843:434 (sep., p. 277) (*Olfersia*). Type-locality: Reunion Islands.

rufipes Macquart, 1848:229 (sep., p. 69) (*Olfersia*). Type-locality: Bourbon Island.

falcinelli Rondani, 1879:23 (*Olfersia*). Type-locality: Malta Island.

maura Bigot, 1885:237 (*Olfersia*). Type-locality: "Algeria".

lividicolor Bigot, 1885:238 (*Olfersia*). Type-locality: "Brazil".

capensis Bigot, 1885:240 (*Olfersia*). Type-locality: South Africa, Cape of Good Hope.

exornata Speiser, 1900:562 (*Olfersia*). Type-locality: Sumatra, D. Tolong.

simillima Speiser, 1904a:337 (*Lynchia*). Type-locality: "Java".

Genus OLFERSIA Leach

Feronia Leach, 1817a:11 (1818:557) (preocc. Latreille, 1817). Type-species, *spinifera* Leach (aut.).

OLFERSIA Leach in Brewster, 1817:162 (nom. nov. for *Feronia* Leach). Type-species, *Feronia spinifera* Leach (Speiser, 1899:202).


Pseudolfersia Coquillett, 1899:336. Type-species, *maculata* Coquillett (mon.) = *fumipennis* (Sahlberg).

AENESCENS Thomson, 1869:610. Type-locality: Indian Ocean, Keeling Island. Distr. — Panama, Cocos Island, Bahama Islands, Puerto Rico, St. Croix, Brazil, Fernando Noronha, South Trinidad Island, Galápagos Islands, Mexico, Isabel Island, Revillagigedo group, Ascension Island; pantropic in the Atlantic, Pacific and Indian Oceans.

erythroptis Bigot, 1885:239. Type-locality: "New Caledonia".

diomedae Coquillett, 1901:379 (*Pseudolfersia*). Type-locality: Galápagos Islands, Albermarle Island.

BISULCATA Macquart, 1847:111 (sep., p. 95), pl. 6, fig. 12. Type-locality: "Chile". Distr. — southern U.S.A. to southern Brazil (São Paulo), Trinidad.



- vulturis* Wulp, 1903:429, pl. 13, figs. 1, 1a. Type-locality: Costa Rica, Río Sucio.
- CORIACEA Wulp, 1903:430, pl. 13, figs. 2, 2a. Type-locality: Guatemala, Mirandilla (lectotype selected by Austen, 1903:265). Distr. — continental Neotropical region, from southern Mexico to Brazil (22° S).
- meleagridis* Lutz, Neiva & Lima, 1915:179, pl. 27, fig. 3 (*Pseudolfersia*). Type-locality: Brazil, Pernambuco; Maranhão, São Luís; "Minas Gerais or Espírito Santo".
- FOSSULATA Macquart, 1843:434 (sep., p. 277). Type-locality: "Brazil". Distr. — Panama, Ecuador, Peru, Brazil, Chile, Cocos Island, Puerto Rico, St. Croix, Galápagos Islands.
- FUMIPENNIS (Sahlberg), 1886:150 (*Lynchia*). Type-locality: "Finland". Distr. — Panama, Ecuador, Peru, Brazil, Chile, Cocos Island, Cuba; Old World, sporadic.
- maculata* Coquillett, 1899:336 (*Pseudolfersia*). Type-locality: U.S.A., Wisconsin.
- mycetifera* Speiser, 1905:359 (*Pseudolfersia*). Type-locality: Arabia. Senafir Island, Red Sea.
- SORDIDA Bigot, 1885:239. Type-locality: "Guatemala". Distr. — U.S.A., Mexico, Guatemala, Panama, Venezuela, Brazil, Bahama Islands, Cuba, Isle of Pines, Jamaica, Hispaniola, St. Croix, Trinidad, Galápagos Islands.
- SPINIFERA (Leach), 1817a:6, 11, pl. 26, figs. 1-3 (1818:557) (*Feronia*). Type-locality: unknown. Distr. — U.S.A., Mexico, Honduras, British Honduras, Turneffe Island, Panama, Brazil, South Trinidad Island, Abrolhos Island, Queimada Grande Island, Alcatrazes Island, Bahama Islands, Cuba, Isle of Pines, Cayman Island, Jamaica, Puerto Rico, St. Croix, Ascension Island, Galápagos Islands; Old World.
- unicolor* Walker, 1849:1144 (*Ornithomyia*). Type-locality: "Jamaica".
- courtillieri* Fairmaire in Courtiller, 1853:196, pl. 15. Type-locality: France, Saumur.
- sulcifrons* Thomson, 1869:611. Type-locality: "Panama".

Subfamily LIPOPTENINAE

Genus NEOLIPOPTENA Bequaert

- NEOLIPOPTENA Bequaert, 1942:47. Type-species, *Lipoptena ferrisi* Bequaert (mon.).
- FERRISI (Bequaert), 1935:170 (*Lipoptena*). Type-locality: U.S.A., California, Mendocino Co. Distr. — Canada, U.S.A., Mexico.
- subulata* Coquillett of Ferris & Cole, 1922:187, misident.

Genus LIPOPTENA Nitzsch

- LIPOPTENA Nitzsch, 1818:310. Type-species, *Hippobosca cervina* Nitzsch (mon.) = *cervi* (Linnaeus).



- Haemobora* Curtis, 1824: pl. 14. Type-species, *pallipes* Curtis (mon.). = *cervi* (Linnaeus).
- Ornithobia* Meigen, 1830:229. Type-species, *pallida* Meigen (mon.) = *cervi* (Linnaeus).
- Alcephagus* Gimmerthal, 1845:152 (unnecessary change of name for *Ornithobia* Meigen). Type-species, *Ornithobia pallida* Meigen (aut.) = *cervi* (Linnaeus).
- GUIMARAESI Bequaert, 1957:504, figs. 102a-d. Type-locality: Brazil, Mato Grosso, São Domingos, Rio das Mortes. Distr. — Brazil.
- MAZAMAE Rondani, 1878:153. Type-locality: "Central and South America". Distr. — From U.S.A. to Argentina, Trinidad.
- mexicana* Townsend, 1897:289 (as *depressa* var.). Type-locality: Mexico, Veracruz, Paso de Telaya.
- conifera* Speiser, 1905:354. Type-locality: "Brazil".
- surinamensis* Bau, 1930:175. Type-locality: Surinam, Paramaribo.
- cervi* American authors, not Linnaeus.

Genus MELOPHAGUS Latreille

- MELOPHAGUS Latreille, 1802:466. Type-species, *Hippobosca ovina* Linnaeus (mon.).
- OVINUS (Linnaeus), 1758:607 (*Hippobosca*). Type-locality: "Europe". Distr. — Palaearctic; introduced and established in the New World.
- hirtella* Olfers, 1816:99. Type-locality: unknown.
- fera* Speiser, 1908:444 (as var.). Type-locality: "Caucasus".
- montanus* Ferris & Cole, 1922:192, figs. 9e, 10 (as ssp.). Type-locality: Alaska-Yukon boundary.
- bolivianus* Bau, 1930:176 (as "form"). Type-locality: Bolivia, Oruro.

LITERATURE CITED

- ALDRICH, J. M.
1923. Notes on the dipterous family Hippoboscidae. *Ins. Insc. Menstr.* 11:75-79.
- AUSTEN, E. E.
1903. Notes on Hippoboscidae in the collections of the British Museum. *Ann. Mag. Nat. Hist.* (7)12:255-266.
- BAU, A.
1922. *Ornithoctona albiventris* nov. spec. und *Ornithoica melaleuca* nov. spec. *Centralbl. Bakt. Parasit.* (2)57:274-279.
1930. *Lipoptena surinamensis* und *Melophagus ovinus bolivianus*, zwei neue Hippobosciden. *Stettin. Ent. Zeitg.* 91(2):175-177.



BEQUAERT, J.

1926. Notes on Hippoboscidae. 1. *Lynchia* Weyenbergh and *Lynchia* Speiser not congeneric. *Psyche* 32(1925):266-277.
1935. The American species of *Lipoptena*. *Bull. Brooklyn Ent. Soc.* 30:170.
1938. Notes on Hippoboscidae. 12. The genus *Microlynchia* Lutz, Neiva and Costa Lima. *Rev. Ent.*, Rio de Janeiro 9:343-349.
1942. A monograph of the Melophaginae, or ked-flies, of sheep, goats, deer and antelopes. *Entomologica Americana* (n.s.) 22:1-220, 19 figs.
- 1943a. Notes on Hippoboscidae. 16. Hippoboscidae from southern Brazil, with the description of a new species of *Lynchia*. *Jour. Parasit.* 29:131-135.
- 1943b. Notes on Hippoboscidae. 18. The genus *Brahypteromyia* Williston, with the description of a new species. *Psyche* 49 (1942): 108-117.
- 1954-1957. The Hippoboscidae or louse-flies (Diptera) of mammals and birds. Part II. Taxonomy, evolution and revision of American genera and species. *Entomologica Americana* 34:1-232, figs. 22-44, 1954; 35:233-416, figs. 45-82, 1955; (1956) 36:417-611, figs. 83-104, 1957.

BERGROTH, E.

1901. Ueber eine auf Eulen schmarotzende Hippoboscidae. *Meddel. Soc. pro Fauna et Flora Fenn.* 27:146-150.

BIGOT, J. M. F.

1885. Diptères nouveaux ou peu connus. 27e partie. XXXV. *Ann. Soc. Ent. France* (6)5:225-246.
1892. Description d'une espèce nouvelle de Diptère parasite de Costa Rica. *The Humming Bird* 2(7):49.

COQUILLETT, D. W.

1899. New genera and species of Nycteribiidae and Hippoboscidae. *Canad. Ent.* 31:333-336, 343.
1901. Papers from the Hopkins-Stanford Galapagos Expedition, 1898-1899. II. Entomological Results. (2). Diptera. *Proc. Wash. Acad. Sci.* 3:371-379.

COURTILLER, A.

1853. Note sur le diptère *Olfersia courtillieri* (Fairmaire). *Ann. Soc. Linn. Départ. Maine-et-Loire* 1:196-197, pl. 15.

CURTIS, J.

1824. *British entomology: Being illustrations and descriptions of the genera of insects found in Great Britain and Ireland* 1: pls. 1-50. London.
1836. *British entomology: Being illustrations and descriptions of the genera of insects found in Great Britain and Ireland* 13: pls. 578-625. London.

DONOSO BARROS, R.

1949. Observación de um caso de ectoparasitismo accidental del hombre, por *Olfersia fossulata* Macquart (1843). *Rev. Méd. Chile* 77(6):166.



DUGÈS, A.

1887. *Adelophis copei*, *Argas sanchezi* y *Ornithomyia villadae*. *La Naturaleza*, México (2) 1(1):18-21, pl. 3.

FALCOZ, L.

1930. Diptères pupipares du Muséum National d'Histoire Naturelle de Paris. Hippoboscidae. *Encycl. Ent. (B) Diptera* 5 (1929):27-54.

FERRIS, G. F.

1923. Observations on the larvae of some Diptera Pupipara with description of a new species of Hippoboscidae. *Parasitology* 15(1): 54-58.

1930. Some New World Hippoboscidae. *Canad. Ent.* 62:62-70.

FERRIS, G. F. & F. R. COLE

1922. A contribution to the knowledge of the Hippoboscidae. *Parasitology* 14(2):178-205.

GIMMERTHAL, B. A.

1845. Bemerkungen über zwei Dipteren-Arten. *Stettin. Ent. Zeitg.* 6:151-153.

GUÉRIN-MÉNEVILLE, F. E.

1835. Pls. 92-104, in his *Iconographie du règne animal de G. Cuvier*. Vol. 2: *Planches des animaux invertébrés, Insectes*, 104 pls. Paris.

HARRIS, T. W.

1833. VIII. Insects, pp. 566-595, in E. Hitchcock, *Report on the geology, mineralogy, botany and zoology of Massachusetts* (Ed. 1), 700 pp., 104 figs. + Atlas, 17 pls. Amherst, Mass.

LATREILLE, P. A.

1802. *Histoire naturelle, générale et particulière, des Crustacés et des Insectes* 3:468 pp. Paris, "An X".

1810. *Considerations générales sur l'ordre naturel des animaux*, 444 pp. Paris.

LEACH, W. E.

- 1817a. *On the genera and species of eproboscideous insects*, 20 pp., 3 pls. Edinburgh, September 1817. (Also in 1818, *Mem. Werner. Nat. Hist. Soc.*, Edinburgh 2(2):547-566, pls. 25-27).

- 1817b. Insecta, pp. 155-164, in D. Brewster, ed., *The Edinburgh encyclopaedia* 12(1):1-376, pls. 330-340 (November, 1817). Edinburgh.

LINNAEUS, C.

1758. *Systema naturae per regna tria naturae*, Ed. 10 1:824 pp. Holmiae (= Stockholm).

LUTZ, A., A. NEIVA & A. DA C. LIMA

1915. Sobre "Pupipara" ou "Hippoboscidae" de aves brasileiras. *Mem. Inst. Oswaldo Cruz* 7(2):173-199, pls. 27-28.

LYNCH ARRIÁZAGA, E.

1881. Diptera, in *Informe oficial de la comisión científica agregada al Estado Mayor General de la expedición al Río Negro (Patagonia)* 1 (Zoología): 88-91, Buenos Aires.

MAA, T. C.

1963. Genera and species of Hippoboscidae (Diptera): types, synonymy, habitats and natural groupings. *Pacific Insects Monograph* 6:1-186, 55 figs.

MACQUART, J.

1835. *Histoire naturelle des insectes, Diptères* 2: 703 pp., 12 pls. Paris.
1840. *Diptères exotiques nouveaux ou peu connus* 2(1):5-135, 21 pls. Paris. (Also published in *Mém. Soc. Roy. des Sci., de l'Agr. et des Arts*, Lille 1840:283-413, 21 pls., 1841).
1843. *Diptères exotiques nouveaux ou peu connus. Mém. Soc. Roy. des Sci., de l'Agr. et des Arts*, Lille 1842:162-460, 36 pls. (Also separately published, pp. 5-304, 36 pls. Paris, 1843).
1846. *Diptères exotiques nouveaux ou peu connus [Ier] Supplément. Mém. Soc. Roy. des Sci., de l'Agr. et des Arts*, Lille (1845) 1844: 133-364, 20 pls. (Also separately published, pp. 5-238, 20 pls., Paris, 1846).
1847. *Diptères exotiques nouveaux ou peu connus. 2^e supplément. Mém. Soc. Roy. des Sci., de l'Agr. et des Arts*, Lille 1846:21-120, 6 pls. (Also separately published, pp. 5-104, 6 pls. Paris, 1847).

MEIGEN, J. W.

1830. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten* 6:iv + 401 pp., pls. 55-66. Hamm.

NITZSCH, C. L.

1818. Die Familien und Gattungen der Thierinsekten (Insecta epizoica) als Prodrömus eine Naturgeschichte derselben. *Mag. d. Ent.*, Halle 3:261-316.

OLFERS, J. F. M. VON

1816. *De vegetativis et animatis corporibus in corporibus animatis reperiundis commentarius* 1:iv+112 pp., 1 pl. Berlin.

OLIVIER, G. A.

1812. Insectes [i. e., Arthropoda, pt. 5], vol. 8, pp. 361-722 (= livr. 77), in *Société de Gens de Lettres, de Savans et d'Artistes, Encyclopédie Méthodique. Histoire Naturelle*, Paris.

PERTY, M.

1833. *Delectus animalium articulorum quae in itinere per Brasiliam annis MDCCCXVII-MDCCCXX jussu et auspiciis Maximiliani Josephi I. Bavariae regis augustissimi peracto collegerunt Dr. J. B. de Spix et Dr. C. F. Ph. de Martius*, pp. 125-224, pls. 25-40. Monachii (= Munich).

PACKARD, A. S.

1869. *Guide to the study of insects and a treatise on those injurious and beneficial to crops*, 702 pp., 651 figs., 10 pls. Salem, Mass.

REED, E. C.

1904. Los dipteros pupíparos de Chile. *Rev. Chil. Hist. Nat.* 8:149-153.

RONDANI, C.

1875. *Muscaria exotica* Musei Civici Januensis. III. Species in Insula Bonae Fortunae (Borneo), Provincia Sarawak, annis 1865-68, lectae a March. J. Doria et Doct. O. Beccari. *Ann. Mus. Civ. Stor. Nat. Genova* 7:421-466.
1878. *Muscaria exotica* Musei Civici Januensis observatae et distincta. IV. *Hippoboscita exotica* non vel minus cognita. *Ann. Mus. Civ. Stor. Nat. Genova* 12:150-170.
1879. *Hippoboscita italica* in familias et genera distributa. *Boll. Soc. Ent. Ital.* 11:3-28.

SAHLBERG, J.

1886. *Lynchia fumipennis* n. sp. en på *Pandion haliaëtus* lefvande hippoboscid. *Meddel. Soc. pro Fauna et Flora Fenn.* 13:149-152, 1 fig.

SAY, T.

1823. Descriptions of dipterous insects of the United States. *Jour. Acad. Nat. Sci. Philad.* 3:9-54, 73-104.
1859. In J. L. LeConte, ed., *The complete writings of Thomas Say on the entomology of North America* 1:421 pp., 54 pls.; 2:814 pp. New York.

SCHUURMANS STEKHOFEN, J. H., JR., I. I. SILVA & P. SAN ROMÁN

1957. Die Fortpflanzungsbiologie von *Pseudolynchia canariensis* (Fam. Hippoboscidae). *Zeitschr. z. Parasitenk.* 17:490-509, 2 figs.

SPEISER, P.

1900. Studien über Hippobosciden. I. *Ann. Mus. Civ. Stor. Nat. Genova* 40:553-562.
- 1902a. Besprechung einiger Gattungen und Arten der Diptera Pupipara. I. *Termész. Füzetek* 25:327-338.
- 1902b. Studien über Diptera Pupipara. *Zeitschr. Syst. Hym. Dipt.* 2:145-180.
- 1902c. Diptera Pupipara, in *Fauna Hawaiiensis* 2(2):86-92.
- 1904a. Studien über Hippobosciden. II. *Ann. Mus. Civ. Stor. Nat. Genova* 41:332-350.
- 1904b. Besprechung einiger Gattungen und Arten der Diptera Pupipara. II. *Ann. Mus. Nat. Hungarici* 2:386-395.
1905. Beiträge zur Kenntnis der Hippobosciden. *Ztschr. Syst. Hym. Dipt.* 5:347-360.
1908. Die geographische Verbreitung der Diptera Pupipara und ihre Phylogenie. *Ztschr. Wiss. Insektenbiol.* 4:241-246, 301-305, 420-427, 437-447.

SWENK, M. H.

1916. Descriptions and records of North American Hippoboscidae. *Jour. New York Ent. Soc.* 24:126-136.

THOMSON, C. G.

1869. 6. Diptera. Species novas descripsit, pp. 443-612, pl. 9 (= h. 12, n° 2), in K. Svenska Vetenskaps-Akademien, *Kongliga svenska fregatten Eugenies resa omkring jorden 2* (Zoologie), [Sec.] 1: *Insekter*:617 pp., 9 pls., Stockholm, "1868".

TOWNSEND, C. H. T.

1897. On a collection of Diptera from the lowlands of the Rio Nautla, in the State of Vera Cruz. *Ann. Mag. Nat. Hist.* (6)20:272-291.

WALKER, F.

1849. *List of the specimens of dipterous insects in the collection of the British Museum* 4:689-1172, London.

WESTWOOD, J. O.

1840. Order XIII. Diptera Aristotle (Antliata Fabricius. Alteriptera Clairv.), pp. 125-128, 129-144, 145-158, in his *An introduction to the modern classification of insects. Synopsis of the genera of British insects*, 158 pp. London.

WEYENBERGH, H.

1881. Dos nuevas especies del grupo de los dípteros pupíparos. *An. Soc. Cient. Argent.* 11(5):193-200.

WIEDEMANN, C. R. W.

1830. *Aussereuropäische zweiflügelige Insekten* 2:xii + 684 pp., pls. 7-10b. Hamm.

WILLISTON, S. W.

1896. A new genus of Hippoboscidae. *Ent. News* 7:184-185.

WULP, F. M. VAN DER

1880. Eenige Diptera van nederlandsch Indië. *Tijdschr. v. Ent.* 23: 155-194.


1903. Fam. Hippoboscidae, pp. 429-432, pl. 13, figs. 1-6, in F. D. Godman & O. Salvin, eds., *Biologia Centrali Americana, Zoologia-Insecta-Diptera* 2:489 pp., 11 figs., 13 pls. London.

INDEX

<i>acarta</i> (Speiser), 1902 (Olfersia), <i>Lynchia</i>	99.6
<i>aenescens</i> Thomson, 1869, <i>Olfersia</i>	99.7
<i>albipennis</i> (Say), 1823 (Olfersia), <i>Lynchia</i>	99.5
<i>albiventris</i> Bau, 1922, <i>Ornithoctona</i>	99.4
<i>Alcephagus</i> Gimmerthal, 1845	99.9
<i>ambigua</i> (Lutz, Neiva & Lima), 1915 (<i>Pseudornithomyia</i>), <i>Ornithomyia</i>	99.2
<i>americana</i> (Leach), 1817 (<i>Feronia</i>), <i>Lynchia</i>	99.5
<i>Anapera</i> Meigen, 1830	99.3
<i>anchineuria</i> Speiser, 1905, <i>Ornithomyia</i>	99.2
<i>anchineuria</i> Speiser of Donoso Barros, 1949, <i>Ornithomyia</i>	99.2
<i>angustifrons</i> (Wulp), 1903 (Olfersia), <i>Lynchia</i>	99.5
<i>anthracina</i> Speiser, 1904, <i>Ornithoctona</i>	99.4
<i>ardeae</i> (Macquart), 1835 (Olfersia), <i>Lynchia</i>	99.5
<i>avicularia</i> authors, <i>Ornithomyia</i>	99.2
<i>beccarina</i> Rondani, 1878, <i>Ornithoica</i>	99.1
<i>bellardiana</i> (Rondani), 1878 (<i>Ornithomyia</i>), <i>Ornithoctona</i>	99.3
<i>bisulcata</i> Macquart, 1847, <i>Olfersia</i>	99.7





<i>bolivianus</i> Bau, 1930, Melophagus	99.9
<i>botauri</i> (Rondani), 1879 (Olfersia), Lynchia	99.5
<i>botaurinorum</i> (Swenk), 1916 (Olfersia), Lynchia	99.5
<i>Brachypteromyia</i> Williston, 1896	99.3
<i>brunnea</i> (Latreille), 1812 (Ornithomyia), Pseudolynchia	99.7
<i>bubonis</i> (Packard), 1869 (Hippobosca), Lynchia	99.5
<i>buteonis</i> (Swenk), 1916 (Ornithomyia), Ornithoctona	99.4
<i>canariensis</i> (Macquart), 1840 (Olfersia), Pseudolynchia	99.7
<i>capensis</i> (Bigot), 1885 (Olfersia), Pseudolynchia	99.7
<i>cervi</i> authors, Lipoptena	99.9
<i>Chelidomyia</i> Rondani, 1879	99.3
<i>chilensis</i> (Guérin-Ménéville), 1835 (Ornithomyia), Ornithoctona ...	99.4
<i>chilensis</i> (Macquart), 1843 (Ornithomyia), Ornithoctona	99.4
<i>chilensis</i> Reed, 1904, Ornithomyia	99.2
<i>chloropus</i> Bergroth, 1901, Ornithomyia	99.2
<i>columbarum</i> Lutz, Neiva & Lima, 1915, Stilbometopa	99.5
<i>confluens</i> authors, Ornithoica	99.1
<i>confluens</i> (Wiedemann), 1830 (Ornithomyia), Ornithoica	99.1
<i>confluenta</i> authors, Ornithoica	99.1
<i>confluenta</i> (Say), 1823 (Ornithomyia), Ornithoica	99.1
<i>conifera</i> Speiser, 1905, Lipoptena	99.9
<i>coriacea</i> Wulp, 1903, Olfersia	99.8
<i>coriacea</i> (Wulp), 1903 (Olfersia), Lynchia	99.5
<i>costaricensis</i> (Swenk), 1916 (Ornithomyia), Ornithoctona	99.4
<i>courtillieri</i> Fairmaire, 1853, Olfersia	99.8
<i>Craterina</i> Olfers, 1816	99.2
<i>crypturelli</i> Bequaert, 1938, Microlynchia	99.6
<i>diomedae</i> (Coquillett), 1901 (Pseudolfersia), Olfersia	99.7
<i>erythrocephala</i> (Leach), 1817 (Ornithomyia), Ornithoctona	99.3
<i>erythrospis</i> Bigot, 1885, Olfersia	99.7
<i>exornata</i> (Speiser), 1900 (Olfersia), Pseudolynchia	99.7
<i>falcinelli</i> (Rondani), 1879 (Olfersia), Pseudolynchia	99.7
<i>falconis</i> (Harris), 1833 (Ornithomyia), Lynchia	99.5
<i>fera</i> Speiser, 1908, Melophagus	99.9
<i>Feronia</i> Leach, 1817	99.7
<i>ferrisi</i> (Bequaert), 1935 (Lipoptena), Neolipoptena	99.8
<i>fossulata</i> Macquart, 1843, Olfersia	99.8
<i>fringillina</i> Curtis, 1836, Ornithomyia	99.2
<i>fulvifrons</i> (Walker), 1849 (Ornithomyia), Stilbometopa	99.4
<i>fumipennis</i> (Sahlberg), 1886 (Lynchia), Olfersia	99.8
<i>furtiva</i> Bequaert, 1955, Microlynchia	99.6
<i>fusca</i> (Macquart), 1846 (Olfersia), Lynchia	99.5
<i>fusca</i> Macquart of Lutz, Neiva & Lima, 1915, Lynchia	99.6
<i>fuscipennis</i> (Macquart), 1835 (Olfersia), Lynchia	99.6
<i>fusciventris</i> (Wiedemann), 1830 (Ornithomyia), Ornithoctona	99.4
<i>galapagoensis</i> Bequaert, 1955, Microlynchia	99.6
<i>gemina</i> (Thomson), 1869 (Ornithomyia), Ornithoctona	99.3
<i>geniculata</i> (Bigot), 1892 (Ornithomyia), Ornithoctona	99.4
<i>guimaraesi</i> Bequaert, 1957, Lipoptena	99.9




<i>Haemobora</i> Curtis, 1824	99.9
<i>haitiensis</i> (Bigot), 1885 (Ornithomyia), Ornithoctona	99.3
<i>hirtella</i> Olfers, 1816, Melophagus	99.9
<i>hoffmanae</i> Bequaert, 1945, Ornithomyia	99.2
holoptera (Lutz, Neiva & Lima), 1915 (Olfersia), Lynchia	99.6
<i>Icosta</i> Speiser, 1905	99.5
<i>impressa</i> (Bigot), 1885 (Olfersia), Stilbometopa	99.5
<i>intertropica</i> (Walker), 1849 (Ornithomyia), Lynchia	99.6
latifacies Bequaert, 1955, Lynchia	99.6
legtersi Bequaert 1955, Stilbometopa	99.5
Lipoptena Nitzsch, 1818	99.8
Lipoptenini	99.8
<i>lividicolor</i> (Bigot), 1885 (Olfersia), Pseudolynchia	99.7
Lynchia Weyenbergh, 1881	99.5
<i>macquartii</i> (Rondani), 1878 (Olfersia), Lynchia	99.6
<i>maculata</i> (Coquillett), 1899 (Pseudolfersia) Olfersia	99.8
<i>maura</i> (Bigot), 1885 (Olfersia), Pseudolynchia	99.7
<i>mazamae</i> Rondani, 1878, Lipoptena	99.9
<i>melaleuca</i> Bau, 1922, Ornithoica	99.1
<i>meleagridis</i> (Lutz, Neiva & Lima), 1915 (Pseudolfersia), Olfersia.	99.8
Melophagus Latreille, 1802	99.9
<i>mexicana</i> (Macquart), 1843 (Olfersia), Lynchia	99.6
<i>mexicana</i> Townsend, 1897, Lipoptena	99.9
Microlynchia Lutz, Neiva & Lima, 1915	99.6
<i>montanus</i> Ferris & Cole, 1922, Melophagus	99.9
<i>mycetifera</i> (Speiser), 1905 (Pseudolfersia), Olfersia	99.8
Myiophthiria Rondani, 1875	99.3
<i>Myocoryza</i> Rondani, 1878	99.3
<i>Myophthiria</i> Rondani, 1875	99.3
<i>nebulosa</i> (Say), 1823 (Ornithomyia), Ornithoctona	99.3
Neolipoptena Bequaert, 1942	99.8
neotropica (Bequaert), 1943 (Brachypteromyia), Myiophthiria ...	99.3
nigra (Perty), 1833 (Hippobosca), Lynchia	99.6
nitens (Bigot), 1885 (Ornithomyia), Ornithoctona	99.4
<i>obliquinervis</i> (Rondani), 1878 (Olfersia), Lynchia	99.6
Olfersia Leach, 1817	99.7
orizabae Bequaert, 1954, Ornithoctona	99.4
<i>Ornithobia</i> Meigen, 1830	99.9
Ornithoctona Speiser, 1902	99.3
Ornithoica Rondani, 1878	99.1
Ornithoicinae	99.1
<i>Ornithomya</i> Latreille, 1802	99.2
Ornithomyia Latreille, 1802	99.2
Ornithomyiinae	99.2
Ornithopertha Speiser, 1902	99.4
<i>Ornithoponus</i> Aldrich, 1923	99.5
ovinus (Linnaeus), 1758 (Hippobosca), Melophagus	99.9
oxycera Falcoz, 1930, Ornithoctona	99.4
<i>Oxypterum</i> Leach, 1817	99.2


<i>pallida</i> Say, 1823, Ornithomyia	99.2
<i>pallidilabris</i> (Rondani), 1878 (Olfersia), Lynchia	99.6
<i>palustris</i> (Wulp), 1903 (Olfersia), Lynchia	99.5
<i>paricella</i> Speiser, 1905, Ornithomyia	99.2
<i>parva</i> Macquart, 1843, Ornithomyia	99.2
<i>parva</i> Macquart of Speiser, 1902, Ornithoctona	99.4
<i>penelopes</i> Weyenbergh, 1881, Lynchia	99.6
<i>peroneura</i> Speiser, 1902, Ornithoica	99.1
<i>pilosula</i> (Wulp), 1903 (Ornithomyia), Ornithoctona	99.4
<i>pirangae</i> (Swenk), 1916 (Ornithomyia), Ornithoctona	99.4
<i>plaumanni</i> Bequaert, 1943, Lynchia	99.6
<i>podopostyla</i> Speiser, 1904, Stilbometopa	99.5
<i>promiscua</i> Ferris of Cole, 1922, Ornithoica	99.1
<i>propinqua</i> (Walker), 1849 (Olfersia), Lynchia	99.5
<i>Pseudolfersia</i> Coquillett, 1899	99.7
<i>Pseudolynchia</i> Bequaert, 1926	99.7
<i>Pseudornithomyia</i> Lutz, Neiva & Lima, 1915	99.2
<i>pusilla</i> (Speiser), 1902 (Lynchia), Microlynchia	99.7
<i>pyrrhocephala</i> (Speiser), 1904 (Ornithomyia), Ornithoctona	99.4
<i>ramphastonis</i> Ferris, 1930, Stilbometopa	99.5
<i>raptatorum</i> (Lutz, Neiva & Lima), 1915 (Olfersia), Lynchia ...	99.6
<i>remota</i> Walker, 1849, Ornithomyia	99.2
<i>robusta</i> (Wulp), 1903 (Ornithomyia), Ornithoctona	99.3
<i>rufipes</i> (Macquart), 1848 (Olfersia), Pseudolynchia	99.7
<i>rufiventris</i> (Bigot), 1885 (Ornithomyia), Lynchia	99.5
<i>scutellaris</i> (Swenk), 1916 (Olfersia), Lynchia	99.5
<i>seguyi</i> Falcoz, 1930, Craterina	99.3
<i>simillima</i> (Speiser), 1904 (Lynchia), Pseudolynchia	99.7
<i>sordida</i> Bigot, 1885, Olfersia	99.8
<i>spinifera</i> (Leach), 1817 (Feronia), Olfersia	99.8
<i>Stilbometopa</i> Coquillett, 1899	99.4
<i>strigilecula</i> (Ferris), 1923 (Ornithomyia), Ornithoctona	99.4
<i>subulata</i> Coquillett of Ferris & Cole, 1922, Neolipoptena	99.8
<i>sulcifrons</i> Thomson, 1869, Pseudolynchia	99.8
<i>surinamensis</i> Bau, 1930, Lipoptena	99.9
<i>synnalaxidis</i> Lynch Arribalzaga, 1881, Ornithomyia	99.2
<i>testacea</i> (Macquart), 1843 (Olfersia), Pseudolynchia	99.7
<i>testacea</i> (Macquart), 1846 (Ornithomyia), Ornithoctona	99.4
<i>unicolor</i> (Walker), 1849 (Ornithomyia), Olfersia	99.8
<i>variegata</i> Bigot, 1885, Ornithomyia	99.2
<i>varipes</i> (Walker), 1849 (Ornithomyia), Ornithoctona	99.4
<i>vicina</i> (Walker), 1849 (Ornithomyia), Ornithoica	99.1
<i>villadae</i> (Dugès), 1887 (Ornithomyia), Lynchia	99.6
<i>vulturis</i> Wulp, 1903, Olfersia	99.8
<i>wolcottii</i> (Swenk), 1916 (Olfersia), Lynchia	99.6





SciELO₉





MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO

Formerly Departamento de Zoologia, Secretaria da Agricultura

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

100

FAMILY STREBLIDAE

RUPERT L. WENZEL

Field Museum of Natural History
Chicago, Illinois

The Streblidae are blood-sucking pupiparous flies and are obligate parasites of bats. All are ectoparasites, except the females of the Old World genus *Ascodipteron*, which embed within the skin of the host and become endoparasitic. The family is primarily tropical in distribution, with a few species in the subtropical and warm temperate zones. The Old and New World Streblidae have no subfamilies in common. Three subfamilies with 23 genera and 94 described species occur in the New World. Two other subfamilies, with 4 genera and 62 species occur in the Old World.


The greatest diversification of genera and species has occurred on families of bats that are endemic to the New World. Their adaptations are reflected in a wide range of form, size and structure. Many species of *Trichobius* are rather generalized calypterate flies, but the species of *Nyctrophilia* are flea-like, while others, like *Strebla*, are polycetenoid. Some are micropterous and flightless. The species of *Paradysehiria* are wingless.

Most of the New World species are restricted to one host or to a few closely related ones. No truly polyxenous species are known.

The females of most New World species deposit what appears to be a soft puparium, lacking any evidence of larval segmentation. This usually hardens and pigments almost immediately after being deposited, but the "tanning" process sometimes begins while it is still within the female. However, the "puparia" of species of the *Trichobius caecus* group exhibit typical larval segmentation and it may be more nearly correct to refer to these particular species as larviparous.

The Streblidae, Nycteribiidae, Hippoboscidae and Brulidae were at one time placed together in a separate section of the Cyclorrhapha, called the Pupipara. While the Streblidae and Nycteribiidae appear to be rather closely related, these two families have little in common with either the Hippoboscidae or the Brulidae.

The most recent papers dealing with taxonomy of New World species are those of Jobling (1929-52), Guimarães (1938-44), Hoffmann (1953) and Wenzel, Tipton & Kiewlicz (1966). Discussions of higher classification are given by Jobling (1936) and Wenzel, Tipton & Kiewlicz (1966). The



only recent world list of genera and species is that of Maa (1965). Host-parasite relationships are discussed by Jobling (1949b), Theodor (1947) and Wenzel, Tipton & Kiewlicz (loc. cit.) and Wenzel & Tipton (1966). Their biology has been discussed by Jobling (loc. cit.), Ross (1961), and Wenzel, Tipton & Kiewlicz (loc. cit.). Their morphology has been treated by Jobling (1929, 1936, 1951), Zeve & Howell (1962-63) and Wenzel, Tipton & Kiewlicz (loc. cit.).

Subfamily NYCTEROPHILIINAE

Genus NYCTEROPHILIA Ferris

NYCTEROPHILIA Ferris, 1916:436. Type-species, *coxata* Ferris (orig. des.). *Nycterophila*, error.

Ref. — Wenzel, Tipton & Kiewlicz, 1966:432 (revision).

COXATA Ferris, 1916:437, fig. 5, pl. 22, fig. 6 (lateral view; wing). Type-locality: U.S.A., California, Santa Margherita River, ex *Macrotus californicus*. Distr. — U.S.A. (Arizona, California), Mexico (Baja California, Colima, Guerrero, Morelos, Oaxaca, Puebla, Sinaloa). Refs. — Hoffmann, 1953:183, 187 (part), pl. 3, fig. 2 (lateral view); Wenzel, Tipton & Kiewlicz, 1966:434, figs. 48B, 50B, 51A (head, thorax; ♂ abdomen; ♀ abdomen).

FAIRCHILD Wenzel, 1966:436, figs. 47B, 49B, 51B (head, thorax; ♂ abdomen; apex ♀ abdomen). Type-locality: Panama, Coclé, Penonomé Caves, ex *Pteronotus suapurensis*. Distr. — Mexico (Sinaloa), Guatemala (Altavcrapaz), Panama (Coclé), Venezuela (Aragua, Dto. Federal).

NATALI Wenzel, 1966:438, figs. 48A, 50A (head, thorax; ♂ abdomen, hindleg). Type-locality: Panama, Canal Zone, San Lorenzo Caves, ex *Natalus stramineus mexicanus*. Distr. — (?) Mexico, (?) Guatemala, Panama (Canal Zone).

?*coxata* Ferris of Hoffmann, 1953:183, 187 (part).

PARNELLI Wenzel, 1966:434, figs. 45A, 46, 47A, 51C (♂ terminalia; lateral view; head, thorax; apex ♀ abdomen). Type-locality: Panama, Canal Zone, Paraiso, ex *Pteronotus parnelli fuscus*. Distr. — Mexico (Colima, Guerrero, Sinaloa, Yucatán), Guatemala (Altavcrapaz, Escuintla, Petén), Nicaragua (Matagalpa), Panama (Canal Zone, Coclé, Colón, Los Santos, Panamá), Colombia (Bolívar), Venezuela (Apure, Bolívar, Carabobo),

Subfamily TRICHOBIINAE

Genus TRICHOBIUS Gervais

TRICHOBIUS Gervais, 1844:14. Type-species, *parasiticus* Gervais (mon.).


Refs. — Kessel, 1925 (synopsis); Jobling, 1938 (revision); Zeve & Howell, 1962-63 (morphology); Wenzel, Tipton & Kiewlicz, 1966 (revision, C. & S. Am. spp.)

Trichobia, error

Strebla, authors (part), not Wiedemann



- Kolenatia* Rondani, 1878:169. Type-species, *Strebila wiedemanni* Kolenati (orig. des.).
- Trichobius* Townsend, 1891:105 (preocc. Gervais, 1844). Type-species, *dugesii* Townsend (orig. des.)
- Kesselia* Curran, 1934:522. Type-species, *pallida* Curran (orig. des.)
- ADAMSI Augustson, 1943:52, pl. 6, figs. 1, 2 (♂ wing, mesonotum). Type-locality: U.S.A., California, Mohave Co., Parker, ex *Macrotus californicus*. Distr. — U.S.A. (Arizona, California), Mexico (Baja California, Colima, Guerrero, Morelos, Oaxaca, Puebla, Sinaloa). Ref. — Ross, 1959:71 (redescri.), fig. 1A-C (dorsum, ♂ wing; head)
- BRENNANI Wenzel, 1966:497, figs. 62D, 74A, 75B (venter of thorax; ♂ mesonotum; ♂ terminalia). Type-locality: Panama, Chiriquí, Casa Tilley, near Cerro Punta, ex *Sturnira ludovici*. Distr. — Panama (Bocas del Toro, Canal Zone, Darién, Chiriquí).
- BEQUAERTI Wenzel, 1966:471, figs. 63A, 66A (♂ terminalia; ♂ mesonotum). Type-locality: Panama, Bocas del Toro, Almirante, ex *Tonatia bidens*. Distr. — Panama (Bocas del Toro).
- CAECUS Edwards, 1918:424. Type-locality: Trinidad, Guaicharo Caves, no host given. Distr. — Trinidad, Venezuela (Apure, Aragua, Bolívar, Carabobo, Dto. Federal, Guarico, Miranda, Monagas, Sucre, Yaracuy). Host. — *Pteronotus parnellii fuscus*. Ref. — Wenzel, Tipton & Kiewliez, 1966:450, figs. 55B, 57A-C (abd. terga I + II; ♂, ♀ terminalia).
- coecus*, error
- CORYNORHINI Cockerell, 1910:59. Type-locality: U.S.A., Colorado, Boulder Co., Great Sphinx Mine, S. of Crisman, ex *Corynorhinus macrotus pallescens*. Distr. — U.S.A. (West Virginia, Kentucky, Arkansas, Kansas, Oklahoma, Colorado, Arizona, Utah, Nevada, California, Oregon, Washington), Mexico (Chihuahua, Guanajuato, Nuevo León). Ref. — Jobling, 1938:366, fig. 1A-D (dorsum, ♀; labium; apex of abdomen, ♂; wing).
- quadrisetosus* Kessel, 1925:15 (as *major* ssp.). Type-locality: "Mexico?"
- COSTALIMAI Guimarães, 1937:660, pl. 3, fig. 10 (mesonotum). Type-locality: Brazil, Pernambuco, Recife, ex undetermined bat. Distr. — Honduras (Tegueigalpa) to Brazil (Pernambuco), Peru (Piura). Host. — *Phyllostomus discolor*. Ref. — Wenzel, Tipton & Kiewliez 1966:471, figs. 63E, 67B (♂ genitalia; mesonotum).
- DIPHYLAE Wenzel, 1966:492, figs. 68B, 73B (♂ terminalia; mesonotum, ♂). Type-locality: Guatemala, Jalapa, San Lorenzo, ex *Diphylla ecaudata*. Distr. — Mexico (Quintana Roo), Guatemala (Chimaltenango, Jalapa), Venezuela (Aragua).
- DUGESII Townsend, 1891:106. Type-locality: Mexico, Guanajuato, ex *Glossophaga soricina*. Distr. — Mexico (Guerrero) to Guyana (Demerara), Peru (Junín). Ref. — Wenzel, Tipton & Kiewliez, 1966:478, figs. 68G-H, 69A (♂ terminalia; mesonotum).
- blandus* Curran, 1935:10, fig. 11 (mesonotum). Type-locality: Panama, Canal Zone, Paraíso, ex *Glossophaga soricina leachii*.



- DUGESIOIDES** Wenzel, 1966:488, figs. 68D, 71 (♂ terminalia; ♂, ♀ mesonotum). Type-locality: Panama, Panamá, Chepo Road, ex *Trachops c. cirrhosus*. Distr. — Guatemala (Chimaltenango) to Brazil (São Paulo), Peru (Loreto).
- DUNNI** Wenzel, 1966:474, figs. 63C, 67A (♂ terminalia; ♂ mesonotum). Type-locality: Panama, Darién, Río Chucunaquc, ex *Molossus bondae*. Distr. — Panama (Darién).
- DYBASI** Wenzel, 1966:469, figs. 63B, 65A (♂ terminalia; ♀ mesonotum). Type-locality: Panama, San Blas, Armila, ex *Tonatia silvicola*. Distr. — Panama (Bocas del Toro), Colombia (Meta), Peru (Piura).
- FURMANI** Wenzel, 1966:490, figs. 68C, 72 (♂ terminalia; ♂, ♀ mesonotum). Type-locality: Peru, Puno, Segrario, ex *Desmodus rotundus*. Distr. — Colombia (Meta, Vaupés), Venezuela (Aragua), Peru (Junín, Puno), Paraguay (San Pedro).
- GALEI** Wenzel, 1966:449, figs. 54, 57J-L (♂ mesonotum; ♀ terminalia; ♂ terminalia). Type-locality: Panama, Canal Zone, San Lorenzo Caves, ex *Natalus stramineus mexicanus*. Distr. — Panama (Canal Zone), Colombia (Córdoba), Venezuela (Aragua, Miranda).
caecus, authors (part), not Edwards.
- HIRSUTULUS** Bequaert, 1933:572, figs. 30 a, b (♂ mesonotum; ♀ scutellum). Type-locality: Mexico, Yucatán, Ticul District, Yocat, ex *Myotis nigricans extremus*. Distr. — Mexico (Yucatán, Tamaulipas). Ref. — Jobling, 1938:369, fig. 3A-D (dorsum, ♀; labium; apex, ♂ abdomen; wing).
- JOBLINGI** Wenzel, 1966:481, figs. 68E, 70 (♂ terminalia; ♂, ♀ mesonotum). Type-locality: Panama, Canal Zone, east of Summit Golf Club, ex *Carollia perspicillata azteca*. Distr. — Mexico (Chiapas, Vera Cruz) to Brazil (Pará, São Paulo), Peru (Junín, Loreto, Puno), Trinidad, Tobago.
blandus, authors (part), not Curran.
dugesii, authors (part), not Townsend.
- JOHNSONAE** Wenzel, 1966:455, figs. 55A, 57G-I (abd. terga I + II; ♂, ♀ terminalia). Type-locality: Panama, Coclé, Penonomé Cave, ex *Pteronotus suapurensis*. Distr. — Panama (Coclé, Chiriquí, Panamá, San Blas), Colombia (Bolívar), Venezuela (Aragua).
- KEENANI** Wenzel, 1966:462, fig. 60A (♂ terminalia). Type-locality: Panama, Bocas del Toro, Almirante, ex *Micronycteris m. megalotis*. Distr. — Panama (Bocas del Toro, Canal Zone, Los Santos), Venezuela (Amazonas).
- LIONYCTERIDIS** Wenzel, 1966:464, fig. 60C (♂ terminalia). Type-locality: Panama, San Blas, Armila, ex *Lionycteris spurrelli*. Distr. — Panama (San Blas), Peru (Cuzco), Venezuela (Bolívar).
- LONCHOPHYLLAE** Wenzel, 1966:461, figs. 59 A-C, 60D, 61A (dorsal view, ♂; labium, apex of abdomen, ♀; ♂ terminalia; wing). Type-locality: Panama, Colón, Buena Vista Caves, ex *Lonchophylla r. robusta*. Distr. — Panama (Bocas del Toro, Canal Zone, Colón, Darién, San Blas), Colombia (Tolima).

- uniformis* Curran, 1935:10 (part), and authors, including Jobling, 1938:380 (part).
- LONGIPES** Rudow, 1871:121 (*Strebla*). Type-locality: None given, ex "*Phyllostoma hastatum*". Distr. — Nicaragua (Jinotega, Matagalpa) to Brazil (Pará), Bolivia (Cochabamba), Peru (Cuzco Huanuco, Junín, Loreto, Piura), Trinidad. Ref. — Wenzel, Tipton & Kiewlicz, 1966:465, figs. 62A, E, 63F, 64 (venter of thorax; ♂ terminalia; ♀, ♂ mesonotum).
- dugesii*, authors (part), not Townsend.
- mixtus* Curran, 1935:10, fig. 10 (mesonotum). Type-locality: Panama, Panamá, Chilibrillo Caves, ex *Phyllostomus hastatus panamensis*.
- MACROPHYLLI** Wenzel, 1966:486, figs. 68F, 69B (♂ terminalia; mesonotum). Type-locality: Panama, Canal Zone, Fort Davis, ex *Macrophyllum maeophyllum*. Distr. — Panama (Canal Zone, San Blas), Venezuela (Apure, Bolívar).
- MAJOR** Coquillett, 1899:334. Type-locality: U.S.A., Florida, Citrus County, Gum Cave. Distr. — U.S.A. (Florida, Kansas, Oklahoma, Texas, Arizona, California), Mexico (Dto. Federal, Guerrero, Michoacán, Morelos, Nuevo León), Guatemala (Chimaltenango). Hosts. — *Myotis* spp. Refs. — Jobling, 1938:368, fig. 2A-D (labium; head & thorax, ♀; ♀ terminalia; wing); Zeve & Howell, 1962:685-694, figs. 1, 4, 7. 1963:2-17, figs. 1, 5, 8, 11, 14-20; 127-138, figs. 1, 4, 7, 10, 11, 14, 17 (morphology).
- MENDEZI** Wenzel, 1966:469, figs. 63D, 65B (♂ terminalia; ♀ mesonotum). Type-locality: Panama, San Blas, Armila, ex *Tonatia minuta*. Distr. — Panama (Bocas del Toro, Los Santos, San Blas), Venezuela (Yaracuy-Carabobo border).
- PALIDUS** (Curran), 1934:522 (*Kesselia*). Type-locality: Guyana, Kartabo, ex *Furipterus horrens*. Distr. — Guyana (Kartabo), Venezuela. Refs. — Jobling, 1936:373, fig. 5A-C (dorsum, ♂; hindleg; wing); Wenzel, Tipton & Kiewlicz, 1966:447, fig. 53 (Jobling's 1936 fig.).
- PARASITICUS** Gervais, 1844:14, pl. 43, fig. Type-locality: "Guiane" ex *Desmodus rufus* (= *rotundus*). Distr. — Mexico (Chiapas, Guerrero, Jalisco, Oaxaca, Puebla, Quintana Roo, San Luis Potosí, Vera Cruz, Yucatán) to Trinidad, Brazil (Amazonas, Pará, Pernambuco), Argentina (Santa Fé) and Peru (Cuzco, Ica, Iquitos, Loreto, Piura). Refs. — Jobling, 1938:381, fig. 11A-C (labium; dorsum, ♀; apex of abdomen, ♂); Wenzel, Tipton & Kiewlicz, 1966:494, figs. 68A, 73A (♀ terminalia; ♂ mesonotum).
- kesseli* Guimarães, 1938:660, fig. 9 (mesonotum) (nom. nov. for *parasiticus* Gervais of Kessel, 1925:15, and Curran 1935:11, in error).
- PHYLLOSTOMAE** Kessel, 1925:16, pl. 1, fig. 9 (mesonotum). Type-locality: Brazil [Santa Catarina], Hansa Humboldt (= Corupá) ex *Phyllostomus* sp. Distr. — Brazil (Santa Catarina). Refs. — Jobling, 1938:376, fig. 7A-D (dorsum, ♀; labium; sternopleura; wing); Wenzel, Tipton & Kiewlicz, 1966:496 ff., fig. 75D (♂ terminalia).
- PSEUDOTRUNCATUS** Jobling, 1939c:164 (nom. nov. for *kesselae* Jobling). Type-locality: Dominica, Basin Well. Distr. — Dominica (Basin Well), Bahamas (Long Island).

- kesselae* Jobling, 1938:377, fig. 8A-D (labium; dorsum, ♀; apex of abdomen, ♂; wing) (precoe. Guimarães, 1938). Type-locality: Dominica, Basin Well, ex *Myotis nigricans*.
- SPARSUS** Kessel, 1925:17, pl. 1, fig. 7; pl. 2, fig. 10 (mesonotum; wing). Type-locality: Panama, Panamá, Chilibrillo River, ex *Chilonycteris rubiginosa fusca* (= *Pteronotus parnelli fuscus*). Distr. — Guatemala (Eseuintla), Panama (Boeas del Toro, Canal Zone, Los Santos, Panamá), Venezuela (Bolívar, Guarico, Miranda, Nueva Esparta). Refs. — Jobling, 1938:373, fig. 6A-D (head, sternopleura; dorsum, ♀; wing; apex of abdomen, ♀); Wenzel, Tipton & Kiewliez, 1966:457, figs. 42, 58 (wing; Jobling's 1938 figs., plus ♂ terminalia).
- SPHAERONOTUS** Jobling, 1939a:494, fig. 4A-C (dorsum, ♀; labium; apex of abdomen, ♂). Type-locality: Mexico, Nuevo León, Cerro Potosí, ex *Leptonycteris nivalis*. Distr. — Mexico (Nuevo León, Guerrero), U.S.A. (Arizona, New Mexico, Texas, Oklahoma).
- TRUNCATUS** Kessel, 1925:17, pl. 1, fig. 4 (mesonotum). Type-locality: "Dominica". Distr. — Dominica, St. Vincent I., Puerto Rico (Aguas Buenas). Refs. — Jobling, 1938:378, fig. 9A-C (dorsal view, ♂; labium; wing; apex of abdomen, ♀).
- UNIFORMIS** Curran, 1935:10 (part), fig. 8 (dorsum, ♀). Type-locality: Panama, Canal Zone, Paraíso ex *Glossophaga soricina leachi*. Distr. — Mexico (Chiapas, Yucatán) to Venezuela (many localities), Guyana (Demerara, Essequibo) and Peru (Lambayeque). Ref. — Wenzel, Tipton & Kiewliez, 1966:459, figs. 60B, 61B (♂ terminalia; wing).
- URODERMAE** Wenzel, 1966:476, figs. 66B, 68I (♀ mesonotum; ♂ terminalia). Type-locality: Panama, Boeas del Toro, Almirante, ex *Uroderma bilobatum*. Distr. — Panama (Boeas del Toro, Darién, Canal Zone, San Blas), Venezuela (Trujillo).
- VAMPYROPIS** Wenzel, 1966:500, figs. 74B, 75C (♀ mesonotum; ♂ terminalia). Type-locality: Panama, Boeas del Toro, near Rancho Grande, ex *Vampyrops vittatus*. Distr. — Panama (Boeas del Toro), Venezuela (Dto. Federal).
- YUNKERI** Wenzel, 1966:453, figs. 56 A-B, 57D-F (dorsum, ♂; labium; ♂, ♀ terminalia). Type-locality: Panama, Canal Zone, Paraíso, ex *Pteronotus parnelli fuscus*. Distr. — Mexico (Guerrero, Jalisco, Sinaloa, Yucatán), Nicaragua (Matagalpa), Guatemala (Eseuintla, Petén, Zacapa), Panama (Boeas del Toro, Canal Zone, Colón, Darién, Los Santos), Peru (Lima).
caecus, authors (part), not Edwards.
- sp.
wiedemanni Kolenati, 1863, part: 96, pl. 15, fig. 36a-e (dorsum; wing; venter) (*Strebla*). Distr. — Cuba. Not Kolenati, 1856.

Genus ANATRICHOBIOUS Wenzel

- ANATRICHOBIOUS** Wenzel, 1966:502. Type-species, *scorzai* Wenzel (orig. des.).
- SCORZAI** Wenzel, 1966:503, figs. 76-78 (dorsum, venter, ♀; ♂ abdomen; ♂ terminalia). Type-locality: Panama, Chiriquí, Cueva Lara, ex

Myotis sp. Distr. — Panama (Chiriquí, Los Santos, Panamá), Colombia (Huila), Venezuela (Aragua, Dto. Federal, Yaracuy-Carabobo border), Peru (Cuzco).

Genus JOBLINGIA Dybas & Wenzel

JOBLINGIA Dybas & Wenzel, 1947:149. Type-species, *schmidtii* Dybas & Wenzel (orig. des.).

SCHMIDTI Dybas & Wenzel, 1947:152, figs. 25-27 (dorsum, ♀; wing; venter of abdomen, ♀). Type-locality: Guatemala, Chimaltenango, Chocoyos, ex *Myotis n. nigricans*. Distr. — Guatemala (Chimaltenango), Panama (Chiriquí). Ref. — Wenzel, Tipton & Kiewlicz, 1966:507, figs. 79A-B, 80 (dorsal view, ♀; wing; ♂ terminalia).

Genus TRICHOBIODES Wenzel

TRICHOBIODES Wenzel, 1966:510. Type-species, *Trichobius perspicillatus* Pessôa & Galvão (orig. des.).

PERSPICILLATUS (Pessôa & Galvão), 1937:1, figs. 1-3 (mesonotum; ventral view, ♀; wing) (*Trichobius*). Type-locality: [Brazil] "Bahia", ex undet. host and *Hemiderma perspicillatum* (= *Carollia perspicillata*). Distr. — Mexico (Oaxaca) to Brazil (Bahia) and Peru (Piura). Ref. — Wenzel, Tipton & Kiewlicz, 1966:511, figs. 81, 82A (♂ mesonotum; ♂ terminalia).

Genus MASTOPTERA Wenzel

MASTOPTERA Wenzel, 1966:512. Type-species, *Aspidoptera minuta* Lima (orig. des.).


GUIMARAESI Wenzel, 1966:514, figs. 82C, 83, 84 (♂ terminalia, dorsal view, ♀; wing; labium; abdomen, ♂, ♀). Type-locality: Panama, Panamá, Chepo Road, ex *Phyllostomus hastatus panamensis*. Distr. — Nicaragua (Jinotega), Costa Rica (Guanacaste), Panama (Bocas del Toro, Canal Zone, Los Santos, Panamá), Colombia (Huila, Bolívar), Venezuela (Yaracuy).

minuta Lima of authors (part), and of Jobling, 1949a: fig. 4 (dorsum, ♂; labium; wing) (*Aspidoptera*).

MINUTA (Lima), 1921:21, pl. 2, fig. 2 (wing) (*Aspidoptera*). Type-locality: Brazil, Mato Grosso, ex *Tonatia amblyotis*. Distr. — Panama (Bocas del Toro, Canal Zone, Los Santos, San Blas), Colombia (Antioquia, Bolívar, Meta, Santander Nte.), Venezuela (Bolívar, Carabobo, Trujillo), Suriname (Nickerie), Brazil (Mato Grosso), Ecuador (Santiago Zamora), Peru (Cuzco, Loreto, Piura), Bolivia (Cochabamba). Refs. — Jobling, 1949a:142, part (record from *Tonatia amblyotis*); Wenzel, Tipton & Kiewlicz, 1966:515, figs. 82B, 85 (♂ terminalia; abdomen, ♂, ♀).

Genus PARATRICHOBIOUS Lima

PARATRICHOBIOUS Lima, 1921:20. Type-species, *Trichobius longicrnis* Ribeiro (orig. des.).



- DUNNI (Curran), 1935:7, fig. 6 (dorsum) (*Speiseria*). Type-locality: Panama [Darién], El Real, ex *Peropteryx canina* or *Uroderma bilobatum*. Distr. — Guatemala (Eseuintla), Panama (Boeas del Toro, Canal Zone, Colón, Darién, San Blas), Venezuela (Bolívar, Dto. Federal, Falcón, Miranda, Sucre, Trujillo, Yaracuy). Refs. — Jobling, 1939:492, fig. 3A-C (dorsum, ♀; apex of abdomen, ♀; venter of head, sternopleura); Wenzel, Tipton & Kiewliez, 1966:527, figs. 90, 91A, 92A (Jobling's 1939a figs. plus tergum IX, ♂ terminalia).
- LONGIERUS (Ribeiro), 1907:236, pl. 25 (dorsum, ♀) (*Trichobius*). Type-locality: Brazil, Guanabara, Rio de Janeiro, Quinta da Boa Vista, ex *Artibeus jamaicensis*. Distr. — Mexico (Chiapas, San Luis Potosí) to Brazil (Rio de Janeiro) and Peru (Loreto). Ref. — Wenzel, Tipton & Kiewliez, 1966:521, fig. 89 (measurements).
- LOWEI Wenzel, 1966:528, figs. 92B, 93 (♀ terminalia; head, thorax, profemora, ♂). Type-locality: Panama, San Blas, Armila, ex *Artibeus cinereus watsoni*. Distr. — Panama (Darién, San Blas), Venezuela (Bolívar).
- SALVINI Wenzel, 1966:532, figs. 95, 95C (head, thorax, profemora, ♀; ♂ terminalia). Type-locality: Panama, Darién, Cerro Tacareuna, ex *Chiroderma salvini*. Distr. — Panama (Darién), Venezuela (Falcón, Dto. Federal, Miranda, Monagas).
- SANCHEZI Wenzel, 1966:530, figs. 92C, 94 (♂ terminalia; ♀ mesonotum). Type-locality: Panama, Darién, Cerro Mali, ex *Enchisthenes hartii*. Distr. — Panama (Darién, Los Santos), Venezuela (Aragua, Dto. Federal).
- sp. (*longicrus* Ribeiro), Wenzel, Tipton & Kiewliez, 1966:523, figs. 86, 88A, 89, 91B (dorsum, ♂; venter of head, sternopleura; ♂ terminalia; measurements; tergum IX). Host. — *Artibeus lituratus palmarum*. Distr. — Panama (Boeas del Toro, Canal Zone), Venezuela (Aragua, Bolívar, Dto. Federal), Trinidad.
- longicrus* Ribeiro of Jobling, 1939:490 (part), fig. 1A-C (dorsum, ♂; venter of head, sternopleura); 1949:316 (part).
- sp. (*longicrus* complex), Wenzel, Tipton & Kiewliez, 1966:524, figs. 87A, 88D, 89 (♂ mesonotum; ♂ terminalia; measurements). Host. — *Vampyropros vittatus*. Distr. — Panama (Boeas del Toro, Darién), Venezuela (Bolívar, Dto. Federal).
- sp. (*longicrus* complex), Wenzel, Tipton & Kiewliez, 1966:524, figs. 87B, 88B, 89 (♂ mesonotum; ♂ terminalia; measurements). Host. — *Artibeus aztecus*. Distr. — Panama (Bocas del Toro, Chiriquí).
- sp. (*longicrus* complex), Wenzel, Tipton & Kiewliez, 1966:527, figs. 88C, 89 (♂ terminalia; measurements). Host. — *Artibeus toltecus*. Distr. — Panama (Boeas del Toro, Chiriquí).
- sp. A, Wenzel, Tipton & Kiewliez, 1966:535, fig. 96A (♂ terminalia). Host. — *Chiroderma villosum jesupi*. Distr. — Panama (San Blas).
- sp. B, Wenzel, Tipton & Kiewliez, 1966:535, fig. 96A (♂ terminalia). Host. — *Vampyropros helleri*. Distr. — Panama (Los Santos), Venezuela (Yaracuy-Carabobo border).

Genus NEOTRICHOBIVS Wenzel & Aitken

NEOTRICHOBIVS Wenzel & Aitken, 1966:536. Type-species, *stenopterus* Wenzel & Aitken (orig. des.) = *delicatus* (Machado-Allison).

DELICATUS (Machado-Allison), 1966-76, figs. 8-11 (wing; scutum and scutellum; sternum II; profemurs, coxa) (*Pterellipsis*). Type-locality: Venezuela, Distrito Federal, Chichiriviche, ex *Vampyressa pusilla thyone*. Distr. — Panama (Bocas del Toro, Los Santos), Colombia (Santander, Valle), Venezuela (Aragua, Bolívar, Dto. Federal, Guarico, Miranda, Sucre, Yaracuy-Carabobo border), Suriname (Nickerie), Trinidad (Sangre Grande), Brazil (Pará).

stenopterus Wenzel & Aitken, 1966:539, figs. 97-99 (dorsum, venter, ♀; apex of abdomen, ♂, ♀; wing). Type-locality: Panama, Bocas del Toro, Almirante, ex *Uroderma bilobatum*. N. syn.

Genus MEGISTOPODA Macquart

MEGISTOPODA Macquart, 1852:332. Type-species, *pilatei* Macquart (mon.). *Megistopodia*, error.

Pterellipsis Coquillett, 1899:333. Type-species, *arana* Coquillett (orig. des.). Refs. — Speiser, 1900:32 ff.; Jobling, 1936:357 ff. (morphology); Maa, 1965:385 (synonymy); Wenzel, Tipton & Kiewlicz, 1966:541-47 (synonymy, revision, key); Machado-Allison, 1966:69-79 (Venezuelan spp.).


ARANEA (Coquillett), 1899:334 (*Pterellipsis*). Type-locality: Jamaica, ex undetermined bats. Distr. — Greater & Lesser Antilles, Mexico (many localities) to Brazil (Pará, São Paulo Santa Catarina), Paraguay (San Pedro), Peru (Huanuco, Lambayeque), Trinidad and Tobago. Refs. — Jobling, 1949:316 ff., fig. 2C (foreleg) and 1952:134, fig. 2A (thorax); Wenzel, Tipton & Kiewlicz, 1966:542, figs. 100A, D (♂ terminalia; wing); Machado-Allison, 1966:70, figs. 1-4 (wing; scutum, scutellum; sternum II; foreleg).

desiderata Speiser, 1900:57, pl. 3, figs. 6-8 (lateral view; wing; antenna). Type-locality: "Brazil", "Cuba", ex "*Phyllostoma* sp."

PILATEI Macquart, 1852:332, pl. 4, fig. 4 (lateral view). Type-locality: Mexico, Teapa, ex undetermined bat. Distr. — Mexico (Teapa). Ref. — Wenzel, Tipton & Kiewlicz, 1966:541-2.

PROXIMA (Séguy), 1926-194, figs. 2-6 (lateral view; antenna; eye; wing; ♂ terminalia) (*Pterellipsis*). Type-locality: Argentina, "Misiones, environs de San Ignacio, Villa Lutecea.", ex unidentified bat. Distr. — Mexico (Chiapas, Jalisco) to Paraguay (San Pedro), Brazil (Pará, Santa Catarina), Argentina (Misiones), and Peru (Madre de Dios). Host. — *Sturnira lilium*. Refs. — Jobling, 1952:134, fig. 4B (thorax); Wenzel, Tipton & Kiewlicz, 1966:543, figs. 100C, 101 (Séguy's figs., plus ♂ terminalia); Machado-Allison, 1966:74, figs. 5-7 (wing; scutum & scutellum; sternum II).

THEODORI Wenzel, 1966:545, fig. 100B (♂ terminalia). Type-locality: Panama, Chiriquí, Cerro Punta, Casa Tilley, ex *Sturnira ludovici*. Distr. — Costa Rica (Cartago), Panama (Bocas del Toro, Chiriquí,



Darién), Venezuela (Aragua, Dto. Federal, Guarico, Merida, Miranda, Monagas).

Genus SPEISERIA Kessel

SPEISERIA Kessel, 1925:19. Type-species, *ambigua* Kessel (orig. des.).

Speisseria, error

Synthesiostrebla Townsend of Jobling, 1939a:487.

Ref. — Wenzel, Tipton & Kiewlicz, 1966:549.

AMBIGUA Kessel, 1925:20, pl. 1, figs. 1, 2 (mesonotum; venter of thorax).

Type-locality: [Brazil] Pernambuco, ex *Vampyrus*. Distr. — Mexico (Chiapas, San Luis Potosí, Tamaulipas) to Brazil (Pará, São Paulo) and Peru (Lambayeque, Loreto), Trinidad. Ref. — Wenzel, Tipton & Kiewlicz, 1966:549, fig. 102A-C (Jobling's 1939a fig., plus ♀ terminalia).

amorphochili Townsend of Jobling, 1939a:488, fig. 1A-C (dorsal view, ♀; labium; sternopleura, apex of abdomen, ♂) (*Synthesiostrebla*), misident.

anduzei Matheson, 1945:191, fig. 1a-e (mesonotum; sternopleura; profemur; palpus; labium) (*Paratrachobius*). Type-locality: Venezuela, San Esteban, ex undetermined bats.

Genus ASPIDOPTERA Coquillett

ASPIDOPTERA Coquillett, 1894:334. Type-species, *buscki* Coquillett (orig. des.).

Lepopteryx Speiser, 1900:53. Type-species, *Lipoptena phyllostomatis* Perty (orig. des.).

Lipoptena, authors (part), not Nitsch.

Refs. — Jobling, 1949a:135-144 (revision); Wenzel, Tipton & Kiewlicz, 1966:551-558 (revision).

BUSCKI Coquillett, 1894:335. Type-locality: Puerto Rico, Bayamon, ex *Artibeus* sp. Distr. — Cuba (Soledad), Puerto Rico (Bayamon), Mexico (Jalisco), Guatemala (Escuintla, Altaverrapaz), Costa Rica (Guanacaste, Puntarenas), Panama (Bocas del Toro, Canal Zone, Darién, Los Santos, Panamá, San Blas), Colombia (Bolívar, Putumayo), Peru (Lambayeque). Ref. — Wenzel, Tipton & Kiewlicz, 1966:555, figs. 104A, C (♀, ♂ terminalia).

phyllostomatis, authors (part), not Perty.

DELATORREI Wenzel, 1966:557, figs. 104B, D (♀, ♂ terminalia). Type-locality: Panama, Los Santos, Guanico, ex *Sturnira lilium parvidens*. Distr. — Mexico (Jalisco, Chiapas), Guatemala (Escuintla), Nicaragua (Matagalpa), Panama (Darién, Los Santos, San Blas), Colombia (Cauca, Putumayo, Tolima), Venezuela (Aragua, Bolívar, Carabobo, Dto. Federal, Falcón, Guarico, Miranda, Monagas, Sucre, Trujillo, Yaracuy).

PHYLLOSTOMATIS (Perty), 1833:190, pl. 37, fig. 16 (dorsal view) (*Lipoptena*). Type-locality: (Ncotype): Brazil, Santa Catarina, Hansa Humboldt (= Corupá), ex *Phyllostomus* sp. Distr. — Brazil (Santa Cata-

rina). Refs. — Jobling, 1949a:136, fig. 1A-C (dorsal view; wing; labium); Wenzel, Tipton & Kiewlicz, 1966:553, fig. 103A-C (Jobling's 1949a figs.).

buscki, authors (part), not Coquillett.

Genus EXASTINION Wenzel

EXASTINION Wenzel, 1966:558. Type-species, *Aspidoptera clovisi* Pessôa & Guimarães (orig. des.).

CLOVISI (Pessôa & Guimarães), 1937:262, figs. 5, 6 (dorsal view, ♂; wing) (*Aspidoptera*). Type-locality: Brazil, São Paulo, Ipiranga, ex *Anoura geoffroyi*. Distr. — Mexico (Chiapas) to Brazil (São Paulo) and Ecuador (Azuay). Refs. — Jobling, 1949:138, fig. 2A-C (dorsal view, with extruded puparium; mouthparts; wing) (*Aspidoptera*); Wenzel, Tipton & Kiewlicz, 1966:560, fig. 1-5A-E (Jobling's 1949 figs. plus head, ♂ terminalia).

phyllostomatis Perty of Bequaert, 1940:418, and of Kessel (part), 1925:26 (*Aspidoptera*), misident.

Genus NOCTILOSTREBLA Wenzel

NOCTILOSTREBLA Wenzel, 1966:560. Type-species, *Lipoptena dubia* Rudow (orig. des.).

Lipoptena, authors (part), not Nitsch.

Lepopteryx Speiser, 1900:53, part.

Aspidoptera, authors (part), not Coquillett.

Ref. — Wenzel, Tipton & Kiewlicz, 1966:560 (revision, key).

AITKENI Wenzel, 1966:567, figs. 107C, 108 (♀ terminalia; dorsal view; labium; wing). Type-locality: Trinidad, Manzanilla, ex *Noctilio l. leporinus*. Distr. — Venezuela (Bolívar, Miranda), Trinidad (Manzanilla, Scotland Bay), B. W. I. (Monos Island), Suriname (Brokopondo, Nickerie), Brazil (Pará), Peru (Loreto).

megastigma Speiser of Jobling (part), 1949a:140, fig. 3A-C (dorsal view; labium; wing), and of Goodwin & Greenhall, 1961:220 (*Aspidoptera*), misident.

DUBIA (Rudow), 1871:122 (*Lipoptena*). Type-locality: "Venezuela", ex *Noctilio dorsatus* (= *N. leporinus*). Distr. — Venezuela. Ref. — Wenzel, Tipton & Kiewlicz, 1966:563.

megastigma Speiser of Bequaert, 1942:88 (*Aspidoptera*), misident.

MAAI Wenzel, 1966:569, figs. 107A, 109 (♂ terminalia; ♂, ♀ abd. sternum II). Type-locality: Panama, Darién, Río Tuirá, ex *Noctilio labialis*. Distr. — Panama (Canal Zone, Darién, Panamá), Venezuela (Apure, Bolívar, Dto. Federal, Falcón-Yaracuy border, Yaracuy).

MEGASTIGMA (Speiser), 1900:54, pl. 3, fig. 2 (dorsal view) (*Lepopteryx*). Type-locality: Unknown, ex *Noctilio dorsatus* (= *N. leporinus*). Distr. — Not known. Ref. — Wenzel, Tipton & Kiewlicz, 1966:564.

TRAUBI Wenzel, 1966:565, figs. 106, 107B, D (♂, ♀ abd. sternum II, terminalia). Type-locality: Panama, Canal Zone, Fort Sherman,

ex *Noctilio leporinus mexicanus*. Distr. — Cuba (Isle of Pines, Havana), Costa Rica (Limón), Panama (Canal Zone, Darién, Panamá), Peru (Tumbes), Colombia (Antioquia, Magdalena), Venezuela (Yaracuy-Carabobo border).

Genus PARADYSCHIRIA Speiser

- PARADYSCHIRIA Speiser, 1900:55. Type-species, *fusca* Speiser (orig. des.).
Ref. — Wenzel, Tipton & Kiewlicz, 1966:571-78 (revision, key).
- FUSCA Speiser, 1900:56, pl. 3, fig. 1 (dorsum, ♀). Type-locality: Colombia, Orocué, ex *Noctilio leporinus*. Distr. — Colombia (Orocué), Venezuela (Bolívar, Miranda, Yaracuy), Trinidad, Suriname (Nickerie), French Guiana (Cayenne), Brazil (Pará). Refs. — Wenzel, Tipton & Kiewlicz, 1966:573, figs. 110C, 111B (♀ abdomen; ♂ abd. terga I + II).
- dubia* Rudow of Guimarães, 1941:217, figs. 1-4 (dorsum, venter; apex of abdomen, ♀, ♂) (re-described); 1944:186, figs. 5-9 (larva), misident.
- dubia*, authors (part), not Rudow.
- lineata*, authors (part), not Kessel.
- LINEATA Kessel, 1925:27. Type-locality: "Cuba", no host given. Distr. — Puerto Rico (Quebrada Cave near San Germán), Cuba (Isle of Pines, Havana), Panama (Bocas del Toro, Canal Zone, Panamá). Ref. — Wenzel, Tipton & Kiewlicz, 1966:574, figs. 110A-B, 111A, 112A-B, 113A (♀ abdomen; ♀ terminalia; ♂ abd. terga I + II; ♂ terminalia).
- PARVULA Falcoz, 1931:267, figs. 4, 5 (dorsum; venter of head, thorax). Type-locality: "Brazil", ex *Dirias albiventer* (= *Noctilio labialis*). Distr. — Colombia (Antioquia), Venezuela (Apurc, Carabobo, Falcón, Guarico, Merida, Yaracuy), Trinidad, Brazil (Pará), Bolivia (Beni). Ref. — Wenzel, Tipton & Kiewlicz, 1966:574.
- dubia*, authors (part), not Rudow.
- PARVULOIDES Wenzel, 1966:575, figs. 110D, 112C-D, 113B, 114A-B (♀ abdomen; ♀, ♂ terminalia; ♂ thorax; abd. terga I + II). Type-locality: Panama, Darién, Río Tuira, ex *Noctilio l. labialis*. Distr. — Nicaragua (Zelaya), Costa Rica (Puntarenas), Panama (Canal Zone, Darién, Los Santos, Panamá), Colombia (Antioquia, Chocó, Cauca, Valle del Cauca).
- dubia* Rudow of Cooper, 1941:126; and of Jobling (part), 1949b:328, misident.

Genus PARASTREBLA Wenzel

- PARASTREBLA Wenzel, 1966:578. Type-species, *handleyi* Wenzel (orig. des.).
- HANDLEYI Wenzel, 1966:579, fig. 116 (thorax; apex ♀ abdomen). Type-locality: Panama, San Blas, Armila, ex *Micronycteris nicefori*. Distr. — Panama (San Blas).

Genus SYNTHESIOSTREBLA Townsend

- SYNTHESIOSTREBLA Townsend, 1913:98. Type-species, *amorphochili* Townsend (orig. des.).

Refs. — Jobling, 1947:39-41, Wenzel, Tipton & Kiewlicz, 1966:429, 649.

AMORPHOCHILI Townsend, 1913:98. Type-locality: Peru, Piura, ex *Amorphochilus schnablii*. Distr. — Peru (Piura, Cuzco). Refs. — Jobling, 1947:40, fig. 1A, B (dorsal view, ♀: venter of head, sternopleura); Wenzel, Tipton & Kiewlicz, 1966: fig. 115 (Jobling's 1947 figs.).

Genus PSEUDOSTREBLA Lima

PSEUDOSTREBLA Lima, 1921:23. Type-species, *ribciroi* Lima (orig. des.). Refs. — Jobling, 1939b:268-273; Wenzel, Tipton & Kiewlicz, 1966:582.

GREENWELLI Wenzel, 1966:582, fig. 118A, C (♂ terminalia; hind tibia). Type-locality: Panama, San Blas, Armila, ex *Tonatia minuta*. Distr. — Panama (San Blas).

RIBEIROI Lima, 1921:23, pl. 2, fig. 4 (wing). Type-locality: Brazil, Mato Grosso, ex *Tonatia amblyotis* (= *silvicola*). Distr. — Panama (San Blas), Colombia (Antioquia, Bolívar, Meta), Ecuador (Oriente), Peru (Loreto), Bolivia (Cochabamba), Brazil (Mato Grosso, Pará). Refs. — Jobling, 1939b:271 (redescr.), fig. 1 (dorsum; labium; venter of thorax); Wenzel, Tipton & Kiewlicz, 1966:582, figs. 117, 118B, D (Jobling's 1939b figs., plus ♂ terminalia, hind tibia).

Genus STIZOSTREBLA Jobling

STIZOSTREBLA Jobling, 1939b:273. Type-species, *longirostris* Jobling (orig. des.).

Ref. — Wenzel, Tipton & Kiewlicz, 1966:586.

LONGIROSTRIS Jobling, 1939b:273, fig. 2A-C (dorsum; labium; venter of head, thorax). Type-locality: Brazil, Rio de Janeiro, ex unknown bat. Distr. — Brazil (Rio de Janeiro), Colombia (Meta). Host. — *Tonatia* sp., prob. *silvicola*. Ref. — Wenzel, Tipton & Kiewlicz, 1966:587, fig. 119 (Jobling's 1939b figs. plus ♂ terminalia).

Genus ELDUNNIA Curran

ELDUNNIA Curran, 1934:479. Type-species, *breviceps* Curran (mon.).

Refs. — Curran, 1935:6; Jobling, 1939b:269; Wenzel, Tipton & Kiewlicz, 1966:586.

BREVICEPS Curran, 1934:479. Type-locality: Panama, Panamá, Chilibrillo Caves, ex *Lonchophylla robusta*. Distr. — Panama (Panamá, San Blas). Refs. — Curran, 1935:6, figs. 7, 9 (dorsum, ♀; venter of head); Jobling, 1949b:316 ff., fig. 2B (posteroventral margin of head); Wenzel, Tipton & Kiewlicz, 1966:588, figs. 39C, 120, 121 (Jobling's 1949b fig.; head & mesonotum; ♂ terminalia).

Subfamily STREBLINAE

Genus STREBLA Wiedemann

STREBLA Wiedemann, 1824:19. Type-species, *Hippobosca vespertilionis* Fabricius (mon.) = *wiedemanni* Kolenati, 1856.

Euctenodes Waterhouse, 1879:310. Type-species, *mirabilis* Waterhouse (mon.).

Refs. — Kessel, 1924:406 (*Euctenodes*); Wenzel, Tipton & Kiewliez, 1966:591-627 (revision).

ALTMANI Wenzel, 1966:623, figs. 123G, 137A (postvertex; ♂ mesonotum). Type-locality: Panama, Canal Zone, Gamboa Road, mine shaft at Coeo Plantation, ex *Lonchorhina aurita aurita*. Distr. — Panama (Canal Zone, Colón, Darién), Colombia (Bolívar, Córdoba), Venezuela (Apure, Aragua, Bolívar, Dto. Federal, Trujillo, Yaracuy).

ALVAREZI Wenzel, 1966:625, figs. 123E-F, 137B (postvertex; ♂ mesonotum). Type-locality: Panama, Canal Zone, ½ mi. SE of Empire Range, ex *Micronycteris megalotis*. Distr. — Guatemala (Eseuintla), El Salvador (La Libertad, Usulután), Panama (Boeas del Toro, Canal Zone, Chiriquí, Panamá, San Blas), Colombia (Meta), Venezuela (Amazonas, Miranda, Yaracuy), Brazil (Rio Uaupés).

CAROLLIAE Wenzel, 1966:619, figs. 122A, 136 (head; ♂ mesonotum). Type-locality: Panama, Canal Zone, Fort Davis, ex *Carollia perspicillata azteca*. Distr. — Mexico (Chiapas) to Brazil (Pará, São Paulo), Peru (Junín), Trinidad.

mirabilis, authors (part), not Waterhouse (*Euctenodes*).

CHRISTINAE Wenzel, 1966:606, figs. 44A, 124E, 131A (♀ terminalia; postvertex; ♂ mesonotum). Type-locality: Panama, San Blas, Armila, ex *Phylloderma stenops*. Distr. — Panama (San Blas), Venezuela (Apure, Bolívar, Carabobo, Falcón, Sucre, Yaracuy).

CONSOEUS Wenzel, 1966:600, figs. 124B, 125A, 128 (postvertex; metatibia; ♀ mesonotum). Type-locality: Trinidad, Guanapo Heights, Tamana Cave, ex *Phyllostomus h. hastatus*. Distr. — Colombia (Meta, Tolima), Venezuela (Bolívar, Carabobo, Amazonas, Miranda, Monagas, Sucre, Trujillo, Yaracuy), Suriname (Suriname, Nickerie), Peru (Cuzco, Loreto, Madre de Dios).

mirabilis, authors (part), not Waterhouse (*Euctenodes*).

DIAEMI Wenzel, 1966:599, figs. 124A, 125C, 127A (postvertex; metatibia; ♂ mesonotum). Type-locality: Panama, San Blas, Armila, ex *Diaemus youngi cypselinus*. Distr. — Panama (Bocas del Toro, San Blas), Colombia (Valle de Cauca, Caquetá), Venezuela (Sucre, Orinoco River), Brazil (Pará), Peru (Loreto).

DIPHYLLAE Wenzel, 1966:613, figs. 124C, 133 (postvertex; ♂ mesonotum). Type-locality: Guatemala, Jalapa, San Lorenzo, 4 mi. NE of Volcán de Jumay, ex *Diphylla ecaudata centralis*. Distr. — Mexico (Quintana Roo, Yucatán, San Luis Potosí), Guatemala (Chimaltenango, Jalapa, Santa Rosa, Sololá).

mirabilis, authors (part), not Waterhouse (*Euctenodes*).

- GALINDOI** Wenzel, 1966:604, figs. 123D, 124F, 125D, 130 (postvertex; antenna, frontoelypeus; metatibia; ♂ mesonotum). Type-locality: Panama, Bocas del Toro, Almirante, ex *Tonatia bidens*. Distr. — Panama (Bocas del Toro), Venezuela (Apure, Bolivar, Carabobo, Yaracuy), Trinidad (St. Patrick), Brazil (Pará).
- HERTIGI** Wenzel, 1966:596, figs. 122B, 125F, 127B (head; metatibia; ♂ mesonotum). Type-locality: Panama, Panamá, Chepo Road, ex *Phyllostomus hastatus panamensis*. Distr. — Guatemala (Suchitipequez) to Brazil (Pará), Peru (Huanuco, Piura), Trinidad. Host. — *Phyllostomus h. panamensis*, *P. discolor*.
- mirabilis*, authors (part), not Waterhouse (*Euctenodes*).
- HOOGSTRAALI** Wenzel, 1966:603, figs. 123H, 126, 129B (postvertex; wing; mesonotum; thorax). Type-locality: Panama, San Blás, Armila, ex *Tonatia minuta*. Distr. — Panama (Bocas del Toro, Los Santos, San Blas).
- KOHLISI** Wenzel, 1966:618, fig. 123C (postvertex). Type-locality: Panama, San Blas, Armila, ex *Tonatia silvicola*. Distr. — Panama (Darién, San Blas), Colombia (Antioquia), Venezuela (Apure, Yaracuy).
- MACHADOI** Wenzel, 1966:607, figs. 124D, 131B (postvertex; ♀ mesonotum). Type-locality: Venezuela, Monagas, Caripe, La Guanote, ex *Micronycteris minuta*. Distr. — Venezuela (Bolívar, Miranda, Monagas, Sucre).
- MIRABILIS** (Waterhouse), 1879:310 (*Euctenodes*). Type-locality: Colombia, without host. Distr. — Panama (Bocas del Toro, Canal Zone, Panamá) to Brazil (Pará), Bolivia (Cochabamba) and Peru (Junín, Loreto). Hosts. — *Phyllostomus hastatus panamensis*, *P. elongatus*, *Trachops c. cirrhosus*. Ref. — Wenzel, Tipton & Kiewlicz, 1966:615, figs. 123A, 125E, 134, 135A (postvertex; metatibia; ♂ mesonotum; ♂ terminalia).
- TONATIAE** (Kessel), 1924:411, figs. 7-9 (dorsal, ventral views; head) (*Euctenodes*). Type-locality: Ecuador, Gualaquiza, ex *Tonatia brasiliensis* (= *amblyotis*). Distr. — Ecuador (Gualaquiza), Peru (Cuzco), Venezuela (Bolívar, Carabobo, Trujillo, Yaracuy). Refs. — Pessoa & Guimarães, 1937:255, figs. 10-12 (antenna); Wenzel, Tipton & Kiewlicz, 1966:602, fig. 129A (mesonotum).
- WIEDEMANNI** Kolenati 1856:46 (nom. nov. for *vespertilionis* Fabricius). Distr. — Mexico (Chiapas, Yucatan) to Brazil (Guanabara, Pará, São Paulo) and Peru (Apurímac, Cuzco, Junín, Loreto, Piura, Puno, San Martín). Host. — *Desmodus rotundus*.
- vespertilionis* Fabricius, 1805:339. 6 (*Hippobosca*). Suppressed by I.C.Z.N., 1936:29. Type-locality (Neotype, Wenzel, Tipton & Kiewlicz, 1966:610): Brazil, Pernambuco ex undetermined bat. Ref. — Wenzel, Tipton & Kiewlicz, 1966:609, figs. 123B, 125B, 132 (post-vertex; metatibia; mesonotum).
- mirabilis*, authors (part), not Waterhouse (*Euctenodes*).
- sp., Wenzel, Tipton & Kiewlicz, 1966:617 (? *mirabilis* ex *Trachops cirrhosus*, part), fig. 135B (♂ terminalia). Distr. — Guatemala (Chimaltenango). Host. — *Trachops cirrhosus coffini*.

Genus *PARAEUCTENODES* Pessôa & Guimarães

PARAEUCTENODES Pessôa & Guimarães, 1937:257. Type-species, *longipes* Pessôa & Guimarães (orig. des.).

Ref. — Wenzel, Tipton & Kiewlicz, 1966:627.

LONGIPES Pessôa & Guimarães, 1937:258, figs. 1-4, 7 (dorsal view; head; mesonotum; antenna). Type-locality: Brazil, São Paulo, Ipiranga, ex *Lonchoglossa ecaudata*. Distr. — Guatemala (Altaverapaz), Nicaragua (Zelaya), Venezuela (Bolívar, Dto. Federal, Carabobo, Sucre), Brazil (São Paulo). Fef. — Wenzel, Tipton & Kiewlicz, 1966:627.

logipes, error

Genus *ANASTREBLA* Wenzel

ANASTREBLA Wenzel, 1966:627. Type-species, *modestini* Wenzel (orig. des.).

Strebla, authors, not Wiedemann.

MATTADENI Wenzel, 1966:631, figs. 138B, 139E, F (mesonotum; head; vertex; ♂ terminalia). Type-locality: Panama, Bocas del Toro, upper Río Changena camp, ex *Anoura cultrata*. Distr. — Panama (Bocas del Toro, Darién), Venezuela (Aragua, Bolívar).

MODESTINI Wenzel, 1966:629, figs. 138A, 139C-D (♂ mesonotum; head; vertex; ♂ terminalia). Type-locality: Panama, Chiriquí, Cerro Punta, Casa Lewis, ex *Anoura geoffroyi lasiopyga*. Distr. — Mexico (Sinaloa), Guatemala (Chimaltenango, Jalapa), El Salvador (Santa Ana), Panama (Chiriquí, Los Santos), Venezuela (Bolívar, Dto. Federal, Guarico, Merida, Miranda), Trinidad (Aripo Cave, Tamana Caves).

NYCTERIDIS Wenzel, 1966:629, figs. 139A-B (head, vertex; ♂ terminalia). Type-locality: Panama, Bocas del Toro, Almirante, ex *Lonchophylla robusta*. Distr. — Panama (Bocas del Toro, Darién).

sp., Wenzel, Tipton & Kiewlicz, 1966:628.

vespertilionis Fabricius of Speiser, 1900:38, pl. 4, fig. 1, 2 (dorsum & venter) (*Strebla*). Host. — *Anoura caudifera* (= *Lonchoglossa ecaudata*). Distr. — Colombia (Popayan), Venezuela (Bolívar, Miranda, Dto. Federal, Miranda), Brazil (Santa Catarina). Refs. — Kessel, 1924:413, figs. 1, 5, 6 (dorsum; head; antenna); 1925:24 (part), pl. 3, fig. 20; pl. 4, fig. 24 (1924 figs.).

sp., Wenzel, Tipton & Kiewlicz, 1966:628, fig. 39D (head).

vespertilionis Fabricius of Jobling, 1929: fig. 4 (head); 1949: fig. 3F (head) (*Strebla*). Distr. — Unknown.

Genus *METELASMUS* Coquillett

METELASMUS Coquillett, 1907:292. Type species, *pseudopterus* Coquillett (orig. des.).

Lemosia Pessôa & Galvão, 1936:243. Type-species, *setosa* Pessôa & Galvão (orig. des.).

Ref. — Jobling, 1936:370; Wenzel, Tipton & Kiewlicz, 1966:634.

PSEUDOPTERUS Coquillett, 1907:292, fig. (dorsum). Type-locality: Paraguay, Sapucay, ex *Artibeus lituratus*. Distr. — Mexico (Chiapas, Jalisco), Guatemala (Escuintla), Nicaragua (Matagalpa), Panama (Bocas del Toro, Canal Zone, Darién, Los Santos), Venezuela (Bolívar, Falcón, Guarico, Miranda, Monagas, Sucre), Suriname (Nickerie), Brazil (Pará, Santa Catarina), Paraguay (Sapucay). Refs. — Jobling, 1936:370 (redeser.), figs. 2, 3A-E (dorsal view, ♂; head; thorax; end of abdomen, ♂; halter); Wenzel, Tipton & Kiewlicz, 1966:634, fig. 140 (Jobling's 1936 figs. of head).

setosa Pessôa & Galvão, 1936:244, figs. 1-4 (venter; head; wing; antenna) (*Lemosia*). Type-locality: Brazil, São Paulo, Mogi das Cruzes.

LITERATURE CITED

AUGUSTSON, G. F.

1943. A new parasitic fly from bats (Diptera: Pupipara). *Bull. S. Calif. Acad. Sci.*, 42:52-53, pl. 6.

BEQUAERT, JOSEPH C.

1933. Contribution to the entomology of Yucatan. In, The Peninsula of Yucatan, medical, biological, meteorological, and sociological studies [by George C. Shattuck and collaborators]. *Publ. Carnegie Inst. Washington*, no. 431 (Chap. 29):547-574, figs. 27-30.

1940. Moscas parasitas Pupipara de Colombia y Panama. *Rev. Acad. Colomb. Ciencias Exactas, Físicas y Naturales, Bogotá*, 3 (12): 414-418.

1942. The Diptera Pupipara of Venezuela. *Bol. Ent. Venez.* 1 (4):79-88.

COCKERELL, THEODORE D. A.

1910. A new dipterous parasite of bats. *Can. Ent.*, 42:59-60.

COOPER, KENNETH W.

1941. Reports from recipients of grants from research funds. Grant 484 (1940). An investigation of the aberrant chromosome behavior in the male germ cells of flies parasitic on tropical bats and vultures. (Diptera, section Pupipara). *Yearbook Amer. Philos. Soc.*, 1941:122-127.

COQUILLETT, DANIEL W.

1899. New genera and species of Nycteribiidae and Streblidae. *Can. Ent.*, 31 (11): 333-336.

1907. Notes and descriptions of Hippoboscidae and Streblidae. *Ent. News*, 18:290-292.

CURRAN, CHARLES H.

- 1934a. The Diptera of Kartabo, Bartica District, British Guiana. *Bull. Amer. Mus. Nat. Hist.*, 66:287-532, 55 figs., 1 map [Streblidae, pp. 521-523].

- 1934b. *The families and genera of North American Diptera*. New York, The Ballou Press. 512 pp., illus. [Family Streblidae — The batflies, pp. 477-479, 7 text figs.].



1935. New species of Nycteribiidae and Streblidae (Diptera). *Amer. Mus. Novit.*, no. 765:1-15, 12 figs.
- DYBAS, HENRY S. & RUPERT L. WENZEL.
1947. A new genus of batflies from Guatemala (Diptera Acalypterae: Streblidae). *Fieldiana, Zool.*, 31:149-154, figs. 25-27.
- EDWARDS, FREDERICK W.
1918. Two new Diptera from Trinidad. *Ann. Mag. Nat. Hist.* (9) 1:424-425.
- FABRICIUS, JOHANN
1805. *Systema Antliatorum secundum ordines, genera, species, adjectis synonymis, locis, observationibus, descriptionibus*. Brunsvigae. xiv + 372 pp., index (30 pp.).
- FALCOZ, M. LOUIS
1931. Matériaux pour la connaissance des Diptères Pupipares. I. *Parasitology*, 23:264-269, 5 text figs.
- FERRIS, GORDON F.
1916. Some ectoparasites of bats. *Ent. News*, 27:433-438, 1 text fig., pls. 22-23.
1923. Observations on the larvae of some Diptera Pupipara, with descriptions of a new species of Hippoboscidae. *Parasitology*, 15:54-58, 4 text figs.
- GERVAIS, FRANÇOIS L. P.
1844. *Atlas de Zoologie; ou, collection de 100 planches comprenant 257 figures d'animaux nouveaux ou peu connus classés d'après la méthode de M. de Blainville. Avec une explication par M. Paul Gervais. Ouvrage complémentaire des dictionnaires et des traités d'Histoire naturelle*. Paris, Gernier Baillière. 32 pp., 100 pls.
- GUIMARÃES, LINDOLPHO R.
1938. Sobre as espécies sul Americanas do genero *Trichobius* (Diptera-Streblidae). *Revista Mus. Paulista*, 23(1937): 651-666, pls. 2, 3.
1941. Notas sobre *Streblidae* I. Sobre a verdadeira identidade de *Trichobius dugesi* Towns. 1891. *Papéis Avulsos Depto. Zool., Secret. Agric., São Paulo*, 1:213-222, 4 figs.
1944. Sobre os primeiros estádios de alguns Dipteros pupíparos. *Ibidem*, 6:181-192, 19 figs., 1 plate.
- HOFFMANN, ANITA
1953. Estado actual del conocimiento de los estreblidos Mexicanos (Diptera: Pupipara). *Mem. Cong. Cient. Mexicano*, 7:175-193, 4 plates.
- INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE
1936. Opinion 128. *Nycteribia*, 1796, PUPIPARA and *Spinturnix*, 1826, ACARINE. *Smith. Misc. Collns.*, 73:28-44 (Republished in *Ops. Int. Comm. Zool. Nomencl.*, 1, Section B: 465-508.
- JOBLING, B.
1929. A comparative study of the head and mouthparts in the Streblidae (Diptera: Pupipara). *Parasitology*, 21:417-445, 6 text figs., pls. 18-20.

1936. A revision of the subfamilies of the Streblidae and the genera of the Streblinae (Diptera Acalypterae) including a redescription of *Metelasmus pseudopterus* Coquillett and a description of two new species from Africa. *Ibidem*, 28 (3):355-380, 5 text figs.
1938. A revision of the genus *Trichobius* (Diptera, Acalypterae, Streblidae). *Ibidem*, 30:358-587, 14 text figs.
- 1939a. On some American genera of the Streblidae and their species, with the description of a new species of *Trichobius* (Diptera, Acalypterae). *Ibidem*, 31:486-497, 4 text figs.
- 1939b. A redescription of *Pseudostrebla ribetoi* Costa Lima and the description of a new genus and species of the Streblidae of Brazil. *Arb. Morph. Taxon. Ent.*, 6 (3):268-275, 2 text figs.
- 1939c. On the African Streblidae (Diptera Acalypterae) including the morphology of the genus *Ascodipteron* Adens. and a description of a new species. *Parasitology* 31(2):147-165, 4 figs.
1947. On *Speiseria ambigua* Kessel and *Synthesiostrebla amorphochili* Townsend, with a redescription of the latter (Diptera, Streblidae). *Proc. Roy. Ent. Soc. Lond. (B)* 16:39-42, 1 text fig.
- 1949a. A revision of the genus *Aspidoptera* Coquillett, with some notes on the larva and the puparium of *A. clovisi*, and a new synonym. (Diptera, Streblidae). *Ibidem*, 18:135-144, 4 text figs.
- 1949b. Host-parasite relationship between the American Streblidae and the bats, with a new key to the American genera and a record of the Streblidae from Trinidad, British West Indies (Diptera). *Parasitology*, 39:315-329, 3 text figs.
1951. A record of the Streblidae from the Philippines and other Pacific Islands, including morphology of the abdomen, host-parasite relationship and the geographical distribution, and the description of new species. *Trans. Roy. Ent. Soc. Lond.*, 102 (4): 211-246, 10 text figs., 2 maps.
1952. Description of two new species of *Ascodipteron* from Africa and one species of *Nycteribosca* from Madagascar (Diptera, Streblidae). *Parasitology*, 42(1):6-135, 4 text figs.

KESSEL, QUINTA C.

1914. Notes on the Streblinae, a subfamily of the Streblidae (Diptera Pupipara). *Parasitology*, 16:405-414, 9 text figs. [With introductory note, pp. 405-406, by Hugh Scott].
1925. A synopsis of the Streblidae of the world. *Jour. N. Y. Ent. Soc.*, 33:11-33, pls. 1-4.

KOLENATI, FRIEDRICH A.

1856. *Die Parasiten der Chiropteren*. Brünn, Rohrer, 1856. 51 pp., 4 pls. [also Dresden, Kuntze, 1857].
1863. Beiträge zur Kenntnis der Phthirio-Myiarien. *Hor. Soc. Ent. Ross.* 2:9-109, 11 pls. [Separate, Petersburg, 1863, 101 pp., 15 pls.]

LIMA, A. DA COSTA

1921. Sobre os Streblideos americanos (Diptera: Pupipara). *Arch. Esc. Sup. Agr. Med. Veter.*, Niterói 5:17-14, pl. 1.



MAA, TSING C.

1965. An interim world list of batflies (Diptera: Nycteribiidae and Streblidae). *Jour. Med. Ent.*, 1:377-386.

MACHADO-ALLISON, CARLOS E.

1966. Notas sobre Streblidae (Diptera) de Venezuela I. Las especies del genero *Pterellipsis* Coquillett. *Acta Biol. Venez.*, 5 (4):69-79, figs. 1-11, tables 1-4.

MACQUART, PIERRE J. M.

1852. Notice sur un nouveau genre de Diptères de la famille des Pupipares, tribu des Phthiromydes, sous le nom de *Megistopoda* (*M. pilatei*). *Ann. Soc. Ent. Fr.* (2) 10:331-333, pl. 4, fig. 4.

MATHESON, ROBERT A.

1945. Descriptions of new species, *Paratrachobius anduzci* and *Nycteribosca franclemonti* (Streblidae, Diptera, Pupipara). *Jour. Parasit.*, 31 (3):191-194, 2 text figs.

PERTY, J. A. MAXIMILIAN

1833. Insecta Brasiliensia, pp. 125-224, pls. 25-40, in his *Delectus animalium articulatum, quae in itinere per Brasiliam, collegerunt Dr. J. B. Spix, et C. F. Ph. de Martius, digessit, descripsit, pingenda curavit M. Perty*. Monachii. iii + 44 + 224 pp., 40 col. pls. [1830-34].

PESSÔA, S. B., & AYROSA L. GALVÃO

1936. Novo genero e nova specie de Streblidae (Dipt. Pupipara) parasita do morcego do Brasil. *Revista Ent.*, 6:242-248, text figs. 1-4.
1937. Descrição de uma nova especie do genero *Trichobius* (Diptera-Streblidae). *Folia clin. biol., São Paulo*, 9:1-4.

PESSÔA, S. B., & LINDOLPHO R. GUIMARÃES

1937. Notas sobre Streblidae (Diptera) com a descrição de um novo genero e duas novas especies. *Ann. Fac. Med. São Paulo*, 12 (2):255-267, 17 figs.
1940. Nota sobre Streblídeos (Diptera) de Moreegos de Mato-Grosso, Brazil. *Arquivos Inst. Biol. São Paulo*, 11:421-426, 6 text figs.

RIBEIRO, ALIPIO DE MIRANDA

1907. Alguns dipteros interessantes. *Arch. Mus. Nac., Rio de Janeiro* 14:231-239, pls. xxiii-xxv.

RONDANI, CAMILLO

1878. Muscaria exotica Musei civici Januensis observata et distincta. Fragmentum IV: Hippoboscita exotica non vel minus cognita. *Ann. Mus. Civ. Stor. Nat. Genova*, 12:150-170, 1 text fig.

ROSS, ANTHONY

1959. A redescription of *Trichobius adamsi* Augustson (Diptera: Streblidae). *Wasmann Jour. Biol.*, 17 (1): pp. 69-74, 1 text fig.
1961. Biological studies on bat ectoparasites of the genus *Trichobius* (Diptera: Streblidae) in North America, North of Mexico. *Ibidem*, 19 (2): pp. 229-246.

RUDOW, F.

1871. Einige Pupiparen auf Chiropteren schmarotzend. *Zeitschr. f. d. gesamt. Naturwiss.*, 37 (n.s. vol. 3): 121-124.

1872. On some Pupipara parasitic upon Chiroptera. *Ann. Mag. Nat. Hist.* (4) 9:407-408. (English version of 1871 paper).

SÉGUY, E.

1926. Diptères exotiques peu connus. *Encycl. Ent.*, (E) 11, *Diptera*, 3:192-196, 6 text figs.

SPEISER, P.

1900. Ueber die Strebliden, Fledermausparasiten aus der Gruppe der pupiparen Dipteren. *Archiv. Naturg.*, 66, Bd. I, Hft. 1:31-70, pls. 3-4, 8 text figs.

THEODOR, OSKAR

1957. Parasitic adaptations and host specificity in pupiparous Diptera. In, *First Symposium on host specificity among parasites of vertebrates* [Ernst Mayr, Ed.]. Institut de Zoologie, Université de Neuchâtel. Pp. 50-63.

TOWNSEND, C. H. TYLER

1891. A remarkable new Hippoboscid from Mexico. *Ent. News*, 2:105-106.

1913. A new genus of Streblidae. *Proc. Ent. Soc. Wash.*, 15:98-99.

WATERHOUSE, CHARLES O.

1879. On the affinity of the genus *Polystenes* Gigl., with a description of a new species. *Trans. R. Ent. Soc. Lond.*, 1879, pt. 4:309-312, pls. 9-10.

WENZEL, RUPERT L.

1966. [Descriptions of new genera and species] in Wenzel, Tipton & Kiewlicz (q. v.).

WENZEL, RUPERT L. & THOMAS H. G. AITKEN

1966. [Description of *Neotrichobius stenopterus* n.g., n.sp.], *Ibidem*: 536-540, text figs. 97-99.

WENZEL, RUPERT L., VERNON J. TIPTON & ALICJA KIEWLICZ

1966. The streblid batflies of Panama (Diptera Calypterac: Streblidae). In R. L. Wenzel and V. J. Tipton (Eds.), *Ectoparasites of Panama*: 405:675, text figs. 38-146, tables 6-10.

WIEDEMANN, CHRISTIAN R. W.

1824. *Analecta entomologica ex Museo regio Hafniae maxime congesta*. Kilian. 60 pp., 1 pl.

ZEVE, VICTOR H., & D. E. HOWELL


1962-63. The comparative external morphology of *Trichobius corynorhini*, *T. major*, and *T. sphaeronotus* (Diptera, Streblidae). Part I. The head. *Ann. Ent. Soc. Amer.*, 55 (1962), pp. 685-694, 11 text figs. Part II. The thorax. *Ibidem*, 56 (1963), pp. 2-17, 37 text figs. Part III. The abdomen. *Ibidem*, pp. 127-138, 19 text figs.

INDEX

<i>adamsi</i> Augustson, 1943, <i>Trichobius</i>	100.3
<i>aitkeni</i> Wenzel, 1966, <i>Noctiliostrebla</i>	100.11
<i>altmani</i> Wenzel, 1966, <i>Strebla</i>	100.14
<i>alvarezi</i> Wenzel, 1966, <i>Strebla</i>	100.14
<i>ambigua</i> Kessel, 1925, <i>Speiseria</i>	100.10
<i>amorphochili</i> Townsend, 1913, <i>Synthesiostrebla</i>	100.13
<i>amorphochili</i> Townsend of Jobling, 1939, <i>Speiseria</i>	100.10
<i>Anastrebla</i> Wenzel, 1966	100.16
<i>Anatrichobius</i> Wenzel, 1966	100.6
<i>anduzei</i> (Matheson), 1945 (<i>Paratrichobius</i>), <i>Speiseria</i>	100.10
<i>aranea</i> (Coquillett), 1899 (<i>Pterellipsis</i>), <i>Megistopoda</i>	100.9
<i>Aspidoptera</i> , authors (part)	100.11
<i>Aspidoptera</i> Coquillett, 1894	100.10
<i>bequaerti</i> Wenzel, 1966, <i>Trichobius</i>	100.3
<i>blandus</i> , authors (part), <i>Trichobius</i>	100.4
<i>blandus</i> Curran, 1935, <i>Trichobius</i>	100.3
<i>brennani</i> Wenzel, 1966, <i>Trichobius</i>	100.3
<i>breviceps</i> Curran, 1934, <i>Eldunnia</i>	100.13
<i>buscki</i> , authors (part), <i>Aspidoptera</i>	100.11
<i>buscki</i> Coquillett, 1894, <i>Aspidoptera</i>	100.10
<i>caccus</i> , authors (part), <i>Trichobius</i>	100.4
<i>caccus</i> Edwards, 1918, <i>Trichobius</i>	100.3
<i>carolliae</i> Wenzel, 1966, <i>Strebla</i>	100.14
<i>christinae</i> Wenzel, 1966, <i>Strebla</i>	100.14
<i>clovisi</i> (Pessôa & Guimarães), 1937 (<i>Aspidoptera</i>), <i>Exastinion</i> ...	100.11
<i>coecus</i> , <i>Trichobius</i> , error	100.3
<i>consocius</i> Wenzel, 1966, <i>Strebla</i>	100.14
<i>corynorhini</i> Cockerell, 1910, <i>Trichobius</i>	100.3
<i>costalimai</i> Guimarães, 1937, <i>Trichobius</i>	100.3
<i>coxata</i> Ferris, 1916, <i>Nycterophilia</i>	100.2
<i>coxata</i> Ferris of Hoffmann (part), <i>Nycterophilia</i>	100.2
<i>delatorrei</i> Wenzel, 1966, <i>Aspidoptera</i>	100.10
<i>delicatus</i> (Machado-Allison), 1966 (<i>Pterellipsis</i>), <i>Neotrichobius</i> ...	100.9
<i>desiderata</i> Speiser, 1900, <i>Megistopoda</i>	100.9
<i>diaemi</i> Wenzel, 1966, <i>Strebla</i>	100.14
<i>diphyllac</i> Wenzel, 1966, <i>Strebla</i>	100.14
<i>diphyllac</i> Wenzel, 1966, <i>Trichobius</i>	100.3
<i>dubia</i> , authors (part), <i>Paradyschiria</i>	100.12
<i>dubia</i> (Rudow), 1871 (<i>Lipoptena</i>), <i>Noctiliostrebla</i>	100.11
<i>dubia</i> Rudow of Cooper, 1941, and Jobling, 1949, <i>Paradyschiria</i> ..	100.12
<i>dubia</i> Rudow of Guimarães, 1941, <i>Paradyschiria</i>	100.12
<i>dugesii</i> , authors (part), <i>Trichobius</i>	100.4, 100.5
<i>dugesii</i> Townsend, 1891, <i>Trichobius</i>	100.3
<i>dugesioides</i> Wenzel, 1966, <i>Trichobius</i>	100.4
<i>dunni</i> (Curran), 1935 (<i>Speiseria</i>), <i>Paratrichobius</i>	100.8
<i>dunni</i> Wenzel, 1966, <i>Trichobius</i>	100.4
<i>dybasi</i> Wenzel, 1966, <i>Trichobius</i>	100.4




Eldunnia Curran, 1934	100.13
Euctenodes Waterhouse, 1879	100.14
Exastinion Wenzel, 1966	100.11
fairchildi Wenzel, 1966, Nycterophila	100.2
furmani Wenzel, 1966, Trichobius	100.4
fusca Speiser, 1900, Paradyschiria	100.12
galei Wenzel, 1966, Trichobius	100.4
galindoi Wenzel, 1966, Strebla	100.15
greennelli Wenzel, 1966, Pseudostrebla	100.13
guimaraesi Wenzel, 1966, Mastoptera	100.7
handleyi Wenzel, 1966, Parastrebla	100.12
hertigi Wenzel, 1966, Strebla	100.15
hirsutulus Bequaert, 1933, Trichobius	100.4
hoogstraali Wenzel, 1966, Strebla	100.15
joblingi Wenzel, 1966, Trichobius	100.4
Joblingia Dybas & Wenzel, 1947	100.7
johnsonae Wenzel, 1966, Trichobius	100.4
kecnani Wenzel, 1966, Trichobius	100.4
kesselae Jobling, 1938, Trichobius	100.6
kesseli Guimarães, 1938, Trichobius	100.5
Kesselia Curran, 1934	100.3
kohlsi Wenzel, 1966, Strebla	100.15
Kolenatia Rondani, 1878	100.3
Lemosia Pessôa & Galvão, 1936	100.16
Lepopteryx Speiser, 1900	100.10
Lepopteryx Speiser (part)	100.11
lineata, authors (part), Paradyschiria	100.12
lineata Kessel, 1925, Paradyschiria	100.12
lionycteridis Wenzel, 1966, Trichobius	100.4
Lipoptena, authors (part)	100.10, 100.11
logipes, error, Paraeuctenodes	100.16
lonchophyllae Wenzel, 1966, Trichobius	100.4
longicrus (Ribeiro), 1907 (Trichobius), Paratrachobius	100.8
longicrus Ribeiro of Jobling, 1939, Paratrachobius	100.8
longipes Pessôa & Guimarães, 1937, Paraeuctenodes	100.16
longipes (Rudow), 1871 (Strebla), Trichobius	100.5
longirostris Jobling, 1939, Stizostrebla	100.13
lowei Wenzel, 1966, Paratrachobius	100.8
maai Wenzel, 1966, Noctiliostrebla	100.11
machadoi Wenzel, 1966, Strebla	100.15
macrophylli Wenzel, 1966, Trichobius	100.5
major Coquillett, 1899, Trichobius	100.5
Mastoptera Wenzel, 1966	100.7
mattadeni Wenzel, 1966, Anastrebla	100.16
megastigma (Speiser), 1900 (Lepopteryx), Noctiliostrebla	100.11
megastigma Speiser of Bequaert, 1942, Noctiliostrebla	100.11
megastigma Speiser of Jobling, 1949 (part), Noctiliostrebla	100.11
Megistopoda Macquart, 1852	100.9




<i>Megistopodia</i> , error	100.9
<i>mendezi</i> Wenzel, 1966, <i>Trichobius</i>	100.5
<i>Metelasmus</i> Coquillett, 1907	100.16
<i>minuta</i> (Lima), 1921 (<i>Aspidoptera</i>), <i>Mastoptera</i>	100.7
<i>minuta</i> Lima of authors (part), <i>Mastoptera</i>	100.7
<i>mirabilis</i> , authors (part), <i>Strebla</i>	100.14, 100.15
<i>mirabilis</i> (Waterhouse), 1879 (<i>Euctenodes</i>), <i>Strebla</i>	100.15
<i>mixtus</i> Curran, 1935, <i>Trichobius</i>	100.5
<i>modestini</i> Wenzel, 1966, <i>Anastrebla</i>	100.16
<i>natali</i> Wenzel, 1966, <i>Nycterophila</i>	100.2
<i>Neotrichobius</i> Wenzel & Aitken, 1966	100.9
<i>Noctiliostrebla</i> Wenzel, 1966	100.11
<i>nycteridis</i> Wenzel, 1966, <i>Anastrebla</i>	100.16
<i>Nycterophila</i> , error	100.2
<i>Nycterophila</i> Ferris, 1916	100.2
<i>Nycterophiliinae</i>	100.2
<i>pallidus</i> (Curran), 1934 (<i>Kesselia</i>), <i>Trichobius</i>	100.5
<i>Paradyschiria</i> Speiser, 1900	100.12
<i>Paraectenodes</i> Pessôa & Guimarães, 1937	100.16
<i>parasiticus</i> Gervais, 1844, <i>Trichobius</i>	100.5
<i>Parastrebla</i> Wenzel, 1966	100.12
<i>Paratrichobius</i> Lima, 1921	100.7
<i>parnelli</i> Wenzel, 1966, <i>Nycterophila</i>	100.2
<i>parvula</i> Falcoz, 1931, <i>Paradyschiria</i>	100.12
<i>parvuloides</i> Wenzel, 1966, <i>Paradyschiria</i>	100.12
<i>perspicillatus</i> (Pessôa & Galvão), 1937 (<i>Trichobius</i>), <i>Trichobioides</i>	100.7
<i>phyllostomac</i> Kessel, 1925, <i>Trichobius</i>	100.5
<i>phyllostomatis</i> , authors, <i>Aspidoptera</i>	100.10
<i>phyllostomatis</i> (Perty), 1833 (<i>Lipoptena</i>), <i>Aspidoptera</i>	100.10
<i>phyllostomatis</i> Perty of Bequaert, 1940 (part), and Kessel, 1925 (part), <i>Exastinion</i>	100.11
<i>pilatci</i> Macquart, 1852, <i>Megistopoda</i>	100.9
<i>proxima</i> (Séguy), 1926 (<i>Pterellipsis</i>), <i>Megistopoda</i>	100.9
<i>pseudopterus</i> Coquillett, 1907, <i>Metelasmus</i>	100.17
<i>Pseudostrebla</i> Lima, 1921	100.13
<i>pseudotruncatus</i> Jobling, 1939, <i>Trichobius</i>	100.6
<i>Pterellipsis</i> Coquillett, 1899	100.9
<i>quadrisetosus</i> Kessel, 1925, <i>Trichobius</i>	100.3
<i>ribeiroi</i> Lima, 1921, <i>Pseudostrebla</i>	100.13
<i>salvini</i> Wenzel, 1966, <i>Paratrichobius</i>	100.8
<i>sanchezi</i> Wenzel, 1966, <i>Paratrichobius</i>	100.8
<i>schmidt</i> Dybas & Wenzel, 1947, <i>Joblingia</i>	100.7
<i>scorzai</i> Wenzel, 1966, <i>Anatrichobius</i>	100.6
<i>setosa</i> (Pessôa & Galvão), 1936 (<i>Lemosia</i>), <i>Metelasmus</i>	100.17
sp., Wenzel, Tipton & Kiewlicz, 1966, <i>Anastrebla</i>	100.16
sp., Wenzel, Tipton & Kiewlicz, 1966, <i>Paratrichobius</i>	100.8
sp., Wenzel, Tipton & Kiewlicz, 1966, <i>Strebla</i>	100.15
sp., <i>Trichobius</i>	100.6
sp. A, Wenzel, Tipton & Kiewlicz, 1966, <i>Paratrichobius</i>	100.8



sp. B, Wenzel, Tipton & Kiewlicz, 1966, <i>Paratrachobius</i>	100.8
sparsus Kessel, 1925, <i>Trachobius</i>	100.6
Speiseria Kessel, 1925	100.10
<i>Speisseria</i> , error	100.10
sphaeronotus Jobling, 1939, <i>Trachobius</i>	100.6
stenopterus Wenzel & Aitken, 1966, <i>Neotrachobius</i>	100.9
Stizostrebla Jobling, 1939	100.13
<i>Strebla</i> , authors	100.16
<i>Strebla</i> , authors (part)	100.2
Strebla Wiedemann, 1824	100.14
Streblinae	100.14
<i>Synthesiostrebla</i> Townsend of Jobling, 1939	100.10
<i>Synthesiostrebla</i> Townsend, 1913	100.12
theodori Wenzel, 1966, <i>Megistopoda</i>	100.9
tonatae (Kessel), 1924 (<i>Euctenodes</i>), <i>Strebla</i>	100.15
traubi Wenzel, 1966, <i>Noctiliostrebla</i>	100.11
<i>Trachobia</i> , error	100.2
Trachobiinae	100.2
Trachobioides Wenzel, 1966	100.7
Trachobius Gervais, 1844	100.2
Trachobius Townsend, 1891	100.3
truncatus Kessel, 1925, <i>Trachobius</i>	100.6
uniformis, authors (part), <i>Trachobius</i>	100.5
uniformis Curran, 1935, <i>Trachobius</i>	100.6
uniformis Curran (part), <i>Trachobius</i>	100.5
uniformis Curran of Jobling, 1938 (part), <i>Trachobius</i>	100.5
urodermae Wenzel, 1966, <i>Trachobius</i>	100.6
vampyropis Wenzel, 1966, <i>Trachobius</i>	100.6
vespertilionis (Fabricius), 1805 (<i>Hippobosca</i>), <i>Strebla</i>	100.15
vespertilionis Fabricius of Jobling, 1929, <i>Anastrebla</i>	100.16
vespertilionis Fabricius of Speiser, 1900, <i>Anastrebla</i>	100.16
wiedemanni Kolenati, 1863 (<i>Strebla</i>), <i>Trachobius</i> (part)	100.6
wiedemanni Kolenati, 1856, <i>Strebla</i>	100.15
yunkeri Wenzel, 1966, <i>Trachobius</i>	100.6









DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

101

FAMILY NYCTERIBIIDAE

LINDOLPHO R. GUIMARÃES

*Departamento de Zoologia, Secretaria da Agricultura
São Paulo, Brazil*

Members of this family are wingless, spider-like flies, and obligatory blood-sucking ectoparasites of bats. As occurs with the two preceding families, all species are larviparous, the larva pupating immediately after it is laid. Practically nothing is known about their biology, and this is especially true with the New World species. Hase (1931) gives some facts on the behaviour of *Basilia myotis* Curran. Guimarães & d'Andretta (1956) give data on the morphology, host-parasite relationships, and geographic distribution of American species, and keys, descriptions, and figures for their identification. American species are chiefly found on bats of the family Vespertilionidae. Theodor (1967) has recently contributed a complete revision of World species.

Genus BASILIA Ribeiro

BASILIA Ribeiro, 1903:177. Type-species, *ferruginea* Ribeiro (orig. des., as gen. n., sp. n.).

Pseudelytromyia Ribeiro, 1907:233. Type-species, *speiseri* Ribeiro (mon.).

Guimaraesia Schuurmans Stekhoven, 1951:109. Type-species, *guimaraesi* Schuurmans Stekhoven (orig. des.).

ANCEPS Guimarães & d'Andretta, 1956:113, figs. 188-192. Type-locality: Colombia, Caquetá, La Tagua, Tres Troncos, Río Caquetá. Distr. — Panama, Colombia, Peru.

ANOMALA Guimarães & d'Andretta, 1956:67, figs. 83, 95-100. Type-locality: Mexico, Chiapas, Huehuetán. Distr. — Mexico, Guatemala.

ANTROZOI (Townsend), 1893:79 (*Nycteribia*). Type-locality: U.S.A., New Mexico, Las Cruces. Distr. — U.S.A., Mexico.

BELLARDII (Rondani), 1878:152 (*Nycteribia*). Type-locality: Mexico, Veracruz, Córdoba (lectotype selected by Guimarães & d'Andretta, 1956:56). Distr. — Mexico.

101.1

- mexicana* Bigot, 1885:245 (*Nycteribia*). Type-locality: unknown [Mexico].
- BEQUAERTI Guimarães & d'Andretta, 1956:37, figs. 18-24, 56, 57. Type-locality: Paraguay, Paraguari, Sapueay. Distr. — Venezuela, Colombia, Paraguay.
- CARTERI Scott, 1936:498, figs. 9-10. Type-locality: Paraguay, Mathlawaiya. Distr. — Bolivia, Paraguay, Brazil, Argentina.
- romanaei* Del Ponte, 1944:119, fig. Type-locality: Argentina, Depto. Pellegrini, La Fragua.
- CONSTRICTA Guimarães & d'Andretta, 1956:80, figs. 86, 112-117, 123, 130, 131, 148. Type-locality: Ecuador, Gualaquiza. Distr. — Ecuador, Peru, Colombia, Venezuela.
- COSTARICENSIS Guimarães & d'Andretta, 1956:106, figs. 146, 164, 169-174, 178. Type-locality: Costa Rica, Guanacaste, Las Cañas. Distr. — Costa Rica.
- CURRANI Guimarães, 1943:257, figs. 1-2. Type-locality: Brazil, São Paulo, São Paulo. Distr. — Brazil, Argentina.
- DUBIA Guimarães & d'Andretta, 1956:102, figs. 150, 154, 160, 167, 168, 177. Type-locality: Brazil, Mato Grosso, Chavantina. Distr. — Peru, Brazil.
- DUNNI Curran, 1935:3, figs. 1-2. Type-locality: Panama, Santa Rosa. Distr. — Panama.
- FERRISI Schuurmans Stekhoven, 1931:217 (nom. nov. for *speiseri* Ribeiro of Ferris). Type-locality: Costa Rica, Sipurio. Distr. — Costa Rica.
- speiseri* Ribeiro of Ferris, 1924:198, pl. 3, misident.
- ferrisi* Scott, 1936:502 (nom. nov. for *speiseri* Ribeiro of Ferris; preocc. Schuurmans Stekhoven, 1931).
- FERRUGINEA Ribeiro, 1903:179, pl. 1, figs. 1-2. Type-locality: Brazil, Minas Gerais, Santana do Sapé. Distr. — Cuba, Panama, Brazil, Paraguay.
- FLAVA (Weyenbergh), 1881:194 (*Nycteribia*). Type-locality: Argentina, Córdoba. Distr. — Argentina.
- FORCIPATA Ferris, 1924:196, figs. 1a-c. Type-locality: U.S.A., California, Los Angeles Co., Covina. Distr. — Canada, U.S.A., Mexico.
- calverti* Fox & Stabler, 1953:22, figs. 1-2. Type-locality: U.S.A., Colorado, Colorado Springs.
- GUIMARAESI (Schuurmans Stekhoven), 1951:113 (*Guimaraesia*; nom. nov. for *bellardii* Rondani of Guimarães). Type-locality: Brazil, São Paulo, São Paulo, Cantarcira. Distr. — Brazil.
- bellardii* Rondani of Guimarães, 1946:62, figs. 73-80, misident.
- HANDLEYI Guimarães, 1966:394, figs. 35A-C. Type-locality: Panama, San Blás, Armila. Distr. — Panama.

- HUGHSEOTTI Guimarães, 1946:77, figs. 94-97. Type-locality: Brazil. Rio Grande do Sul, São Lourenço. Distr. — Brazil.
- JUQUIENSIS Guimarães, 1946:73, figs. 89-93. Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- MIRANDARIBEIROI Guimarães, 1942:146, fig. 1. Type-locality: Brazil, São Paulo, São Paulo. Distr. — Brazil.
- MIMONI Theodor & Peterson, 1964:107, figs. 1-6. Type-locality: Peru, Loreto, Rio Yavari. Distr. — Peru.
- MYOTIS Curran, 1935:3, figs. 3-5. Type-locality: Panama, Tapia. Distr. — Guatemala, Panama, Colombia, Peru, Venezuela, Guyana. Ref. — Hase, 1931 (behaviour).
- bellardii* Rondani of Hase, 1931:220, figs. 1-17, misident.
- NEAMERICANA Schuurmans Stekhoven, 1951:102, figs. 1-3. Type-locality: Argentina, La Rioja, El Tieson. Distr. — Argentina.
- ORTIZI Machado-Allison, 1964:455, figs. 1-6. Type-locality: Venezuela, Bolívar, Serranía de Nuria. Distr. — Venezuela.
- PERUVIA Guimarães & d'Andretta, 1956:84, figs. 87, 118-122, 125, 128, 149. Type-locality: Peru, Cuzco, Balceadero, Río Nushiseaya. Distr. — Peru.
- PIZONYEHUS Scott, 1939:168, pl. 16. Type-locality: Mexico, Ángel de la Guardia I., Gulf of California. Distr. — Mexico.
- PLAUMANNI Scott, 1940:58, figs. 1-2. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil, Argentina, Paraguay. Ref. — Guimarães, 1946:38, figs. 31, 39, 40, 44 (as *mexicana* ssp.).
- mexicana* Bigot of Guimarães, 1946:28, figs. 21-29, 32-38, 43, misident.
- RONDANI Guimarães & d'Andretta, 1956:50, figs. 39-40, 42, 44, 47-51, 53, 55, 79. Type-locality: Guatemala, Esequintla, Finca los Areos. Distr. — U.S.A., Mexico, Guatemala, Honduras.
- RUGOSA Schuurmans Stekhoven, 1942:96, figs. 1-6 (1952:96). Type-locality: Peru, Ayacucho, Sivia. Distr. — Peru.
- SILVAE (Brèthes), 1913:297, unnumbered text-fig. (*Cyclopodia*). Type-locality: Chile, Santiago. Distr. — Chile.
- SPEISERI (Ribeiro), 1907:233, pl. 23, 24, figs. 2-4 (*Pseudelytromyia*). Type-locality: Brazil, Guanabara, Rio de Janeiro, Quinta da Boa Vista. Distr. — Brazil, Paraguay.
- TIPTONI Guimarães, 1966:396, figs. 36A-C. Type-locality: Panama, Bocas del Toro near Changuinola. Distr. — Panama.
- TRAVASSOSI Guimarães, 1938:183, pl. 1, figs. 1-3. Type-locality: Brazil, Pernambuco, Tapera. Distr. — Brazil.
- WENZELI Guimarães & d'Andretta, 1956:42, figs. 25-33, 57, 80. Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Panama, Venezuela, Colombia.



Genus *HERSHKOVITZIA* Guimarães & d'Andretta

- HERSHKOVITZIA* Guimarães & d'Andretta, 1956:127. Type-species, *primitiva* Guimarães & d'Andretta (orig. des.).
- COECA* Theodor, 1967:346, figs. 578-582. Type-locality: unknown (probably from Central or South America).
- INAEQUALIS* Theodor, 1967:348, figs. 585-594. Type-locality: Peru, Loreto. Distr. — Peru.
- PRIMITIVA* Guimarães & d'Andretta, 1956:130, figs. 213-232. Type-locality: Colombia, Bolívar, Socorré, Río Sinú. Distr. — Colombia.

LITERATURE CITED

EIGOT, J. M. F.

1885. Diptères nouveaux ou peu connus. 27^e partie, XXXV: Famille des Anomalocerati (mihi) (Coriacea, Pupipara, Nycteribiidae, auctor). *Ann. Soc. Ent. France* (6)5:225-246.

BRÊTHES, J.

1913. Une nouvelle espèce de diptère pupipare du Chile. *Bol. Mus. Nac. Chile* 5(2):297-299, fig.

CURRAN, C. H.

1935. New species of Nycteribiidae and Streblidae (Diptera). *Amer. Mus. Nov.* 765:1-11, 2 figs.

DEL PONTE, E.

1944. *Basilia romanaï*, nobis (Diptera, Nycteribiidae) de la Argentina. *An. Inst. Med. Regional, Tucumán* 1(1):117-128, 3 figs.

FERRIS, G. F.

1924. The New World Nycteribiidae (Diptera Pupipara). *Ent. News* 35(6):191-199, fig. 1, pl. 3.

FOX, R. M. & R. M. STABLER

1953. *Basilia calverti* n. sp. (Diptera: Nycteribiidae) from the interior long-legged bat. *Jour. Parasit.* 39(1):22-27, 2 figs.

GUIMARÃES, L. R.

1938. Sobre uma nova espécie de Nycteribiidae (Diptera-Pupipara). *Livro Jubilar Prof. L. Travassos*, Rio de Janeiro, pp. 183-184, pl. 1, figs. 1-3.
1942. Nova espécie do gênero *Basilia* (Nycteribiidae-Diptera) do Brasil. *Papéis Avulsos Dep. Zool. S. Paulo* 2:145-149, fig. 1.
1943. Mais uma nova espécie sul-americana de Nycteribiidae (Diptera). *Papéis Avulsos Dep. Zool. S. Paulo* 3:257-260, 1 fig.
1946. Revisão das espécies sul-americanas do gênero *Basilia* (Diptera — Nycteribiidae). *Arg. Zool.*, São Paulo 5:1-87, figs. 1-98, 1 map.

1966. Nycteribiid batflies from Panama (Diptera: Nycteribiidae), pp. 393-404, figs. 35-37, in R. L. Wenzel & V. J. Tipton, eds., *Ectoparasites of Panama*, xii + 861 pp., 154 figs. Chicago.

GUIMARÃES, L. R. & M. A. V. D'ANDRETTA

1956. Sinopse dos Nycteribiidae (Diptera) do Novo Mundo. *Arq. Zool., São Paulo* 10(1):1-184, 232 figs., 3 tables, 5 maps.

HASE, A.

1931. Über die Lebensgewohnheiten einer Fledermausfliege in Venezuela: *Basilia bellardii* Rondani (Fam. Nycteribiidae-Diptera pupipara). *Zeitschr. Parasitenk.* 3(2):220-257, 7 figs.

MACHADO-ALLISON, C. E.

- (1964). Un nuevo Nycteribiidae (Diptera) de Venezuela. *Acta Zool. Venez.* 3(29):455-459, 6 figs. [Printed 31 Dec., 1963; distributed in 1964].

RIBEIRO, A. M.

1903. *Basilia ferruginea* gênero novo e espécie nova da família das nycteribias. *Arch. Mus. Nac.*, Rio de Janeiro 12:175-179, pl. 1.
1907. Alguns dípteros interessantes. *Arch. Mus. Nac.*, Rio de Janeiro 14:231-239, pls. 23-25.

RONDANI, C.

1878. *Muscaria exotica* Musei Civici Januensis observata et distincta. *Ann. Mus. Civ. Stor. Nat. Genova* 12:150-169.

SCHUURMANS STEKHOVEN, J. H., JR.


1931. Eine seltene, ungenügend beschriebene *Basilia* Art (Diptera Pupipara) aus Venezuela. *Zeitschr. Parasitenk.* 3(2):205-219, figs. 1-6.
1942. Pupiparen (Dipt.). *Beiträge zur Fauna Perus* 3:91-101, figs. 1-6 (Reprinted 1952).
1951. Algunas especies del genero "*Basilia*" Ribeiro y creación del nuevo genero "*Guimarãesia*". *Acta Zool. Lilloana* 12:551-561, figs. 1-3.

SCOTT, H.

1936. Descriptions and records of Nycteribiidae (Diptera Pupipara), with a discussion of the genus *Basilia*. *Linnean Soc. Jour. Zool.* 39(nº 267):479-505, 11 figs.
1939. A new species of Nycteribiidae (Diptera Pupipara) from islands in the Gulf of California. *Allan Hancock Pacific Exp.* 2(10):167-170, pl. 16, figs. 1-3.
1940. Nycteribiidae from southern Brazil (Diptera). *Proc. Ent. Soc. London* 9(4):57-61, 4 figs.

THEODOR, O.

1967. *An illustrated catalogue of the Rothschild collection of Nycteribiidae (Diptera) in the British Museum (Natural History), with keys and short descriptions for the identification of sub-families, genera, species and subspecies*, viii + 506 pp., 898 figs., 6 maps, 5 pls. London.



THEODOR, O. & B. V. PETERSON

1964. On some new species of Nycteribiidae (Diptera: Pupipara).
Great Basin Nat. 24:107-115, 13 figs.

TOWNSEND, C. H. T.


1893. A nycteribiid from a New Mexico bat. *Jour. New York Ent. Soc.* 1(2):79-80.

WEYENBERGH, D. H.

1881. Dos nuevas especies del grupo de los dípteros pupíparos. *An. Soc. Cient. Argent.* 11:193-200.

INDEX

anceps Guimarães & d'Andretta, 1956, Basília	101.1
anomala Guimarães & d'Andretta, 1956, Basília	101.1
antrozoi (Townsend), 1893 (Nycteribia), Basília	101.1
Basília Ribeiro, 1903	101.1
bellardii (Rondani), 1878 (Nycteribia), Basília	101.1
bellardii Rondani of Guimarães, 1946, Basília	101.2
bellardii Rondani of Hase, 1931, Basília	101.3
bequaerti Guimarães & d'Andretta, 1956, Basília	101.2
calverti Fox & Stabler, 1953, Basília	101.2
carteri Scott, 1936, Basília	101.2
coeca Theodor, 1967, Hershkovitzia	101.4
constricta Guimarães & d'Andretta, 1956, Basília	101.2
costaricensis Guimarães & d'Andretta, 1956, Basília	101.2
currani Guimarães, 1943, Basília	101.2
dubia Guimarães & d'Andretta, 1956, Basília	101.2
dunni Curran, 1935, Basília	101.2
ferrisi Schuurmans Stekhoven, 1931, Basília	101.2
ferrisi Scott, 1936, Basília	101.2
ferruginea Ribeiro, 1903, Basília	101.2
flava (Weyenbergh), 1881 (Nycteribia), Basília	101.2
forcipata Ferris, 1942, Basília	101.2
guimaraesi (Schuurmans Stekhoven), 1951 (Guimaraesia), Basília	101.2
Guimaraesia Schuurmans Stekhoven, 1951	101.1
handleyi Guimarães, 1966, Basília	101.2
Hershkovitzia Guimarães & d'Andretta, 1956	101.4
hughscotti Guimarães, 1946, Basília	101.3
inaequalis Theodor, 1967, Hershkovitzia	101.4
juquiensis Guimarães, 1946, Basília	101.3



<i>mexicana</i> (Bigot), 1885 (Nycteribia), Basília	101.2
<i>mexicana</i> Bigot of Guimarães, 1946, Basília	101.3
<i>mirandaribeiroi</i> Guimarães, 1942, Basília	101.3
<i>mimoni</i> Theodor & Peterson, 1964, Basília	101.3
<i>myotis</i> Curran, 1935, Basília	101.3
 <i>neamericana</i> Schuurmans Stekhoven, 1951, Basília	101.3
 <i>ortizi</i> Machado-Allison, 1964, Basília	101.3
 <i>peruvia</i> Guimarães & d'Andretta, 1956, Basília	101.3
<i>pizonychus</i> Scott, 1939, Basília	101.3
<i>plaumanni</i> Scott, 1940, Basília	101.3
<i>primitiva</i> Guimarães & d'Andretta, 1956, Hershkovitzia	101.4
<i>Pseudelytromyia</i> Ribeiro, 1907	101.1
 <i>rondanii</i> Guimarães & d'Andretta, 1956, Basília	101.3
<i>romanai</i> Del Ponte, 1944, Basília	101.2
<i>rugosa</i> Schuurmans Stekhoven, 1942, Basília	101.3
 <i>silvae</i> (Brèthes), 1913 (Cyclopodia), Basília	101.3
<i>speiseri</i> (Ribeiro), 1907 (Pseudelytromyia), Basília	101.3
<i>speiseri</i> Ribeiro of Ferris, 1924, Basília	101.3
 <i>tiptoni</i> Guimarães, 1966, Basília	101.3
<i>travassosi</i> Guimarães, 1938, Basília	101.3
 <i>wenzeli</i> Guimarães & d'Andretta, 1956, Basília	101.3





MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO

(Formerly Departamento de Zoologia, Secretaria da Agricultura, São Paulo)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

102

FAMILY CALLIPHORIDAE¹

MAURICE T. JAMES


*Department of Entomology, Washington
State University, Pullman, Washington*

The family Calliphoridae is of medical, veterinary, and sanitary importance because of the facts that larvae may breed in decaying materials or may cause myiasis in man or animals, and that the adults may transmit pathogens mechanically from putrefactive material to human food.

Some species, such as *Phaenicia sericata*, *P. cuprina*, and *Calliphora vicina* are found over a large part of the world; others are quite narrowly restricted in their distribution, for example, the species of *Phaenicia* (*Viridinsula*) that occur only on the Galápagos Islands. Four subfamilies occur in the Neotropical Region; two of these, the Mesembrinellinae and the Toxotarsinae, are found nowhere else in the World. The Polleniinae, which occur in the Nearctic Region, are, so far, not known from the Neotropical, though the widespread *Pollenia rudis* (Fabricius) and some of the species of *Melanodexia* that occur in California may ultimately be recorded from Mexico. It is also quite likely, when the fauna of northern Mexico becomes better known, that some of the Calliphorinae and, possibly, Chrysomyinae that are known from the southwestern United States, may extend into Mexico.

The work of Shannon (1926), though now outdated, may serve as a starting point for the study of the Neotropical fauna. Hall's (1948) monograph of the blow flies of North America is the standard guide for identifying these insects in Mexico, Central America, and northwestern South America, as well as in the Nearctic Region. Other works of more general value are those of Smart (1937) on the fauna of Patagonia and South Chile, and Aldrich (1922, 1925) and Mello (1967) on the Mesembrinellinae. References dealing with more restricted groups are cited in the proper places. Our knowledge of the larvae of the Neotropical species is very poor; Hall (1948) will serve as a partial guide to these.

¹Scientific paper 3377, College of Agriculture, Washington State University. Work was conducted under project 9043.



An interesting fossil record from South America is the discovery of a number of empty puparia of an undetermined species embedded in the skull of a giant ground sloth, *Megatherium* probably *medinae* Philippi, excavated south of Pica, Tarapacá, Chile (James, in press).

A distinct difficulty involved in cataloguing the Calliphoridae of the Neotropical Region arises from the question of the identity of so many species that have been described under such generic names as *Lucilia*, *Calliphora*, *Somomyia*, and *Musca*. The types of many of these species will have to be examined in order to determine the specific identity, and some of these are lost or in so poor a condition as to be indeterminate. Many metallic muscids have been referred to calliphorid genera, and it is possible that some overlooked Neotropical calliphorids described as *Musca*, *Pyrellia*, etc., may belong to the Calliphoridae.

Subfamily MESEMBRINELLINAE

Genus ALBUQUERQUEA Mello *

- ALBUQUERQUEA Mello, 1967:10. Type-species, *latifrons* Mello (orig. des.).
LATIFRONS Mello, 1967:10. Type-locality: Brazil, Rio de Janeiro, Petrópolis.
Distr. — Brazil (Rio de Janeiro).

Genus EUMESEMBRINELLA Townsend

- EUMESEMBRINELLA Townsend, 1931:69. Type-species, *Musca quadrilincata* Fabricius (mon.).
DORSIMACULA (Aldrich), 1922:20 (*Mesembrinella*). Type-locality: "Brazil".
Distr. — Brazil.
INCOMPLETA (Curran), 1934b:470 (*Mesembrinella*). Type-locality: Guyana (British Guiana), Kartabo. Distr. — Guyana. Description indicates that this species probably is to be placed here.
LANEI Mello, 1967:26. Type-locality: Brazil, Amapá, Serra do Navio.
Distr. — Brazil (Amapá).
PAUCISETA (Aldrich), 1922:18 (*Mesembrinella*). Type-locality: Brazil, Pará. Distr. — Brazil (Pará, Bahia, Espírito Santo, Rio de Janeiro).
QUADRILINEATA (Fabricius), 1805:286 (*Musca*). Type-locality: "South America". Distr. — Venezuela, Guyana, Brazil (Pará, Amazonas, Mato Grosso).
flavipennis Macquart, 1843:391 (separate, p. 134) (*Ochromyia*).
Type-locality: Brazil, Pará.
RANDA (Walker), 1849:852 (*Dexia*). Type-locality: "Brazil". Distr. — Guyana, French Guiana, Surinam, Brazil (Pará).
quadrilineata Fabricius of Surcouf, 1919.

* The author's name was incorrectly spelled "Melo" in his publication on the Mesembrinellinae (Mello, 1967).

Genus HUASCAROMUSCA Townsend

- HUASCAROMUSCA Townsend, 1918:155. Type-species, *cruciata* Townsend (mon.) = *xanthorrhina* (Bigot).
- Promesembrinella* Hall, 1948:65. Type-species, *Mesembrinella semiflava* Aldrich (orig. des.).
- AENEIVENTRIS (Wiedemann), 1830:376 (*Dexia*). Type-locality: "Brazil". Distr. — Panama, Colombia, Peru, Brazil (Guanabara).
tibialis Aldrich, 1922:16 (*Mesembrinella*). Type-locality: Panama, Porto Bello. N. SYN.
- GRAJAUENSIS Mello, 1967:43. Type-locality: Brazil, Guanabara, Rio de Janeiro, Grajaú. Distr. — Brazil (Guanabara).
- NIGRIFRONS (Bigot), 1878:39 (*Ochromyia*). Type-locality: "Brazil". Distr. — Ecuador, Peru, Brazil (Rio de Janeiro, Espírito Santo).
purpurata Aldrich, 1922:16 (*Mesembrinella*). Type-locality: Brazil, Espírito Santo.
- SEMIFLAVA (Aldrich), 1925:14 (*Mesembrinella*). Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica.
- UNISETA (Aldrich), 1925:13 (*Mesembrinella*). Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica.
- VOGELSANGI Mello, 1967:46. Type-locality: Venezuela, Aragua. Distr. — Venezuela.
- XANTHORRHINA (Bigot), 1888:clxxx (*Calliphora*). Type-locality: "Mexico". Distr. — Southern Mexico to Peru and Venezuela.
chrysorhea Brauer, 1895:596 (*Mesembrinella*). Type-locality: "Venezuela".
cruciata Townsend, 1918:156. Type-locality: Peru, Jaen, Huascaray Ridge.
bequaerti Séguy, 1925a:195 (*Mesembrinella*). Type-locality: "Peru".
decrepita Séguy, 1925a:195 (*Mesembrinella*). Type-locality: "Peru".

Genus LANEELLA Mello

- LANEELLA Mello, 1967:3. Type-species, *Mesembrinella brunripes* Surcouf (orig. des.).
- BRUNNIPES (Surcouf), 1919:78 (*Mesembrinella*). Type-locality: "Bolivia". Distr. — Bolivia, Surinam, Brazil (Guanabara, Espírito Santo, São Paulo).

Genus MESEMBRINELLA Giglio-Tos


- MESEMBRINELLA Giglio-Tos, 1893:4. Type-species, *quadrilineata* Fabricius (orig. des., misident.) = *bellardiana* Aldrich.
- Mesembrinella*, subg. *Mesembolia* Aldrich, 1925:10. Type-species, *bellardiana* Aldrich (orig. des.).
- ABACA (Hall), 1948:68 (*Huascaromusca*). Type-locality: Panama, Canal Zone, Barro Colorado Island. Distr. — Guatemala to Panama.



- APOLINARIS Séguy, 1925a:196. Type-locality: "Colombia". Distr. — Colombia.
- BATESI Aldrich, 1922:15. Type-locality: Brazil, "Amazon River". Distr. — Peru, Brazil (Bahia, Espírito Santo, Rio de Janeiro).
- BELLARDIANA Aldrich, 1922:21. Type-locality: "Brazil". Distr. — Mexico, Ecuador, Guyana, Bolivia, Brazil (Amazonas, Pará, Mato Grosso, Goiás, Minas Gerais, Espírito Santo, Guanabara, São Paulo, Santa Catarina), Paraguay.
- quadrilineata* Fabricius of Giglio-Tos, 1893:4.
- BICOLOR (Fabricius), 1805:291 (*Musca*). Type-locality: "South America". Distr. — Mexico (north to Nayarit), to Brazil (south to São Paulo) and Paraguay.
- socors* Walker, 1861:311 (*Calliphora*). Type-locality: "Mexico".
- bicolor* Giglio-Tos, 1893:4. Type-localities: Mexico: Chinantla, Oaxaca, and Playa Vicente (cotypes).
- aeneiventris* authors, not Wiedemann.
- CONFUSA Séguy, 1925a:196. Type-locality: "Peru". Distr. — Peru.
- CYANEICINETA (Surcouf), 1919:69 (*Ochromyia*). Type-locality: "Brazil". Distr. — Brazil.
- FACIALIS Aldrich, 1922:17. Type-locality: Costa Rica, Higuito. Distr. — British Honduras, Guatemala, Canal Zone, Panama.
- FLAVICRURA Aldrich, 1925:16. Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica.
- FORMOSA Aldrich, 1932:25. Type-locality: Guatemala, Pochuta, Santa Emilia. Distr. — Guatemala.
- FULVIFES Aldrich, 1922:22. Type-locality: Brazil, Espírito Santo. Distr. — Brazil (Espírito Santo).
- FUSCICOSTA Séguy, 1925a:195. Type-locality: French Guiana, St-Jean du Maroni. Distr. — French Guiana.
- PEREGRINA Aldrich, 1922:22. Type-locality, Brazil, Espírito Santo. Distr. — Guyana to Brazil (Minas Gerais, Rio de Janeiro, Guanabara, São Paulo).
- PICTIPENNIS Aldrich, 1922:11. Type-locality: Bolivia, Yungas de la Paz. Distr. — Bolivia.
- SEMIHYALINA Mello, 1967:73. Type-locality: Brazil, Espírito Santo, Parque Sooretama. Distr. — Brazil (Espírito Santo, Rio de Janeiro).
- SPICATA Aldrich, 1925:13. Type-locality, Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica.
- TRANSPOSITA Séguy, 1925a:196. Type-locality: (French) "Guiana". Distr. — French Guiana.
- UMBROSA Aldrich, 1922:12. Type-locality, Costa Rica, Tucurrique. Distr. — Costa Rica, Colombia, Ecuador.

Unplaced species

- BENOISTI Séguy, 1925a:196 (*Ochromyia*). Type-locality: "French Guiana".
- FUSCIPENNIS Macquart, 1843:293 (separate, p. 135) (*Ochromyia*). Type-locality, Brazil, Pará. Probably *Mesembrinella*: See Aldrich, 1922:23.



GIGAS Macquart, 1846:324 (separate, p. 196) (*Ochromyia*). Type-locality: said to be Brazil, but probably mislabelled: probably African and may be a bengaline.

PURPURATA Séguéy, 1925a:195 (*Mesembrinella*). Type-locality, "Brazil". Homonym of *purpurata* Aldrich.

Subfamily CHRYSOMYINAE

Tribe CHRYSOMYINI

Genus CALLITROGOPSIS Townsend

CALLITROGOPSIS Townsend, 1935:70. Type-species, *costalis* Townsend (orig. des.). Ref. — Mello, 1965 (revision).

COSTALIS Townsend, 1935:70. Type-locality: Brazil, Pernambuco, Tapera. Distr. — Brazil (Amapá, Pará, Pernambuco, Goiás, Espírito Santo, Rio de Janeiro, São Paulo). Ref. — Mello, 1965 (redescr.).

Genus CHLOROPROCTA Wulp

CHLOROPROCTA Wulp, 1896:296. Type-species, *semiviridis* Wulp (mon.) = *fuscipennis* (Macquart).

FUSCIPENNIS (Macquart), 1851:223 (separate, p. 250) (*Lucilia*). Type-locality: Brazil, Bahia. Distr. — U.S.A. (Texas) to Peru and Brazil (Bahia) chiefly along the Atlantic coast.

semiviridis Wulp, 1896:296. Type-locality: Mexico, Northern Yucatán.

IDIOIDEA (Robineau-Desvoidy), 1830:445 (*Chrysomya*). Type-locality: "Brazil". Distr. — Southern Mexico (Quintana Roo, Yucatán) to Paraguay; Cuba.

purpurea Walker, 1852:337 (*Musca*). Type-locality: "South America".

Genus COCHLIOMYIA Townsend

Callitroga Brauer, 1883:74 [also Brauer & Bergenstamm, 1893:194 (1893:106), in generic synonymy]. Type-species, *Musca macellaria* Fabricius (Hall, 1948:126). Based on a name originally cited in synonymy, hence invalid.

COCHLIOMYIA Townsend, 1915:646. Type-species, *Musca macellaria* Fabricius (orig. des.). Ref. — Garcia, 1952 (revision).

Compsomyia authors, in part, not Rondani.

Paralucilia authors, not Brauer & Bergenstamm.

ALDRICHI Del Ponte, 1938:274. Type-locality: Bahama Islands, San Salvador Is. Distr. — U.S.A. (Florida Keys), Bermuda, Bahamas, Cuba, Puerto Rico, Virgin Islands.

FONTANAI Garcia, 1952:71. Type-locality: Argentina, Capital Federal, Belgrano. Distr. — Argentina (La Plata).



- HOMINIVORAX (Coquerel), 1858:173 (*Lucilia*). Type-locality: "Guiana".
 Distr. — U.S.A. through West Indies and Central and South America to Argentina and Chile. The primary screw-worm.
?infesta Philippi, 1861:513 (*Calliphora*). Type-locality: Chile, Santiago.
?anthropophaga Conil, 1878:71 (*Calliphora*). Type-locality: Argentina, Córdoba.
fulvobarbata Bigot, 1887:598 (*Somomyia*). Type-locality: "Mexico".
homicida Brauer, 1899:525 (*Compsomyia*). Type-locality: French Guiana, Cayenne.
americana Cushing & Patton, 1933:539. Type-locality: U.S.A., Texas, Menard.
macellaria authors, not Fabricius.
- MACELLARIA (Fabricius), 1775:776 (*Musca*). Type-locality: "West Indies".
 Distr. — U.S.A. South to Argentina and Chile; throughout West Indies; Galápagos Islands. The secondary screw-worm fly.
taniaria Wiedemann, 1830:406 (*Musca*), lapsus.
laniaria Wiedemann, 1830:683 (*Musca*; correction of name *taniaria*).
 Type-locality: "Brazil".
affinis Robineau-Desvoidy, 1830:445 (*Chrysomya*). Type-locality: "Brazil".
alia Robineau-Desvoidy, 1830:447 (*Chrysomya*). Type-locality: "Brazil". Sabrosky (1953) says "du nord de la Capit. de St. Paul" (north of the State of São Paulo).
?decora Robineau-Desvoidy, 1830:448 (*Chrysomya*). Type-locality: "Antilles".
hyacinthina Robineau-Desvoidy, 1830:450 (*Chrysomya*). Type-locality: "South America".
lepada Robineau-Desvoidy, 1830:448 (*Chrysomya*). Type-locality: "Brazil". Sabrosky (1953) says "du nord de la Capit. de St. Paul" (north of the State of São Paulo).
placii Robineau-Desvoidy, 1830:448 (*Chrysomya*). Type-locality: "Antilles".
socia Robineau-Desvoidy, 1830:447 (*Chrysomya*). Type-locality: "Brazil". Sabrosky (1953) says "du nord de la Capit. St. Paul" (north of the State of São Paulo).
?tibialis Robineau-Desvoidy, 1830:446 (*Chrysomya*). Type-locality: Jamaica, Port-au-Prince.
viridula Robineau-Desvoidy, 1830:445 (*Chrysomya*). Type-locality: "Brazil". Sabrosky (1953) says "du midi de la Capit. de Goyaz" (southern State of Goiás).
durvillei Macquart, 1843:299 (separate, p. 142) (*Lucilia*). Type-locality: Peru, Payta.
violacea Macquart, 1843:285 (separate, p. 128) (*Calliphora*). Type-locality: Given as "Africa", but Sabrosky (1953) says the label on the type is "Mr. Gay-Chile".
phauda Walker, 1849:896 (*Musca*). Type-locality: Galápagos Islands.
cyanicolor Rondani, 1850:178 (*Lucilia*; nom. nov. for *violacea* Macquart). Type-locality: "Africa" (in error).
rubrifrons Macquart, 1851:223 (separate, p. 250) (*Lucilia*). Type-locality: Argentina, Buenos Aires.

- tibialis* Macquart, 1851:215 (separate, p. 242) (*Calliphora*). Type-locality: Brazil, Rio de Janeiro.
- fasciata* Walker, 1852:337 (*Musca*). Type-locality: "Brazil".
- turbida* Walker, 1852:336 (*Musca*). Type-locality: "Jamaica".
- curvipes* Thomson, 1869:544 (*Lucilia*). Type-locality: Brazil, Rio de Janeiro.
- picicrus* Thomson, 1869:543 (*Lucilia*). Type-locality: "Panama".
- porticola* Thomson, 1869:544 (*Lucilia*). Type-locality: Peru, Callao.
- quadrisignata* Thomson, 1869:544 (*Lucilia*). Type-locality: Galápagos Islands.
- aztequina* Bigot, 1877:252 (*Somomyia*). Type-locality: "Mexico".
- flavigena* Bigot, 1877:249 (*Somomyia*). Type-locality: "Mexico".
- iridicolor* Bigot, 1887:clxxxi (*Somomyia*). Type-locality: "Cuba".
- tristriata* Verhuell, 1850:273 (*Calliphora*). Type-locality: "Surinam".
- lynchi* Lahille, 1915:12, fig. 6c, pl. 2, figs. 3, 5 (*Chrysomyia*). Type-locality: Argentina, Chaco.
- MINIMA Shannon, 1926:124. Type-locality: West Indies, Santo Domingo, San Francisco Mts. Distr. — Cuba, Santo Domingo, Puerto Rico.

Genus HEMILUCILIA Brauer

- HEMILUCILIA Brauer, 1895:598. Type-species, *Musca segmentaria* Fabricius (mon.). Ref. — Mello, 1959 (revision).
- RENOISTI Séguy, 1925b:440. Type-locality: "French Guiana". Distr. — Venezuela to Peru and Brazil (Goiás, Paraná, Minas Gerais).
- parva* Shannon, 1926:125. Type-locality: Brazil, "Amazon".
- FLAVIFACIES (Engel), 1931:138 (*Strongyloneura*). Type-locality: Bolivia, "Nord-Chiquitos". Distr. — Mexico (Quintana Roo) to Ecuador, Brazil (south to Rio Grande do Sul) and Paraguay; Trinidad.
- fuscanipennis* authors, not Macquart.
- idioidea* authors, not Robineau-Desvoidy.
- SEGMENTARIA (Fabricius), 1805:292 (*Musca*). Type-locality: "South America". Distr. — Mexico (Quintana Roo) to Chile and Brazil (south to Santa Catarina).
- femorata* Walker, 1861:310 (*Calliphora*). Type-locality: "Mexico".
- nubipennis* Rondani, 1848:77 (*Lucilia*). Type-locality: "Brazil".
- semidiaphana* Rondani, 1850:177 (*Mya*). Type-locality: Brazil, São Paulo, São Sebastião Is.
- TOWNSENDI Shannon, 1926:125. Type-locality: Peru, Yahuar Mayo. Distr. — Peru.
- townsendia*, lapsus

Genus MYIOLUCILIA Hall

- MYIOLUCILIA Hall, 1948:109. Type-species, *Musca lyrcæa* Walker (orig. des.).
- FULVINOTA (Bigot), 1877:251 (*Somomyia*). Type-locality: "Mexico". Distr. — Mexico to Brazil (Espírito Santo) and Chile.
- fulvipes* authors, not Macquart.
- desvoidyi* Hough, 1900:208 (*Chrysomyia*). Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).
- braueri* Townsend, 1916:11 (*Paralucilia*). Type-locality: "Chile".



- LYRCEA (Walker), 1849:873 (*Musca*). Type-locality: Uruguay, Montevideo. Distr. — Mexico (Campeche) to Chile and Argentina (Tucumán).
verena Walker, 1849:874 (*Musca*). Type-locality: "Venczucla".
caruca Walker, 1849:877 (*Musca*). Type-locality: "Chile".
gamelia Walker, 1849:878 (*Musca*). Type-locality: Uruguay, Montevideo.
callipes Bigot, 1877:249 (*Somomya*). Type-locality: "Mexico".


Genus PARALUCILIA Brauer & Bergenstamm

- PARALUCILIA Brauer & Bergenstamm, 1891:391 (separate pagination, 87; may have priority). Type-species, *Calliphora fulvipes* Macquart (mon.) = *fulvicrura* (Robineau-Desvoidy). Ref. — Mello, 1968.
Compsomyiops Townsend, 1918:153. Type-species, *Calliphora fulvipes* Macquart (orig. des.) = *fulvicrura* (Robineau-Desvoidy).
ALVARENGAI Mello, 1968:188. Type-locality: Bolivia, La Paz, El Alto. Distr. — Bolivia.
APOLINARI Mello, 1968:185. Type-locality: Colombia, Bogotá. Distr. — Colombia.
AREQUIPENSIS Mello, 1968:187. Type-locality: Peru, Arequipa. Distr. — Peru.
BOLIVIANA Mello, 1968:190. Type-locality: Bolivia, La Paz, Calacoto. Distr. — Bolivia.
FULVICRURA (Robineau-Desvoidy), 1830:446 (*Chrysomyia*). Type-locality: Uruguay, Montevideo. Distr. — Mexico to Chile and Argentina (Rio Negro); Juan Fernandez Is.; Easter Island. Ref. — Lopes & Albuquerque, 1955 (description).
affinis authors, in part.
rufipes Macquart, 1843:286 (separate, p. 129) (*Calliphora*). Type-locality: "America" (original description has "Java", in error; cf. Sabrosky, 1953).
fulvipes Macquart, 1843:289 (separate, p. 132) (*Calliphora*). Type-locality: "Chile".
?annulipes Philippi, 1861:544 (*Calliphora*). Type-locality: Chile, Santiago.
taeniaria Thomson, 1869:544 (*Lucilia*). Type-locality: Argentina, Buenos Aires.
NIGRIPES Mello, 1968:184. Type-locality: Argentina, Buenos Aires, Boulogne. Distr. — Argentina, Bolivia, Chile.
WHEELERI (Hough), 1899:284 (*Chrysomyia*). Type-locality: U.S.A., California, Monterey. Distr. — U.S.A. (Washington to Colorado) to Bolivia.

Tribe PHORMIINI

Genus PHORMIA Robineau-Desvoidy

- PHORMIA Robineau-Desvoidy, 1830:465. Type-species, *Musca regina* Meigen (Robineau-Desvoidy, 1849:v).



Euphormia Townsend, 1919:542. Type-species, *Musca regina* Meigen (orig. des.).

REGINA (Meigen), 1826:58 (*Musca*). Type-locality: "Germany". Distr. — Eurasia; North America; Mexico (south to Mexico City, Baja California).

Subfamily CALLIPHORINAE

Tribe LUCILIINI

Genus BLEPHARICNEMA Macquart

BLEPHARICNEMA Macquart, 1843:283 (separate, 126). Type-species, *splendens* Macquart (mon.). This genus probably should be placed in a separate tribe.

SPLENDENS Macquart, 1843:284 (separate, p. 127). Type-locality: "unknown"; (Townsend, 1937, says "Colombia"). Distr. — Colombia, Ecuador, Peru, Bolivia.

auriceps Walker, 1858:209 (*Cynomyia*). Type-locality: Ecuador, Quito.

Genus BUFOLUCILIA Townsend

BUFOLUCILIA Townsend, 1919:542. Type-species, *Lucilia bufonivora* Moniez (orig. des.) Ref. — James, 1953 (key).

Xerophilophaga Enderlein, 1933:120. Type-species, *Onesia lucilioides* Wulp = *silvarum* (Meigen).

SILVARUM (Meigen), 1826:53 (*Musca*). Type-locality: Austria. Distr. — Eurasia, Northern Africa, North America south to Mexico (D.F.).

lucilioides Wulp, 1836:288 (*Onesia*). Type-locality: Mexico, Mexico City.


Genus LUCILIA Robineau-Desvoidy

LUCILIA Robineau-Desvoidy, 1830:452. Type-species, *Musca caesar* Linnaeus (Macquart, 1834:162; separate, p. 26).

ILLUSTRIS (Meigen), 1826:54 (*Musca*). Type-locality: Germany. Distr. — Eurasia; North America south to northern Mexico.

Genus PHAENICIA Robineau-Desvoidy

PHAENICIA Robineau-Desvoidy, 1863:750. Type-species, *concinna* Robineau-Desvoidy (Townsend, 1916:8) = *sericata* (Meigen). Refs. — Aubertin, 1933 (*Lucilia*, in part, revision); Mello, 1961 (Brazilian species).



Subgenus PHAENICIA Robineau-Desvoidy

- CAERULEVIRIDIS (Macquart), 1855:133 (separate, p. 113) (*Lucilia*). Type-locality: U.S.A., Maryland, Baltimore. Distr. — eastern United States; Cuba.
- oculata Townsend, 1908:122 (*Lucilia*). Type-locality, U.S.A., Kentucky, Cumberland Gap; type series partly from Cuba.
- CLUVIA (Walker), 1849:885 (*Musca*). Type-locality: "West Indies". Distr. — Southeastern United States; Cuba, Puerto Rico.
- CUPRINA (Wiedemann), 1830:654 (*Musca*). Type-locality: "China". Distr. — almost World-wide, tropics and warmer temperate regions; southern U.S.A. to Uruguay and northern Argentina. Listed in the Nearctic catalog as *pallescens* (Shannon).
- pallescens Shannon, 1924:78 (*Lucilia*). Type-locality: U.S.A., North Carolina, Wilmington.
- pseudosericata Gaminara, 1930:1264, 1931:1266 (*Lucilia*). Type-locality: "Uruguay".
- EXIMIA (Wiedemann), 1819:53 (*Musca*). Type-locality: "Brazil". Distr. — southwestern U.S.A. (Texas) to Argentina and Chile; West Indies, widespread. The common greenbottle fly of tropical America.
- ochricornis Wiedemann, 1830:408 (*Musca*), in part. Type-locality: "Brazil".
- ?putrida Fabricius, 1775:775 (*Musca*). Type-locality: "South America".
- smaragdula Robineau-Desvoidy, 1830:462 (*Lucilia*). Type-locality: "Brazil".
- ?paraensis Macquart, 1843:299 (separate, p. 142) (*Lucilia*). Type-locality: Brazil, Pará.
- ruficornis Macquart, 1846:198 (separate, p. 326) (*Lucilia*). Type-locality: "Colombia".
- punctipennis Macquart, 1848:216 (separate, p. 56) (*Lucilia*). Type-locality: "Brazil".
- subrectineuris Macquart, 1851:223 (separate, p. 250) (*Lucilia*). Type-locality: Brazil, Minas Gerais.
- insularis Walker, 1852:340 (*Musca*). Type-locality: "West Indies".
- amazona Bigot, 1877:255 (*Somomya*). Type-locality: "Brazil".
- mutabilis Bigot, 1877:248 (*Somomya*). Type-locality: "Mexico".
- pueblensis Bigot, 1877:250 (*Somomya*). Type-locality: "Mexico".
- oreoquina Bigot, 1877:253 (*Somomya*). Type-locality: "Brazil".
- hirtiforceps Shannon, 1926:133 (*Lucilia*). Type-locality: Panama, Canal Zone, Ancon.
- mera Shannon & Del Ponte, 1926:40 (*Lucilia*). Type-locality: Argentina, Jujuy, San Pedro de Jujuy.
- primaveris Shannon & Del Ponte, 1926:40 (*Lucilia*). Type-locality: Argentina, Buenos Aires, San Isidro.
- IBIS (Shannon), 1926:132 (*Lucilia*). Type-locality: Peru, Huadquiña. Distr. — Peru.
- JAPUHYRENSIS Mello, 1961:274. Type-locality: Brazil, Rio de Janeiro, Angra dos Reis, Japuiba. Distr. — Brazil (Rio de Janeiro).

- MEXICANA (Macquart), 1843:300 (separte, p. 143) (*Lucilia*). Type-locality: "Mexico". Distr. — Southwestern U.S.A., Mexico, Guatemala; Hall (1948) says south to Brazil.
- PROBLEMATICA (Johnson), 1913:448 (*Lucilia*). Type-locality: West Indies, Bermuda. Distr. — "West Indies", uncommon, according to Hall (1948).
- PURPURESCENS (Walker), 1837:355 (*Musca*). Type-locality: Brazil, Santa Catarina. Distr. — Southern Mexico to Argentina.
- ?*brunnicornis* Macquart, 1843:299 (separte, p. 142) (*Lucilia*). Type-locality: "Mexico".
- violacea* Macquart, 1847:99 (separte, p. 83) (*Lucilia*). Type-locality: "Mexico".
- pallidibasis* Bigot, 1877:247 (*Somomyia*). Type-locality: "Mexico".
- ?*praescia* Giglio-Tos, 1893:4 (*Lucilia*). Type-locality: Mexico, Tampico.
- ocularis* Shannon, 1926:131, 132 (*Lucilia*; *oculatis*, p. 132, lapsus). Type-locality: Costa Rica, San Mateo, Higuito.
- RICA (Shannon), 1926:132 (*Lucilia*). Type-locality: West Indies, Antigua. Distr. — Antigua, Puerto Rico, ?Haiti.
- ?*azurea* Robineau-Desvoidy, 1830:455 (*Lucilia*). Type-locality: Not stated; probably South American, according to Aubertin (1933).
- ?*semiviolacea* Bigot, 1877:46 (*Somomyia*). Type-locality: "Puerto Rico".
- ?*soulouquina* Bigot, 1877:47 (*Somomyia*). Type-locality: "Haiti".
- SERICATA (Meigen), 1826:53 (*Musca*). Type-locality: "Germany". Distr. — Almost world-wide; southern Canada to Argentina. The common greenbottle fly. Synonyms are numerous, but none are for certain based on Neotropical types, unless *brunnicornis* (Macquart, 1843), is this species, as Mello (1961) states.

Subgenus VIRIDINSULA Shannon

- VIRIDINSULA Shannon, 1926:131 (as genus). Type-species, *Musca pionia* Walker (orig. des.). Ref. — James, 1966.
- DECEPTOR (Curran), 1934a:166 (*Viridinsula*). Type-locality: Galápagos Islands, North Seymour (Baltra). Distr. — Galápagos Islands, Cocos Island.
- PIONIA (Walker), 1849:880 (*Musca*). Type-locality: Galápagos Islands. Distr. — Galápagos Islands.
- SETOSA James, 1966:479. Type-locality: Galápagos Islands, Darwin Island. Distr. — Galápagos Islands.

Tribe CALLIPHORINI

Genus ALDRICHINA Townsend

- Aldrichella* Rohdendorf, 1931:177 (preocc. Vaughan, 1903). Type-species, *Calliphora grahami* Aldrich (orig. des.).



ALDRICHINA Townsend, 1934:111 (nom. nov. for *Aldrichella* Rohdendorf).
Type-species, *Calliphora grahami* Aldrich (aut.).

GRAHAMI (Aldrich), 1930:1 (*Calliphora*). Type-locality: China, Szechuen, Chuan Chien and Mt. Omei. Distr. — Asia (widespread), Europe (Russia), western United States, Mexico south to Mexico City.

Genus CALLIPHORA Robineau-Desvoidy

CALLIPHORA Robineau-Desvoidy, 1830:433. Type-species, *Musca vomitoria* Linnaeus (orig. des.). Ref. — James, 1953 (key to Nearctic species); Mello, 1962 (Brazilian species).

COLORADENSIS Hough, 1899:286. Type-locality: U.S.A., Colorado. Distr. — Alaska to Mexico (San Martin and Baja California).

LOPESI Mello, 1962:270. Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil (Rio de Janeiro, São Paulo, Rio Grande do Sul).

PERUVIANA Robineau-Desvoidy, 1830:438. Type-locality: "Peru". Distr. — Mexico (Puebla), Costa Rica, Colombia, Ecuador, Peru, Bolivia.
? *semiatria* Schiner, 1868:308. Type-locality: "Colombia". N. SYN.

nigribasis Macquart, 1851:215 (separate, p. 242). Type-locality, "Colombia".

irazuana Townsend, 1908:118. Type-locality: Costa Rica, Irazu. Synonymy by Shannon, 1926, appears to be correct.

VICINA Robineau-Desvoidy, 1830:435. Type-locality: U.S.A., Pennsylvania, Philadelphia. Distr. — Holarctic; Alaska to Mexico; Uruguay, Argentina, and Chile south to Tierra del Fuego.
erythrocephala authors, not De Geer.

Genus EUCALLIPHORA Townsend

EUCALLIPHORA Townsend, 1908:118. Type-species, *Calliphora latifrons* Hough (mon.) = *lilaea* (Walker).

ARTA Hall, 1948:287. Type-locality: Mexico, San Luis Potosí. Distr. — Mexico.


LILAEA (Walker), 1849:894 (*Musca*). Type-locality: Canada, Hudson's Bay, Albany River, St. Martin's Falls. Distr. — Southern Canada to Mexico (Mexico); Hawaii.

Genus METALLICOMYIA Röder

Chalcomyia Röder, 1886:267 (preocc. Williston, 1885). Type-species, *elegans* Röder (mon.).

METALLICOMYIA Röder, 1886:307 (nom. nov. for *Chalcomyia* Röder). Type-species, *Chalcomyia elegans* Röder (aut.).

ELEGANS (Röder), 1886:268 (*Chalcomyia*). Type-locality: Ecuador, Rioshamba. Distr. — Ecuador.



Subfamily TOXOTARSINAE

Genus CALLYNTROPYGA Enderlein

CALLYNTROPYGA Enderlein, 1940:644. Type-species, *selkirki* Enderlein (mon.) = *humeralis* (Walker).

HUMERALIS (Walker), 1837:348 (*Stomoxys*). Type-locality, Chile, Concepción. Distr. — Chile, Juan Fernández Islands. Refs. — Smart, 1937; Lopes & Albuquerque, 1955.

selkirki Enderlein, 1940:645. Type-locality: Chile, Juan Fernández Islands.

Genus CHLOROBACHYCOMA Townsend

CHLOROBACHYCOMA Townsend, 1918:155. Type-species, *splendida* Townsend (orig. des.).

SPLENDIDA Townsend, 1918:155. Type-locality: Peru, Oroya. Distr. — Peru.

Genus KUSCHELOMYIA Lopes

KUSCHELOMYIA Lopes, 1961:455. Type-species, *ambrosiana* Lopes (orig. des.)

AMBROSIANA Lopes, 1961:456. Type-locality: Chile, Santo Ambrosio Is. Distr. — Chile.

Genus NETA Shannon

NETA Shannon, 1926:123. Type-species, *Phryssopoda splendens* Macquart 1851 (orig. des.) = *chilensis* (Walker).

Callyntropus Enderlein, 1930:70. Type-species, *Phryssopoda splendens* Macquart (mon.) = *chilensis* (Walker).

CHILENSIS (Walker), 1837:354 (*Musca*). Type-locality: "Chile". Distr. — Bolivia, Peru, Chile, Argentina (south to Straits of Magellan). *magellanica* Macquart, 1843:288 (separate, p. 131) (*Calliphora*). Type-locality: Chile, Strait of Magellan.

peruviana Robineau-Desvoidy of Macquart, 1851:216 (separate, p. 243) (*Calliphora*). Locality: Peru, Payta.


ortogesa Walker, 1849:834 (*Sarcophaga*). Type-locality: "Chile". *splendens* Macquart of Macquart, 1851:204 (separate, p. 231) (*Phryssopoda*).

calogaster Bigot, 1877:246 (*Somomyia*). Type-locality: Argentina, Buenos Aires, La Plata.

paytensis Townsend, 1893:36 (*Calliphora*; nom. nov. for *Calliphora peruviana* Robineau-Desvoidy of Macquart).

Genus SARCONESIA Bigot

SARCONESIA Bigot, 1857:301. Type-species, *Sarcophaga chlorogaster* Wiedemann (orig. des.).



- CHLOROGASTER (Wiedemann), 1830:359 (*Sarcophaga*). Type-localities: Uruguay, Montevideo, and Argentina, Buenos Aires, La Plata. Distr. — Peru, Brazil (Paraná), Paraguay, Uruguay, Argentina (south to Tierra del Fuego), Chile; Juan Fernández Islands, Easter Island. Ref. — Lopes & Albuquerque, 1955.
- minor* Enderlein, 1940:646 (as subspecies). Type-locality: Chile, Juan Fernández Islands.
- ?DICHROA (Schiner), 1868:313 (*Sarcophaga*). Type-locality: "Chile".
- ?NIGROCYANEA (Walker), 1837:351 (*Sarcophaga*). Type-locality: "Chile".
- PROERNA (Walker), 1849:835 (*Sarcophaga*). Type-locality: Uruguay, Montevideo. Distr. — Uruguay. Listed as synonym of *chlorogaster* by Smart (1937); Lopes & Albuquerque (1955) say this is not so.
- VERSICOLOR Bigot, 1857:302. Type-locality: "Argentina". Distr. — Argentina (L. Nahuel Huapi).

Genus SARCONESIOMIMA Lopes & Albuquerque

- SARCONESIOMIMA Lopes & Albuquerque, 1955:104. Type-species, *bicolor* Lopes & Albuquerque (orig. des.).
- BICOLOR Lopes & Albuquerque, 1955:105. Type-locality: Chile, Santiago. Distr. — Chile (Atacama, Coquimbo, Santiago); Juan Fernández Islands.

Genus SARCONESIOPSIS Townsend

- SARCONESIOPSIS Townsend, 1918:156. Type-species, *caerulea* Townsend (orig. des.) = *chilensis* (Macquart).
- CHILENSIS (Macquart), 1843:288 (separate, p. 131) (*Calliphora*). Type-locality: "Chile". Distr. — Colombia, Ecuador, Peru, Bolivia, Chile (south to Cautin).
- splendens* Macquart, 1843:253 (separate, p. 96) (*Phrissopoda*). Type-locality stated to be "Africa", also from Chile (former apparently by mistake). Not *splendens* (Macquart, 1851).
- fuscipennis* Macquart, 1843:267 (separate, p. 110) (*Cynomyia*). Type-locality: "Chile".
- incerta* Walker, 1852:334 (*Musca*). Type-locality: "Colombia".
- nitens* Bigot, 1877:244 (*Somomya*). Type-locality: "Colombia".
- castanipes* Bigot, 1877:245 (*Somomya*). Type-locality: Ecuador, Quito.
- caerulea* Townsend, 1918:156. Type-locality: Peru, Oroya.

Genus TOXOTARSUS Macquart

- TOXOTARSUS Macquart, 1851:211 (separate, p. 238). Type-species, *rufipalpis* Macquart (mon.).
- Trixoneura* Shannon, 1926:121. Type-species, *Agria fuscipennis* Macquart (orig. des.).

FUSCIPENNIS (Macquart), 1843:266 (separate, p. 109) (*Agria*). Type-locality: "Brazil or Chile". Distr. — Chile (Brazilian type record probably incorrect).

RUFIPALPIS Macquart, 1851:211. Type-locality: "Chile". Distr. — Chile (Coquimbo, Santiago, Chiloé Island). Ref. — Lopcs, 1962.

Unplaced species

ACUTANGULA Rondani, 1863:28 (*Somomya*). Type-locality: "South America".

AMERICANA Macquart, 1846:321 (separate, p. 193) (*Idia*). Type-locality: "Colombia".

AMERICANA Schiner, 1868:311 (*Onesia*). Type-locality: "Chile".

ARGENTINA Bigot, 1877:254 (*Somomya*). Type-locality: Argentina, Buenos Aires. Bigot says this might be the female of his *S. gratiosa*.

BIVITTATA Jaennicke, 1867:378 (*Onesia*). Type-locality: "Chile".

CLEMENTEI Ichcs, 1906: (*Calliphora*). Type-locality: Argentina, Chaco Austral.

DESVOIDYI Jaennicke, 1867:378 (*Cynomyia*). Type-locality: "Chile".

FASCIATA Macquart, 1846:322 (separate, p. 194) (*Rhynchomyia*). Type-locality: "Colombia".

FULVICORNIS Robineau-Desvoidy, 1830:462 (*Lucilia*). Type-locality: "Brazil".

FUSCIPENNIS Jaennicke, 1867:376 (*Calliphora*). Type-locality: "Brazil".

GRATIOSA Bigot, 1877:253 (*Somomya*). Type-locality: Argentina, Buenos Aires. See *argentina* Bigot.

HYACINTHINA Robineau-Desvoidy, 1830:462 (*Lucilia*). Type-locality: "Brazil".

brasiliensis Townsend, 1893:36 (nom. nov. for *Lucilia hyacinthina* Robineau-Desvoidy, 1830:462, not *Chrysomya hyacinthina* Robineau-Desvoidy, 1830:450). Type-locality: "Brazil".

INCISURALIS Macquart, 1843:304 (separate, p. 147) (*Lucilia*). Type-locality: said to be "Brazil", but according to Séguy (1928) it is African.

LIMBIPENNIS Macquart, 1843:283 (separate, p. 126) (*Rhynchomyia*). Type-locality: "America".

LITTORALIS Blanchard, 1937:380 (*Lucilia*). Type-locality: "Argentina".

LUTEICORNIS Jaennicke, 1867:375 (*Lucilia*). Type-locality: "Venezuela".

MARGINATA Macquart, 1843:304 (separate, p. 147) (*Lucilia*). Type-locality: said to be "Brazil or Chile" but according to Séguy it is African.

melanaria Wulp, 1896:295 (*Calliphora*). Type-locality: Mexico, Guerrero, Amula (Not Calliphoridae — see Tachinidae, Theresiini).

MONTEVIDENSIS Bigot, 1877:46 (*Somomyia*). Type-locality: Uruguay, Montevideo.

MUSCARIA Jaennicke, 1867:378 (*Onesia*). Type-locality: "Chile".

NIGRICEPS Macquart, 1843:300 (separate, p. 143) (*Lucilia*). Type-locality: "Mexico".

PERUVIANA Robineau-Desvoidy, 1830:455 (*Lucilia*). Type-locality: "Peru".

- PHACOPTERA Wulp, 1882:88 (*Calliphora*). Type-locality: "Chile".
- PHILIPPIANA Rondani, 1863:29 (*Somomyia*) (as variety of "*S. chilensis* Macq." or a separate species. Type-locality: not given.
- PRAEPES Giglio-Tos, 1893:3 (*Calliphora*). Type-locality: "Mexico".
- PRINCEPS Rondani, 1848:17 (*Lucilia*). Type-locality: "Brazil".
- QUIETA Giglio-Tos, 1893:4 (*Lucilia*). Type-locality: Mexico, no locality indicated.
- RUFIPALPIS Macquart, 1851:216 (separate, p. 243) (*Calliphora*) (not *C. rufipalpis* Macquart, 1843). Type-locality: "America"; may be South America.
- americana* Rondani, 1863:39 (*Somomyia*). Change of name for "*rufipalpis* Macq. — Dipt. exot.", presumably this species rather than *rufipalpis* Macquart, 1843.
- TRANSMARINA Rondani, 1863:29 (*Somomyia*). Type-locality: "South America".
- VARIPALPIS Macquart, 1843:298 (separate, p. 141) (*Lucilia*). Type-locality: "Brazil".

LITERATURE CITED

ALDRICH, J. M.

1922. The Neotropical musicoid genus *Mesembrinella* Giglio-Tos and other testaceous muscoid flies. *Proc. U. S. Nat. Mus.* 62(11):1-24.
1925. New Diptera or two-winged flies in the United States National Museum. *Proc. U. S. Nat. Mus.* 66(18):1-36.
1930. New two-winged flies of the family Calliphoridae from China. *Proc. U. S. Nat. Mus.* 78(1):1-5 (illus.).
1932. A new genus and two new species of muscoid flies from Guatemala. *Proc. Entomol. Soc. Washington* 34:23-25.

AUBERTIN, D.

1933. Revision of the genus *Lucilia* R. D. (Diptera, Calliphoridae). *J. Linn. Soc. London* 38:389-436.

EIGOT, J. M.

1857. Diptères nouveaux provenant du Chili. *Ann. Soc. Entomol. France* 5:277-308, illus.
1877. Diptères nouveaux ou peu connus. 7^e partie, IX: Genre *Somomyia* (Rondani) *Lucilia* (Rob.-Desv.). *Calliphora*, *Phormia*, *Chrysomyia* (id.); 8^e partie, X: Genre *Somomyia* (Rondani). *Calliphora*, *Melinda*, *Mufetia*, *Lucilia*, *Chrysomyia* (alias *Microchrysa* Rond.) Robineau-Desvoidy. *Ann. Soc. Entomol. France* 5(7):35-48, 243-259.
1878. Diptères nouveaux ou peu connus. 9^e partie, XII: Genus *Phumosa*, *Pyrellia*, *Cosmina*, *Ochromyia* et *Curtonevra*. *Ann. Soc. Entomol. France* 5(8):31-40.
1888. (Diagnoses abrégées de quelques Diptères nouveaux, provenant de l'Amerique de Nord). *Bul. Ann. Soc. Entomol. France* 6(7): clxxx-clxxxii (1887).

BLANCHARD, E. E.

1937. Descripciones y anotaciones de dípteros argentinos. *An. Soc. Cienc. Argentina* 126:345-386, illus.

BRAUER, F.

1883. Die Zweiflügler der Kaiserlichen Museums zu Wien. III. *Denkschr. K. Akad. Wiss. Wien, Math.-Nat. Cl.* 47:1-100, illus. Also published separately.
1889 Beiträge zur Kenntniss der Muscaria Schizometopa. *Sitz. K. Akad. Wiss. Wien, Math.-Nat. Cl. Abt. 1*, 108:495-529.
1895. Bemerkungen zu einigen neuen Gattungen der Muscarien und Deutung einiger Original-Exemplare. *Sitz. K. Akad. Wiss. Wien, Math.-Nat. C.* 104:582-604, illus.

BRAUER, F., & J. E. VON BERGENSTAMM,

1891. Die Zweiflügler des Kaiserlichen Museums zu Wien. V. Vorarbeiten zu einer Monographie der Muscaria Schizometopa (exclusive Anthomyidac). Pars II. *K. Akad. Wiss., Wien, Math.-Nat. Cl. Denk.* 58:305-446, illus. Also published separately, 142 pp.

CONIL, P. A.

1878. In Lesbini, C., H. Weyenbergh, & P. A. Conil. Études sur la Myiasis. *Actas Acad. Nac. Sci. Buenos Aires* 3:43-98, illus.

COQUEREL, C.

1858. Note sur les larvæ appartenant à une espèce nouvelle de Diptère (*Lucilia hominivorax*) développées dans les frontaux de l'homme à Cayenne. *Ann. Soc. Entomol. France* 3(6):171-176, illus.

CURRAN, C. H.

- 1934a. The Templeton Crocker Expedition of the California Academy of Sciences, 1932. No. 13. Diptera. *Proc. Calif. Acad. Sci.* 4(21): 147-172, illus.
1934b. The Diptera of Kartabo, Bartica District, British Guiana. *Bul. Amer. Mus. Nat. Hist.* 66:287-532.

CUSHING, E. C. & W. S. PATTON

1933. Studies on the higher Diptera of medical and veterinary importance. *Cochliomyia americana* sp. nov., the screw-worm fly of the New World. *Ann. Trop. Med. Parasitol.* 27:539-551, illus.

DEL PONTE, E.

1938. Las especies argentinas del genero *Cochliomyia* T. T. (Dipt. Musc.). *Rev. Entomol.* 8:273-281, illus.

ENDERLEIN, G.

1930. Diptero-logische Studien. XX. *Deut. Entomol. Ztschr.* 1930: 65-71.
1933. Neue paläarktische Calliphoridae, darunter Schneckenparasiten (Dipt.). *Mitt. Deut. Entomol. Gesell.* 4:120-128.
1940. Die Dipterenfauna der Juan Fernández-Inseln und der Oster-Inseln. In Skottsberg, *Nat. Hist. Juan Fernández and Easter Island, (Zool.)* 3(5):643-680, illus.

ENGEL, O.

1931. Die Ausbeute der deutschen Chaco-Expedition 1925/26. Diptera. *Konowia* 10:131-154.

FABRICIUS, J. C.

1775. *Systema entomologiae, sistens insectorum classes, ordines, species adiectis synonymis, locis, descriptionibus, observationibus*. 832 pp. Flensburg et Lipsiae (Flensburg and Leipzig).
1805. *Systema antliatorum secundum ordines, genera, species*. 373 pp. Brunsvigae (Brunswick).

GAMINARA, A.

1930. Clasificacion de algunos muscoideos uruguayos (Muscidae y Calliphoridae). *An. Fac. Med. Montevideo* 1930(1929-1930): 1235-1282. Also *Arch. Soc. Biol. Montevideo*, 1931, suppl. 5:1237-1280; illus.

GARCIA, MIGUEL

1952. Consideraciones generales sobre el genero *Cochliomyia* Townsend, 1916 y descripcion de *C. fontana* n. sp. (Diptera Calliphoridae). *Publ. Inst. Regional Entomol. Sanitaria*, no. 1-3(1948-1950): 68-80, illus.

GIGLIO-TOS, E.

1893. *Ditteri del Messico*. Pt. 4, 74 pp., illus. Torino.

HALL, D. G.

1948. *The blowflies of North America*. Thomas Say Foundation, Entomological Society of America. Vol. 4, 477 pp., illus.

HOUGH, G. DE N.

1899. Synopsis of the Calliphorinae of the United States. *Zool. Bul.* 2:283-290, illus.
1900. South American Muscidae in the collection of S. W. Williston. *Kansas Univ. Quarterly*, 9:203-232, illus.

ICHES, L.

1906. Sobre cinco dipteros nuevos del Chaco Austral. *Bol. Minist. Agric.*, Buenos Aires 6(5):262-273.

JAENNICKE, F.

1867. Neue exotische Dipteren. *Abhandl. Senckenb. Naturf. Gesell.* 6:311-408, illus.

JAMES, M. T.

1953. Notes on the distribution, systematic position, and variation of some Calliphorinae, with particular reference to the species of western North America. *Proc. Entomol. Soc. Washington* 55: 143-148.
1966. The blow flies of the Galapagos Islands (Diptera: Calliphoridae). *Proc. Calif. Acad. Sci.* 34:475-482, illus.

JOHNSON, C. W.

1913. The dipteran fauna of Bermuda. *Ann. Entomol. Soc. Amer.* 6: 443-452, illus.

LAHILLE, F.

1915. Nota sobre la ura y otras larvas daniñas de dipteros. *Bol. Dir. Gen. Ganad. Min. Agric.*, Buenos Aires, pp. 1-18, 7 figs., 2 pls.

LOPES, H. DE S.

1961. Sôbre um nôvo gênero de Calliphoridae de Ilha de Santo Ambrósio, Chile (Diptera). *Rev. Brasil Biol.* 21:455-459, illus.
1962. Sôbre dois Calliphoridae do Chile (Diptera). *Rev. Brasil. Biol.* 22:421-426, illus.

LOPES, H. DE S. & D. DE O. ALBUQUERQUE

1955. Los insectos de las islas Juan Fernandez. 22. Calliphoridae et Sarcophagidae (Diptera). *Rev. Chilena Entomol.* 4:95-119, illus.

MACQUART, J.

1834. Insectes diptères du nord de la France. Athéricières: Créophiles, Oestrides, Myodaires, Conopsaires, Scénopiniens, Céphalopsides. *Mém. Soc. Roy. Sci., Agric., Arts Lille* 1833:137-368, illus. Also published separately (vol. 5) 232 pp.
1843. Diptères exotiques nouveaux ou peu connus. *Mém. Soc. Roy. Sci., Agric., Arts Lille* (1842):162-460, illus. Also published as separate, 2(3):5-304.
1846. Diptères exotiques nouveaux ou peu connus. 1er. Supplément. *Mém. Soc. Roy. Sci., Agric., Arts, Lille* (1845) (1844):133-364, illus. Also published as Supplément I, pp. 133-364.
1847. Diptères exotiques nouveaux ou peu connus. 2e Supplément. *Mém. Soc. Roy. Sci., Agric., Arts, Lille* (1846):21-120, illus. Also published as separate, Supplément II, pp. 5-104.
1848. Diptères exotiques nouveaux ou peu connus. Suite de 2me. supplément (3e supplement). *Mém. Soc. Roy. Sci., Agric., Arts, Lille* (1847):161-237, illus. Also published as Supplément III, pp. 1-77.
1851. Diptères exotiques nouveaux ou peu connus. Suite de 4e supplément. *Mém. Soc. Roy. Sci., Agric., Arts, Lille* (1850):134-294, illus. Also published as Supplément IV (part), pp. 161-309, 324-336.
1855. Diptères exotiques nouveaux ou peu connus. 5e. supplement. *Mém. Soc. Roy. Sci., Agric., Arts, Lille* (1854):25-156, illus. Also published as separate, Supplément V, pp. 5-136.

MEIGEN, J. W.

1826. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten*. Vol. 5, xii + 512 pp., illus.

MELLO, J. A. S. N. DE

1959. Contribuição ao estudo do gênero "*Hemilucilia*" Brauer, 1895 (Calliphoridae, Diptera). *Rev. Brasil. Biol.* 19:429-439, illus.

MELLO, R. P. DE

1961. Contribuição ao estudo do gênero *Phaenicia* (R.D., 1863) (Diptera, Calliphoridae). *Mem. Inst. Oswaldo Cruz* 59:259-278, illus.
1962. Contribuição ao estudo do gênero *Calliphora* R.D., 1830 (Diptera, Calliphoridae). *Mem. Inst. Oswaldo Cruz* 60:263-274, illus.
1965. Contribuição ao estudo do gênero *Callitrogopsis* Townsend (Diptera, Calliphoridae). *Studia Entomol.* 8:435-440, illus.

1967. Contribuição ao estudo dos Mesembrinellinae sul-americanos (Dipt. Calliphoridae). *Studia Entomol.* 10:1-80, illus.
1968. Contribuição ao estudo do gênero "*Paralucilia*" Brauer & Bergenstamm, 1891 (Diptera, Calliphoridae). *Rev. Brasil. Biol.* 28: 177-192, illus.

PHILIPPI, R. A.

1861. Beschreibung einer neuen Fliege deren Larven in der Nase und Stirnhöle einer Frau gelebt haben. *Z. Ges. Naturw.* 17:513-515.

ROBINEAU-DESVOIDY, J. B.

1830. Essai sur les Myodaires. (Paris) Inst. de France, (Cl. des) Sci. Math. et Phys., Acad. Roy. des Sci., *Mém. présentés par divers Savans* (ser. 2), 2:1-813.
1849. (Note on *Lucilia dispar* and *Phormia regina*). *Ann. Soc. Entomol. France* (2) 7(Bul.):iv-v.
1863. *Histoire naturelle des Diptères des environs de Paris*. Vol. 2, 920 pp.

RÜDER, V. VON

1886. Dipteren von den Cordilleren in Columbien. *Stettiner Entomol. Z.* 47:257-270; 307.

ROHDENDORF, B. B.

1931. Calliphorinen-Studien. IV. (Dipt.). Eine neue Calliphorinen-Gattung aus Ostsibirien. *Zool. Anz.* 95:175-177 (illus.).

RONDANI, C.

1848. Esame di varie specie d'insetti ditteri brasiliani. *Studi Entomol.* (Turin) 1:63-112, illus.
1850. Osservazioni sopra alquante specie di esapodi ditteri del Museo Torinese. *Nuovi Ann. Sci. Nat. Bologna* 3(2):165-197, illus.
1863. *Diptera exotica revisa et annotata*. 99 pp., illus. Modena. Also 1864. *Arch. Zool. Anat. fis.* (Modena) (1863) 3:1-99, illus.

SABROSKY, C. W.


1953. The scientific name of the screw-worm, with a note on *Paralucilia fulvicrura* (Diptera, Calliphoridae). *Proc. Entomol. Soc. Washington* 55:36-38.

SCHINER, I. R. (J. R.)

1868. Diptera (Art. 1), 388 pp., illus.. In Wüllerstorff-Urbair, B. von, *Reise der österreichischen Fregatte Novara*. *Zool.*, Vol. 2, Abt. 1, Sect. B. Wien.

SÉGUY, E.

- 1925a. Espèces nouvelles du genre *Mesembrinella* G-T. *Encycl. Entomol. Diptera, Ser. B2*, 2:195-196.
- 1925b. Étude sur quelques Calliphorines testacés rares ou peu connus. *Bul. Mus. Hist. Nat. Paris* 1925:438-441.
1928. Études sur les mouches parasites. Tome I. Conopides, Oestridas et Calliphorines de l'Europe occidentale. *Encycl. Entomol. Diptera, Ser. B2*, 9:1-251, illus.



SHANNON, R. C.

1924. Nearctic Calliphoridae, Luciliini. *Insecutor Inscitiae Menstruus*, 12:67-81.

1926. Synopsis of the American Calliphoridae (Diptera). *Proc. Entomol. Soc. Washington* 28:115-139.

SHANNON, R. C. & E. DEL PONTE

1926. Sinopsis parcial de los Muscoideos Argentinos. *Rev. Inst. Bacteriol.* 4(5):1-48, illus.

SMART, J.

1937. Calliphoridae, in British Museum (Natural History): *Diptera of Patagonia and South Chile* 7(3):376-384, illus.

SURCOUF, J. M. R.

1919. Revision des Muscidae testaceae. *Nouv. Arch. Mus. Hist. Nat. Paris* 5:27-124 (1914).

THOMSON, C. G.

1869. 6. Diptera. Species nova descripsit. Pp. 443-614, illus. In K. Svenska Vetenskaps-Akademien, *Kongliga svenska fregatten Eugenies resa omkring jorden. Pt. 2: Zoologie (sec.) 1: Insekter*, 617 pp. Stockholm, "1868".

TOWNSEND, C. H. T.

1893. Catalogue of the described South American species of Calyptrae Muscidae. *Ann. N. Y. Acad. Sci.* 7:1-44.

1908. The taxonomy of the muscoidean flies, including descriptions of new genera and species. *Smithsonian Misc. Coll.* 51(2 [publ. 1803]):1-138.

1915. A new generic name for the screw-worm fly. *J. Washington Acad. Sci.* 5:644-646.

1916. Designation of muscoid genotypes, with new genera and species. *Insecutor Inscitiae Menstruus* 4:4-12.

1918. New muscoid genera, species and synonymy. *Insecutor Instittiae Menstruus* 6:151-156.

1919. New genera and species of muscoid flies. *Proc. U. S. Nat. Mus.* 56:541-592.

1931. Notes on American oestromuscoid types. *Rev. Entomol.* 1:65-104, 157-183.

1934. Muscoid notes and descriptions. *Rev. Entomol.* 4:110-112.

1935. New Muscoid genera, mainly from the Neotropical region. *Rev. Entomol.* 5:68-74.

VERHUELL, Q. M. R.

1850. Mededeeling der metamorphose van cene in Guyana voorkomende Vliegensoort. *Tijdschr. wis- en natuurk. Wetensch.* 3: 273-275, fig.

WALKER, F.

1837. Descriptions of the insects collected by Captain P. P. King, R. N., F.R.S., in the survey of the Straits of Magellan. *Trans. Linnacan Soc. London* 17:331-359.


1849. *List of the specimens of dipterous insects in the collection of the British Museum.* Vol. 4, pp. 689-1172. London.



1852. *Insecta Saundersiana, or characters of undescribed insects in the collection of W. W. Saunders. Diptera*, 1:253-414, illus. London, "1856".
1858. Characters of undescribed Diptera in the collection of W. W. Saunders (continuation). *Trans. Entomol. Soc. London* 2(5):190-235.
1861. Characters of undescribed Diptera in the collection of W. W. Saunders. *Trans. Entomol. Soc. London*. 2(5):268-334.
- WIEDEMANN, C. R. W.
1819. Brasilianische Zweiflügler. *Zool. Mag.* (Wiedemann's), 1:40-56.
- 1830...*Aussereuropäische zweiflügelige Insekten* 2:xii + 684 pp., illus.
- WULP, F. M. VAN DER
1882. Remarks on certain American Diptera in the Leyden Museum and descriptions of nine new species. *Leyden Mus. Notes* 4:73-92.
1836. Supplement, pp. 289-304, in Godman, F. D., & Salvin, O., eds., *Biologia Centrali-Americana. Zoologia-Insecta-Diptera*, 1:378 pp., illus.

INDEX


abaca (Hall), 1948 (Huascaromusca), Mesembrinella	102.3
acutangula Rondani, 1863 (Somomya)	102.15
aeneiventris (Wiedemann), 1830 (Dexia), Huascaromusca.	102.3
aeneiventris authors, Mesembrinella	102.4
affinis authors, Paralucilia	102.8
affinis (Robineau-Desvoidy), 1830 (Chrysomya), Cochliomyia ..	102.6
Albuquerque Mello, 1967	102.2
aldrichi Del Ponte, 1938, Cochliomyia	102.5
Aldrichella Rohdendorf, 1931	102.11
Aldrichina Townsend, 1934	102.11, 102.12
alia (Robineau-Desvoidy), 1830 (Chrysomya), Cochliomyia	102.6
alvarengai Mello, 1968, Paralucilia	102.8
amazona (Bigot), 1877 (Somomya), Phaenicia	102.10
ambrosiana Lopes, 1961, Kuscheiomyia	102.13
americana Cushing & Patton, 1933, Cochliomyia	102.6
americana Macquart, 1846 (Idia)	102.15
americana Rondani, 1863 (Somomya)	102.16
americana Schiner, 1868 (Onesia)	102.15
annulipes (Philippi), 1861 (Caliphora), Paralucilia	102.8
anthropophaga (Conil), 1878 (Calliphora) Cochliomyia	102.6
apolinari Mello, 1968, Paralucilia	102.8
apolinaris Séguy, 1925, Mesembrinella	102.4
arequipensis Mello, 1968, Paralucilia	102.8
argentina Bigot, 1877 (Somomya)	102.15
arta Hall, 1948, Eucalliphora	102.12
auriceps (Walker), 1858 (Cynomyia), Blepharicnema	102.9
aztequina (Bigot), 1877 (Somomya), Cochliomyia	102.7
azurea (Robineau-Desvoidy), 1830 (Lucilia) Phaenicia	102.11



batesi Aldrich, 1922, Mesembrinella	102.4
bellardiana Aldrich 1922, Mesembrinella	102.4
benoisti Séguy, 1925, Hemilucilia	102.7
benoisti Séguy, 1925 (Ochromyia)	102.4
bequaerti (Séguy), 1925 (Mesembrinella), Huascaromusca	102.3
bicolor (Fabricius), 1805 (Musca), Mesembrinella	102.4
bicolor Giglio-Tos, 1893, Mesembrinella	102.4
bicolor Lopes & Albuquerque, 1955, Sarconesiomima	102.14
bivittata Jaenicke, 1867 (Onesia)	102.15
Blepharicnema Macquart, 1843	102.9
boliviana Mello, 1968, Paralucilia	102.8
brasiliensis Townsend, 1893 (Lucilia)	102.15
braueri (Townsend), 1916 (Paralucilia), Myiolucilia	102.7
brunnicornis (Macquart), 1843 (Lucilia), Phaenicia	102.11
brunnipes (Surcouf), 1919 (Mesembrinella), Lancella	102.3
Bufolucilia Townsend, 1919	102.9
<i>cacrulea</i> Townsend, 1918, Sarconesiopsis	102.14
caeruleiviridis (Macquart), 1855 (Lucilia), Phaenicia	102.10
callipes (Bigot), 1877 (Somomyia), Myiolucilia	102.8
Calliphora Robineau-Desvoidy 1839	102.12
Calliphorinae	102.9
Calliphorini	102.11
Callitroga Brauer, 1883	102.5
Callitrogopsis Townsend, 1935	102.5
Callyntropus Enderlein, 1930	102.13
Callyntropyga Enderlein, 1940	102.13
calogaster (Bigot), 1877 (Somomyia), Neta	102.13
caruca (Walker), 1849 (Musca), Myiolucilia	102.8
castanipes (Bigot), 1877 (Somomyia) Sarconesiopsis	102.14
Chalcomyia Röder, 1886	102.12
chilensis (Macquart) 1843 (Calliphora) Sarconesiopsis	102.14
chilensis (Walker), 1837 (Musca), Neta	102.13
Chlorobrachycoma Townsend, 1918	102.13
chlorogaster (Wiedemann), 1930 (Sarcophaga), Sarconesia	102.14
Chloroprocta Wulp. 1896	102.5
Chrysomyinae	102.5
Chrysomyiini	102.5
chrysorhea (Brauer), 1895 (Mesembrinella), Huascaromusca ...	102.3
clementei Ichas, 1906 (Calliphora)	102.15
cluvia (Walker), 1849 (Musca), Phaenicia	102.10
Cochliomyia Townsend, 1915	102.5
coloradensis Hough, 1895, Calliphora	102.12
Compsomyia authors	102.5
Compsomyiops Townsend, 1918	102.8
confusa Séguy, 1925, Mesembrinella	102.4
costalis Townsend, 1935, Callitrogopsis	102.5
cruciata Townsend, 1918, Huascaromusca	102.3
cuprina (Wiedemann), 1830 (Musca), Phaenicia	102.10
curvipes (Thomson), 1869 (Lucilia), Cochliomyia	102.7
cyaneicincta (Surcouf), 1919 (Ochromyia), Mesembrinella	102.4
cyanicolor (Rondani), 1850 (Lucilia), Cochliomyia	102.6



<i>deceptor</i> (Curran), 1934 (Viridinsula), Phaenicia	102.11
<i>decora</i> (Robineau-Desvoidy), 1930 (Chrysomya), Cochliomyia ..	102.6
<i>decrepita</i> (Séguy), 1925 (Mesembrinella), Huascaromusca	102.3
<i>desvoidyi</i> (Hough), 1900 (Chrysomyia), Myiolucilia	102.7
<i>desvoidyi</i> Jaenicke, 1867 (Cynomyia)	102.15
<i>dichroa</i> (Schiner), 1868 (Sarcophaga), Sarconesia	102.14
<i>dorsimacula</i> (Aldrich), 1922 (Mesembrinella), Eumesembrinella .	102.2
<i>durvillei</i> (Macquart), 1843 (Lucilia), Cochliomyia	102.6
<i>elegans</i> (Röder), 1886 (Chalcomyia), Metallicomyia	102.12
<i>erythrocephala</i> , authors, Calliphora	102.12
<i>Eucalliphora</i> Townsend, 1908	102.12
<i>Eumesembrinella</i> Townsend, 1931	102.2
<i>Euphormia</i> Townsend, 1919	102.9
<i>eximia</i> (Wiedemann), 1819 (Musca), Phaenicia	102.10
<i>facialis</i> Aldrich, 1922, Mesembrinella	102.4
<i>fasciata</i> Macquart, 1846 (Rhynchomyia)	102.15
<i>fasciata</i> (Walker), 1852 (Musca), Cochliomyia	102.7
<i>femorata</i> (Walker), 1861 (Calliphora), Hemilucilia	102.7
<i>flavicrura</i> Aldrich, 1922, Mesembrinella	102.4
<i>flavifacies</i> (Engel), 1931 (Strongyloncura), Hemilucilia	102.7
<i>flavigena</i> (Bigot), 1877 (Somomya), Cochliomyia	102.7
<i>flavipennis</i> (Macquart), 1843 (Ochromyia), Eumesembrinella	102.2
<i>fontanai</i> Garcia 1952, Cochliomyia	102.5
<i>formosa</i> Aldrich, 1932, Mesembrinella	102.4
<i>fulvicornis</i> Robineau-Desvoidy, 1830 (Lucilia)	102.15
<i>fulvicrura</i> (Robineau-Desvoidy), 1830 (Chrysomyia), Paralucilia .	102.8
<i>fulvinota</i> (Bigot), 1877 (Somomya), Myiolucilia	102.7
<i>fulvipes</i> Aldrich, 1922, Mesembrinella	102.4
<i>fulvipes</i> authors, Myiolucilia	102.7
<i>fulvipes</i> (Macquart), 1843 (Calliphora) Paralucilia	102.8
<i>fulvobarbata</i> (Bigot), 1887 (Somomya), Cochliomyia	102.6
<i>fuscipennis</i> authors, Hemilucilia	102.7
<i>fuscipennis</i> (Macquart), 1851 (Lucilia), Chloroprocta	102.5
<i>fuscicosta</i> Séguy, 1925, Mesembrinella	102.4
<i>fuscipennis</i> Jaenicke, 1867 (Calliphora)	102.15
<i>fuscipennis</i> Macquart, 1843 (Ochromyia)	102.4
<i>fuscipennis</i> (Macquart), 1843 (Cynomyia), Sarconesiopsis	102.14
<i>fuscipennis</i> (Macquart), 1843 (Agria), Toxotarsus	102.15
<i>gamelia</i> (Walker), 1849 (Musca), Myiolucilia	102.8
<i>gigas</i> Macquart, 1846 (Ochromyia)	102.5
<i>grahami</i> (Aldrich), 1930 (Calliphora), Aldrichina	102.12
<i>grajauensis</i> Mello, 1967, Huascaromusca	102.3
<i>gratiosa</i> Bigot, 1877 (Somomya)	102.15
<i>hirtiforceps</i> (Shannon), 1926 (Lucilia), Phaenicia	102.10
<i>Hemilucilia</i> Brauer, 1895	102.7
<i>homicida</i> (Brauer), 1899 (Compsomyia), Cochliomyia	102.6
<i>hominivorax</i> (Coquerel), 1858 (Lucilia), Cochliomyia	102.6
<i>Huascaromusca</i> Townsend, 1918	102.3



humeralis (Walker), 1837 (Stomoxys), Callyntropyga	102.13
hyacinthina (Robineau-Desvoidy), 1830 (Chrysomya), Cochliomyia	102.6
hyacinthina Robineau-Desvoidy, 1830 (Lucilia)	102.15
ibis (Shannon), 1926 (Lucilia), Phaenicia	102.10
idioidea authors, Hemilucilia	102.7
idioidea (Robineau-Desvoidy), 1830 (Chrysomya), Chloroprocta ..	102.5
illustris (Meigen), 1826 (Musca), Lucilia	102.9
incerta (Walker), 1852 (Musca), Sarconesiopsis	102.14
incisuralis Macquart 1943 (Lucilia)	102.15
incompleta (Curran), 1934 (Mesembrinella), Eumesembrinella ..	102.2
infesta (Philippi), 1861 (Calliphora), Cochliomyia	102.6
insularis (Walker), 1852 (Musca), Phaenicia	102.10
irazuana Townsend, 1908, Calliphora	102.12
iridicolor (Bigot), 1887 (Somomyia), Cochliomyia	102.7
japuhybensis Mello, 1961, Phaenicia	102.10
Kuschelomyia Lopcs, 1961	102.13
lanitaria (Wiedemann), 1830 (Musca), Cochliomyia	102.6
Lancella Mello, 1967	102.3
lanci Mello, 1967, Eumesembrinella	102.2
latifrons Mello, 1967, Albuquerquea	102.2
lepida (Robineau-Desvoidy), 1830 (Chrysomya), Cochliomyia	102.6
lilaea (Walker), 1849 (Musca), Eucalliphora	102.12
limbidipennis Macquart, 1843 (Rhynchomyia)	102.15
littoralis Blanchard, 1937 (Lucilia)	102.15
lopesi Mello, 1962, Calliphora	102.12
Lucilia Robineau-Desvoidy, 1830	102.9
Luciliini	102.9
lucilioides (Wulp), 1896 (Onesia), Bufolucilia	102.9
luteicornis Jaennicke, 1867 (Lucilia)	102.15
lynchi (Lahille), 1915 (Chrysomya), Cochliomyia	102.7
lyrcea (Walker), 1849 (Musca), Myiolucilia	102.8
macellaria authors, Cochliomyia	102.6
macellaria (Fabricius), 1775 (Musca), Cochliomyia	102.6
magellanica (Macquart), 1843 (Calliphora), Neta	102.13
marginata Macquart, 1843 (Lucilia)	102.15
melanaria Wulp, 1896 (Calliphora)	102.15
mera (Shannon & Del Ponte), 1926 (Lucilia), Phaenicia	102.10
Mesembolia Aldrich, 1925	102.3
Mesembrinella Giglio-Tos, 1893	102.3
Mesembrinellinae	102.2
Metallicomyia Röder, 1886	102.12
mexicana (Macquart), 1843 (Lucilia), Phaenicia	102.11
minima Shannon, 1926, Cochliomyia	102.7
minor Enderlein, 1940, Sarconesla	102.14
montevideensis Bigot, 1877 (Somomyia)	102.15
muscaria Jaennicke, 1867 (Onesia)	102.15
mutabilis (Bigot), 1877 (Somomyia), Phaenicia	102.10
Myiolucilia Hall, 1948	102.7




<i>Neta</i> Shannon, 1926	102.13
<i>nigribasis</i> Macquart, 1851, <i>Calliphora</i>	102.12
<i>nigriceps</i> Macquart, 1843 (<i>Lucilia</i>)	102.15
<i>nigrifrons</i> (Bigot), 1878 (<i>Ochromyia</i>), <i>Huascaromusca</i>	102.3
<i>nigripes</i> Mello, 1968, <i>Paralucilia</i>	102.8
<i>nigrocyanca</i> (Walker), 1837 (<i>Sarcophaga</i>), <i>Sarconesia</i>	102.14
<i>nitens</i> (Bigot), 1877 (<i>Somomya</i>), <i>Sarconesiopsis</i>	102.14
<i>nubipennis</i> (Rondani), 1848 (<i>Lucilia</i>), <i>Hemilucilia</i>	102.7
<i>oehricornis</i> (Wiedemann), 1830 (<i>Musca</i>), <i>Phaenicia</i>	102.10
<i>ocularis</i> (Shannon), 1926 (<i>Lucilia</i>), <i>Phaenicia</i>	102.11
<i>oculata</i> (Townsend), 1908 (<i>Lucilia</i>), <i>Phaenicia</i>	102.10
<i>oculatis</i> , error, <i>Phaenicia</i>	102.11
<i>oreoquina</i> (Bigot), 1877 (<i>Somomya</i>), <i>Phaenicia</i>	102.10
<i>ortogesa</i> (Walker), 1849 (<i>Sarcophaga</i>), <i>Neta</i>	102.13
<i>pallasecns</i> (Shannon), 1924 (<i>Lucilia</i>), <i>Phaenicia</i>	102.10
<i>pallidibasis</i> (Bigot), 1877 (<i>Somomya</i>), <i>Phaenicia</i>	102.11
<i>Paralucilia</i> Brauer & Bergenstamm, 1891	102.8
<i>Paralucilia</i> , authors	102.5
<i>paraensis</i> (Macquart), 1843 (<i>Lucilia</i>), <i>Phaenicia</i>	102.10
<i>parva</i> Shannon, 1926, <i>Hemilucilia</i>	102.7
<i>pauciseta</i> (Aldrich), 1922 (<i>Mesembrinella</i>), <i>Eumesembrinella</i> ...	102.2
<i>paytensis</i> (Townsend), 1893 (<i>Calliphora</i>), <i>Neta</i>	102.13
<i>peregrina</i> Aldrich, 1922, <i>Mesembrinella</i>	102.4
<i>peruviana</i> Robineau-Desvoidy of Macquart, 1851, <i>Neta</i>	102.13
<i>peruviana</i> Robineau-Desvoidy, 1830 (<i>Lucilia</i>)	102.15
<i>peruviana</i> Robineau-Desvoidy, 1830 (<i>Calliphora</i>)	102.12
<i>phacoptera</i> Wulp, 1882 (<i>Calliphora</i>)	102.16
<i>Phaenicia</i> Robineau-Desvoidy, 1863	102.9, 102.10
<i>phanda</i> (Walker), 1849 (<i>Musca</i>), <i>Cochliomyia</i>	102.6
<i>philippiana</i> Rondani, 1863 (<i>Somomya</i>)	102.16
<i>Phormia</i> Robineau-Desvoidy, 1830	102.8
<i>Phormiini</i>	102.8
<i>pictierus</i> (Thomson), 1869 (<i>Lucilia</i>), <i>Cochliomyia</i>	102.7
<i>pictipennis</i> Aldrich, 1922, <i>Mesembrinella</i>	102.4
<i>pionia</i> (Walker), 1849 (<i>Musca</i>) <i>Phaenicia</i>	102.11
<i>placii</i> (Robineau-Desvoidy), 1830 (<i>Chrysomya</i>), <i>Cochliomyia</i> ...	102.6
<i>porticola</i> (Thomson), 1869 (<i>Lucilia</i>), <i>Cochliomyia</i>	102.7
<i>praepes</i> Giglio-Tos, 1893 (<i>Calliphora</i>)	102.16
<i>praescia</i> (Giglio-Tos), 1893 (<i>Lucilia</i>), <i>Phaenicia</i>	102.11
<i>primaveris</i> (Shannon & Dei Ponte), 1926 (<i>Lucilia</i>), <i>Phaenicia</i> ..	102.10
<i>princeps</i> Rondani, 1848 (<i>Lucilia</i>)	102.16
<i>problematica</i> (Johnson), 1913 (<i>Lucilia</i>), <i>Phaenicia</i>	102.11
<i>procrna</i> (Walker), 1849 (<i>Sarcophaga</i>), <i>Sarconesia</i>	102.14
<i>Promesembrinella</i> Hall, 1948	102.3
<i>pseudoseriata</i> (Gaminara), 1930 (<i>Lucilia</i>), <i>Phaenicia</i>	102.10
<i>pueblensis</i> (Bigot), 1877 (<i>Somomya</i>), <i>Phaenicia</i>	102.10
<i>punctipennis</i> (Macquart), 1848 (<i>Lucilia</i>), <i>Phaenicia</i>	102.10
<i>purpurata</i> (Aldrich), 1922 (<i>Mesembrinella</i>), <i>Huascaromusca</i> ...	102.3
<i>purpurata</i> Séguy, 1925 (<i>Mesembrinella</i>)	102.5

<i>purpurca</i> (Walker), 1852 (Musca), Chloroprocta	102.5
<i>purpurescens</i> (Walker), 1837 (Musca), Phaenicia	102.11
<i>putrida</i> (Fabricius), 1775 (Musca), Phaenicia	102.10
<i>quadrilincata</i> (Fabricius), 1805 (Musca), Eumesebrinella	102.2
<i>quadrilineata</i> Fabricius of Giglio-Tos, 1893 Mesembrinella	102.4
<i>quadrilineata</i> Fabricius of Surcouf, 1919, Eumesebrinella	102.2
<i>quadrisignata</i> (Thomson), 1869 (Lucilia), Cochliomyia	102.7
<i>quicta</i> Giglio-Tos, 1893 (Lucilia)	102.16
<i>randa</i> (Walker), 1849 (Dexia), Eumesebrinella	102.2
<i>regina</i> (Meigen), 1826 (Musca), Phormia	102.9
<i>rica</i> (Shannon), 1926 (Lucilia), Phaenicia	102.11
<i>rubrifrons</i> (Macquart), 1851 (Lucilia), Cochliomyia	102.6
<i>ruficornis</i> (Macquart), 1846 (Lucilia), Phaenicia	102.10
<i>rufipalpis</i> (Macquart), 1843 (Calliphora)	102.16
<i>rufipalpis</i> Macquart, 1851, Toxotarsus	102.15
<i>rufipes</i> (Macquart), 1843 (Calliphora), Paralucilia	102.8
<i>Sarconesia</i> Bigot, 1857	102.13
<i>Sarconesiomima</i> Lopes & Albuquerque, 1955	102.14
<i>Sarconesiopsis</i> Townsend, 1918	102.14
<i>segmentaria</i> (Fabricius), 1805 (Musca); Hemilucilia	102.7
<i>selkirki</i> Enderlein, 1940, Callitropypa	102.13
<i>semitra</i> Schiner, 1868, Calliphora	102.12
<i>semidiphana</i> (Rondani), 1850 (Mya), Hemilucilia	102.7
<i>semiflava</i> (Aldrich), 1925 (Mesembrinella), Huascaromusca	102.3
<i>semihyalina</i> Mello, 1967, Mesembrinella	102.4
<i>semiviridis</i> Wulp, 1896, Chloroprocta	102.5
<i>semiviolacea</i> (Bigot), 1877 (Somomyia), Phaenicia	102.11
<i>sericata</i> (Meigen), 1826 (Musca), Phaenicia	102.11
<i>setosa</i> James, 1966, Phaenicia	102.11
<i>silvarum</i> (Meigen), 1826 (Musca), Bufolucilia	102.9
<i>smaragdula</i> (Robineau-Desvoidy), 1830 (Lucilia), Phaenicia	102.10
<i>socors</i> (Walker), 1861 (Calliphora), Mesembrinella	102.4
<i>socia</i> (Robineau-Desvoidy), 1830 (Chrysomya), Cochliomyia	102.6
<i>soulouquina</i> (Bigot), 1877 (Somomyia), Phaenicia	102.11
<i>spicata</i> Aldrich, 1925, Mesembrinella	102.4
<i>splendens</i> Macquart, 1843, Blepharicnema	102.9
<i>splendens</i> (Macquart), 1851 (Phryssopoda), Neta	102.13
<i>splendens</i> (Macquart), 1843 (Phryssopoda), Sarconesiopsis	102.14
<i>splendida</i> Townsend, 1918, Chlorobrachycoma	102.13
<i>subrectineuris</i> (Macquart), 1851 (Lucilia), Phaenicia	102.10
<i>taeniaria</i> (Thomson), 1869 (Lucilia), Paralucilia	102.8
<i>taniaria</i> (Wiedemann), 1830 (Musca), Cochliomyia, error	102.6
<i>tibialis</i> (Aldrich), 1922 (Mesembrinella), Huascaromusca	102.3
<i>tibialis</i> (Macquart), 1851 (Calliphora), Cochliomyia	102.7
<i>tibialis</i> (Robineau-Desvoidy), 1830 (Chrysomya), Cochliomyia ...	102.6
<i>townsendi</i> Shannon, 1926, Hemilucilia	102.7
<i>townsendia</i> , error, Hemilucilia	102.7
<i>Toxotarsinac</i>	102.13



<i>Toxotarsus</i> Macquart, 1851	102.14
<i>transmarina</i> Rondani, 1863 (<i>Somomyia</i>)	102.16
<i>transposita</i> Séguy, 1925, <i>Mesembrinella</i>	102.4
<i>tristriata</i> (Verhuel), 1850 (<i>Calliphora</i>), <i>Cochliomyia</i>	102.7
<i>Trixoneura</i> Shannon, 1926	102.14
<i>turbida</i> (Walker), 1952 (<i>Musca</i>), <i>Cochliomyia</i>	102.7
<i>umbrosa</i> Aldrich, 1922, <i>Mesembrinella</i>	102.4
<i>uniseta</i> (Aldrich), 1925 (<i>Mesembrinella</i>), <i>Huascaromusca</i>	102.3
<i>varipalpis</i> Macquart, 1843 (<i>Lueilia</i>)	102.16
<i>vercna</i> (Walker), 1849 (<i>Musca</i>), <i>Myiolucilia</i>	102.8
<i>versicolor</i> Bigot, 1857, <i>Sarconesia</i>	102.14
<i>vicina</i> Robineau-Desvoidy, 1830, <i>Calliphora</i>	102.12
<i>violacea</i> (Macquart), 1843 (<i>Calliphora</i>), <i>Cochliomyia</i>	102.6
<i>violacea</i> (Macquart), 1847 (<i>Lueilia</i>), <i>Phaenicia</i>	102.11
<i>Viridinsula</i> Shannon, 1926	102.11
<i>viridula</i> (Robineau-Desvoidy), 1830 (<i>Chrysomya</i>), <i>Cochliomyia</i> ..	102.6
<i>vogelsangi</i> Mello, 1967, <i>Huascaromusca</i>	102.3
<i>wheeleri</i> (Hough), 1899 (<i>Chrysomyia</i>), <i>Paralucilia</i>	102.8
<i>xanthorrhina</i> (Bigot), 1888 (<i>Calliphora</i>), <i>Huascaromusca</i>	102.3
<i>Xerophilophaga</i> Enderlein, 1933	102.9



DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

103

FAMILY SARCOPHAGIDAE

HUGO SOUZA LOPES¹*Seção de Entomologia, Instituto Oswaldo Cruz
Rio de Janeiro, Guanabara
Brasil*

As in many other families of Diptera, the Sarcophagidae have been studied only with reference to the external characters; the male genitalia was used only for the characterization of species. However, after the important publication of Roback on the structures of the penis of many species (1954) I am convinced that an accurate study of the male and female genitalia, and of the larvae, will be necessary for the establishment of a true phylogenetic system for the Sarcophagidae.

With basis on those characters, four subfamilies are provisionally considered for the Neotropical species (Miltogramminae, Macronychiinae, Paramacronychiinae, and Sarcophaginae); the Sarcophaginae are also provisionally divided into six tribes (Microcerellini, Notochaetini, Tephromyiini, Raviniini, Sarcophagulini, and Sarcophagini). Genera are arranged phylogenetically within each tribe. The family is represented, in the area covered by this Catalogue, by 144 genera and 622 species.

Aldrich (1916) included some Neotropical species in his revision of the North American fauna, and published a very good contribution to the knowledge of the American species, redescribing those of the earlier authors (1930). Townsend (1927) considered the Neotropical genera in his study of the "humid tropical America" Muscoidea, examined the types of some species (1931a, 1931b), and referred to all known genera in his Manual of Myiology (1935b, 1937b, 1938). Hall published valuable contributions concerning the sarcophagid fauna of Panama (1933), and Patagonia and South Chile (1937b). Curran & Walley (1934) have dealt with Guyanan species, and Blanchard (1934, 1942c) with those of Argentina. Lopes published a paper on the female genital apparatus (1941a) and a morphological study of the larvae (1943). Dodge (1965d, 1965e) studied the fauna of the Bahamas and Jamaica.

Habits and special information referring to subfamilies, tribes or single species are added under each of these taxa, in this catalogue.

1 Partially supported by the Conselho Nacional de Pesquisas, Brasil.

Subfamily MILTOGRAMMINAE

MILTOGRAMMINAE Brauer & Bergenstamm, 1889:8 (76), 47 (112) (as Milto-grammidac).

Larvac of the known species of the subfamily feed on provisions of wasps' nests. Those of *Amobia*, in Brazil, live in nests of Trypoxylionidae, Sphecidae, and Eumenidae.

Genus AMOBIA Robineau-Desvoidy

AMOBIA Robineau-Desvoidy, 1830:96. Type-species, *conica* Robineau-Desvoidy (mon.) = *signata* (Meigen).

Pachyophthalmus Brauer & Bergenstamm, 1889:117 (49). Type-species, *Tachina signata* Meigen (mon.).

Subgenus SARCOMACRONYCHIA Townsend

SARCOMACRONYCHIA Townsend, 1892f:100 (as genus). Type-species, *unica* Townsend (orig. des.) = *erythrura* (Wulp).

AURIFACIES (Séguy), 1931:782, figs. 1-2 (*Pachyophthalmus*) (larva III, nest of *Scelifron* or *Trypoxylon*). Type-locality: Venezuela, Puerto La Cruz. Distr. — Venezuela.

ERYTHRURA (Wulp), 1890:89 (*Miltogramma*). Type-locality: Mexico, Presidio. Distr. — U.S.A., Mexico.

unica Townsend, 1892f:101 (*Sarcomacronychia*). Type-locality: U.S.A., New Mexico.

sarcophagoides Townsend, 1892g:165 (*Sarcomacronychia*). Type-locality: U.S.A., New Mexico.

FLORIDENSIS (Townsend), 1892c:80 (*Pachyophthalmus*). Type-locality: "U.S.A." Distr. — U.S.A., Cuba, Jamaica, Puerto Rico, Panama, Peru.

trypoxylonis Townsend, 1893b:165 (*Sarcomacronychia*). Type-locality: U.S.A., Ohio.

trivittata Townsend, 1912a:363 (*Sarcomacronychia*). Type-locality: Peru, Piura. Ref. — Allen, 1926:10 (type examined).

ssp. ANTILLARUM Richards, 1935:212 (*Pachyophthalmus*). Type-locality: Trinidad, Tunapuma (nest of *Trypoxylon paliditarse* Sauss.). Distr. — Trinidad.

ORNATICAUDA (Aldrich), 1928:9 (*Pachyophthalmus*). Type-locality: Chile, Perales. Distr. — Chile.

PRIDEI (Richards), 1936:89 (*Pachyophthalmus*). Type-locality: Paraguay, Chaco (nest of *Trypoxylon*). Distr. — Paraguay.

SARCOPIAGINA (Wulp), 1890:90 (*Miltogramma*). Type-locality: Mexico, Vera Cruz. Distr. — Mexico.

Genus EUMACRONYCHIA Townsend

EUMACRONYCHIA Townsend, 1892f:98. Type-species, *decens* Townsend (orig. des.).

Ref. — Reinhard, 1965b (revision).

- ARMILLA Reinhard, 1965b:344. Type-locality: U.S.A., Nevada, Mt. Rose. Distr. — U.S.A., Mexico.
- BORREGOENSIS Reinhard, 1965b:344. Type-locality: U.S.A., New Mexico, Las Cruces. Distr. — U.S.A., Mexico.
- DECENS Townsend, 1892f:99. Type-locality: U.S.A., New Mexico, Las Cruces. Distr. — U.S.A., Mexico.
- DUPLICATA Reinhard, 1944:57. Type-locality: U.S.A., New Mexico, Albuquerque. Distr. — U.S.A., Mexico.
- NIGRICORNIS Allen, 1926:90. Type-locality: U.S.A., Ohio, Sugar Grove. Distr. — U.S.A.
- ssp. TRICOSA Reinhard, 1965b:341. Type-locality: U.S.A., Texas, College Station. Distr. — U.S.A., Mexico.
- PERSONELLA Reinhard, 1965b:399. Type-locality: U.S.A., California, Antioch. Distr. — U.S.A., Mexico.
- STERNALIS Allen, 1926:89. Type-locality: U.S.A., Texas, Galveston. Distr. — U.S.A., Mexico, Guatemala, Nicaragua.
- prolixa Reinhard, 1939:67. Type-locality: U.S.A., Texas, Galveston.
- STIGMOSA Reinhard, 1965b:346. Type-locality: Mexico, Sonora, Guaymas. Distr. — Mexico.

Genus HILARELLA Rondani

- HILARELLA Rondani, 1856a:70. Type-species, *Miltogramma hilarella* Zetterstedt (taut.).
- HILARELLA (Zetterstedt), 1844:1212 (*Miltogramma*). Type-locality: "Europe". Distr. — Europe; U.S.A., Mexico.
- zetterstedti Rondani, 1856a:70 (*Miltogramma*). Type-locality: "Europe".

Genus METOPIA Meigen

- METOPIA Meigen, 1803:280. Type-species, *Musca leucocephala* Rossi (as Panzer, mon.) = *argyrocephala* (Meigen).

Subgenus METOPIA Meigen

- ARGYROCEPHALA (Meigen), 1803:280 (*Tachina*). Type-locality: "Europe". Distr. — Palaearctic, Nearctic; Mexico, Cuba, Jamaica, Peru.
- leucocephala Rossi, 1790:306 (*Musca*; preocc. Villers, 1789). Type-locality: "Italy".
- meridiana Townsend, 1912a:361. Type-locality: Peru, Piura. Ref. — Allen, 1926:49 (type examined).
- PAUCISETA Dodge, 1966b:701. Type-locality: Peru, Tingo Maria. Distr. — Peru.
- PERPENDICULARIS Wulp, 1890:115, pl. 3, figs. 18, 18a. Type-locality: "Mexico". Distr. — Mexico.

Subgenus ALLENANICIA Townsend

- Parametopia* Townsend, 1916b:619 (preocc. Reitter, 1884). Type-species, *morrisoni* Townsend (orig. des.) = *lateralis* (Macquart).



ALLENANICIA Townsend, 1935a:68 (as genus). Type-species, *Metopia sinipalpis* Allen (orig. des.).

LATEROPILI Allen, 1926:57. Type-locality: Cuba, Havana. Distr. — U.S.A., Cuba.

Genus JUQUIANICIA Townsend

JUQUIANICIA Townsend, 1934a:112. Type-species, *juquiana* Townsend (orig. des.).

JUQUIANA Townsend, 1934a:112. Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.

Genus OPSIDIA Coquillett

OPSIDIA Coquillett, 1895:102. Type-species, *gonioides* Coquillett (orig. des.).
Opsidiopsis Townsend, 1919b:544. Type-species, *oblata* Townsend (orig. des.).

Neopsidia Townsend, 1928:144. Type-species, *Opsidia metopioides* Allen (orig. des.).

Goniogramma Townsend, 1931b:316. Type-species, *herbsti* Townsend (orig. des.) = *intonsa* Aldrich.

INTONSA Aldrich, 1928:8. Type-locality: Chile, Santiago. Distr. — Chile.
herbsti Townsend, 1931b:317 (*Goniogramma*). Type-locality: Chile, Concepción.

METOPIOIDES Allen, 1926:44. Type-locality: Panama, Porto Bello. Distr. — Panama.

VITTATA Reinhard, 1965a:79. Type-locality: "West Indies". Distr. — West Indies.

Genus RUPUNUNIA Townsend

RUPUNUNIA Townsend, 1935a:74. Type-species, *polita* Townsend (orig. des.).

POLITA Townsend, 1935a:74. Type-locality: Guyana, Rupununi. Distr. — Guyana.

Genus ANICIA Robineau-Desvoidy

ANICIA Robineau-Desvoidy, 1863:99. Type-species, *sabulosa* Robineau-Desvoidy (Coquillett, 1910:506) = *campestris* (Fallén).

BRASILLANA Townsend, 1929:565. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil.

Genus SENOTAINIA Macquart

SENOTAINIA Macquart, 1846:295 (sep., p. 167) (as *Senetainia*, pp. 295 and 167; as *Senotaina*, pp. 364 and 236). Type-species, *rubriventris* Macquart (orig. des.).

Euselenomyia Townsend, 1912a:364. Type-species, *peruviansis* Townsend (orig. des.) = *trilineata* (Wulp).

Eusenotainia Townsend, 1915b:22. Type-species, *Hilarella rufiventris* Coquillett (orig. des.).

Microsenotainia Townsend, 1916b:618. Type-species, *Senotainia nana* Coquillett (orig. des.).

- Ref. — Allen, 1926:16 (revision).
- BRASILIENSIS (Townsend), 1929:365 (*Euselcnomyia*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil.
- FLAVICORNIS (Townsend), 1891:355 (*Miltogramma*). Type-locality: U.S.A., Illinois. Distr. — U.S.A., Cuba.
- similis* Townsend, 1891:357 (*Miltogramma*). Type-locality: U.S.A., Illinois.
- FULVICORNIS (Wulp), 1890:89 (*Miltogramma*). Type-locality: Mexico, Guerrero. Distr. — Mexico. Ref. — Allen, 1926:21 (type examined).
- LITORALIS Allen, 1924:90. Type-locality: U.S.A., Mississippi. Distr. — U.S.A., Bahamas.
- NANA (Wulp), 1892:195 (*Miltogramma*). Type-locality: Mexico, Guerrero. Distr. — Mexico. Ref. — Allen, 1926:21.
- RUBRIVENTRIS Macquart, 1846:295 (sep., p. 167). Type-locality: U.S.A., Texas, Galveston. Distr. — U.S.A., Jamaica, Puerto Rico.
- decisa* Townsend, 1892c:81 (*Miltogramma*). Type-locality: U.S.A., Florida.
- TRILINEATA (Wulp), 1890:89 (*Miltogramma*). Type-locality: Mexico, Presidio. Distr. — U.S.A., Mexico, Bahamas, Jamaica, Nicaragua, Costa Rica, Peru. Ref. — Allen, 1926:32.
- americanus* Brauer & Bergenstamm, 1891:57 (361) (*Arrenopus*). Type-locality: U.S.A., Georgia.
- argentifrons* Townsend, 1891:357 (*Miltogramma*). Type-locality: U.S.A., Illinois.
- cinerascens* Townsend, 1891:358 (*Miltogramma*). Type-locality: U.S.A., Illinois.
- peruviansis* Townsend, 1912a:364 (*Euselenomyia*). Type-locality: Peru, Piura.

Genus SPHENOMETOPA Townsend

- SPHENOMETOPA Townsend, 1908:64. Type-species, *Araba nebulosa* Coquillett (orig. des.).
- Euaraba* Townsend, 1915a:20. Type-species, *Araba tergata* Coquillett (orig. des.).
- Arabiopsis* Townsend, 1915d:285. Type-species, *cocklei* Townsend (orig. des.).
- NEBULOSA (Coquillett), 1902:200 (*Araba*). Type-locality: Mexico, Chihuahua, Sierra Madre. Distr. — U.S.A., Mexico. Ref. — Allen, 1926:66.
- TERGATA (Coquillett), 1895:103 (*Araba*). Type-locality: U.S.A., Illinois. Distr. — U.S.A., Mexico. Ref. — Allen, 1926:64.

Subfamily MACRONYCHIINAE

- MACRONYCHIINAE Brauer & Bergenstamm, 1889:8 (76), 49 (117); Rohdendorf, 1937:48.

Some species of this subfamily live in the nests of Hymenoptera, but "the species with a sharp piercing sternotheca evidently inject their maggots within the integument of caterpillars or other insects" (Townsend, 1935:213).

Genus MACRONYCHIA Rondani

MACRONYCHIA Rondani, 1859:229, 239 (as *Macronichia*, p. 229). Type-species, *Tachina agrestis* Fallén (mon.).

Amobiopsis Townsend, 1915b:20. Type-species, *Amobia aurata* Coquillett (orig. des.).

CONFUNDENS (Townsend), 1915b:20 (*Amobiopsis*). Type-locality: U.S.A., New Hampshire. Distr. — U.S.A., Mexico.

ORNATA (Townsend), 1917c:221 (*Amobiopsis*). Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Distr. — Brazil.

Genus ITAMOBIA Townsend

ITAMOBIA Townsend, 1927a:224. Type-species, *ornata* Townsend (orig. des.).

AURIFRONS (Hall), 1937b:351, figs. 61a-b (*Macronychia*). Type-locality: Argentina, Misiones. Distr. — Argentina.

ORNATA Townsend, 1927a:224, 321. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil. Ref. — Lopes, 1936a:851, figs. 10-13 (paratype examined).

Genus DOLICHAMOBIA Townsend

DOLICHAMOBIA Townsend, 1915c:408. Type-species, *auromaculata* Townsend (orig. des.).

AUROMACULATA Townsend, 1915c:408. Type-locality: Peru, Oroya. Distr. — Peru.

Genus NEOPHYTODES Townsend

NEOPHYTODES Townsend, 1931b:317. Type-species, *lindneri* Townsend (orig. des.).

LINDNERI Townsend, 1931b:318. Type-locality: Argentina, Chaco, San José. Distr. — Argentina.

Subfamily PARAMACRONYCHIIDAE

PARAMACRONYCHIIDAE Brauer & Bergenstamm, 1889:8 (76), 47 (115); Rohdendorf, 1937:46.

Nothing is known about the habits of the Neotropical species of this subfamily. The larvae of the Palaeartic and Nearctic species are parasitic or live on dead insects or in pustules in the skin of mammals. The species of the Oriental genus *Goniophyto* are collected only on sea beaches. All species considered here show a median transverse series of bristles in the apparently first genital tergite of the male.

Genus CATTASOMA Reinhard

CATTASOMA Reinhard, 1947:97. Type-species, *mediocre* Reinhard (orig. des.).

FESTINANS Reinhard, 1947:98, fig. 4. Type-locality: U.S.A., Texas, Marathon. Distr. — U.S.A., Mexico.

Genus GALAPAGOMYIA Bischof

GALAPAGOMYIA Bischof, 1904:100. Type-species, *Microcerella steindachneri* Brauer & Bergenstamm (orig. des.).

Microcerella Macquart of Brauer & Bergenstamm, 1891:85 (389).

STEINDACHNERI (Brauer & Bergenstamm), 1891:85 (389) (*Microcerella*).
Type-locality: Galápagos Is. Distr. — Galápagos Is.

Genus PANAMAPHYTO Townsend

PANAMAPHYTO Townsend, 1931b:314. Type-species, *Miltogramma biseta* Thomson (orig. des.).

BISETA (Thomson), 1869:542 (*Miltogramma*). Type-locality: "Panama".
Distr. — Panama. Ref. — Townsend, 1931a:314.

Subfamily SARCOPHAGINAE

SARCOPHAGINAE Macquart, 1835:29 (as Sarcophagini).

Tribe MICROCERELLINI

MICROCERELLINI Séguy, 1928:99, 108; Roback, 1954:58 (as Hypopeltina).

Species of this tribe show bare or pubescent arista. Living specimens of all the species I have collected have green, instead of red, eyes and the larvae of some were reared in decaying animal matter. Most of the Neotropical species live in temperate regions or on high mountains. *Doringia*, *Parahybopelta*, and *Scopaediscus* were reared from Acrididae and Bulimulidae.

Genus MICROCERELLA Macquart

MICROCERELLA Macquart, 1851:209 (236). Type-species, *rufomaculata* Macquart (orig. des.).

RUFOMACULATA Macquart, 1851:209 (236), pl. 22, fig. 1. Type-locality: "Chile". Distr. — Chile. Refs. — Séguy, 1926:184; Townsend, 1931a:72 (type examined).

SARCOPHAGINA Thomson, 1869:541. Type-locality: "Chile". Distr. — Chile.

Genus AULACOPHYTO Townsend

AULACOPHYTO Townsend, 1919a:158. Type-species, *auromaculata* Townsend (orig. des.).

Ref. — Hall, 1937b:350, 360.

AUROMACULATA Townsend, 1919a:158. Type-locality: Peru, Oroya. Distr. — Peru, Chile. Paratype examined.

RUSCA Hall, 1937b:350, 361. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.

sp. Dodge, 1966b:675, figs. 1a-b. Locality: Chile, Longuen.

Genus MICROCERELLAMYIA Hall

MICROCERELLAMYIA Hall, 1937b:350, 354. Type-species, *retusa* Hall (orig. des.).



RETUSA Hall, 1937b:350, 354, figs. 63a-d. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.

Genus EUPARAPHYTO Townsend

EUPARAPHYTO Townsend, 1912a:359. Type-species, *alpina* Townsend (orig. des.).

ALPINA Townsend, 1912a:359. Type-locality: Peru, Tirapata. Distr. — Peru.

AUROMACULATA Townsend, 1919a:157. Type-locality: Peru, Oroya. Distr. — Peru.

Genus ITIOPHYTO Hall

ITIOPHYTO Hall, 1937b:350, 355. Type-species, *engelii* Hall (orig. des.).

ENGELI Hall, 1937b:350, 356, figs. 64a-b. Type-locality: Chile, Tanuncie. Distr. — Chile.

Genus EUMICROCERELLA Hall

EUMICROCERELLA Hall, 1938a:253. Type-species, *duca* Hall (orig. des.).

DUCA Hall, 1938a:253, fig. 1. Type-locality: Argentina, Mendoza. Distr. — Argentina.

Genus PARAHYDOPELTA Blanchard

PARAHYDOPELTA Blanchard, 1955:30. Type-species, *salavini* Blanchard (orig. des.).

SALAVINI Blanchard, 1955:30, fig. 2 (ex Acrididae). Type-locality: Argentina, Patagonia. Distr. — Argentina.

nana Enderlein, 1928:40 (*Dasypygia*). *Nomen nudum*.

Genus AUSTROHARTIGIA Townsend

AUSTROHARTIGIA Townsend, 1937a:115 (March, 1937). Type-species, *magellanic* Townsend (orig. des.) = *spinigena* Rondani.

Phacosarcodexia Hall, 1937b:350, 362 (April 25, 1937). Type-species, *aldrichi* Hall (orig. des.) = *spinigena* Rondani.

EDWARDSI (Hall), 1937b:351, 364, figs. 68a-d (*Phacosarcodexia*). Type-locality: Chile, Perales. Distr. — Chile.

SPINIGENA (Rondani), 1846:26 (*Sarcophaga*). Type-locality: Chile, Valdivia. Distr. — Chile, Argentina. Ref. — Hall, 1937b:350, 363.

ruficauda Aldrich, 1928:10 (*Brachicoma*; preocc. Wulp, 1890). Type-locality: Chile, Angol.

magellanica Townsend, 1937a:116. Type-locality: Chile, Punta Arenas. Paratype examined.

aldrichi Hall, 1937b:348, 350, 362, figs. 67a-c (*Phacosarcodexia*; nom. nov. for *ruficauda* Aldrich). Type-locality: Chile, Angol.

SPINOSA (Hall), 1937b:351, 365 (*Phacosarcodexia*). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Chile, Argentina.

SUBANDINA (Blanchard), 1938:377, fig. 6 (*Dorinia*). Type-locality: Argentina, Mendoza. Distr. — Argentina.

Genus ALACCOPROSOPA Townsend

- ALACCOPROSOPA Townsend, 1934b:202. Type-species, *apicalis* Townsend (orig. des.).
- Mimophytomyia* Hall, 1937b:350, 357. Type-species, *chilensis* Hall (orig. des.) = *apicalis* Townsend.
- APICALIS Townsend, 1934b:202. Type-locality: Chile, Concepción. Distr. — Chile, Argentina. Ref. — Lopes, 1938e:565, pl. 4, figs. 5-7.
- chilensis* Hall, 1937b:350, 357, figs. 65a-d (*Mimophytomyia*). Type-locality: Argentina, Rio Negro, Lake Nahuel Huapi.
- MALLOCHI (Hall), 1937b:350, 359, figs. 66a-c (*Mimophytomyia*). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina, Chile.

Genus XANTHOBACHYCOMA Townsend

- XANTHOBACHYCOMA Townsend, 1927a:232. Type-species, *analís* Townsend (orig. des.).
- ANALIS Townsend, 1927a:232, 365. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil (Minas Gerais, Espírito Santo, Rio de Janeiro, São Paulo, Santa Catarina).
- bicoloricauda* Enderlein, 1928c:151 (*Mesothysia*). Type-locality: "Brazil".
- WEYRAUCHI Lopes, 1954:69, figs. 14-18. Type-locality: Peru, Atocongo, near Lima. Distr. — Peru.
- WYGODZINSKYI Lopes, 1954:67, figs. 8-13. Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil.

Genus TOWNSENDIMYIA Prado & Fonseca

- TOWNSENDIMYIA Prado & Fonseca, 1932b:167. Type-species, *Stephanostoma argenteum* Prado & Fonseca (orig. des.) = *halli* (Engel).
- HALLI (Engel), 1931:151, fig. 20 [*Sarcophaga* (*Bellieria*)]. Type-locality: Bolivia, n. Chiquitos. Distr. — Bolivia, Brazil (Minas Gerais, São Paulo), Argentina.
- georgina* Wiedemann of Mattos, 1919:53, pl. 3 (*Sarcophaga*), misident. *argenteum* Prado & Fonseca, 1932a:35, fig. 6 (*Stephanostoma*). Type-locality: Brazil, São Paulo.

Genus DORINGIA Weyenbergh

- DORINGIA Weyenbergh, 1875:85. Type-species, *Nemorea acridiorum* Weyenbergh (orig. des.).
- ACRIDIORUM (Weyenbergh), 1875:85 (*Nemorea*) (ex Acrididae). Type-locality: "Argentina". Distr. — Argentina. Ref. — Crouzel, 1950:291, figs. (larva I).
- minuta* Lahille, 1907:82, 86, figs. 8-11 (*Sarcophaga*) (ex Acrididae). Type-locality: "Argentina".
- ERYTHROPYGA (Lopes), 1936b:71, figs. 1-2 (*Xenopopia*). Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil (Rio de Janeiro, Guanabara, São Paulo). Ref. — Lopes, 1940:950 (ex *Bulimulus tenuissimus*).

Genus XENOPPIA Townsend

XENOPPIA Townsend, 1915b:20. Type-species, *hypopygialis* Townsend (orig. des.).

Camptopyga Aldrich, 1916:41. Type-species, *aristata* Aldrich (orig. des.)
= *hypopygialis* Townsend.

HYPOPYGIALIS Townsend, 1915b:20. Type-locality: U.S.A., Georgia. Distr.
— U.S.A., Mexico.

intermedia Townsend of Coquillett, 1897:132 (*Sarcotachinella*), misident.

aristata Aldrich, 1916:42, figs. 10a-b (*Camptopyga*). Type-locality:
U.S.A., Texas, Alta Loma.

Genus SCOPAEDISCUS Blanchard

SCOPAEDISCUS Blanchard, 1939:802. Type-species, *muchni* Blanchard (orig. des.).

MUEHNI Blanchard, 1939:802, fig. 2 (ex *Oeceticus kirbyi*). Type-locality:
Argentina, San Miguel. Distr. — Argentina.

aristata Aldrich of Engel, 1931:141, fig. 6 (*Camptopyga*), misident.

Genus SAROTHROMYIOPS Townsend

SAROTHROMYIOPS Townsend, 1917b:191, 195. Type-species, *cinctus* Townsend
(orig. des.) = *inoa* (Walker).

Prosthetocirca Townsend, 1917b:192, 195. Type-species, *cana* Townsend
(orig. des.) = *inoa* (Walker).

INOA (Walker), 1849:832 (*Sarcophaga*). Type-locality: Galápagos Is.
Distr. — Galápagos Is. Refs. — Austen, 1907:342; Cole, 1927: fig.
197.

cinctus Townsend, 1917b:196. Type-locality: Galápagos Is., Culpepper I.

cana Townsend, 1917b:196 (*Prosthetocirca*). Type-locality: Galápagos
Is., Narborough I. Ref. — Townsend, 1931:78 (type examined).

Genus CHRYSAGRIA Townsend

CHRYSAGRIA Townsend, 1935a:72. Type-species, *duodecimpunctata* Townsend
(orig. des.).

DUODECIMPUNCTATA Townsend, 1935a:72. Type-locality: Brazil, Pernambuco.
Tapera. Distr. — Brazil.

Genus CATHETERONYCHIA Townsend

CATHETERONYCHIA Townsend, 1927a:230. Type-species, *chactosa* Townsend
(orig. des.).

CHACTOSA Townsend, 1927a:230, 295. Type-locality: Brazil, São Paulo,
Itaquaquecetuba. Distr. — Brazil (São Paulo).

Genus OPSOPHYTOPSIS Townsenu

OPSOPHYTOPSIS Townsend, 1918a:163. Type-species, *insularis* Townsend
(orig. des.).

INSULARIS Townsend, 1918a:163. Type-locality: Galápagos Is., Albermale I. Distr. — Galápagos Is.

Genus NEOPHYTO Townsend

NEOPHYTO Townsend, 1908:55. Type-species, *Phyto setosa* Coquillett (orig. des.).

Phytodes Coquillett, 1910a:127. Type-species, *hirculus* Coquillett (orig. des.).

INCONISMUS (Reinhard), 1957:99 (*Phytodes*). Type-locality: Mexico, San Luis Potosí, El Maíz. Distr. — Mexico.

Genus CHAPINIOLA Dodge

CHAPINIOLA Dodge, 1965c:253. Type-species, *impressa* Dodge (orig. des.).

IMPRESSA Dodge, 1965e:253, figs. 1-3. Type-locality: Colombia, Guasea. Distr. — Colombia.

Tribe NOTOCHAETINI, new

All species included here may be parasitic. Larvae of the studied species do not show a complete elypeal arch. Some species of *Notochaeta* are parasitic on Oligochaeta, Amphibia, and Lacertilia. *Johnsonia* was reared from *Succinea* (Mollusea). Adult *Gymnocamptops* and *Lepidodexia* were found on bark of trees.

Genus NOTOCHAETA Aldrich

NOTOCHAETA Aldrich, 1916:52. Type-species, *subpolita* Aldrich (orig. des.) = *fuscianalis* (Wulp).

Micronotochaeta Townsend, 1927a:233. Type-species, *costalis* Townsend (orig. des.).

Actenella Enderlein, 1928a:16. Type-species, *fiebrigi* Enderlein (orig. des.).

Udamoctis Enderlein, 1928a:25. Type-species, *bogotana* Enderlein (orig. des.).

Anolisimya Dodge, 1955:183. Type-species, *blakeae* Dodge (orig. des.). N. SYN.

ALDRICH Lopes, 1942a:362, figs. 1-4 (ex Oligochaeta). Type-locality: Brazil, Guanabara, Rio de Janeiro, Grajaú. Distr. — Brazil (Minas Gerais, Rio de Janeiro, Guanabara, Santa Catarina).

AUGUSTA Aldrich, 1925a:23. Type-locality: Panama, Canal Zone, Corozal. Distr. — Panama. Ref. — Lopes, 1946a:521, figs. 18-23.

BLAKEAE (Dodge), 1955:183, figs. 1-7 (*Anolisimya*) (ex *Anolis carolinensis*). Type-locality: U.S.A., North Carolina. Distr. — U.S.A. (North Carolina, Florida). N. COMB.

BOGOTANA (Enderlein), 1928a:25 (*Udamoctis*). Type-locality: Colombia, Bogotá. Distr. — Colombia.

BUFONIVORA Lopes & Vogelsang, 1953:140, figs. 1-9 (ex Amphibia). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.

COGNATA (Walker), 1852:325 (*Sarcophaga*). Type-locality: "South America". Distr. — South America. Ref. — Aldrich, 1930:18 (type examined).

- COMATA Aldrich, 1925a:22. Type-locality: Costa Rica, La Suiza. Distr. — Costa Rica.
- CONFUSA Lopes, 1946a:537, figs. 56-63. Type-locality: Brazil, Guanabara, Rio de Janeiro, Grajaú. Distr. — Brazil (Minas Gerais, Guanabara).
- COSTALIS (Townsend), 1927a:233, 330 (*Micronotochacta*). Type-locality: Peru, Chaquimayo R. Distr. — Peru.
- CYANEIVENTRIS Lopes, 1946a:542, figs. 66-68. Type-locality: Brazil, Rio de Janeiro, Angra dos Reis. Distr. — Brazil (Rio de Janeiro, Guanabara).
- DIMIDIATA (Wiedemann), 1830:360 (*Sarcophaga*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Aldrich, 1930:4, fig. 25.
- DISTINCTA Lopes, 1947:378, figs. 4-8. Type-locality: Brazil, Amazonas, Manaus. Distr. — Brazil.
- DIVERSA Lopes, 1946a:528, figs. 31-38. Type-locality: Brazil, Guanabara, Rio de Janeiro, Grajaú. Distr. — Brazil (Guanabara).
- DIVERSINERVIS (Wulp), 1895:268, 1896:274, pl. 7, fig. 8 (*Sarcophaga*). Type-locality: Mexico, Tabasco. Distr. — Mexico. Ref. — Aldrich, 1930:31 (type examined).
- FIEBIGI (Enderlein), 1928a:17 (*Actocella*). Type-locality: "Paraguay". Distr. — Paraguay. Refs. — Townsend, 1931a:78 (type examined).
- FUMIPENNIS Lopes, 1946a:525, figs. 24-30. Type-locality: Brazil, Guanabara, Rio de Janeiro, Jardim Botânico. Distr. — Brazil (Guanabara, Paraná).
- FUSCIANALIS (Wulp), 1895:268, 1896:275 (*Sarcophaga*). Type-locality: "Mexico". Distr. — Mexico, Guatemala, Guyana, Brazil (Pará). Ref. — Aldrich, 1930:31 (type examined).
- subpolita* Aldrich, 1916:53, fig. 14. Type-locality: "Guatemala".
- IGNOTA Lopes, 1947:380, figs. 9-13. Type-locality: Brazil, Amazonas, Rio Negro, Umarituba. Distr. — Brazil.
- MICROPYGA (Wulp), 1895:268, 1896:279 (*Sarcophaga*). Type-locality: Mexico, Guerrero. Distr. — Mexico. Ref. — Aldrich, 1930:34 (type examined).
- OBSCURA Lopes, 1950a:354, 360, figs. 13-18. Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- PARVA Lopes, 1946a:519, figs. 12-17. Type-locality: Brazil, Pará, Belém, Utinga. Distr. — Brazil.
- PLUMIGERA (Wulp), 1895:268, 1896:273, pl. 7, figs. 6-6a (*Sarcophaga*). Type-locality: Mexico, Tabasco. Distr. — Mexico. Ref. — Aldrich, 1930:31 (type examined).
- RUSTICA Lopes, 1950a:354, figs. 12-14. Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- TOWNSENDI Aldrich, 1925a:22. Type-locality: Peru, Jaén, Huascaray Ridge. Distr. — Peru.
- UNISETA Lopes, 1950a:353, 357, figs. 8-11. Type-locality: Ecuador, Baños. Distr. — Ecuador.
- VIRIDIS Lopes, 1950a:353, 355, figs. 1-7. Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.

Genus EUFLETCHERIMYIA Townsend

- EUFLETCHERIMYIA Townsend, 1934a:111. Type-species, *aurea* Townsend (orig. des.).

AUREA Townsend, 1934a:111. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil.

Genus MONONOTOCHAETA Townsend

MONONOTOCHAETA Townsend, 1935a:73. Type-species, *myersi* Townsend (orig. des.).

MYERSI Townsend, 1935a:73. Type-locality: "Trinidad". Distr. — Trinidad.

Genus EUNOTOCHAETA Townsend

EUNOTOCHAETA Townsend, 1934b:203. Type-species, *amazonica* Townsend (orig. des.).

AMAZONICA Townsend, 1934b:203. Type-locality: Brazil, Pará, Rio Tapajós. Distr. — Brazil.

Genus DISCOMYIOPHORA Townsend

DISCOMYIOPHORA Townsend, 1927a:233. Type-species, *aurata* Townsend (orig. des.).

AURATA Townsend, 1927a:233, 302. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil.

Genus SISASARCOPHAGA Townsend

SISASARCOPHAGA Townsend, 1928:143. Type-species *metallica* Townsend (orig. des.).

METALLICA Townsend, 1928:144. Type-locality: Peru, Hualлага R., Sisa. Distr. — Peru.

Genus ORODEXIA Townsend

ORODEXIA Townsend, 1927a:227, 341. Type-species, *ornata* Townsend (orig. des.) = *opima* (Wiedemann).

OPIMA (Wiedemann), 1830:364 (*Sarcophaga*). Type-locality: "Brazil". Distr. — Brazil (Rio de Janeiro, São Paulo). Ref. — Aldrich, 1930:6, fig. 21 (type examined).

ornata Townsend, 1927a:341. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Ref. — Lopes, 1936a:850, figs. 8-9 (type examined).

Genus GEJSKESIA Lopes

GEJSKESIA Lopes, 1945a:295. Type-species, *brevigaster* Lopes (orig. des.).

BREVIGASTER Lopes, 1945a:295, figs. 1-6. Type-locality: "Surinam". Distr. — Surinam.

Genus HARPAGOPYGA Aldrich

HARPAGOPYGA Aldrich, 1916:61. Type-species, *Sarcophaga diversipes* Coquillett (orig. des.).

Pachygraphiops Townsend, 1934b:204. Type-species, *pacta* Townsend (orig. des.).

Ref. — Lopes, 1951a:275 (revision).

- ALEIDA Lopes, 1951a:287, figs. 40-43 (female, larva I). Type-locality: Brazil, São Paulo, Ilha Sêca. Distr. — Brazil.
- ALBILURTA Dodge, 1965c:506. Type-locality: Jamaica, Portland. Distr. — Jamaica.
- ANGUSTIVENTRIS Curran & Walley, 1934:477, fig. 34. Type-locality: Guyana, Kartabo. Distr. — Guyana. Ref. — Lopes, 1951a:283, figs. 27-32 (female, larva I).
- ATRATA Dodge, 1965a:505. Type-locality: Jamaica, Manchester. Distr. — Jamaica.
- CUBANA Lopes, 1951a:279, figs. 7-13. Type-locality: Cuba, Oriente, Cristo. Distr. — Cuba.
- diversipes* Coquillett of Aldrich, 1916:62 (in part).
- DISSIMILIS Dodge, 1965c:505 (female). Type-locality: Jamaica, St. Andrew. Distr. — Jamaica.
- DIVERSIPES (Coquillett), 1900:255 (*Sarcophaga*) (female, ex *Laphygma*). Type-locality: Puerto Rico, Fajardo. Distr. — Puerto Rico.
- DOWNESI Dodge, 1965d:485, figs. 12-14. Type-locality: Bahamas, Nassau. Distr. — Bahamas.
- diversipes* Coquillett of Aldrich, 1916: fig. 19 (in part).
- FLAVOMILOSA Lopes, 1951a:285, figs. 33-39 (female, larva I). Type-locality: Brazil, Goiás, Goiânia. Distr. — Brazil.
- NIGRIBIMBO Dodge, 1965c:505. Type-locality: Jamaica, Portland. Distr. — Jamaica.
- PACTA (Townsend), 1934b(June):204 (*Pachygraphiops*). Type-locality: Brazil, Pará, Boa Vista, Rio Tapajós. Distr. — Trinidad, Guyana, Brazil.
- divergens* Curran & Walley, 1934 (30 July):477, fig. 32. Type-locality: Guyana, Kartabo. Ref. — Lopes, 1951a:281, figs. 15-26 (male, female, larva I).
- RUFIPES Dodge, 1965d:485, fig. 15. Type-locality: Bahamas, Grand Bahama I. Distr. — Bahamas.
- VILLIPES Dodge, 1965c:506. Type-locality: Jamaica, St. Andrew. Distr. — Jamaica.
- sp. Dodge, 1965c:507. Locality: Jamaica, Cinchona.
- diversipes* Coquillett of Lopes, 1951a:276, figs. 1-6, misident.

Genus GYMNOCAMPTOPS Townsend

- GYMNOCAMPTOPS Townsend, 1927a:222. Type-species, *griseescens* Townsend (orig. des.).
- Agriochaeta* Enderlein, 1928a:25. Type-species, *aurcostrigata* Enderlein (orig. des.).
- Agriochaetops* Townsend, 1934b:204. Type-species, *Dexia fervens* Wiedemann (orig. des.).
- AUREOSTRIGATA (Enderlein), 1928a:26 (*Agriochaeta*). Type-locality: Brazil, Espírito Santo. Distr. — Brazil.
- FERVENS (Wiedemann), 1830:382 (*Dexia*). Type-locality: "Brazil". Distr. — Brazil. Refs. — Aldrich, 1927c:4 (type examined); Townsend, 1931a:81 (type examined).
- GRISESCENS Townsend, 1927a:222, 311. Type-locality: Brazil, São Paulo, São Paulo, Cantareira. Distr. — Brazil (Rio de Janeiro, Guanabara, São Paulo).



- OCHRISTRIGA (Enderlein), 1928c:149 (*Agriochacta*). Type-locality: "Brazil".
Distr. — Brazil.
- RUBRIVENTRIS (Macquart), 1851:262 (sep., p. 235) (*Sarcophaga*). Type-locality: Argentina, Corrientes. Distr. — Argentina. Ref. — Aldrich, 1930:23 (type examined).

Genus LEPIDODEXIA Brauer & Bergenstamm

- LEPIDODEXIA Brauer & Bergenstamm, 1891:69 (373), 75 (379). Type-species, *tetraptera* Brauer & Bergenstamm (mon.).
- Raimondia* Townsend, 1917a:47. Type-species, *uruhuasi* Townsend (orig. des.).
- Xylocampta* Townsend, 1927a:233. Type-species, *sarcophagina* Townsend (orig. des.).
Ref. — Lopes, 1951b:641 (revision).
- ANTENNATA (Dodge), 1966b:697, figs. 6a-c (*Petriana*). Type-locality: Peru, 40 miles w. of Cusco. Distr. — Peru.
- APOLINARI Lopes, 1951b:644, figs. 6-9. Type-locality: Colombia, Choachi. Distr. — Colombia, Venezuela.
- nigricornis* Dodge, 1966b:696, fig. 6, L (*Petriana*; apud Dodge, in litt.). Type-locality: "Venezuela". N. SYN.
- CINGULATA Lopes, 1962a:254, figs. 1-4. Type-locality: Colombia, Cundinamarca. Distr. — Colombia.
- CURRANI Lopes, 1951b:646, figs. 10-16. Type-locality: Ecuador, Bolívar, Hda. Talahua. Distr. — Ecuador.
- DISTINCTA Lopes, 1951b:655, figs. 39-43. Type-locality: Ecuador, Azuay, Cuenca. Distr. — Ecuador.
- CRISEA Lopes, 1951b:651, figs. 23-29. Type-locality: Venezuela, Caracas. Distr. — Venezuela.
- NIGROPILOSA Lopes, 1951b:649, figs. 17-22. Type-locality: Ecuador, Imbabura, Chichocha. Distr. — Ecuador.
- SARCOPHAGINA (Townsend), 1927a:233, 266 (*Xylocampta*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil (Minas Gerais, Rio de Janeiro, São Paulo, Paraná, Santa Catarina). Refs. — Lopes, 1936a:848, figs. 3-7 (paratype examined); Lopes, 1943:157, figs. 70-71 (larva I).
- SQUAMATA (Walker), 1852:279 (*Tachina*). Type-locality: "Colombia". Distr. — Colombia. Refs. — Austen, 1907:334 (type examined); Townsend, 1931a:81 (type examined).
- TETRAPTERA Brauer & Bergenstamm, 1891:75 (379). Type-locality: "Venezuela". Distr. — Venezuela. Ref. — Aldrich, 1929a:31, fig. 2 (type examined).
- URUHUASI (Townsend), 1917a:47 (*Raimondia*). Type-locality: Peru, San Gabón R. Distr. — Peru. Ref. — Aldrich, 1929a:33 (male and female).

Genus XYLOCAMPTOPSIS Townsend

- XYLOCAMPTOPSIS Townsend, 1927a:221. Type-species, *teffeensis* Townsend (orig. des.).
- TEFFEENSIS Townsend, 1927a:221, 367. Type-locality: Brazil, Amazonas, Tefé. Distr. — Brazil (Amazonas).

Genus ASILODEXIA Townsend

- ASILODEXIA Townsend, 1927b:279. Type-species, *asiliformis* Townsend (orig. des.).
Shannomyiella Lopes, 1938d:433. Type-species, *elegans* Lopes (orig. des.)
= *asiliformis* Townsend.
ASILIFORMIS Townsend, 1927b:279. Type-locality: Paraguay, Santa Trinidad.
Distr. — Brazil, Paraguay.
elegans Lopes, 1938d:433, pl. 1, figs. 1-8 (*Shannomyiella*). Type-locality: Brazil, Mato Grosso, Maracaju.

Genus PETRIANA Lopes

- PETRIANA Lopes, 1945c:453. Type-species, *brevirostris* Lopes (orig. des.).
BREVIROSTRIS Lopes, 1945c:453, figs. 1-21. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Santa Catarina, São Paulo).

Genus JOHNSONIA Coquillett

- JOHNSONIA Coquillett, 1895:316. Type-species, *elegans* Coquillett (orig. des.).
BIVITTATA Curran, 1928:95. Type-locality: Puerto Rico, Aibonito. Distr. — U.S.A., Puerto Rico.
ELEGANS Coquillett, 1895:316. Type-locality: U.S.A., Florida, St. Augustine. Distr. — U.S.A., Mexico.
SETOSA Aldrich, 1916:33, fig. 5. Type-locality: "Peru". Distr. — Peru.

Genus STHENOPYGA Aldrich

- STHENOPYGA Aldrich, 1916:59. Type-species, *globosa* Aldrich (orig. des.)
= *rufitibia* (Wulp).
Camptopsis Townsend, 1918a:162. Type-species *miamensis* Townsend (orig. des.).
RUFITIBIA (Wulp), 1895:270, pl. 7, fig. 3 (*Sarcophaga*). Type-locality: Mexico, Guerrero. Distr. — U.S.A., Mexico.
globosa Aldrich, 1916:59, fig. 18. Type-locality: U.S.A., Louisiana, Opelousas.

Genus PACHYGRAPHIA Brauer & Bergenstamm

- PACHYGRAPHIA Brauer & Bergenstamm, 1891:68 (372), 75 (379). Type-species, *Dexia virgata* Wiedemann (Townsend, 1916d:8).
FRONTALIS Townsend, 1928:366. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil.
VIRGATA (Wiedemann), 1830:382 (*Dexia*). Type-locality: "Brazil". Distr. — Brazil. Refs. — Aldrich, 1927c:4 (type examined); Townsend, 1931a:80 (type examined).

Genus ABACANTHA Hall

- ABACANTHA Hall, 1938b:173. Type-species, *zygax* Hall (orig. des.).
ZYGAX Hall, 1938b:173, fig. 3. Type-locality: Venezuela, Maracay. Distr. — Venezuela.

Genus BOETTCHERIMIMA Lopes

BOETTCHERIMIMA Lopes, 1950b:706. Type-species, *hypopygialis* Lopes (orig. des.).

HYPOPYGIALIS Lopes, 1950b:706, figs. 85-93. Type-locality: Ecuador, Azuay, Cuenca. Distr. — Ecuador.

Genus EMDENIMYIA Lopes

EMDENIMYIA Lopes, 1946c:117. Type-species, *Dexomyiophora myersi* Curran (orig. des.).

banksi Dodge, 1966b:683 (only in key).

BISERIATA Dodge, 1966b:681, figs. 3a-c. Type-locality: "Venezuela". Distr. — Venezuela.

MYERSI (Curran), 1938:7 (*Dexomyiophora*). Type-locality: Guyana, Barabuirna (ex *Castnia licoides*). Distr. — Guyana. Ref. — Lopes, 1946c:118, figs. 1-8.

SPINOSA Lopes, 1948:556, figs. 1-5. Type-locality: Mexico, Guerrero, Ayotzinapa. Distr. — Mexico, Brazil (São Paulo).

VAURIAE Dodge, 1963:240, figs. 5-7. Type-locality: Mexico, Tamazunchale. Distr. — Mexico.

XANTHOPIORINA Dodge, 1967:125, fig. C. Type-locality: "Venezuela". Distr. — Venezuela.

Tribe TEPHROMYIINI

TEPHROMYIINI Townsend, 1919:544; Townsend, 1935:109 (as Myiorhinini); Roback, 1954:83 (as Servasiina).

The species included in this tribe are parasitic on Acrididae, Mantodea or Colcoptera, or sometimes live on dead insects. The glans does not show styli, but a large terminal opening.

Genus OPSOPHYTO Townsend

OPSOPHYTO Townsend, 1915b:23. Type-species, *Sarcophaga opifera* Coquillett (orig. des.).

OPIFERA (Coquillett), 1892:22 (*Sarcophaga*) (ex Acrididae). Type-locality: U.S.A., California. Distr. — U.S.A., Mexico. Ref. — Aldrich, 1916:119, fig. 48.

Genus PROTODEXIA Townsend

PROTODEXIA Townsend, 1912b:117. Type-species, *synthetica* Townsend (orig. des.) = *hunteri* (Hough).

ALEUAPHAGA Blanchard, 1942b:343, fig. 2 (ex *Aleuas lineatus*). Type-locality: Argentina, Buenos Aires. Distr. — Argentina.

AUSTRALIS Blanchard, 1942b:344, fig. 3 (ex *Dichroplus elongatus*). Type-locality: Argentina, Rio Negro. Distr. — Argentina.

CROUZELLI Blanchard, 1942b:347, fig. 4 (ex *Eleochoira veridicata*). Type-locality: Argentina, José C. Paz. Distr. — Argentina.

DEAURATA Blanchard, 1942b:349, fig. 5 (ex *Zoniopoda* sp.). Type-locality: Argentina, Córdoba. Distr. — Argentina.

- DIPONTHI Blanchard, 1942b:351, fig. 6 (ex *Diponthus* sp.). Type-locality: Argentina, Córdoba. Distr. — Argentina.
- HUNTER (Hough), in Hunter, 1898:207, fig. (*Sarcophaga*) (ex *Melanophus differentialis*). Type-locality: U.S.A., Kansas. Distr. — Canada, U.S.A., Mexico, Brazil.
- synthetica* Townsend, 1912b:117. Type-locality: U.S.A., Massachusetts.
- acnigma* Reinhard, 1947:103 (*Sarcophaga*). Type-locality: U.S.A. New York.
- LIEBERMANNI Blanchard, 1955:33, fig. 3 (ex *Dichroplus maculipennis*). Type-locality: Argentina, Patagonia. Distr. — Argentina, Chile.
- OGLOBLINI Blanchard, 1942b:353, fig. 7 (ex *Anchochroma ogloblini*). Type-locality: Argentina, San Luis. Distr. — Argentina.
- QUAESITA (Hall), 1937b:366 (*Sarcophaga*) (ex grasshoppers). Type-locality: Chile. Angol. Distr. — Chile. Ref. — Dodge, 1966b:677 (type examined).

Genus MANTIDOPHAGA Townsend

- MANTIDOPHAGA Townsend, 1919a:160. Type-species, *stagnomontis* Townsend (orig. des.).
- Helicobiella* Blanchard, 1939:837. Type-species, *sanchezi* Blanchard (orig. des.) = *blandita* (Brèthes).
- BLANDITA (Brèthes), 1920:282 (*Sarcophaga*). Type-locality: "South America". Distr. — Brazil, Argentina.
- sanchezi* Blanchard, 1939:837, fig. 13 (*Helicobiella*) (ex Mantodea). Type-locality: Argentina, Catamarca.
- mantidophaga* Blanchard, 1942b:340, fig. 1 (*Helicobiella*) (ex Mantodea). Type-locality: Argentina, José C. Paz.


Genus ACRIDIOPHAGA Townsend

- ACRIDIOPHAGA Townsend, 1917a:46. Type-species, *Sarcophaga aculcata* Aldrich (orig. des.).
- Amblycoryphencs* Townsend, 1918a:161. Type-species, *Sarcophaga amblycoryphae* Coquillett (orig. des.).
- ANGUSTIFRONS (Aldrich), 1916:142, fig. 63 (*Sarcophaga*). Type-locality: U.S.A., New Mexico, Socorro. Distr. — U.S.A., Mexico.
- ARTEGAI (Blanchard), 1939:840, fig. 14 (*Opsophyto*). Type-locality: Argentina, Santa Fé. Distr. — Argentina (ex *Schistocerca paranensis* and *Trigonophymus arrogans*). Ref. — Salavin, 1958:299, figs. p's. 1-3 (habits, larvae I & II).
- CARIDEI (Brèthes), 1906:299, figs. 1-3 (*Sarcophaga*). Type-locality: "Argentina". Distr. — Brazil (Rio Grande do Sul) (ex *Acrididae*). Argentina. Refs. — Blanchard, 1942c:167, fig. 15; Crouzel, 1944:239, 12 figs. (larva I).
- acridiorum* Weyenbergh of Conil, 1879:426, pls. 5-6 (*Nemorea*), and of Lahille, 1907:38, 50, 67, pls. 1-3 (*Sarcophaga*) (larvae, pupa, habits), misident.
- DOLOSA (Dodge), 1966b:675, fig. 1c (*Blacusoripha*). Type-locality: Chile, Casa Blanca. Distr. — Chile.

- EBI (Hall), 1937b:369 (*Sarcophaga*). Type-locality: Chile, Perales. Distr. — Chilc. Ref. — Dodge, 1966b:675, fig. 1 D-F (cx *Dichroplus* spp.).
- FAZI (Hall), 1937b:368, figs. 69 A-B (*Sarcophaga*). Type-locality: Chilc, Perales. Distr. — Chilc.
- GUADALUPENSIS (Arnaud), 1963:124, figs. 1-7 (*Blaesoxipha*). Type-locality: Mexico, Guadalupe Is. Distr. — Mexico.
- LAMBENS (Wiedemann), 1830:365 (*Sarcophaga*). Type-locality: "Americae Meridionalis Insulae". Distr. — West Indies. Ref. — Townsend. 1931a:74 (type examined).
- LOROSA (Hall), 1937b:367 (*Sarcophaga*). Type-locality: Chile, Santiago. Distr. — Chilc.
- NEUQUENENSIS (Blanchard), 1942b:356, fig. 8 (*Tephromyella*). Type-locality: Argentina, Neuquén. Distr. — Argentina (cx *Dichroplus elongatus* G.T.).
- OBERRANS (Wulp), 1896:283 (*Sarcophaga*). Type-locality: Mexico, Guerrero. Distr. — Mexico. Ref. — Aldrich, 1930:34, fig. 14 (type examined).
- OBSCURIOR (Lahille), 1907:91, fig. 16 (as *Sarcophaga acridiorum* var.). Type-locality: "Argentina". Distr. — Argentina.
- UNCATA (Wulp), 1896:277 (*Sarcophaga*). Type-locality: Mexico, Guerrero. Distr. — Mexico. Ref. — Aldrich, 1930:34, fig. 14 (type examined).
- tridens* Wulp, 1896:281 (*Sarcophaga*). Type-locality: Mexico, Guerrero.
- marginata* Aldrich, 1916:136, fig. 58 (*Sarcophaga*). Type-locality: U.S.A., Indiana, La Fayette.

Genus ACANTHODOTHECA Townsend

- ACANTHODOTHECA Townsend, 1918a:159. Type-species, *Sarcophaga prohibita* Aldrich (orig. des.).
- Eleodimyia* Townsend, 1918a:160. Type-species, *Sarcophaga eleodis* Aldrich (orig. des.).
- Sarcophodexia* Townsend, 1918a:161. Type-species, *Sarcophaga hamata* Aldrich (orig. des.).
- Notochaetopsis* Townsend, 1918a:162. Type-species, *Sarcophaga masculina* Aldrich (orig. des.).
- Hamatomyia* Blanchard, 1939:816. Type-species, *denicri* Blanchard (mon.).
- ACRIDIOPIAGOIDES Lopes & Downs, 1951:588, figs. 53-59. Type-locality: Brazil, São Paulo, São Paulo, Ipiranga. Distr. — U.S.A., Mexico, Brazil.
- ALCEDO Aldrich, 1916:132, fig. 55. Type-locality: U.S.A. Indiana, La Fayette. Distr. — U.S.A., Mexico.
- ssp. DENIERI Blanchard, 1939:816, fig. 6 (as sp.). Type-locality: Argentina, Santa Fé. Distr. — Argentina, Brazil.
- ssp. MEXICANA Lopes & Downs, 1951:586, figs. 46-49. Type-locality: Mexico, Morelos, Cuernavaca. Distr. — Mexico.
- ssp. MINENSE Lopes & Downs, 1951:587, figs. 54-57. Type-locality: Brazil, Minas Gerais, Serra do Cipó. Distr. — Brazil (Minas Gerais).
- AZTECA Lopes & Downs, 1951:578, figs. 20-23. Type-locality: Mexico, Bosenchevc. Distr. — Mexico.



- BOXI Hall, 1938a:258, fig. 5. Type-locality: Argentina, Tucumán. Distr. — Argentina.
- DAMPFI (Hall), 1937a:213, fig. 26 (*Sarcophaga*). Type-locality: Mexico, Jalisco, La Barea. Distr. — Mexico.
- EXUBERANS Lopes & Downs, 1951:578, figs. 24-28. Type-locality: Brazil, São Paulo, São José dos Campos. Distr. — Brazil (São Paulo).
- INORNATA Lopes & Downs, 1951:573, figs. 5-7. Type-locality: Brazil, Goiás, Camplinas. Distr. — Brazil (Goiás).
- ssp. PAUCISETA Lopes & Downs, 1951:575, figs. 9-13. Type-locality: Brazil, São Paulo, São José dos Campos. Distr. — Brazil (São Paulo).
- LANEI (Lopes), 1938c:336, figs. 1-3 (*Sarcophaga*). Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil (Minas Gerais, São Paulo, Santa Catarina), Argentina.
- wagneri Blanchard, 1939:812, fig. 5 (*Sarcophodexia*). Type-locality: "Argentina".
- PROHIBITA (Aldrich), 1916:133, fig. 56 (*Sarcophaga*). Type-locality: U.S.A., Western Kansas. Distr. — U.S.A., Mexico.
- RIMOSA (Hall), 1938b:172, fig. 2 (*Sarcophaga*). Type-locality: Venezuela, Maracay. Distr. — Venezuela.
- RUDIS (Aldrich), 1916:125, fig. 51 (*Sarcophaga*) (ex *Ligyrrus gibbosus* DeGeer, Coleoptera, Dynastidae). Type-locality: U.S.A., Indiana, La Fayette. Distr. — U.S.A., Brazil (Rio Grande do Norte, Mato Grosso, São Paulo).

Genus SPECIOSIA Roback

- SPECIOSIA Roback, 1954:87. Type-species, *Fletcherimyia speciosa* Lopes (orig. des.).
- SPECIOSA (Lopes), 1946b:121, figs. 1-6 (*Fletcherimyia*). Type-locality: Mexico, Chapultepec. Distr. — Mexico. Ref. — Roback, 1954:87, figs. 411-412.

Genus CICATRICIA Dodge

- CICATRICIA Dodge, 1965e:503. Type-species, *jamaicorum* Dodge (orig. des.).
- JAMAICORUM Dodge, 1965e:503, figs. 4-8. 8a. Type-locality: "Jamaica". Distr. — Jamaica.

Genus KURTOMYIA Roback

- KURTOMYIA Roback, 1954:84. Type-species, *Sarcophaga postilla* Reinhard (orig. des.) = *californica* (Parker).
- CALIFORNICA (Parker), 1918:32, fig. (*Sarcophaga*). Type-locality: U.S.A., California. Distr. — U.S.A., Mexico.
- postilla Reinhard, 1947:111 (*Sarcophaga*). Type-locality: U.S.A., California. Ref. — Roback, 1954:84, figs. 408-410.
- CESSATOR (Aldrich), 1916:84, fig. 31 (*Sarcophaga*). Type-locality: U.S.A., California, Pine Lake. Distr. — U.S.A., Mexico.

Genus NEOTEPHROMYIELLA Blanchard

- NEOTEPHROMYIELLA Blanchard, 1955:28. Type-species, *chubutina* Blanchard (orig. des.).

CHUBUTINA Blanchard, 1955:28, fig. 1 (ex Acrididae). Type-locality: Argentina, Chubut, Cushamen. Distr. — Argentina.

Tribe RAVINIINI

RAVINIINI Rohdendorf, 1937:49; Roback, 1954:72 (as Raviniina).

Most of the genera here considered are characterized by the fusion of theca and paraphallus (without articulation) and by the presence of a pair of plates representing female's tergite VIII. The larvae of the reared species feed commonly on human faeces. *Oxysarcoderia* has the largest number of Neotropical species.

Genus RAVINIA Robineau-Desvoidy

RAVINIA Robineau-Desvoidy, 1863:434. Type-species, *Sarcophaga haematodes* Meigen (orig. des.) = *pernix* (Harris).

Punasarcophaga Townsend, 1915c:408. Type-species, *auromaculata* Townsend (orig. des.).

Euravinia Townsend, 1917b:191. Type-species, *Ravinia communis* Parker (orig. des.) = *querula* (Walker).

Miltoravinia Townsend, 1917b:191. Type-species, *Sarcophaga planifrons* Aldrich (orig. des.).

Triosarcophaga Townsend, 1917b:191. Type-species, *Sarcophaga aurigena* Townsend (orig. des.).

ADDENTATA (Hall), 1929:71 (*Sarcophaga*; nom. nov. for *minuta* Hall). Type-locality: Mexico, Puebla. Distr. — Mexico.

minuta Hall, 1928:335, fig. 1 (*Sarcophaga*; preocc. Robineau-Desvoidy, 1830). Type-locality: Mexico, Puebla.

AUREOPYGA (Hall), 1928:339, fig. 10 (*Sarcophaga*). Type-locality: Chile, Angol. Distr. — Brazil (Rio Grande do Sul), Chile, Argentina.

AURIGENA (Townsend), 1912a:357 (*Sarcophaga*). Type-locality: Peru, Piura. Distr. — Peru.

AUROMACULATA (Townsend), 1915e:408 (*Punasarcophaga*). Type-locality: Peru, Oroya. Distr. — Peru. Ref. — Hall, 1928:342, fig. 14.

BELFORTI (Prado & Fonseca), 1932a:39, fig. 7 (*Euravinia*). Type-locality: Brazil, São Paulo. Distr. — Colombia, Trinidad, Brazil (Mato Grosso, Goiás, Minas Gerais, Rio de Janeiro, Guanabara, São Paulo, Paraná), Paraguay, Argentina.

tessellata Fabricius of Mattos, 1919:67 (*Sarcophaga*; as Wiedemann, error), misident.

COVAI (Dodge), 1965c:255, figs. 11-12 (*Boetia*). Type-locality: "Venezuela". Distr. — Venezuela.

FLORIDENSIS (Aldrich), 1916:249, fig. 117 (*Sarcophaga*). Type-locality: U.S.A., Florida. Distr. — U.S.A., Bahamas.

LHERMINIERI (Robineau-Desvoidy), 1830:339 (*Myophora*). Type-locality: U.S.A., Carolina. Distr. — U.S.A., Mexico, Bermudas; Hawaii.

anxia Walker, 1849:818 (*Sarcophaga*). Type-locality: "North America".

rediviva Walker, 1849:823 (*Sarcophaga*). Type-locality: Canada, Ontario.

pallinervis Thomson, 1869:535 (*Sarcophaga*). Type-locality: Hawaii, Honolulu.

- OBSCURIPES (Hall), 1928:335, fig. 2 (*Sarcophaga*). Type-locality: Mexico. Sierra Madre. Distr. — Mexico.
- OCHREACEA (Aldrich), 1916:255, fig. 121 (as *Sarcophaga communis* ssp.). Type-locality: U.S.A., Mississippi, Hamburg. Distr. — U.S.A. Guatemala.
- OLLANTAYTAMBENSIS (Hall), 1928:341, fig. 13 (*Sarcophaga*). Type-locality: Peru, Ollantaytambo. Distr. — Peru.
- PUSIOLA (Wulp), 1895:268; 1896:278 (*Sarcophaga*). Type-locality: Mexico, Mexico City. Distr. — Mexico.
- QUADRIVITTATA (Maequart), 1843:259 (sep., p. 102), pl. 12, fig. (*Sarcophaga*). Type-locality: Brazil, Pará. Distr. — Brazil. Ref. — Aldrich, 1930:22 (type examined).
- SUETA (Wulp), 1895:268; 1896:281 (*Sarcophaga*). Type-locality: Mexico, Guerrero. Distr. — Mexico.
- TACITURO Roback, 1952:48, figs. 6 A-C. Type-locality: Mexico, Michoacán. Distr. — Mexico.

Genus ANDINORAVINIA Townsend

- ANDINORAVINIA Townsend, 1917b:190, 193, 195. Type-species, *rufipes* Townsend (orig. des.). Ref. — Lopes, 1962:165 (revision).
- Engelina* Enderlein, 1928a:16. Type-species, *Sarcophaga xanthopyga* Wulp (orig. des; misident.) = *rufipes* Townsend.
- MEINCKE Blanchard, 1939:835, fig. 12. Type-locality: Argentina, Córdoba. Distr. — Argentina.
- RUFIPES Townsend, 1917b:195. Type-locality: Peru, Huariaca. Distr. — Peru, Colombia, Ecuador, Chile. Ref. — Lopes, 1962b:167, figs 1-11.
- xanthopyga* Wulp of Enderlein, 1928a:16 (*Engelina*), misident.
- ssp. INTERMEDIA Lopes, 1962b:169, figs. 12-18. Type-locality: Colombia, Nariño.
- ssp. COLUMBIANA Lopes, 1962b:170, figs. 19-22. Type-locality: Colombia, Nariño.

Genus CHAETORAVINIA Townsend

- CHAETORAVINIA Townsend, 1917b:190, 193, 195. Type-species, *Helicobia quadrisetosa* Coquillett (orig. des.) = *derelicta* (Walker).
- ALMEIDA Lopes, 1946d:227, figs. 1-5. Type-locality: Brazil, Mato Grosso, Salobra. Distr. — Brazil (Mato Grosso, Minas Gerais, São Paulo, Paraná).
- BARROI Dodge, 1956:97. Type-locality: Cuba, Sierra Maestra, Loma del Gato. Distr. — Cuba.
- CAMPFI Lopes, 1946b:135, figs. 36-40. Type-locality: Mexico, Chapultepec. Distr. — Mexico.
- DERELICTA (Walker), 1852:322 (*Sarcophaga*). Type-locality: "U.S.A." Distr. — U.S.A., Mexico. Ref. — Aldrich, 1930:17 (type examined).
- quadrisetosa* Coquillett, 1901:17 (*Helicobia*). Type-locality: U.S.A., Washington, D. C. Ref. — Hall, 1928:246, fig. 18.
- EFFRENATA (Walker), 1861:309 (*Sarcophaga*). Type-locality: "Mexico". Distr. — U.S.A., Mexico, Colombia, Jamaica, Bahamas. Ref. — Aldrich, 1930:20, 30.

- xanthopyga* Wulp, 1895:272 (*Sarcophaga*). Type-locality: "Mexico".
conjugens Wulp, 1895:272 (*Sarcophaga*). Type-locality: "Mexico".
adamsi Hall, 1928:345, fig. 17 (*Sarcophaga*). Type-locality: "Jamaica".
 ERRABUNDA (Wulp), 1895:268, 1896:278 (*Sarcophaga*). Type-locality: Mexico.
 Tabasco. Distr. — Mexico, U.S.A. Ref. — Aldrich, 1930:33 (type
 examined).
reinhardii Hall, 1928:246, fig. 20 (*Sarcophaga*). Type-locality: U.S.A.,
 Texas.
 GLOBULUS (Aldrich), 1916:293, fig. 145 (*Sarcophaga*). Type-locality: Cuba.
 Havana. Distr. — Cuba, Puerto Rico.
 TRIVITTATA (Townsend), 1927:220, 295 (*Sarcophaga*). Type-locality: Brazil,
 São Paulo, Itaquaquecetuba. Distr. — Bolivia, Brazil (Mato
 Grosso, Minas Gerais, São Paulo, Santa Catarina), Argentina.
 Ref. — Lopes, 1932:46, pl. 3 (paratype examined).
townsendi Engel, 1931:149, fig. 18 [*Sarcophaga* (*Chactoravina*)].
 Type-locality: Bolivia, northern Chiquitos.
 VAGABUNDA (Wulp), 1895:270, pl. 7, fig. 4 (*Sarcophaga*). Type-locality:
 "Mexico". Distr. — Mexico. Ref. — Lopes, 1946b:132, figs. 23-26.

Genus OXYVINIA Dodge

- OXYVINIA Dodge, 1966b:692. Type-species, *piliventris* Dodge (orig. des.).
 EXCISA (Lopes), 1950a:361, figs. 19-26 (*Dexosarcophaga*). Type-locality:
 Brazil, Guanabara, Rio de Janeiro, Grajaú. Distr. — Peru, Brazil
 (Mato Grosso, Rio de Janeiro, Guanabara).
 GRATA (Lopes), 1953a:44, figs. 9-12 (*Dexosarcophaga*). Type-locality:
 Ecuador, Azuay, Cuenca. Distr. — Ecuador.
 PILIVENTRIS Dodge, 1966b:692, figs. 5C-E. Type-locality: Venezuela, near
 Caracas, Sierro Ávila. Distr. — Venezuela.
 XANTHOPHORA (Schiner), 1868:313 (*Sarcophaga*). Type-locality: "Vene-
 zuela". Distr. — Venezuela. Ref. — Aldrich, 1930:25, fig. 23 (type
 examined).
 WICHARTI (Lopes), 1953a:46, figs. 13-21 (*Dexosarcophaga*). Type-locality:
 Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil.

Genus HYBOPYGIA Enderlein

- HYBOPYGIA Enderlein, 1928a:40. Type-species, *Sarcophaga chilensis* Mac-
 quart (orig. des.) = *varia* (Walker).
Amesothysus Enderlein, 1928a:41. Type-species, *chilensis* Enderlein (orig.
 des.) = *minalis* (Wiedemann).
 AURA (Hall), 1937b:349, 372 (*Sarcophaga*, nom. nov. for *chilensis* Engel).
 Type-locality: Bolivia, northern Chiquitos. Distr. — Bolivia,
 Brazil.
chilensis Enderlein of Engel, 1931:148, fig. 17 (*Amesothysus*), mis-
 ident.
 TERMINALIS (Wiedemann), 1830:366 (*Sarcophaga*). Type-locality: "Brazil".
 Distr. — Brazil, Argentina, Chile. Ref. — Lopes, 1946c:62.
chilensis Enderlein, 1928a:41 (*Amesothysus*). Type-locality: "Chile".
varia Walker of Engel, 1931:147, fig. 15a [*Sarcophaga* (*Hybopygia*)].
 misident. (In part; Argentinian specimen).
complicata Hall, 1937b:371, fig. 70 (*Sarcophaga*). Type-locality:
 "Chile".
pseudovarioria Blanchard, 1939:806, fig. 3. Type-locality: Argentina,
 Córdoba.

- VARIA (Walker), 1837:353 (*Sarcophaga*). Type-locality: Uruguay, Gorrito Is., off Maldonado. Distr. — Brazil, Uruguay, Argentina, Chile, Juan Fernandez Is.; New Zealand.
- flavicostata* Macquart, 1843:261 (1843:104), pl. 13, fig. 4 (*Sarcophaga*). Type-locality: Chile, Concepción. Ref. — Aldrich, 1930:22 (type examined).
- chilensis* Macquart, 1843:261 (sep., p. 104), pl. 11, fig. 6 (*Sarcophaga*). Type-locality: "Chile". Ref. — Aldrich, 1930:22 (type examined).
- truncata* Schiner, 1868:314 (*Sarcophaga*). Type-locality: "Chile". Ref. — Aldrich, 1930:26 (type examined).
- nobilis* Thomson, 1869:536 (*Sarcophaga*). Type-locality: "South America". Ref. — Aldrich, 1930:28 (type examined).
- milleri* Johnston & Tiegs, 1922:185, fig. 1 (*Sarcophaga*). Type-locality. "New Zealand".

Genus APELOPHYL A Hall

- APELOPHYL A Hall, 1938a:356. Type-species, *marina* Hall (orig. des.).
- MARINA Hall, 1938a:257, fig. 4. Type-locality: Argentina, Mendoza. Distr. — Argentina.

Genus OXYSARCODEXIA Townsend

- OXYSARCODEXIA Townsend, 1917b:191. Type-species, *Sarcophaga peltata* Aldrich (orig. des.).
- Dasyproctia* Enderlein, 1928a:23. Type-species, *Sarcophaga praevolans* Wulp (orig. des.; misident.) = *ochripyga* (Wulp).
- ADMIXTA (Lopes), 1933b:156, fig. 3 (*Sarcophaga*). Type-locality: Brazil, Rio de Janeiro, Angra dos Reis. Distr. — Brazil (Golás, Rio de Janeiro, São Paulo), Argentina.
- AFFICTA (Wulp), 1895:269, 1896:286 (*Sarcophaga*). Type-locality: "Mexico". Distr. — Mexico, Panama, Colombia. Ref. — Aldrich, 1930:36, fig. 18 (type examined).
- AMOROSA (Schiner), 1868:314 (*Sarcophaga*). Type-locality: "Brazil". Distr. — Mexico, Colombia, Guyana, Brazil (Mato Grosso, Minas Gerais, Rio de Janeiro, Guanabara). Ref. — Aldrich, 1930:26, fig. 26 (type examined).
- xanthosoma* Aldrich of Curran & Walley, 1934:489, fig. 29 (*Sarcophaga*), misident.
- ANGOLENSIS (Hall), 1937b:374, fig. 71 (*Sarcophaga*). Type-locality: Chile, Angol. Distr. — Chile.
- ANGRENSIS (Lopes), 1933b:153, fig. 1 (*Sarcophaga*). Type-locality: Brazil, Rio de Janeiro, Angra dos Reis. Distr. — Guyana, Brazil (Pará, Mato Grosso, Golás, Rio de Janeiro, Guanabara).
- kartabo* Curran & Walley, 1934:487, fig. 43 (*Sarcophaga*). Type-locality: Guyana, Kartabo.
- ARTIGASI Dodge, 1966b:684, fig. 3G. Type-locality: Chile, Santiago. Distr. — Chile.
- AUGUSTA Lopes, 1946e:84, figs. 42-44. Type-locality: Brazil, Guanabara, Rio de Janeiro, Méier. Distr. — Brazil (Rio de Janeiro, Guanabara), Argentina.

- AUREICEPS (Macquart), 1855:128 (sep., p. 108) (*Sarcophaga*). Type-locality: "South America". Distr. — South America. Ref. — Aldrich, 1930:24 (type examined, female).
- AURIFINIS (Walker), 1852:325 (*Sarcophaga*). Type-locality: "Brazil". Distr. — Guyana, Brazil (Pará, Ceará, Pernambuco, Paraíba, Mato Grosso, Goiás, Minas Gerais, Rio de Janeiro, Guanabara, São Paulo, Paraná, Santa Catarina, Rio Grande do Sul), Paraguay, Argentina. Ref. — Aldrich, 1930:18, fig. 3 (type examined).
- tessellata* Fabricius of Wiedemann, 1830:363 (*Sarcophaga*), misident. Ref. — Townsend, 1931a:74 (type examined).
- compta* Wiedemann of Mattos, 1919:63, pl. 3 (*Sarcophaga*), misident.
- aurigena* Townsend of Engel, 1931:146, fig. 12 (*Trirosarcophaga*), misident.
- neotropica* Prado & Fonseca, 1932a:36, fig. 4. Type-locality: Brazil, São Paulo.
- AVUNCULA (Lopes), 1933b:156, fig. 4 (*Sarcophaga*). Type-locality: Brazil, Guanabara, Rio de Janeiro, Manguinhos. Distr. — Mexico, Brazil (Goiás, Minas Gerais, Rio de Janeiro, Guanabara, São Paulo, Santa Catarina), Paraguay, Argentina.
- BAKERI (Aldrich), 1916:270, fig. 129 (*Sarcophaga*). Type-locality: Cuba, Havana. Distr. — U.S.A., Mexico, Cuba, Bahama Is. (Turk & Caicos), Jamaica, Puerto Rico, Brazil (Bahia, Goiás).
- BICOLOR Lopes, 1946c:127, figs. 155-158. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil (Rio de Janeiro, São Paulo).
- BIKINI Dodge, 1966b:685, figs. 3H-J. Type-locality: "Chile". Distr. — Chile.
- CARVALHOI Lopes, 1946c: 92, figs. 59-62. Type-locality: Brazil, Minas Gerais, Cordisburgo. Distr. — Guyana, Brazil (Mato Grosso, Minas Gerais).
- CHAETOPYGIALIS (Williston), 1896:366 (*Sarcophaga*). Type-locality: West Indies, St. Vincent. Distr. — St. Vincent, Jamaica. Ref. — Aldrich, 1916:110, fig. 45 (type examined).
- CONFUSA Lopes, 1946c:96, figs. 83-88. Type-locality: Brazil, Rio de Janeiro, Miguel Pereira. Distr. — Brazil (Minas Gerais, Rio de Janeiro, Guanabara, São Paulo, Paraná), Argentina.
- COROLLA Dodge, 1965e:508, figs. 12-14. Type-locality: Jamaica, St. Andrew. Distr. — Jamaica.
- CUERNAVACA Dodge, 1966b:685, figs. 4 L-M. Type-locality: Mexico, Morelos, Cuernavaca. Distr. — Mexico.
- CULMINATA (Aldrich), 1916:289, fig. 139 (*Sarcophaga*). Type-locality: Puerto Rico, Mayaguez. Distr. — Puerto Rico, Brazil (Minas Gerais, Guanabara, Rio de Janeiro, São Paulo, Paraná, Santa Catarina, Rio Grande do Sul), Paraguay, Argentina.
- meridionalis* Blanchard, 1942c:146, fig. 6 (preocc. Engel, 1931). Type-locality: Argentina, Delta del Paraná.
- culminiforceps* Dodge, 1966b:687. Type-locality: Brazil, Rio de Janeiro.
- CYANIFORCEPS (Hall), 1933:282, fig. 23 (*Sarcophaga*). Type-locality: Panama, Barro Colorado Is. Distr. — Panama.
- DIANA (Lopes), 1933b:154, fig. 2 (*Sarcophaga*). Type-locality: Brazil, Rio de Janeiro. Distr. — Mexico, Brazil (Rio de Janeiro, Guanabara, Paraná), Paraguay, Argentina.
- DORISAE Dodge, 1965e:508, figs. 15-16. Type-locality: Jamaica, Manchester. Distr. — Jamaica.

- EDWARDSI Lopes, 1946e:123, figs. 146-149. Type-locality: Colombia, Gorgona Is. Distr. — Colombia.
- FAVORABILIS (Lopes), 1935b:318, figs. 5-6 (*Sarcophaga*). Type-locality: Brazil, Rio de Janeiro, Angra dos Reis. Distr. — Brazil.
- FLAVIFRONS (Maequart), 1846:319 (sep., p. 191) (*Sarcophaga*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Aldrich, 1930:23 (type examined).
- FLUMINENSIS Lopes, 1946e:104, figs. 103-105. Type-locality: Brazil, Guanabara. Rio de Janeiro, Grajaú. Distr. — Brazil.
- FRINGIDEA (Curran & Walley), 1934:488, fig. 40 (*Sarcophaga*). Type-locality: Guyana, Georgetown. Distr. — Guyana, Brazil (Amazonas, Pará, Minas Gerais). Ref. — Lopes, 1946e:77, figs. 2-6, 23-26.
- lopesi Dodge, 1966b:687 (unnecessary nom. nov. for *fringidea* Curran & Walley of Lopes, 1946a:77).
- GRANDIS Lopes, 1946e:82, figs. 39-41. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil (Rio de Janeiro, Guanabara, Paraná).
- ILLATA (Lopes), 1938:340, pl. 4, figs. 1-2 (*Sarcophaga*). Type-locality: Brazil, Goiás, Campinas. Distr. — Brazil.
- INSOLITA Lopes, 1946e:89, figs. 53-54. Type-locality: Guyana, Moraballi Creek. Distr. — Guyana.
- INTONSA (Curran & Walley), 1934:489, fig. 42 (*Sarcophaga*). Type-locality: Guyana, Kartabo. Distr. — Guyana.
- MAJOR Lopes, 1946e:88, figs. 50-52. Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil (Rio de Janeiro, Guanabara).
- MERIDIONALIS (Engel), 1931:146, fig. 14 (*Sarcophaga*, as *ventricosa* var.). Type-locality: Argentina, San José. Distr. — Bolivia, Argentina.
- MITIFICA Lopes, 1953a:50, figs. 26-30. Type-locality: Venezuela, Caracas. Distr. — Venezuela.
- MODESTA Lopes, 1946e:129, fig. 162. Type-locality: Brazil, Guanabara, Rio de Janeiro, Gávea. Distr. — Brazil (Minas Gerais, Guanabara, São Paulo).
- MOLITOR (Curran & Walley), 1934:488, fig. 38 (*Sarcophaga*). Type-locality: Guyana, Kartabo. Distr. — Guyana.
- NEIVAI (Mattos), 1919:78, fig., pl. 3 (*Sarcophaga*). Type-locality: Brazil, São Paulo. Distr. — Brazil.
- OCCULTA Lopes, 1946e:112, figs. 122-124. Type-locality: Brazil, Guanabara, Rio de Janeiro, Coreovado. Distr. — Brazil (Rio de Janeiro, Guanabara).
- OCHRIPTYGA (Wulp), 1895:269; 1896:85 (*Sarcophaga*). Type-locality: Mexico, Guerrero. Distr. — U.S.A., Mexico, Guatemala, Costa Rica, Panama, Colombia, Peru. Ref. — Aldrich, 1930:30 (type examined).
- australis Aldrich, 1916:282, fig. 135 (*Sarcophaga*). Type-locality: "Peru".
- OMISSA Lopes, 1946b:140, figs. 47-50. Type-locality: Mexico, Chapultepec. Distr. — Mexico.
- ORBITALIS Dodge, 1966b:686, figs. 5A-B. Type-locality: Martinique, Mt. Péló. Distr. — Martinique.
- PALLISTERI Dodge, 1966b:686, figs. 3K-N. Type-locality: Peru, Machu-Picchu. Distr. — Peru.

- PARVA Lopes, 1946e:97, figs. 77-82. Type-locality: Brazil, Guanabara, Rio de Janeiro, Grajaú. Distr. — Brazil (Mato Grosso, Rio de Janeiro, Guanabara), Argentina.
- PAULISTANENSIS (Mattos), 1919:72, fig., pl. 3 (*Sarcophaga*). Type-locality: Brazil, São Paulo. Distr. — Brazil (Minas Gerais, Rio de Janeiro, São Paulo, Paraná, Rio Grande do Sul), Argentina.
- amarali* Prado & Fonseca, 1932a:35, fig. 1. Type-locality: Brazil, São Paulo.
- deipontei* Blanchard, 1939:809, fig. 4. Type-locality: "Argentina".
- PELTATA (Aldrich), 1916:316, fig. 103 (*Sarcophaga*). Type-locality: Puerto Rico, Mayaguez. Distr. — U.S.A., Mexico, Cuba, Puerto Rico, Guadalupe, Bahamas, Santa Lucia, Jamaica.
- PERNETA (Walker), 1861:308 (*Sarcophaga*). Type-locality: "Mexico". Distr. — Mexico. Ref. — Aldrich, 1930:20, fig. 6 (type examined).
- PLAUMANNI Lopes, 1946e:102, figs. 93-96. Type-locality: Brazil, São Paulo, Eugênio Lefèvre. Distr. — Brazil (São Paulo, Paraná).
- PLEBEJA Lopes, 1946b:142, figs. 51-52. Type-locality: Mexico. Cuernavaca. Distr. — U.S.A., Mexico.
- RIOGRANDENSIS Lopes, 1946c:103, figs. 100-102. Type-locality: Brazil, Rio Grande do Sul, São João de Montenegro. Distr. — Brazil (Rio de Janeiro, Rio Grande do Sul).
- SARCINATA Lopes, 1953a:48, figs. 22-25. Type-locality: Colombia, Medellin. Distr. — Colombia.
- SIMPLICOIDES (Lopes), 1953b:156, fig. 5 (*Sarcophaga*). Type-locality: Brazil, Guanabara, Rio de Janeiro, Méier. Distr. — Brazil.
- TAITENSIS (Schiner), 1868:314 (*Sarcophaga*). Type-locality: "Tahiti". Distr. — Tahiti, Samoa, Fidji; Galápagos. Refs. — Aldrich, 1929:256, fig. 6.
- obtusifrons* Thomson, 1869:536 (*Sarcophaga*). Type-locality: "Galápagos". Ref. — Townsend, 1931a:77.
- auricauda* Enderlein, 1928a:23 (*Dasyproctia*; *nomen nudum*). Locality: "Samoa".
- TIMIDA (Aldrich), 1916:283, fig. 136 (*Sarcophaga*). Type-locality: Guatemala, Puerto Barrios. Distr. — Guatemala, Honduras, Panama, Brazil (Pará, Maranhão, Rio de Janeiro, Guanabara).
- sanguisuga* Hall, 1933:283, fig. 24 (*Sarcophaga*). Type-locality: Panama, Canal Zone. Ref. — Dodge, 1966b:692 (type examined by Downes).
- TITUBATA Lopes, 1945c:459, figs. 26-31. Type-locality: Brazil, Paraná, Foz do Iguaçu. Distr. — Brazil (Paraná, Santa Catarina).
- ssp. FRATERNA Lopes, 1946b:143, figs. 56-59. Type-locality: Mexico, Cuernavaca. Distr. — Mexico.
- TRIVIALIS (Wulp), 1895:268; 1896:277 (*Sarcophaga*). Type-locality: "Mexico". Distr. — U.S.A., Mexico. Ref. — Aldrich, 1930:32, fig. 11 (type examined).
- VENTRICOSA (Wulp), 1895:268; 1896:274 (*Sarcophaga*). Type-locality: Mexico, Guerrero. Distr. — Canada, U.S.A., Mexico. Ref. — Aldrich, 1930:31 (type examined).
- tenuiventris* Wulp, 1895:268; 1896:282 (*Sarcophaga*). Type-locality: "Mexico". Ref. — Aldrich, 1930:34 (type examined).
- assidua* Walker of Aldrich, 1916:285, fig. 137 (*Sarcophaga*), misident.



- VESICA (Hall), 1933:283, fig. 26 (*Sarcophaga*). Type-locality: Panama, Canal Zone, Aneon. Distr. — Panama, Ecuador.
- VILLOSA Lopes, 1945e:458, figs. 22-25. Type-locality: Brazil, Pará. Distr. — Brazil.
- XANTHIOSOMA (Aldrich), 1916:274, fig. 131 (*Sarcophaga*). Type-locality: Guatemala, Los Amates. Distr. — Guatemala, Panama, Brazil (Minas Gerais, Rio de Janeiro, Guanabara), Argentina.
- ZAYASI Dodge, 1956:100, figs. 6-12. Type-locality: Cuba, Lomas de Camoa. Distr. — Cuba, Bahamas.

Genus OROSARCOPHAGA Townsend

- OROSARCOPIAGA Townsend, 1927a:231. Type-species, *ornata* Townsend (orig. des.).
- ORNATA Townsend, 1927a:231, 342. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil.

Genus NEPHOCHAETOPTERYX Townsend

- NEPHOCHAETOPTERYX Townsend, 1934b:203. Type-species, *pallidiventris* Townsend (orig. des.).
- Sarcohehobia* Blanchard, 1939:795. Type-species, *clegans* Blanchard (orig. des.).
- AFFINIS Lopes, 1936b:88, figs. 24-25. Type-locality: Brazil, São Paulo, Birigui. Distr. — Brazil.
- ANGUSTIFRONS Lopes, 1942a:187, pl. 20, figs. 4-6. Type-locality: Brazil, Guanabara, Rio de Janeiro, Grajaú. Distr. — Brazil.
- CALIDA (Wiedemann), 1830:366 (*Sarcophaga*). Type-locality: "Brazil". Ref. — Lopes, 1936b:81, 83.
- CYANEIVENTRIS Lopes, 1936b:86, figs. 22-23. Type-locality: Brazil, São Paulo, Alto da Serra (= Paranapiacaba). Distr. — Brazil.
- ELEGANS (Blanchard), 1939:795 (*Sarcohehobia*). Type-locality: Argentina, Misiones. Distr. — Argentina.
- FLAVIPALPIS Lopes, 1936b:85, figs. 19-21. Type-locality: Brazil, Guanabara, Rio de Janeiro, Jardim Botânico. Distr. — Brazil (Guanabara, São Paulo).
- FUSCIPENNIS Lopes, 1941b:361, figs. 1-3. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- MOLINAI Lopes, 1942a:186, pl. 20, figs. 1-3. Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil.
- ORBITALIS (Curran & Walley), 1934:478, fig. 35 (*Sarcophaga*). Type-locality: Guyana, Kartabo. Distr. — Guyana, Brazil. Ref. — Lopes, 1936b:85, figs. 15-16.
- PALLIDIVENTRIS Townsend, 1934b:203. Type-locality: Brazil, Pará, Rio Tapajós, Boa Vista. Distr. — Brazil (Pará, Rio de Janeiro). Ref. — Lopes, 1936b:83, figs. 13-14.
- SUBAURATA (Engel), 1931:142, fig. 8 (*Bercaca*). Type-locality: N. Argentina, San José. Distr. — Argentina.
- TRAVASSOSI Lopes, 1938b:279, pl. 1, figs. 1-4. Type-locality: Brazil, Guanabara, Rio de Janeiro, Gávea. Distr. — Brazil.

Tribe SARCOPHAGULINI

SARCOPHAGULINI Roback, 1954:59 (as Sarcophagulina).

The species of Sarcophagulini show proclinate fronto-orbital bristles in the male and the penis is very similar in the species of all genera. *Tricharaca* and *Sarothromyia* are found only on sea or river beaches.

Genus SARCOPHAGULA Wulp

SARCOPHAGULA Wulp, 1887:173. Type-species, *Musca occidua* Fabricius (Coquillett, 1910:602). Ref. — Lopes, 1955b:589 (revision).

Pseudosarcophagula Townsend, 1927a:220. Type-species, *brasiliensis* Townsend (orig. des.).

Desvoidyella Enderlein, 1928a:21. Type-species, *aurata* Enderlein (orig. des.) = *brasiliensis* (Townsend).

BRASILIENSIS (Townsend), 1927a:220, 354 (*Pseudosarcophagula*). Type-locality: Brazil, Amazonas, Tefé. Distr. — Brazil, Paraguay.

aurata Enderlein, 1928a:21 (*Desvoidyella*). Type-locality: "Paraguay".

CANUTA Wulp, 1896:289, pl. 8, fig. 12. Type-locality: "Mexico". Distr. — Mexico, Cuba, Jamaica, Dominica, Honduras, Colombia, Ecuador, Brazil, Paraguay. Ref. — Lopes, 1955b:594, figs. 18-29.

INDONATA Lopes, 1955b:597, figs. 36-38. Type-locality: "Surinam". Distr. — Surinam, Brazil (Mato Grosso), Paraguay.

MACROPHTHALMA Lopes, 1955b:599, figs. 39-45. Type-locality: Brazil, Guanabara, Rio de Janeiro, Gávea. Distr. — Brazil (Guanabara, Rio de Janeiro).

OBSOLETA (Wiedemann), 1830:367 (*Sarcophaga*). Type-locality: "West Indies". Distr. — West Indies.

OCCIDUA (Fabricius), 1794:315 (*Musca*). Type-locality: "America meridionalis Insulis". Distr. — U.S.A., West Indies, Central America, Galápagos, Ecuador, Colombia, Venezuela, Bolivia, Guyana, Brazil (Amazonas, Pará, Maranhão, Ceará, Mato Grosso, Goiás, Rio de Janeiro, Guanabara, São Paulo), Paraguay, Argentina, Chile. Ref. — Aldrich, 1930:2 (type examined).

sugens Wiedemann, 1830:367 (*Sarcophaga*). Type-locality: "Brazil". Ref. — Aldrich, 1930:12 (type examined).

parvula Wiedemann, 1830:368 (*Sarcophaga*). Type-locality: "Brazil". Ref. — Aldrich, 1930:12 (type examined).

parva Walker, 1852:321 (*Sarcophaga*). Type-locality: Brazil, Pará. Ref. — Aldrich, 1930:17 (type examined).

despecta Thomson, 1869:540 (*Sarcophaga*). Type-locality: Ecuador, Isla Puná. Ref. — Aldrich, 1930:29 (type examined).

imbecilla Wulp, 1896:289. Type-locality: "Mexico".

PERUANA Townsend, 1912a:358. Type-locality: Peru, Piura. Distr. — Peru.

PUELLA (Williston), 1896:361, pl. 12, fig. 120bis (*Sarcophiloides*). Type-locality: West Indies, St. Vincent. Distr. — West Indies.

TENUIS Wulp, 1896:289. Type-locality: "Mexico". Distr. — Mexico. Ref. — Lopes, 1955b:596, figs. 30-35.

Genus BAHAMIOLA Dodge

BAHAMIOLA Dodge, 1965d:481. Type-species, *orbitalis* Dodge (orig. des.).
ORBITALIS Dodge, 1965d:481, figs. 8-10. Type-locality: Bahamas, Grand
Bahama Is. Distr. — Bahamas.

Genus SARCOFAHRTIOPSIS Hall

- SARCOFAHRTIOPSIS Hall, 1933:261. Type-species, *Sarcofahrtia capitata* Curran
(mon.).
Cataphyto Townsends, 1935a:72. Type-species, *euncata* Townsends (orig.
des.).
BAUMHOVERI Dodge, 1965d:491, figs. 38, 39. Type-locality: Bahamas, Grand
Bahama Is. Distr. — Bahamas.
CAPITATA (Curran), 1928:96, fig. 33 (*Sarcofahrtia*). Type-locality: Puerto
Rico, Mayaguez. Distr. — Puerto Rico, Panama.
CUNEATA (Townsends), 1935a:72 (*Cataphyto*). Type-locality: Brazil, Per-
nambuco, Tapera. Distr. — Trinidad, Brazil (Pernambuco, Rio
de Janeiro). Ref. — Lopes, 1936:89, figs. 26-28.
DIEMBROMA Dodge, 1965c:511, fig. 27. Type-locality: Jamaica, St. Andrew.
Distr. — Jamaica.
FARRI Dodge, 1965c:511, fig. 25. Type-locality: Jamaica, St. Andrew. Distr.
— Jamaica, Costa Rica.
JAMAICENSIS Dodge, 1965c:513, fig. 22. Type-locality: Jamaica, St. Andrew.
Distr. — Jamaica.
PATERNA Dodge, 1965c:513, figs. 26, 29-35. Type-locality: U.S.A., Florida
Keys. Distr. — U.S.A., Cuba, Puerto Rico.

Genus PHAGITA Lopes

PHAGITA Lopes, 1941b:362. Type-species, *natiusecula* Lopes (orig. des.).
NATIUSECULA Lopes, 1941b:363, figs. 4-7. Type-locality: Colombia, Gorgona
Is. Distr. — Colombia.

Genus CRYPTOSARCOPHILA Townsends

CRYPTOSARCOPHILA Townsends, 1931b:314. Type-species, *chaetosa* Townsends
(orig. des.).
CHAETOSA Townsends, 1931b:315. Type-locality: Argentina, Mendoza. Distr.
— Argentina.

Genus SAROTHIROMYIA Brauer & Bergenstamm

SAROTHIROMYIA Brauer & Bergenstamm, 1891:61 (365). Type-species, *Sar-
cophila femoralis* Schiner (mon.).
ENGELI (Dodge), 1966b:699, fig. 6G (*Tricharea*). Type-locality: Bolivia,
Fortín Esteros. Distr. — Bolivia.
femoralis Schiner of Engel, 1931:141, misident.
FEMORALIS (Schiner), 1868:315 (*Sarcophila*). Type-locality: "Brazil".
Distr. — West Indies, Panama, Brazil (Guanabara, São Paulo).

Refs. — Aldrich, 1916:37, figs. 7, 7a; Dodge, 1965e:516 (type examined).

brevispina Thomson, 1869:539 (*Sarcophaga*). Type-locality: Brazil, Guanabara, Rio de Janeiro. Ref. — Aldrich, 1930:29 (type examined).

SIMPLEX Aldrich, 1916:39 (as *femoralis* ssp.). Type-locality: U.S.A., Florida. Distr. — U.S.A. (Florida), Bahamas.

Genus TRICHARAEA Thomson

TRICHARAEA Thomson, 1869:540. Type-species, *scatophagina* Thomson (mon.) = *brevicornis* (Wiedemann).

Mallonotum Aldrich, 1925b:466. Type-species, *Sarcophaga brevicornis* Wiedemann (orig. des.).

Leucoravinia Townsend, 1928:143. Type-species, *quadraticephala* Townsend (orig. des.) = *brevicornis* (Wiedemann).

Euleucomyia Townsend, 1934b:202. Type-species, *australis* Townsend (orig. des.) = *brevicornis* (Wiedemann).

Xenagria Blanchard, 1939:799. Type-species, *mallogaster* Blanchard (orig. des.) = *brevicornis* (Wiedemann).

BREVICORNIS (Wiedemann), 1830:299 (*Sarcophaga*). Type-locality: Uruguay, Montevideo. Distr. — Peru, Brasil (Guanabara, Amazon Valley), Chile, Argentina, Uruguay, Australia. Ref. — Aldrich, 1925:467 (type examined).

albicans Walker, 1858:204 (?*Dezia*). Type-locality: Brazil, Amazon Valley. Ref. — Aldrich, 1930:28 (type examined).

scatophagina Thomson, 1869:541. Type-locality: Brazil, Guanabara, Rio de Janeiro. Ref. — Townsend, 1931a:73 (type examined).

canescens Thomson, 1869:539 (*Sarcophaga*). Type-locality: Brazil, Guanabara, Rio de Janeiro. Ref. — Aldrich, 1930:28 (type examined).

quadraticephala Townsend, 1928:143 (*Leucoravinia*). Type-locality: Peru, Huallaga R.

pellita Enderlein, 1928:42 (*Liopygia*; *nomen nudum*).

australis Townsend, 1934:302 (*Euleucomyia*). Type-locality: Brazil, Rio de Janeiro.

mallogaster Blanchard, 1939:800, fig. 1 (*Xenagria*). Type-locality: Argentina, Adrogué.

bilineata Dodge, 1966b:700. Type-locality: Peru, Tingo Maria.

penai Dodge, 1965a:90. Type-locality: Chile, Maullin.

Tribe SARCOPHAGINI

SARCOPHAGINI Macquart, 1835:219.

Many groups of genera are here included, some of them poorly known. Groups of genera like *Peckia*, *Paraphrissopoda*, *Euboettcheria* and *Sarcodexia*; *Helicobia* and *Punaphyto*; *Raviniopsis*, *Tripanurga*, *Argoravinia* and *Gigantotheca*; *Airypel*, *Cuculomyia* and *Titanogrypa*; *Boettcheria* and *Boettia*, will be separated in tribes at a near future, but their limits have to be more accurately known. *Malacophagula* was described as parasite of *Bulimulus*; *Helicobia*, *Udamopyga* and

Malacophagomyia were reared from dead molluscs; *Sarcodexia* is commonly found on dead Arthropods; *Arachnidomyia* was reared from spider cocoons; *Dexosarcophaga* breeds in nests of termites and ants, or in galleries of some boring insects; *Neobellieria* lives in nests of Vespidae; *Boettcheria* has been referred from Lepidoptera and Hymenoptera larvae; *Lopostomyia* and *Sarcophagulopsis* were reared from Acrididae; *Sarcophodexiops* was found in Alabama (Lepidoptera); *Ravinlops* was found near *Bembix* nests; *Sarcophahrtimys* is found only on fresh water vegetation. Larvae of *Cuculomyia* and *Sarconetva* are venomous, destroying all larvae of other species in the same environment. Species of *Parasarcophaga* were introduced from Palaearctic or Oriental regions.

Genus GLUTOXYS Aldrich

GLUTOXYS Aldrich, 1929b:2. Type-species, *elegans* Aldrich (orig. des.).
ELEGANS Aldrich, 1929b:2. Type-locality: "Santo Domingo". Distr. — Santo Domingo.

Genus DISCOHELICOBIA Dodge

DISCOHELICOBIA Dodge, 1965e:484. Type-species, *annae* Dodge (orig. des.).
ANNAE Dodge, 1965e:484, fig. 11. Type-locality: Turks and Caicos Is., South Caicos Is. Distr. — Turks and Caicos Is.

Genus DEXOMYIOPHORA Townsend

DEXOMYIOPHORA Townsend, 1927a:233. Type-species, *facialis* Townsend (orig. des.).
FACIALIS Townsend, 1927a:233, 301. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil (São Paulo). (Allotype examined).
HINEI (Aldrich), 1930:31 (*Notochacta*). Type-locality: Guatemala, Puerto Barrios. Distr. — Guatemala.
plumigera Wulp of Aldrich, 1916:55, fig. 15 (*Notochacta*), misident.

Genus OROBRACHYCOMA Townsend

OROBRACHYCOMA Townsend, 1927a:230. Type-species, *ornata* Townsend (orig. des.).
ORNATA Townsend, 1927a:230, 341. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil.

Genus MELANOPHYTO Townsend

MELANOPHYTO Townsend, 1916c:304. Type-species, *macrens* Townsend (orig. des.).
MAERENS Townsend, 1916c:304. Type-locality: Mexico, Mexico City. Distr. — Mexico.

Genus SARCOFAHRTIMYIA Hall

SARCOFAHRTIMYIA Hall, 1937:350, 353. Type-species, *tenta* Hall (orig. des.).

TENTA Hall, 1937b:353, fig. 62. Type-locality: Argentina, Misiones, Bonpland. Distr. — U.S.A., Jamaica, Guatemala, Brazil (Roraima, Minas Gerais, Rio de Janeiro, Guanabara), Argentina.

Genus PARAMINTHO Brauer & Bergenstamm

PARAMINTHO Brauer & Bergenstamm, 1891:62(366). Type-species, *modulata* Wulp (sub. mon.).

MODULATA Wulp, 1895:265. Type-locality: Mexico, Guerrero. Distr. — Mexico. Ref. — Townsend, 1938:211.

Genus RAFAELIA Townsend

RAFAELIA Townsend, 1917a:45. Type-species, *rufiventris* Townsend (orig. des.). Ref. — Lopes, 1953e:225 (revision).

ACANTHIOPTERA (Wulp), 1895:267, 271, pl. 7, fig. 3 (*Sarcophaga*). Type-locality: Mexico, Guerrero. Distr. — Mexico. Ref. — Aldrich, 1930:29, fig. 7 (type examined).

RUFIVENTRIS Townsend, 1917a:45. Type-locality: Mexico, Vera Cruz, San Rafael. Distr. — U.S.A., Mexico.

rufiventris Wiedemann of Aldrich, 1916:150, fig. 69 (*Sarcophaga*). misident.

Genus MALACOPHAGULA Bequaert

MALACOPHAGULA Bequaert, 1925:202. Type-species, *neotropica* Bequaert (orig. des.).

NEOTROPICA Bequaert, 1925:208, fig. 1. (ex *Bulimulus*). Type-locality: Brazil, Pará, Belém. Distr. — Brazil (Pará, Goiás).

Genus EMBLEMASOMA Aldrich

EMBLEMASOMA Aldrich, 1916:56. Type-species, *erro* Aldrich (orig. des.).

ERRO Aldrich, 1916:56, figs. 16, 16a, 16b. Type-locality: U.S.A., New Jersey. Distr. — U.S.A., Brazil.

Genus PESSOAMYIA Lopes

PESSOAMYIA Lopes, 1938e:333. Type-species, *prosternalis* Lopes (orig. des.).

PROSTERNALIS Lopes, 1938e:333, pl. 1, figs. 1-4. Type-locality: Brazil, Rio de Janeiro, Angra dos Reis, Jussara. Distr. — Brazil.

Genus SARCODEXIA Townsend

SARCODEXIA Townsend, 1892a:105. Type-species, *sternodontes* Townsend (orig. des.).

Cricobrachia Enderlein, 1928a:19. Type-species, *anitsiana* Enderlein (orig. des.) = *sternodontes* (Townsend).

NOTATA Lopes, 1953e:230, figs. 16-19 (ex dead lepidopterous larva). Type-locality: Venezuela, near Caracas. Distr. — Venezuela.

STERNODONTES Townsend, 1892a:106 (ex dead arthropods). Type-locality: "Jamaica". Distr. — U.S.A., Jamaica, Puerto Rico, Tobago, Bahamas, St. Vincent, Guyana, Colombia, Bolivia, Brazil, Argentina, Paraguay.

- piophila* Neiva & Gomes, 1913:17 (*Sarcophaga*). Type-locality: Brazil, Rio de Janeiro.
- freirei* Mattos, 1919:75, fig., pl. 3 (*Sarcophaga*). Type-locality: Brazil, São Paulo.
- anitsiana* Enderlein, 1928a:19 (*Cricobrachia*). Type-locality: Paraguay, Asunción.
- tessellata* Enderlein, 1928a:42 (*Liopygia*; *nomen nudum*).
- butantani* Prado & Fonseca, 1932a:35, fig. 3 (*Ctenoprosballia*). Type-locality: Brazil, São Paulo.
- diminuta* Blanchard, 1942c:170, fig. 16 (as *Cricobrachia anitsiana* ssp.) Type-locality: "Argentina".

Genus EUBOETTCHERIA Townsend

- EUBOETTCHERIA Townsend, 1927a:231. Type-species. *australis* Townsend (orig. des.).
- Ctenollioproctia* Enderlein, 1928a:27. Type-species, *venusta* Enderlein (orig. des.) = *australis* Townsend.
- Ctenoprosballia* Enderlein, 1928a:28. Type-species, *Sarcophaga afficta* Wulp (orig. des.; misident.) = *enderleini* Townsend.
- Neosarcodexia* Blanchard, 1942a:60. Type-species, *deaurata* Blanchard (orig. des.) = *florencioi* (Prado & Fonseca).
- ABRUPTA Lopes, 1955a:85, figs. 7-14. Type-locality: Peru. Callao. Distr. — Peru.
- AEQUATA (Wulp), 1895:269, 1896:286 (*Sarcophaga*). Type-locality: Mexico. Guerrero. Distr. — Mexico. Ref. — Aldrich, 1930:36, fig. 17 (type examined).
- garleppi* Enderlein, 1928c:150 (*Udamoctis*). Type-locality: "Bolivia"
- ANATINA Lopes, 1953a:42, fig. 5-8. Type-locality: Ecuador. Puerto Grande, Isla Puna. Distr. — Ecuador.
- ANGULLA (Curran & Walley), 1934:484, fig. 36 (*Sarcophaga*). Type-locality: Guyana, Kartabo. Distr. — Guyana, Brazil (Mato Grosso, Rio de Janeiro, Guanabara, São Paulo), Argentina.
- ASINOMA Hall, 1938a:259, fig. 6. Type-locality: Ecuador. Oña. Distr. — Ecuador.
- AUSTRALIS Townsend, 1927a:306. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil (São Paulo, Santa Catarina, Rio Grande do Sul), Paraguay, Argentina. Ref. — Lopes, 1934:45, pls. 1-2 (paratype examined).
- tessellata* Fabricius, 1805:285 (*Musca*; not Fabricius, 1794), misident. Ref. — Townsend, 1931a:74 (type examined).
- plinthopyga* Wiedemann of Mattos, 1919:71, fig., pl. 3 (*Sarcophaga*), misident.
- venusta* Enderlein, 1928a:27 (*Ctenollioproctia*). Type-locality: "Paraguay". Ref. — Townsend, 1931a:78 (type examined).
- humeralis* Prado & Fonseca, 1932a:38, fig. 8 (*Stephanostoma*). Type-locality: Brazil, São Paulo.
- EIFURCATA (Dodge), 1965:487, figs. 24-26 (*Peckia*). Type-locality: Bahamas. New Providence Is. Distr. — Bahamas.
- BUETHNI Dodge, 1965e:509, figs. 18-20. Type-locality: Jamaica, Papine Distr. — Jamaica.

- COLLUSOR (Curran & Walley), 1934:485, fig. 45 (*Sarcophaga*). Type-locality: Guyana, Kartabo. Distr. — Trinidad, Guyana, Bolivia, Brazil (Mato Grosso, Guanabara, Paraná, Santa Catarina), Argentina.
- CUBENSIS (Dodge), 1965d:488, fig. 27 (*Peckia*). Type-locality: Cuba, Havana, Lomas de Camoa. Distr. — Cuba.
- ENDERLEINI (Townsend), 1931a:76 (*Ctenoprosballia*). Type-locality: "Colombia". Distr. — Colombia.
- afficta* Wulp of Enderlein, 1928a:28 (*Sarcophaga*), misident.
- naides* Lopes, 1941b:377, figs. 19-22. Type-locality: "Colombia".
- EPIMELIA (Lopes), 1938b:284, pl. 2, figs. 3-4 (*Sarcophaga*). Type-locality: Brazil, São Paulo. Distr. — Brazil (São Paulo).
- FLORENCIOI (Prado & Fonseca), 1932a:36, fig. 2 (*Ctenoprosballia*). Type-locality: Brazil, São Paulo. Distr. — Brazil (São Paulo, Santa Catarina), Argentina.
- xanthophora* Schiner of Mattos, 1919:59, fig., pl. 3 (*Sarcophaga*), misident.
- dentifera* Hall, 1937a:208, fig. 22 (*Sarcophaga*). Type-locality: Argentina, Iguazu.
- deaurata* Blanchard, 1942a:60, fig. 1 (*Neosarcodexia*) (ex *Alabama argilacea*). Type-locality: "Argentina".
- NICASIA Lopes, 1941b:375, figs. 11, 12, 18. Type-locality: Jamaica, Mona. Distr. — Jamaica.
- PERCUSSA (Lopes), 1938c:337, pl. 3, figs. 6, 8 (*Sarcophaga*). Type-locality: Haiti, Port-au-Prince. Distr. — Haiti.
- PERLITA Lopes, 1953a:41, figs. 1-4. Type-locality: Guyana, Shudihar. Distr. — Guyana.
- SUBDUCTA (Lopes), 1935a:41, figs. 6-7 (*Sarcophaga*). Type-locality: Brazil, Guanabara, Rio de Janeiro, Corcovado. Distr. — Brazil.
- TREJOSI Lopes, 1955a:83, figs. 4-6. Type-locality: Costa Rica, San José. Distr. — Costa Rica.
- TRIDENTATA (Hall), 1937a:209, fig. 23 (*Sarcophaga*). Type-locality: "Brazil". Distr. — Brazil.
- VOLUCRIS (Wulp), 1895:269, 1896:285 (*Sarcophaga*). Type-locality: Mexico, Atoyac. Distr. — U.S.A., Mexico. Ref. — Aldrich, 1930:35, fig. 16 (type examined).

Genus PECKIA Robineau-Desvoidy

- PECKIA Robineau-Desvoidy, 1830:335. Type-species, *imperialis* Robineau-Desvoidy (Coquillett, 1910:585) = *praeceps* (Wiedemann).
- Phrissopoda* Macquart, 1835:222. Type-species, *Sarcophaga praeceps* Wiedemann (orig. des.).
- PRAECEPT (Wiedemann), 1830:355 (*Sarcophaga*). Type-locality: "Cuba". Distr. — Mexico, Cuba, Haiti, San Domingos. Ref. — Aldrich, 1916:23, fig. 1.
- imperialis* Robineau-Desvoidy, 1830:335. Type-locality: "Cuba".
- SATANICA Dodge, 1965d:489, fig. 36. Type-locality: Bahamas, Cat Is. Distr. — Bahamas.

Genus PARAPHRISSOPODA Townsend

- PARAPHRISSOPODA Townsend, 1915a:118. Type-species, *Peckia lamanensis* Robineau-Desvoidy (orig. des.) = *gulo* (Fabricius).

- Chrysostomyia* Townsend, 1931b:315. Type-species, *Sarcophaga chrysostoma* Wiedemann (orig. des.).
- Paradiscochacta* Blanchard, 1939:834. Type-species, *Adiscochacta enderleini* Engel (orig. des.).
- ADOLENDA (Lopes), 1935a:40, figs. 4-5 (*Sarcophaga*). Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil.
- AMOENA (Aldrich), 1916:207, fig. 97 (*Sarcophaga*). Type-locality: "Dominica". Distr. — Dominica.
- AURIBARBATA (Townsend), 1912a:357 (*Sarcophaga*). Type-locality: Peru, Piura. Distr. — Peru.
- CAPITATA (Aldrich), 1916:209, fig. 98 (*Sarcophaga*). Type-locality: Puerto Rico, Mayaguez. Distr. — Puerto Rico, Virgin Is., Panama.
- CHRYSSOSTOMA (Wiedemann), 1830:356 (*Sarcophaga*). Type-locality: St. Thomas. Distr. — St. Thomas, Tobago, Jamaica, Mexico, Nicaragua, Panama, Trinidad, Guyana, Brazil, Argentina. Ref. — Aldrich, 1924:210 (type examined).
- townsendi* Prado & Fonseca, 1932a:37, fig. 5 (*Chrysostoma*). Type-locality: Brazil, São Paulo.
- clotho* Curran & Walley, 1934:482, fig. 41 (*Sarcophaga*; precoc. Brèthes, 1920). Type-locality: Guyana, Kartabo.
- impura* Curran & Walley, 1934:493 (*Sarcophaga*; as *clotho* ssp.). Type-locality: Guyana, Kartabo.
- bergi* Blanchard, 1939:831 fig. 11 (*Chrysostomyia*). Type-locality: Argentina, Misiones.
- CHRYSSOSTOMATA (Hall), 1933:267, fig. 9 (*Sarcophaga*). Type-locality: Panama, Canal Zone, Miraflores. Distr. — Panama.
- CONCINATA (Williston), 1896:364 (*Sarcophaga*). Type-locality: St. Vincent. Distr. — St. Vincent (Grenadines). Ref. — Aldrich, 1930:37 (type examined).
- otiosa* Williston, 1896:364 (*Sarcophaga*). Type-locality: St. Vincent. Ref. — Aldrich, 1930:37, fig. 20 (type examined).
- CRAIGI (Dodge), 1964:52, figs. 2-4 [*Peckia* (*Chrysostomyia*)]. Type-locality: Mexico, Revillagigedo Is. Distr. — Mexico.
- ECUATORIANA Lopes, 1958:223, figs. 29, 38, 43, 50, 62, 82. Type-locality: Ecuador, Isla Puná, Puerto Grande. Distr. — Ecuador.
- ENDERLEINI (Engel), 1931:145, fig. 10 (*Adiscochacta*). Type-locality: Bolivia, northern Chiquitos. Distr. — Bolivia, Argentina, Brazil (Mato Grosso, São Paulo).
- GULO (Fabricius), 1805:283 (*Musca*). Type-locality: "South America". Distr. — Mexico, Costa Rica, Colombia, Panama, Honduras, Ecuador, Guyana, Brazil. Ref. — Aldrich, 1930:2 (type examined).
- lamanensis* Robineau-Desvoidy, 1830:335 (*Peckia*). Type-locality: French Guiana, La Mana.
- hirtipes* Walker, 1852:321 (*Sarcophaga*). Type-locality: "Colombia". Ref. — Aldrich, 1930:17.
- wiedemanni* Aldrich, 1916:193, fig. 88 (*Sarcophaga*). Type-locality: Colombia, Santa Marta.
- ILLIFERA (Aldrich), 1916:210, fig. 99 (*Sarcophaga*). Type-locality: U.S.A., Florida, Miami. Distr. — U.S.A., Puerto Rico, Bahama, Trinidad.
- HIRSUTA (Hall), 1933:268, fig. 10 (*Sarcophaga*). Type-locality: Panama, Canal Zone, Miraflores. Distr. — Panama.

- HONDURANA (Dodge), 1965d:490, fig. 34 (*Peckia*). Type-locality: British Honduras, Rio Grande. Distr. — British Honduras.
- INCERTA (Walker), 1852:324 (*Sarcophaga*). Type-locality: "Jamaica". Distr. — Jamaica. Ref. — Aldrich, 1930:18; Dodge, 1965c:510 (type examined).
- IRWINI (Dodge), 1966b:693, figs. 5I-K (*Peckia*). Type-locality: Salvador, Quezaltepeque. Distr. — Salvador.
- KEYENSIS (Dodge), 1965d:490, fig. 35 (*Peckia*). Type-locality: U.S.A., Florida Keys. Distr. — U.S.A., ?Costa Rica.
- LITHIOGASTER (Curran & Walley), 1934:482, fig. 49 (*Sarcophaga*). Type-locality: Guyana, Kartabo. Distr. — Guyana, Trinidad, Tobago.
- LUTZI Lopes, 1958:224, figs. 44, 62, 80. Type-locality: Brazil, Guanabara. Rio de Janeiro, Mangueiros. Distr. — Brazil (Pará, Rio de Janeiro, Guanabara).
- NEPHELE Lopes, 1941b:272, figs. 13-15. Type-locality: "Trinidad". Distr. — Trinidad.
- NIGRICAUDA (Dodge), 1964:51, fig. 1 [*Peckia* (*Chrysostomia*)]. Type-locality: Mexico, Revillagigedo Is. Distr. — Mexico.
- PENATA (Wulp), 1895:269; 1896:284 (*Sarcophaga*). Type-locality: Mexico, Yucatán. Distr. — Mexico, Trinidad, Ecuador, Bolivia, Brazil (Ceará, Rio de Janeiro).
- PLUMIPES (Robineau-Desvoidy), 1830:336 (*Peckia*). Type-locality: "Haiti". Distr. — Mexico, Haiti.
- RETROCITA (Hall), 1933:280, fig. 20 (*Sarcophaga*). Type-locality: Panama, Corozal. Distr. — Panama, Guyana, Brazil (Pará, Roraima, Rio de Janeiro).
- SETIFACIES Lopes, 1945d:450, figs. 1-7. Type-locality: Colombia, Gorgona Is. Distr. — Colombia.
- SPECTABILIS (Aldrich), 1916:212, fig. 100 (*Sarcophaga*). Type-locality: Bahama, Nassau. Distr. — Bahama, Cuba, Hispaniola.
- UNCINATA (Hall), 1933:271, fig. 11 (*Sarcophaga*). Type-locality: Panama, Canal Zone, Corozal. Distr. — Panama, Tobago, Peru, Brazil (Rio de Janeiro).
- URCEOLA (Hall), 1933:273, fig. 13 (*Sarcophaga*). Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.
- VILLEGASI (Dodge), 1966b:693, figs. 5G-H (*Peckia*). Type-locality: Venezuela, Tacarigua. Distr. — Venezuela.

Genus PATTONELLA Enderlein

- PATTONELLA Enderlein, 1928a:38. Type-species, *magnifica* Enderlein (orig. des.).
- Ref. — Lopes, 1941b:365 (key to spp.).
- MAGNIFICA Enderlein, 1928a:39. Type-locality: "Paraguay". Distr. — Paraguay, Brazil (Guanabara, Santa Catarina).
- OCCIDENTALIS (Thomson), 1869:532 (*Sarcophaga*). Type-locality: Peru, Callao. Distr. — Mexico, Guatemala, Santa Lucia, Honduras, Tobago, Panama, Guyana, Ecuador, Peru, Brazil (Pará, Goiás, Mato Grosso, Guanabara, São Paulo). Ref. — Aldrich, 1930:27 (type examined).

- cotyledonea* Aldrich, 1916:187, fig. 86 (*Sarcophaga*). Type-locality: Brazil, Rio de Janeiro.
- PALLIDIPILOSA (Curran & Wailely), 1934:483, fig. 47 (*Sarcophaga*). Type-locality: Guyana, Kartabo. Distr. — Guyana.
- RESONA (Lopes), 1935:320, figs. 7-8 (*Sarcophaga*). Type-locality: Brazil, São Paulo. Distr. — Brazil (Rio de Janeiro, São Paulo, Santa Catarina), Argentina.
- dangelo* Blanchard, 1938:375, fig. 5. Type-locality: Argentina, Corrientes.
- SMARTI Lopes, 1941b:367, figs. 8, 10. Type-locality: Guyana, Essequibo R. Distr. — Guyana, Brazil (Pará).

Genus SQUAMATOIDES Curran

- SQUAMATOIDES Curran, 1927:2. Type-species, *trivittata* Curran (orig. des.).
- STANLI Dodge, 1966b:699, figs. 6J-K. Type-locality: Argentina, Misiones. Distr. — Argentina.
- TRIVITTATUS Curran, 1927:3, fig. 3. Type-locality, Brazil, Mato Grosso, Chapada. Distr. — Brazil (Mato Grosso, Goiás, Minas Gerais, São Paulo).
- splazi* Lopes, 1933a:69, pls. 4-5 (*Sarcophaga*). Type-locality: Brazil, Goiás, Campinas.

Genus ADISCOCHAETA Enderlein

- ADISCOCHAETA Enderlein, 1928a:44. Type-species, *abnormis* Enderlein (orig. des.). Ref. — Townsend, 1931a:78.
- Squamata* Curran, 1928:94 (*nomen nudum*, only in key).
- ABNORMIS Enderlein, 1928a:45. Type-locality: "Paraguay". Distr. — Mexico, Panama, Puerto Rico, Colombia, Venezuela, Guyana, Brazil (Pará, Mato Grosso, Minas Gerais, Rio de Janeiro, São Paulo), Paraguay, Argentina. Ref. — Engel, 1931:145, fig. 11; Townsend, 1931a:78 (type examined).

Genus SOCORROMYIA Dodge

- SOCORROMYIA Dodge, 1964:49. Type-species, *blodgeti* Dodge (orig. des.).
- BLODGETI Dodge, 1964:50, figs. 5-8. Type-locality: Mexico, Revillagigedo Is. Distr. — Mexico.

Genus PECKIAMYIA Dodge

- PECKIAMYIA Dodge, 1966b:695. Type-species, *pecorensis* Dodge (orig. des.).
- ABNORMALIS (Hall), 1937a:211, fig. 25 (*Sarcophaga*). Type-locality: "Colombia". Distr. — Colombia, Guyana, Brazil (Rio de Janeiro, Guanabara).
- CALX (Hall), 1933:279, fig. 19 (*Sarcophaga*). Type-locality: Panama, Tobago Is. Distr. — Panama.
- EXPUNCTA (Lopes), 1935a:38, figs. 1-3 (*Sarcophaga*). Type-locality: Brazil, Rio de Janeiro, Angra dos Reis. Distr. — Brazil (Rio de Janeiro, Guanabara).
- MINUTIPENIS (Hall), 1937a:210, fig. 24 (*Sarcophaga*). Type-locality: Brazil, Mato Grosso, Chapada. Distr. — Brazil.

PECORENSIS Dodge, 1966b:695, figs. 5M-0. Type-locality: Panama, Pacora. Distr. — Panama.

Genus ABAPA Dodge

- ABAPA Dodge, 1965d:479. Type-species, *Sarcophaga currani* Hall (orig. des.).
- BELLULA Dodge, 1965d:481, fig. 4. Type-locality: Bahamas, Exuma Cays. Distr. — Bahamas.
- CURRANI (Hall), 1930:423, fig. 2 (*Sarcophaga*). Type-locality: Cuba, Portio E' de Mora. Distr. — Cuba.
- ELEUTHERAE (Dodge), 1965d:492, fig. 40 (*Sarcophaga*). Type-locality: Bahamas, Eleuthera Is. Distr. — Bahamas.
- PEDATA (Aldrich), 1916:180, fig. 82 (*Sarcophaga*). Type-locality: Cuba. Havana. Distr. — Cuba.
- PEDATULLA (Dodge), 1965d:488, fig. 28 (female only) (*Peckia*). Type-locality: Bahamas, Grand Bahama Is. Distr. — Bahamas.
- SETIFORCEPS (Hall), 1930:422, fig. 1 (*Sarcophaga*). Type-locality: Cuba. Punta E' de Mora. Distr. — Cuba.
- SULCATA (Dodge), 1965d:489, fig. 29 (*Peckia*) (female only). Type-locality: Bahamas, Grand Bahama Is. Distr. — Bahamas.
- TURKENSIS Dodge, 1965d:481. Type-locality: Turk and Caicos Is., Long Cay, south of Grand Turk Is. Distr. — Caicos and Turk Is.

Genus PTILOCNEMA Enderlein

- PTILOCNEMA Enderlein, 1928a:43. Type-species, *miniaticauda* Enderlein (orig. des.).
- MINIATICAUDA Enderlein, 1928a:43. Type-locality: Brazil, Santa Catarina. Distr. — Brazil.

Genus DIENCHAETA Enderlein

- DIENCHAETA Enderlein, 1928a:16. Type-species, *Sarcophaga turbata* Enderlein (orig. des.). = *enderleini* Townsend.
- ENDERLEINI Townsend, 1913a:76. Type-locality: "Colombia". Distr. — Colombia.
- turbata* Wulp of Enderlein, 1928a:16, misident. Ref. — Townsend, 1931a: 76 (specimen examined).

Genus HELICOBIA Coquillett

- HELICOBIA Coquillett, 1895:317. Type-species, *Sarcophaga helcis* Townsend (orig. des.) = *rapax* (Walker). Ref. — Lopes 1939c:497 (revision).
- Helicobiopsis* Townsend, 1927a:230. Types-species, *aurescens* Townsend (orig. des.).
- Notochaetophyto* Hall, 1933:258. Type-species, *resinata* Hall (orig. des.).
- Anaravinia* Townsend, 1934b:202. Type-species, *hondurana* Townsend (orig. des.) = *luginucula* (Hall).
- Oxyhelicobia* Blanchard, 1942a:63. Type-species, *chacoana* Blanchard (orig. des.).
- AURESCENS (Townsend), 1927a:230, 312 (*Helicobiopsis*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil, Argentina. Ref. — Lopes, 1936a:844, figs. 1, 2 (paratype examined).

- parvula* Lahille, 1907:87-89, fig. 13 (*Sarcophaga*; preocc. Wiedemann, 1830). Type-locality: Argentina.
- lahillei* Blanchard, 1939:842 (*Opsophyto?*; unnecessary nom. nov. for *parvula* Lahille). Type-locality: Argentina.
- BETHIAE Dodge, 1965d:487, fig. 20 (female). Type-locality: Turk and Caicos Is., West Caicos Is. Distr. — Turk and Caicos Is.
- DIPLAGIATA Dodge, 1966b:683, figs. 3D-E. Type-locality: Peru, Callao. Distr. — Ecuador, Peru.
- BORGMEIERI Lopes, 1939c:510, figs. 7, 16, 17, 46, 51. Type-locality: Brazil, Goiás, Campinas. Distr. — Brazil.
- DEBILIS (Wulp), 1895:268, 1896:279 (*Sarcophaga*). Type-locality: "Mexico" Distr. — Mexico. Ref. — Aldrich, 1930:33, fig. 12 (type examined).
- EDWARDSI (Hall), 1937b:351, 370 (*Sarcophaga*). Type-locality: Argentina, Bonpland. Distr. — Argentina.
- GIOVANNOLI Dodge, 1965d:486, figs. 18, 21. Type-locality: Bahama, Exuma Cays. Distr. — Bahama Is.
- HAYDENI Dodge, 1965d:486, figs. 17, 22. Type-locality: Bahama Is., North Bimini Is. Distr. — Bahama Is.
- HIERINGI Lopes, 1939c:513, figs. 4, 14, 15, 36. Type-locality: Brazil, Rio de Janeiro. Distr. — Guyana, Brazil (Paraíba, Rio de Janeiro, Guanabara).
- LAGINUCULA (Hall), 1933:256, fig. 1 (*Johnsonia*). Type-locality: Panama, Canal Zone. Distr. — Costa Rica, British Honduras, Panama.
- hondurana* Townsend, 1934b:202 (*Anaravinia*). Type-locality: British Honduras, Manatee. Ref. — Dodge, 1966b:684, fig. 3 (type examined).
- LUBERA (Brèthes), 1920:282 (*Sarcophaga*). Type-locality: Brazil, Rio Grande do Sul. Distr. — Brazil.
- MORIONELLA (Aldrich), 1930:31 (*Sarcophaga*). Type-locality: Cuba, Havana. Distr. — Mexico, Bahamas Is., Bermudas, Cuba, Jamaica, Puerto Rico, Virgin Is., Brazil, Argentina.
- surrubea* Wulp of Aldrich, 1916:154, fig. 72 (*Sarcophaga*). misident.
- chacoana* Blanchard, 1942a:63, fig. 2 (*Oxyhelicobia*) (ex *Alabama argillacea*). Type-locality: Argentina, Chaco.
- NEGLECTA Lopes, 1946b:127, figs. 17-22. Type-locality: Mexico, Chapultepec. Distr. — Mexico.
- PILIFERA Lopes, 1939c:508, figs. 1, 25, 26, 37, 43, 45, 54. Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil (Goiás, Guanabara), Argentina (Misiones).
- PILIPLEURA Lopes, 1939c:514, figs. 2, 8, 18, 39. Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil (Goiás, Minas Gerais, Rio de Janeiro, Guanabara), Argentina (Misiones).
- PROVIDENCIA Dodge, 1965d:486, fig. 9. Type-locality: Bahamas, New Providence Is. Distr. — Bahama Is.
- RABEI Dodge, 1965d:486. Type-locality: Turk and Caicos Is., South Caicos Is. Distr. — Turk and Caicos Is.
- RAFAX (Walker), 1849:818 (*Sarcophaga*). Type-locality: "North America". Distr. — U.S.A., Mexico, B. Honduras, Panama, Cuba, Puerto Rico, Jamaica, Bermudas, Brazil, Argentina (reared from dead Molluscs and Insects). Ref. — Aldrich, 1930:15 (type examined).
- genalis* Thomson, 1869:539 (*Sarcophaga*). Type-locality: "Brazil". Ref. — Aldrich, 1930:29 (type examined).

- Helicis* Townsend, 1892:220 (*Sarcophaga*). Type-locality: U.S.A., Ohio. Ref. — Aldrich, 1916:158, fig. 74.
- RESINATA* (Hall), 1933:259, fig. 2 (*Notochactophyto*). Type-locality: Panama. Canal Zone, Barro Colorado Is. Distr. — Panama.
- SETINERVIS* Lopes, 1939c:515, figs. 13, 19, 32. Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil.
- STELLATA* (Wulp), 1895:267, 272 (*Sarcophaga*). Type-locality: "Mexico". Distr. — Mexico. Ref. — Aldrich, 1930:30 (type examined); Lopes, 1948:563, figs. 15-16.
- SURRUBEA* (Wulp), 1895:267, 1896:273 (*Sarcophaga*). Type-locality: Mexico. Tabasco. Distr. — Mexico. Ref. — Aldrich, 1930:30, fig. 8 (type examined).

Genus PUNAPHYTO Townsend

- PUNAPHYTO* Townsend, 1915c:410. Type-species, *tridens* Townsend (orig. des.).
- TRIDENS* Townsend, 1915c:410. Type-locality: Peru, High Andes. Distr. — Peru, Ecuador.

Genus PROMAYOA Dodge

- PROMAYOA* Dodge, 1966b:697. Type-species, *peculiaris* Dodge (orig. des.).
- PECULIARIS* Dodge, 1966b:697, figs. 6D-F. Type-locality: Surinam, Paramaribo. Distr. — Surinam.

Genus STHENOPYGOPSIS Townsend

- STHENOPYGOPSIS* Townsend, 1935a:73. Type-species, *subcylindrica* Townsend (orig. des.).
- SUBCYLINDRICA* Townsend, 1935a:73. Type-locality: "Trinidad". Distr. — Trinidad.

Genus DEXOSARCOPHAGA Townsend

- DEXOSARCOPHAGA* Townsend, 1917c:221. Type-species, *transita* Townsend (orig. des.).
- Sarcomyia* Roback, 1954:64. Type-species, *Sarcophaga scelestia* Hall (orig. des.) = *transita* Townsend.
- Farrimyia* Dodge, 1965e:504. Type-species, *ruthae* Dodge (orig. des.).
- ALMEDI* (Lopes), 1938c:346, pl. 6, figs. 1-5 (*Sarcophaga*). Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil.
- BIDENTATA* Dodge, 1966b:680, figs. 2H-K. Type-locality: West Indies, Trinidad. Distr. — Trinidad.
- CHAETOSA* Blanchard, 1939:849, fig. 18. Type-locality: Argentina, Corrientes. Distr. — Argentina.
- EGREGIA* (Hall), 1933:276, fig. 17 (*Sarcophaga*). Type-locality: Panama, Canal Zone. Distr. — Panama, Trinidad.
- GLOBULOSA* Lopes, 1946c:129, figs. 24-27. Type-locality: Brazil, Rio de Janeiro. Distr. — Guyana, Brazil (Mato Grosso, Bahia, Rio de Janeiro, Guanabara).
- ITAQUA* Dodge, 1966b:681, figs. 2L-M. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil.
- MEGISTA* (Hall), 1933:278, fig. 18 (*Sarcophaga*). Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.
- METAMASHI* Lopes, 1946c:127, figs. 18-23 (ex *Metamasius*). Type-locality: Guyana, Berbice. Distr. — Guyana.

- QUECETUBA Dodge, 1966b:681, figs. 2N-0. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil.
- PETUSA (Hall), 1933:282, fig. 22 (*Sarcophaga*). Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama, Colombia (Gorgona Is).
- RUTHIAE (Dodge), 1965e:501, figs. 10, 11 (*Farrimyia*). Type-locality: "Jamalea". Distr. — Jamalea.
- TERMITARIA (Lopes), 1939a:63, figs. 1-2 (*Sarcophaga*) (ex nest of *Armitermes*). Type-locality: Brazil, São Paulo, Caçapava. Distr. — Brazil.
- TRANSITA Townsend, 1917e:221. Type-locality: Brazil, Mato Grosso, Chapada. Distr. — U.S.A., Guyana, Brazil.
- scelesta* Hall, 1931b:285, fig. 4 (*Sarcophaga*). Type-locality: U.S.A., Texas.
- steinbachii* Enderlein, 1928:22 (*Tetradiscalis*; *nomen nudum*). Type-locality: Bolivia, Sara.
- excitans* (Curran & Walley), 1934:486, fig. 44 (*Sarcophaga*). Type-locality: Guyana, Kartabo.
- adunca* Hall, 1933:275, fig. 15 (*Sarcophaga*; as *scelesta* ssp.). Type-locality: "Panama".

Genus ARACHNIDOMYIA Townsend

- ARACHNIDOMYIA Townsend, 1934a:111. Type-species, *Sarcophaga davidsoni* Coquillett (orig. des.).
- DAVIDSONI (Coquillett), 1892:24 (*Sarcophaga*). Type-locality: U.S.A., California. Distr. — U.S.A., Mexico, West Indies. Ref. — Roback, 1954:65, figs. 70-72.
- GUYANENSIS Lopes, 1946c:125, figs. 13-17 (ex cocoon of *Argiope*). Type-locality: Guyana, Berbice. Distr. — Guyana.
- INSULARIS Lopes, 1946c:123, figs. 9-12 (ex egg-sacs of Spider). Type-locality: "Haiti". Distr. — Haiti.

Genus BOETTCHERIA Parker

- BOETTCHERIA Parker, 1914:65. Type-species, *latisterna* Parker (orig. des.). Ref. — Lopes, 1950b:687.
- Parasarcocxia* Townsend, 1917b:192, 195. Type-species, *Sarcophaga parkeri* Aldrich (orig. des.).
- Winthemiola* Dodge, 1967:126. Type-species, *calceata* Dodge (orig. des.).
- ARNAUDI Lopes, 1950b:692, figs. 20-25. Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- AUREIFERA Lopes, 1950b:690, figs. 6-15. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (São Paulo, Santa Catarina).
- CALCEATA (Dodge), 1967:127, figs. A.B. (*Winthemiola*). Type-locality: "Brazil". Distr. — Peru, Brazil.
- CIMBICIS (Townsend), 1892d:126 (*Sarcophaga*). Type-locality: U.S.A., South Dakota. Distr. — U.S.A., Mexico (ex Lepidoptera and Hymenoptera larvae).
- fernaldi* Parker, 1914:72, fig. 47. Type-locality: "U.S.A."
- CUBANA Lopes, 1950b:697, figs. 45-50. Type-locality: Cuba, Oriente, Guaro Dam. Distr. — Cuba.

- DENTATA Dodge, 1966b:677, figs. 1J-K. Type-locality: El Salvador, Cerro Verde. Distr. — El Salvador.
- MARSTONI Dodge, 1966b:677, figs. 1H-I. Type-locality: Brazil, Rio de Janeiro. Distr. — Brasil.
- MEXICANA Lopes, 1950b:702, figs. 67-71. Type-locality: Mexico, Bosencheve. Distr. — Mexico.
- MUNDELLI Blanchard, 1939:825, fig. 9. Type-locality: Argentina, Tucumán. Distr. — Argentina.
- PARKERI (Aldrich), 1916:78, fig. 27 (*Sarcophaga*). Type-locality: "Jamaica". Distr. — Jamaica.
- PERUVIANA Lopes, 1950b:691, figs. 16-19. Type-locality: Peru, Junin, Chanchamayo. Distr. — Peru.
- PRAEVLANS (Wulp) 1895:268, 1896:276, pl. 7, fig. 8 (*Sarcophaga*). Type-locality: Mexico, Guerrero. Distr. — U.S.A., Mexico, Costa Rica, Colombia. Ref. — Aldrich, 1930:32, figs. 9, 9a.
- podexaureus* Enderlein, 1928b:149 (*Robineanella*). Type-locality "Colombia".
- siccana* Reinhard, 1947:117, fig. 20 (*Sarcophaga*). Type-locality: U.S.A., Arizona.
- PYRRHOPYGA (Hall), 1933:265, fig. 7 (*Sarcophaga*). Type-locality: Panama, Barro Colorado, Canal Zone. Distr. — Panama.
- RETROVERSA (Lopes), 1935:44, figs. 10-11 (*Sarcophaga*). Type-locality: Brazil, Rio de Janeiro, Corcovado. Distr. — Brazil.
- SIMILIS Lopes, 1946b:126, figs. 12-16. Type-locality: Mexico, Cuernavaca. Distr. — Mexico.

Genus BOETIA Dodge

- BOETIA Dodge, 1965c:255. Type-species, *curiosa* Dodge (orig. des.).
- CURIOSA Dodge, 1965c:255, figs. 7-10. Type-locality: Peru, Cuzco. Distr. — Peru.
- FUSCIPENNIS Dodge, 1967:123, figs. E-F. Type-locality: Venezuela, Kaol?. Distr. — Venezuela.

Genus LIPOPTILOCNEMA Townsend

- LIPOPTILOCNEMA Townsend, 1934a:111. Type-species, *lanei* Townsend (orig. des.).
- Lahillemyia* Blanchard, 1939:818. Type-species, *Sarcophaga margareti* Lahille (orig. des.).
- Parapeltopyga* Blanchard, 1939:845. Type-species, *liguloides* Blanchard (orig. des.) = *lanei* Townsend.
- CRISPINA (Lopes), 1938b:282, pl. 1, fig. 5, pl. 2, figs. 1-2 (*Sarcophaga*). Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil (Rio de Janeiro, Guanabara, São Paulo).
- CRISPULA (Lopes), 1938b:281, pl. 1, figs. 2, 3, 6 (*Sarcophaga*). Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil (Guanabara, São Paulo).
- KOEHLERI (Blanchard), 1939:847, fig. 17 (*Parapeltopyga*). Type-locality: Argentina, Salta. Distr. — Brazil (Rio Grande do Sul), Argentina.
- LANEI Townsend, 1934a:111. Type-locality: Brazil, São Paulo, São Paulo, Pinheiros. Distr. — Brazil (São Paulo, Santa Catarina, Rio

Grande do Sul), Argentina. Ref. — Lopes, 1942c:297, figs. 1-3. *liguloides* Blanchard, 1939:845, fig. 16 (*Parapeltopyga*). Type-locality: Argentina, Salta.

MARGARETI (Lahille), 1907:92-94, fig. 17 (*Sarcophaga*) (ex Acriidae). Type-locality: Argentina. Distr. — Argentina. Ref. — Blanchard, 1939:819, fig. 7 (*Lahillemyia*).

MISELLA (Lopes), 1938c:335, figs. 4-6 (*Sarcophaga*). Type-locality: Brazil, Goiás, Campinas. Distr. — Brazil.

SALOBRENSIS Lopes, 1942c:301, figs. 4-9. Type-locality: Brazil, Mato Grosso, Salobra. Distr. — Brazil.

Genus MALACOPHAGOMYIA Lopes

MALACOPHAGOMYIA Lopes, 1966:316. Type-species, *Sarcophaga filamenta* Dodge (orig. des.).

FILAMENTA (Dodge), 1963:240, fig. 4 (*Sarcophaga*). Type-locality: "Surinam". Distr. — Surinam, Bolivia, Brazil (Pará, Rio de Janeiro). Ref. — Lopes, 1966:316, figs. 1-18.

Genus LOPESIMYIA Blanchard

LOPESIMYIA Blanchard, 1954:9. Type-species, *Sarcophaga albicans* Lahille (orig. des.) = *stallengi* Lahille.

STALLENGI (Lahille), 1907:89, figs. 14-15 (*Sarcophaga*). Type-locality: "Argentina". Distr. — Argentina.

albicans Lahille: 1907:91 (*Sarcophaga*; as *stallengi* ssp.). Type-locality: "Argentina". Ref. — Blanchard, 1945:40, figs. a-g.; Salavin, 1958:301 (ex dead Acriidae).

Genus UDAMOPYGA Hall

UDAMOPYGA Hall, 1938a:255. Type-species, *smagra* Hall (orig. des.). Ref. — Lopes, 1940:931 (revision); Dodge, 1965b:679 (revision).

MEROKELLYMYIA Blanchard, 1939:843. Type-species, *nodositas* Blanchard (orig. des.) = *setigena* (Enderlein).

APOLINARI (Lopes), 1938c:347, pl. 6, figs. 6-7 (*Sarcophaga*). Type-locality: "Colombia". Distr. — Colombia.

BARTICA (Curran & Walley), 1934:485, fig. 48 (*Sarcophaga*). Type-locality: Guyana, Kartabo. Distr. — Guyana.

BATEMANI Dodge, 1965b:681, figs. 1G-I. Type-locality: Mexico, Linares, Nuevo León. Distr. — Mexico.

BONIATO Dodge, 1965b:680, figs. 1E-F. Type-locality: Cuba, Boniato. Distr. — Cuba.

CUBANA Lopes, 1940:946, figs. 7, 8, 37, 38, 51. Type-locality: Cuba, Soledad. Distr. — Cuba.

DIVERSA Lopes, 1940:948, figs. 4-5, 33, 34. Type-locality: Brazil, Guanabara, Rio de Janeiro, Méier. Distr. — Brazil (Guanabara).

MALACOPHAGA Lopes, 1940:944, figs. 1-3, 25-26, 49 (ex *Thaumastus*). Type-locality: Brazil, Guanabara, Rio de Janeiro, Grajaú. Distr. — Brazil (Goiás, Guanabara, Santa Catarina).

- NEIVAI Lopes, 1940:942, figs. 13-15, 29, 45-48 (ex *Thaumastus*). Type-locality: Brazil, Guanabara Rio de Janeiro, Grajaú. Distr. — Brazil (Guanabara).
- PERCITA (Lopes), 1938a:344, pl. 5, figs. 5-7 (*Sarcophaga*). Type-locality: Brazil, Guanabara Rio de Janeiro, Méier. Distr. — Brazil (Guanabara, São Paulo).
- PROVEETA (Lopes), 1939b:122, figs. 3-4 (*Sarcophaga*). Type-locality: Argentina, Misiones. Distr. — Argentina.
- SETIGENA (Enderlein), 1928b:129 (*Udamoctis*) (ex *Ampullaria* and *Strophocheilus*). Type-locality: Paraguay, Santa Trinidad. Distr. — Brazil (Goiás, Rio Grande do Sul) Paraguay, Argentina. Ref. — Hall, 1938:256, fig. 3 (type examined).
- ogloblini Lopes, 1938b:119, figs. 1-2 (*Sarcophaga*). Type-locality: Argentina, Misiones.
- nodositas Blanchard, 1939:843, fig. 15 (*Merokellymyia*). Type-locality: Argentina, Misiones.
- SMAGRA Hall, 1938:255, fig. 2. Type-locality: Argentina, Tucumán. Distr. — Argentina.

Genus CHLOROSARCOPHAGA Townsend

- CHLOROSARCOPHAGA Townsend, 1919b:543. Type-species, *cochliomyia* Townsend (orig. des.).
- AZUREA (Enderlein), 1928c:148 (*Robineanella*). Type-locality: "Colombia". Distr. — Colombia.
- BICOLOR Dodge, 1967:124, fig. D. Type-locality: "Venezuela". Distr. — Venezuela.
- CALLIPHORINA (Enderlein), 1928c:147 (*Robineanella*). Type-locality: "Paraguay". Distr. — Bolivia, Brazil, Paraguay. Ref. — Dodge, 1966b:680, figs. 2C-G (type examined).
- COCHLIOMYIA Townsend, 1919b:543. Type-locality: Panama, Canal Zone, Upper Chagres R. Distr. — Panama.

Genus CHLORONESIA Townsend

- CHLORONESIA Townsend, 1912a:360. Type-species, *andina* Townsend (orig. des.).
- ANDINA Townsend, 1912a:360. Type-locality: Ecuador, Cordillera Occidental. Distr. — Ecuador.
- ARAGUA Dodge, 1966b:678, figs. 1L-M; 2A-B. Type-locality: Venezuela, Maracay?. Distr. — Venezuela.

Genus HYSTRICOCNEMA Townsend

- HYSTRICOCNEMA Townsend, 1919a:160. Type-species, *Sarcophaga robusta* Aldrich (orig. des.) = *plinthopyga* (Wiedemann).
- PLINTHOPYGA (Wiedemann), 1830:360 (*Sarcophaga*). Type-locality: St. Thomas Is. Distr. — U.S.A., Mexico, Guatemala, West Indies, Galápagos, Guyana, Brazil.
- tripartita Wulp, 1895:269; 1896:284 (*Sarcophaga*). Type-locality: Mexico, Jalisco. Ref. — Aldrich, 1930:35 (type examined).
- robusta Aldrich, 1916:268, fig. 128 (*Sarcophaga*). Type-locality: U.S.A., Texas.

quadriforceps Curran, 1932:365, fig. (*Sarcophaga*). Type-locality: Galápagos, Floreana.

khalili Salen, 1935:41, figs. 10-11 (*Sarcophaga*). Type-locality "Egypt" (wrong label).

Genus SABINATA Parker

SABINATA Parker, 1921:112. Type-species, *catalina* Parker (orig. des.) = *villipes* (Wulp).

VILLIPES (Wulp), 1895:269, pl. 7, fig. 2 (*Sarcophaga*). Type-locality: Mexico, Guerrero. Distr. — U.S.A., Mexico.

catalina Parker, 1921:112, figs. 1-5. Type-locality: U.S.A., Arizona.

Genus RAVINIOPSIS Townsend

RAVINIOPSIS Townsend, 1918a:160. Type-species, *aurca* Townsend (orig. des.).

Sarcophagla Curran, 1928:102. Type-species, *candida* Curran (orig. des.).

Pachygraphomyia Hall, 1933:259. Type-species, *spinosa* Hall (orig. des.).

AUREA Townsend, 1918a:160. Type-locality: Peru, Piura. Distr. — Peru.

CANDIDA (Curran), 1928:102, fig. 34 (*Sarcophagla*). Type-locality: Puerto Rico, Santurce. Distr. — Puerto Rico, Jamaica.

SPINOSA (Hall), 1933:260, fig. 3 (*Pachygraphomyia*). Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama, Guyana, Brazil (Pará, Guanabara, São Paulo, Paraná).

Genus ARGORAVINIA Townsend

ARGORAVINIA Townsend, 1917b:190, 193, 195. Type-species, *Sarcophaga argentea* Townsend (orig. des.) = *rufliventris* (Wiedemann).

RUFLIVENTRIS (Wiedemann), 1830:362 (*Sarcophaga*). Type-locality: "Brazil". Distr. — U.S.A., Mexico, Jamaica, Puerto Rico, Guatemala, Panama, Honduras, Guyana, Peru, Bolivia, Brazil, Argentina. Ref. — Aldrich, 1930:5, fig. 27 (type examined).

modesta Wiedemann, 1830:363 (*Sarcophaga*). Type-locality: "Brazil". Ref. — Aldrich, 1930:7 (type examined).

argentea Townsend, 1912a:358 (*Sarcophaga*). Type-locality: Peru, Piura. Ref. — Townsend, 1918b:20.

fissa Aldrich, 1916:290, fig. 140 (*Sarcophaga*). Type-locality: Honduras, Pt. Cortez.

sanctijosephi Engel, 1931:150, fig. 19 [*Sarcophaga* (?*Plerretia*)]. Type-locality: Bolivia, Chiquitos, San José.

guyanica Curran & Walley, 1934:479, fig. 33 (*Helicobia*). Type-locality: Guyana, Kartabo.

Genus TRIPANUEGA Brauer & Bergenstamm

TRIPANUEGA Brauer & Bergenstamm, 1891:63 (367). Type-species, *Sarcophaga albicans* Wiedemann [Brauer & Bergenstamm, 1893:77 (165)].

ALBICANS (Wiedemann), 1830:363 (*Sarcophaga*). Type-locality: "Brazil". Distr. — Mexico, Costa Rica, Venezuela, Peru, Bolivia, Brazil (Pará, Goiás, Minas Gerais, Rio de Janeiro, São Paulo, Paraná), Argentina. Ref. — Aldrich, 1930:8, fig. 24 (type examined).

TERMITOPHILA Silvestri, 1903:185 (from termite nest). Type-locality: "America Meridionalis". Distr. — South America.

Genus ZYGASTROPYGA Townsend

ZYGASTROPYGA Townsend, 1917b:191, 194, 195. Type-species, *aurea* Townsend (orig. des.).

SULCATA (Aldrich), 1916:223, fig. 106, 106a (*Sarcophaga*). Type-locality: U.S.A., Texas, Crystal City. Distr. — U.S.A., Mexico.

Genus GIGANTOTHECA Townsend

GIGANTOTHECA Townsend, 1917b:192, 194, 195. Type-species, *galapagensis* Townsend (orig. des.) = *violenta* (Walker).

ISLA (Curran), 1934:169, fig. 3 (*Sarcophaga*). Type-locality: Galápagos Is. Distr. — Galápagos Is.

VIOLENTA (Walker), 1849:826 (*Sarcophaga*). Type-locality: Galápagos Is. Distr. — Galápagos Is. Ref. — Aldrich, 1930:16, fig. 2, 2a (type examined).

Genus AIRYPEL Dodge

AIRYPEL Dodge, 1965:502. Type-species, *molluscoperta* Dodge (orig. des.) = *cryptopyga* (Lopes).

CRYPTOPYGA (Lopes), 1956a:208, figs. 1-10 (*Titanogrypa*). Type-locality: Brazil, Mato Grosso, Salobra. Distr. — Brazil (Mato Grosso). Bolivia, Guyana, Jamaica.

molluscoperta Dodge, 1965:502, figs. 1-3. Type-locality: Jamaica. Mandeville.

Genus CUCULOMYIA Roback

CUCULOMYIA Roback, 1954:83. Type-species, *Sarcophaga pedunculata* Hall (orig. des.).

LARVICIDA (Lopes), 1935c:470, figs. 1-6 (*Sarcophaga*) (larvae I — III. habits). Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil (Rio Grande do Norte, Rio de Janeiro).

LUCULENTA (Lopes), 1935a:209 (*Sarcophaga*; nom. nov. for *despecta* Lopes). Type-locality: Brazil, Rio de Janeiro, Angra dos Reis. Distr. — Brazil.

despecta Lopes, 1935a:42, figs. 8-9 (*Sarcophaga*; preocc. Thomson. 1869). Type-locality: Brazil, Rio de Janeiro, Angra dos Reis.

PEDUNCULATA (Hall), 1931:284, fig. 3 (*Sarcophaga*). Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico.

PLACIDA (Aldrich), 1925:24, fig. 1b (*Sarcophaga*). Type-locality: Panama. Canal Zone, Ancon. Distr. — Panama.

Genus TITANOGRYPHA Townsend

TITANOGRYPHA Townsend, 1917a:44. Type-species, *Sarcophaga alata* Aldrich (orig. des.).

Acanthotheca, subg. *Lepyria* Roback, 1954:89. Type-species, *Sarcophaga alata* Aldrich (orig. des.).

ALATA (Aldrich), 1916:109, fig. 44 (*Sarcophaga*, as *melampyga* var.). Type-locality: U.S.A., Florida, Miami. Distr. — U.S.A., Cuba, Bahamas.

RUBRICASTER Dodge, 1965d:495, figs. 49, 57. Type-locality: Bahamas, Grand Bahama Is. Distr. — Bahamas.

Genus TULAEPODA Townsend

TULAEPODA Townsend, 1917b:190, 195. Type-species, *Sarcophaga pervillosa* Aldrich (orig. des.).

ABDOMINALIS Dodge, 1965d:496, figs. 50, 53. Type-locality: Bahamas, Fortune Is. Distr. — Bahamas.

CAROLAE Dodge, 1965d:496, fig. 55. Type-locality: Turk and Caicos Is. Long Cay. Distr. — Turk and Caicos Is.

INCHOATA (Aldrich), 1916:92 (*Sarcophaga*, as *pervillosa* var.). Type-locality: Bahamas, Nassau. Distr. — Bahamas.

PERVILLOSA (Aldrich), 1916:92, fig. 36 (*Sarcophaga*). Type-locality: Cuba, Havana. Distr. — Cuba.

Genus SARCONEIVA Lopes

SARCONEIVA Lopes, 1940:949. Type-species, *Sarcophaga larvivorax* Lopes (orig. des.).

AFFINIS Lopes, 1948:559, figs. 6-12. Type-locality: Mexico, Yucatán, Peto. Distr. — Mexico.

FIMBRIATA (Aldrich), 1916:293, fig. 141 (*Sarcophaga*). Type-locality: Jamaica, Portland. Distr. — Jamaica.

HIRERINGI (Lopes), 1935b:317, figs. 3-4 (*Sarcophaga*). Type-locality: Brazil, Pernambuco, Pajeú R. Distr. — Brazil.

LARVIVORAX (Lopes), 1935c:475, figs. 7-16 (*Sarcophaga*). Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil. Ref. — Lopes, 1940:949, 950, fig. 50 (ex *Bradybaena*).

Genus VILLEGASIA Dodge

VILLEGASIA Dodge, 1966b:700. Type-species, *difficilis* Dodge (orig. des.). *DIFFICILIS* Dodge, 1966b:700, figs. 6H-I. Type-locality: Venezuela, Tacarigua. Distr. — Venezuela.

POSTUNCINATA (Hall), 1933:275, fig. 16 (*Sarcophaga*). Type-locality: Panama, Canal Zone, Patilla Point. Distr. — Panama.

Genus STEATOPYGA Dodge

STEATOPYGA Dodge, 1965c:254. Type-species, *mirabilis* Dodge (orig. des.). *MIRABILIS* Dodge, 1965c:254, figs. 4-6. Type-locality: "Colombia". Distr. — Colombia.

Genus SARCOPODEXTIOPSIS Blanchard

SARCOPODEXTIOPSIS Blanchard, 1942a:96. Type-species, *tucumana* Blanchard (orig. des.).

TUCUMANA Blanchard, 1942a:96, fig. 3 (ex *Alabama*). Type-locality: Argentina, Tucumán. Distr. — Argentina.

Genus SARCODEXIOPSIS Townsend

SARCODEXIOPSIS Townsend, 1917b:190. 193. 195. Type-species, *Sarcophaga biseriata* Aldrich (orig. des.).

Aphelomyia Roback, 1954:81. Type-species, *Sarcophaga welchi* Hall (orig. des.).

AMPULLULA (Engel), 1931:142, fig. 7 (*Bercaea*). Type-locality: N. Argentina, Lapango. Distr. — Brazil (São Paulo), Argentina.

BISERIATA (Aldrich), 1916:153, fig. 71 (*Sarcophaga*). Type-locality: Cuba, Havana. Distr. — Cuba.

SALUTARIS Dodge, 1965d:491, fig. 37. Type-locality: Bahamas, Grand Bahama Is. Distr. — Bahamas.

WELCHI (Hall), 1930:2, fig. 3 (*Sarcophaga*). Type-locality: Cuba, Punta Alegre. Distr. — U.S.A., Cuba, Bahamas.

Genus MICROPLAGIA Townsend

MICROPLAGIA Townsend, 1915c:437. Type-species, *nitens* Townsend (orig. des.).

NITENS Townsend, 1915c:437. Type-locality: Peru, Lima. Distr. — Peru.

Genus GLAUCOSARCOPIAGA Townsend

GLAUCOSARCOPIAGA Townsend, 1917a:158. Type-species, *knabi* Townsend (orig. des.).

KNABI Townsend, 1917a:158 (nom. nov. for *villipes* Wulp of Aldrich). Type-locality: Mexico, Vera Cruz. Distr. — Mexico.

villipes Wulp of Aldrich, 1916:178 (*Sarcophaga*), misident.

Genus CARINOCLYPEUS Dodge

CARINOCLYPEUS Dodge, 1965d:483. Type-species, *creameri* Dodge (orig. des.).

CREAMERI Dodge, 1965d:483, figs. 58-59. Type-locality: Bahamas, Nassau. Distr. — Bahamas.

Genus TOLUCAMYIA Dodge

TOLUCAMYIA Dodge, 1965c:256. Type-species, *marstoniana* Dodge (orig. des.).

CAVAGNAROI Dodge, 1965c:259, figs. 16-18. Type-locality: Guatemala, Panajachel. Distr. — Guatemala.

MARSTONIANA Dodge, 1965c:256, figs. 13-15. Type-locality: Mexico, 20 mi. E. of Toluca. Distr. — Mexico.

Genus SARCOPHAGULOPSIS Blanchard

SARCOPHAGULOPSIS Blanchard, 1939:798. Type-species, *trigonophynt* Blanchard (orig. des.).

TRIGONOPHYNI Blanchard, 1939:798 (ex *Acerididae*). Type-locality: "Argentina". Distr. — Argentina (Bahia Blanca, La Colona, Pampa).

Genus BERCAEA Robineau-Desvoidy

BERCAEA Robineau-Desvoidy, 1863:421. Type-species, *Musca haemorrhoidalis* Fallén (orig. des.).

HAEMORRHODALIS (Fallén), 1816:236 (*Musca*). Type-locality: "Europa". Distr. — Nearly Cosmopolitan. Ref. — Boettcher, 1913:10, fig. 37.

Genus PARASARCOPHAGA Johnston & Tlegs

Sarcophaga, subg. PARASARCOPHAGA Johnston & Tlegs, 1921:86. Type-species, *omega* Johnston & Tlegs (orig. des.) = *knabi* (Parker).

Subgenus THOMSONEA Rohdendorf

Parasarcophaga, subg. THOMSONEA Rohdendorf, 1937:247. Type-species, *Sarcophaga barbata* Thomson (orig. des.) = *argyrostoma* (Robineau-Desvoidy).

Probelliaria Blanchard, 1942:149. Type-species, *Sarcophaga barbata* Thomson (orig. des.) = *argyrostoma* (Robineau-Desvoidy).

ARGYROSTOMA (Robineau-Desvoidy), 1830:340 (*Myophora*). Type-locality: "Cap de Bonne Esperance". Distr. — Cosmopolitan. Ref. — Séguy, 1911:77, figs. 65-70 (type examined).

barbata Thomson, 1869:533 (*Sarcophaga*). Type-locality: Hawaii, Honolulu.

falculata Pandellé, 1896:185 (*Sarcophaga*). Type-locality: "France".

argentina Brèthes, 1916:142 (*Sarcophaga*). Type-locality: "Argentina".

henseliana Enderlein, 1928e:152 (*Mesothyrsta*). Type-locality: Brazil. Rio Grande do Sul, Porto Alegre.

Subgenus JANTIA Rohdendorf

Parasarcophaga, subg. JANTIA Rohdendorf, 1937:251. Type-species, *Sarcophaga securifera* Villeneuve (orig. des.) = *crassipalpis* (Macquart).

CRASSIPALPIS (Macquart), 1839:112 (*Sarcophaga*). Type-locality: "Canary Is". Distr. — Europe, N. Africa, Canary Is., N. America, Asia, Australia; Argentina.

securifera Villeneuve in Becker, 1908:123, fig. 38 (*Sarcophaga*). Type-locality: Canary Is., Teneriffe.

Subgenus LIOPYGIA Enderlein

LIOPYGIA Enderlein, 1928a:41 (as genus). Type-species, *Musca ruficornis* Fabricius (orig. des.).

RUFICORNIS (Fabricius), 1794:314 (*Musca*). Type-locality: "East Indies". Distr. — Oriental Region, Hawaii; Brazil.

Subgenus LIOSARCOPHAGA Rohdendorf

Parasarcophaga, subg. LIOSARCOPHAGA Rohdendorf, 1937:205. Type-species, *Sarcophaga madeirensis* Schiner (orig. des.).

SARRACENIOIDES (Aldrich), 1916:227, fig. 108 (*Sarcophaga*; as *tuberosa* ssp.).
Type-locality: U.S.A., Indiana, La Fayette. Distr. — N. America, Mexico.

Genus NEOBELLIERIA Blanchard

NEOBELLIERIA Blanchard, 1939:822. Type-species, *brothesi* Blanchard (orig. des.) = *offecta* (Lopes).

Sapromyia Roback, 1954:64. Type-species, *Sarcophaga bullata* Parker (orig. des.).

OFFECTA (Lopes), 1938:342, pl. 5, figs. 1-4 (*Sarcophaga*; as *polistensis* ssp.).
Type-locality: Brazil, São Paulo, São Paulo, Cantareira. Distr. — Brazil (Guanabara, Rio de Janeiro, São Paulo, Santa Catarina), Argentina.

brothesi Blanchard, 1939:822, fig. 8 (ex *Polistes versicolor* L.).
Type-locality: Argentina, Buenos Aires.

TRIPLASIA (Wulp), 1895:269, 1896:283, pl. 7, fig. 10 (*Sarcophaga*). Type-locality: Mexico, Guerrero. Distr. — U.S.A., Mexico.

dissidia Parker, 1917:157, fig. 7 (*Sarcophaga*, as *fulvipes* ssp.).
Type-locality: U.S.A., New York, Niagara Falls.

Genus WOHLFAHRTIOPSIS Townsend

WOHLFAHRTIOPSIS Townsend, 1917a:45. Type-species, *Sarcophaga johnsoni* Aldrich (orig. des.).

Scarabacophaga Townsend, 1918a:160. Type-species, *Sarcophaga utilis* Aldrich (orig. des.).

Petrosarcophaga Townsend, 1919b:543. Type-species, *arizonica* Townsend (orig. des.) = *bishoppi* (Aldrich).

BEERI (Dodge), 1966a:89 [*Sarcophaga* (*Scarabacophaga*)]. Type-locality: Mexico, Cholula. Distr. — Mexico.

BISHOPPI (Aldrich), 1916:258, fig. 123 (*Sarcophaga*). Type-locality: U.S.A., Texas, Crystal City. Distr. — U.S.A., Mexico.

arizonica Townsend, 1919b:544 (*Petrosarcophaga*). Type-locality: U.S.A., Arizona. Ref. — Townsend, 1938:53.

KESSELLI (Dodge), 1966a:92, fig. 2 [*Sarcophaga* (*Scarabacophaga*)]. Type-locality: U.S.A., Arizona, Yavapai. Distr. — U.S.A., Mexico.

UNPLACED BUT WELL CHARACTERIZED SPECIES OF SARCOPHAGIDAE.

Some of these species are well characterized by illustrations, but not sufficiently known to be included in the present genera; some of them may even constitute new genera.

ACCURATA Hall, 1933:264, fig. 6 (*Sarcophaga*). Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.

ALBISIGNUM Aldrich, 1916:111, fig. 46 (*Sarcophaga*). Type-locality: Guatemala, Los Amates. Distr. — Guatemala.

BIFRONS Walker, 1852:327 (*Sarcophaga*). Type-locality: "South America". Distr. — South America. Ref. — Aldrich, 1930:18, fig. 4 (type examined).

- CAMURA Hall, 1938b:171, fig. 1 (*Sarcophaga*). Type-locality: Venezuela, Maracay. Distr. — Venezuela.
- CASSIDIFERA Engel, 1931:152, fig. 21 [*Sarcophaga* (*Paraphrissopoda*)]. Type-locality: Bolivia, El Cairo. Distr. — Panama, Bolivia, Brazil (Pará, Rio de Janeiro, Guanabara), Argentina.
- losquí Blanchard, 1939:828, fig. 10 (*Paraphrissopoda*). Type-locality: Argentina, Misiones.
- CHIROTHECA Hall, 1933:266, fig. 8 (*Sarcophaga*). Type-locality: Panama. Canal Zone, Barro Colorado Is. Distr. — Panama.
- DECLINATA Hall, 1933:263, fig. 4 (*Sarcophaga*). Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.
- DELECTA Wulp, 1895:268; 1896:282 (*Sarcophaga*). Type-locality: Mexico, Guerrero. Distr. — Mexico. Ref. — Aldrich, 1930:34, fig. 13 (type examined).
- DOWNSIANA Dodge, 1963:239, figs. 1-3 (*Sarcophaga*). Type-locality: Venezuela, Caracas, Cerro Ávila. Distr. — Venezuela.
- ELABORATA Hall, 1933:281, fig. 21 (*Sarcophaga*). Type-locality: Panama, Corozal. Distr. — Panama.
- GLADYSAE Dodge, 1965d:493, fig. 45 (*Sarcophaga*). Type-locality: Bahamas, Exuma Cays. Distr. — Bahamas.
- HOSPES Aldrich, 1916:156, fig. 73 (*Sarcophaga*). Type-locality: Guyana, Bartica. Distr. — Guyana.
- MENTOR Curran & Walley, 1934:487, fig. 46 (*Sarcophaga*). Type-locality: Guyana, Kartabo. Distr. — Guyana.
- MICROPYGIALLIS Williston, 1896:363 (*Sarcophaga*). Type-locality: St. Vincent. Distr. — St. Vincent, Jamaica. Ref. — Aldrich, 1930:37, fig. 19 (Type examined).
- PARALIS Lopes, 1938c:341, pl. 4, figs. 3-4 (*Sarcophaga*). Type-locality: Brazil, São Paulo, Valinhos. Distr. — Brazil.
- RUDICOPAGES Hall, 1933:264, fig. 5 (*Sarcophaga*). Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.
- TAURUS Aldrich, 1916:82, fig. 30 (*Sarcophaga*). Type-locality: Puerto Rico, Naguabo. Distr. — Puerto Rico.
- TURBATA Wulp, 1895:268; 1896:276 (*Sarcophaga*). Type-locality: "Mexico". Distr. — Mexico. Ref. — Aldrich, 1930:32, fig. 10 (type examined).
- VANVOASTI Dodge, 1965d:494, fig. 42, 43 (*Sarcophaga*). Type-locality: Bahamas, Exuma Cays. Distr. — Bahamas.
- WILLIAMSII Curran, 1934:170, fig. 4 (*Sarcophaga*). Type-locality: Galápagos Is. Charles Is. Distr. — Galápagos Is.

UNRECOGNIZED SPECIES

- ADVENA Walker, 1852:324 (*Sarcophaga*). Type-locality: "Brazil".
- ARTICULATA Hall, 1933:284 (*Sarcophaga*). Type-locality: Panama, Canal Zone, Los Cascadas.
- AURIMACULATA Vlummer & Soukoup, 1940:367 (*Sarcophaga*). Type-locality: Peru, Puno.

- BONPLANDI Hall, 1937b:372 (*Sarcophaga*). Type-locality: Argentina, Misiones, Bonpland.
- BRASILIENSIS Robineau-Desvoidy, 1830:338 (*Myophora*). Type-locality: Brazil, Goiás.
- BRULLEI Macquart, 1835:223 (*Phrissopoda*). Type-locality: "South America".
- CAYENNENSIS Macquart, 1843:105 (*Sarcophaga*). Type-locality: French Guiana, Cayenne.
- CHRYSELLA Robineau-Desvoidy, 1830:339 (*Myophora*). Type-locality: "Brazil".
- CHRYSTURA Rondani, 1864:25 (*Sarcophaga*). Type-locality: "South America".
- CIRCUMCISA Rondani, 1848:76 (*Sarcophaga*). Type-locality: "Brazil".
- CLOTHO Brèthes, 1920:283 (*Sarcophaga*) (ex *Schistocerca*). Type-locality: Brazil, Rio Grande do Sul.
- COMTA Wiedemann, 1830:365 (*Sarcophaga*). Type-locality: "Brazil".
- CONCLAUSA Walker, 1861:309 (*Sarcophaga*). Type-locality: "Mexico". Ref. — Aldrich, 1930:20 (type examined).
- CONTERMINA Walker, 1852:327 (*Sarcophaga*). Type-locality: "Brazil".
- CUBENSIS Macquart, 1843:263 (sep., p. 106), pl. 12, fig. 6 (*Sarcophaga*). Type-locality: "Cuba".
- CUBENSIS Robineau-Desvoidy, 1830:342 (*Myophora*). Type-locality: "Cuba".
- DECEDENS Walker, 1857:207 (*Sarcophaga*). Type-locality: "Colombia".
- DESPENSA Walker, 1861:309 (*Sarcophaga*). Type-locality: "Mexico". Ref. — Aldrich, 1930:20 (type examined).
- DIVERSIMACULATA Macquart, 1848:214 (sep., p. 54), pl. 6, fig. 4 (*Sarcophaga*). Type-locality: "Brazil". Ref. — Aldrich, 1930:23 (type examined).
- EDNAE Dodge, 1965d:492, fig. 44 (*Sarcophaga*). Type-locality: Bahamas, Exuma Cays.
- FERVIDA Robineau-Desvoidy, 1830:341 (*Myophora*). Type-locality: "Haiti".
- FLAVEOLA Robineau-Desvoidy, 1830:339 (*Myophora*). Type-locality: Brazil, Paraná, Guaratuba.
- FLAVICEPS Macquart, 1843:260 (sep., p. 103), pl. 13, fig. 3 (*Sarcophaga*). Type-locality: "Brazil". Ref. — Aldrich, 1930:22 (type examined).
- FORTIPES Walker, 1861:310 (*Sarcophaga*). Type-locality: "Haiti".
- FULVICRUS Rondani, 1850b:365 (*Sarcophaga*; nom. nov. for *rufipes* Macquart). Type-locality: "Chile".
- rufipes* Macquart, 1843:260 (sep., p. 103) (*Sarcophaga*, preocc. Wiedemann, 1830). Type-locality: "Chile".
- FULVIFES Macquart [1843:262 (sep., p. 105), pl. 12, fig. 5 (*Sarcophaga*)], of authors.
- FULVIVITTATA Walker, 1852:326 (*Sarcophaga*). Type-locality: "South America".
- GRISOFLAVESCENS Robineau-Desvoidy, 1830:383 (*Gesneria*). Type-locality: French Guiana, Cayenne.
- HUMBOLDTI Robineau-Desvoidy, 1830:338 (*Myophora*). Type-locality: "Peru".
- IMMANIS Walker, 1849:815 (*Sarcophaga*). Type-locality: "Honduras".
- INGENS Walker, 1849:816 (*Sarcophaga*). Type-locality: unknown. Ref. — Austen, 1907:342.
- INJUNCTA Walker, 1857:208 (*Sarcophaga*). Type-locality: "Brazil".
- INNOTA Walker, 1861:308 (*Sarcophaga*). Type-locality: "Mexico". Ref. — Aldrich, 1930:20 (type examined).

- INTERMUTANS Walker, 1861:308 (*Sarcophaga*). Type-locality: "Mexico".
Ref. — Aldrich, 1930:20 (type examined).
- JEJUNA Walker, 1849:811 (*Sarcophaga*). Type-locality: "Venezuela".
- MACULATA Macquart, 1843:254 (sep., p. 97), pl. 11, fig. 4 (*Phrissopoda*).
Type-locality: French Guiana, Cayenne.
- MINIMA Vimmer & Soukoup, 1940:368 (*Agria*). Type-locality: Peru, Puno.
- MINIMA Vimmer & Soukoup, 1940:368 (*Sarcophila*). Type-locality: Peru.
Cuzco.
- MUSCOIDES Bigot, 1857:817 (*Sarcophaga*). Type-locality: "Cuba".
- NOTATA Macquart, 1855:128 (sep., p. 108) (*Sarcophaga*). Type-locality:
"South America".
- ORALIS Rondani, 1868:26 (*Sarcophaga*). Type-locality: Brazil, Bahia.
- PATRICIAE Dodge, 1965d:493, fig. 46 (*Sarcophaga*). Type-locality: Bahamas,
East Bimini Is.
- PHILIPPI Rondani, 1864:24 (*Sarcophaga*). Type-locality: Chile, Valdivia.
- PHOENICRUS Wiedemann, 1830:365 (*Sarcophaga*). Type-locality: "Brazil"
- PIGMEA Rondani, 1850b:365 (*Sarcophaga*). Type-locality: Ecuador, Rio
Napo.
- PROERNA Walker, 1849:835 (*Sarcophaga*). Type-locality: Uruguay, Montevideo.
- PUDICA Rondani, 1850a:174 (*Sarcophaga*). Type-locality: Brazil, São Paulo,
São Sebastião Is.
- PUSILLA Bigot, 1857:817 (*Sarcophaga*; preocc. Wiedemann, 1830). Type-
locality: "Cuba".
- RUBELLA Wiedemann, 1830:357 (*Sarcophaga*). Type-locality: "Antigua".
Ref. — Aldrich, 1930:2 (type examined).
- RUBRIANALIS Macquart, 1851:262 (sep., p. 235), (*Sarcophaga*). Type-locality:
"Chile". Ref. — Aldrich, 1930:24 (type examined).
- RUFICRURA Rondani, 1864:25 (*Sarcophaga*). Type-locality: "Equatorial
America".
- RUFIPALPIS Macquart, 1843:102 (*Sarcophaga*). Type-locality: "Brazil".
- SUBSERICANS Walker, 1857:207 (*Sarcophaga*). Type-locality: "South America".
- SURINAMENSIS Wiedemann, 1830:366 (*Sarcophaga*). Type-locality: "Surinam".
- TRIGONOMACULATA Macquart, 1843:263 (sep., p. 106), pl. 12, fig. 2 (*Sarcophaga*).
Type-locality: "Mexico".
- TRIVITTATA Macquart, 1843:262 (sep., p. 105), pl. 12, fig. 3 (*Sarcophaga*).
Type-locality: "Cuba".
- VARIPES Rondani, 1850b:364 (*Sarcophaga*). Type-locality: Ecuador, Rio
Napo.
- VITTATA Walker, 1837:353 (*Sarcophaga*). Type-locality: unknown.

LITERATURE CITED


ALDRICH, J. M.

1916. *Sarcophaga and allies in North America*. [Vol. 1], 302 pp.,
16 pls. In Entomological Society of America, Thomas Say
Foundation. Lafayette, Ind.

1924. Notes on some types of American muscoid Diptera in the collection of the Vienna Natural History Museum. *Ann. Ent. Soc. Amer.* 17:209-218.
- 1925a. New Diptera or two-winged flies in the United States National Museum. *Proc. U. S. Nat. Mus.* 66 (art. 18):1-36, 1 fig.
- 1925b. Notes on some types of American muscoid Diptera in the collection of the Vienna Natural History Museum (cont.). *Ann. Ent. Soc. Amer.* 18:456-469, 1 fig. (cont.).
- 1927a. Notes on Muscoid synonymy. *Bull. Brooklyn Ent. Soc.* 22:18-25.
- 1927b. The dipterous parasites on the migratory locust of Tropical America — *Schistocerca paranensis* Burmeister. *J. Econ. Ent.* 20:588-593.
- 1927c. Redescription of types of American muscoid flies in the Vienna Natural History Museum, with incidental notes. *Proc. U.S. Nat. Mus.* 72 (art. 7):1-35, 3 figs.
1928. New Diptera or two-winged flies from South America. *Proc. U.S. Nat. Mus.* 74 (art. 1):1-25, 2 figs.
- 1929a. Further studies of types of American muscoid flies in the collection of the Vienna Natural History Museum. *Proc. U.S. Nat. Mus.* 74 (art. 19):1-34, 2 figs.
- 1929b. New genera and species of muscoid flies. *Proc. U.S. Nat. Mus.* 76 (art. 15):1-13.
1930. Notes on the types of American two-winged flies of the genus *Sarcophaga* and a few related forms. described by the early authors. *Proc. U.S. Nat. Mus.* 78 (art. 12):1-39, 3 pls.
- ALLEN, H. W.
1926. North American species of two-winged flies belonging to the tribe Miltogrammini. *Proc. U.S. Nat. Mus.* 68(9):1-105, 5 pls.
- ARNAUD, P. H.
1963. Records of Diptera from Guadalupe Island, Mexico. Second paper. *Ent. News* 74(5):117-129, 7 figs.
- AUSTEN, E. E.
1907. The synonymy and generic position of certain species of Muscidae (sens. lat.) in the collection of the British Museum, described by the late Francis Walker. *Ann. Mag. Nat. Hist.* (7) 19:326-347.
- BECKER, T.
1908. Dipteren der Kanarischen Inseln. *Mitt. Zool. Mus. Berlin* 4:1-180, 4 pls.
- BEQUAERT, J.
1925. The arthropod enemies of molluscs with description of a new dipterous parasite from Brazil. *J. Paras.* 11:201-212, 1 fig.
1926. A dipterous parasite of a snail from Brazil with an account of the arthropod enemies of mollusks, in Medical report of the Hamilton Rice Seventh Expedition to the Amazon, in conjunction with the Department of Tropical Medicine of Harvard University, 1924-1925. Part II. Medical and economic entomology. *Harvard Univ., Inst. Trop. Biol. and Med. Contrib.* 4:155-257, 9 figs., pls. 61-67.

- BIGOT, J. M.
1857. Ordre des Diptères, Latr., in R. de la Sagra: *Historia física, política y natural de la Isla de Cuba* 7:783-829, 1 pl.
- BISCHOF, J.
1904. Beitrag zur Kenntnis der Muscaria Schizometopa. *Verh. zool-bot. Ges. Wien* 54:79-101.
- BLANCHARD, E.
1852. Diptera, in C. Gay: *Historia física y política de Chile, Zoología* 7:327-468. Paris.
- BLANCHARD, E. E.
1933. Parasitos animales de la langosta. *Bol. Min. Agric. Buenos Aires* 34:247-266, 5 pls.
1938. Descripciones y anotaciones de dípteros argentinos. *An. Soc. Cient. Argent.* 126:345-386, 8 figs.
1939. Los sarcófágidos argentinos, contribución a su conocimiento *Physis*, Buenos Aires 17:791-856, 19 figs.
1940. Apuntes sobre muscoideos argentinos. *An. Soc. Cient. Argent.* 129:224-246.
1942a. Parasitos de *Alabama argilacea* Hbn. en la Republica Argentina. Estudio Preliminar. *An. Soc. Cient. Argent.* 134:54-63, 94-128, 17 figs.
1942b. Nuevos dípteros y himenópteros de la Republica Argentina. *Rev. Soc. Ent. Argent.* 11:340-379, 17 figs.
1942c. Los sarcophagidos argentinos, segunda contribución a su conocimiento. *Physis*, Buenos Aires 19(52):133-172, 16 figs.
1954. "*Lopesimyia*", un nuevo genero de sarcófagido argentino (Diptero). *Neotropica* 1(1):9-12, 1 fig.
1955. Nuevos sarcófagidos argentinos (Diptera). *Rev. Soc. Ent. Argent.* 18:27-34, 3 figs.
- BRAUER, F. & J. E. VON BERGENSTAMM
1889. Die Zweiflügler des kaiserlichen Museums zu Wien. IV. Vorarbeiten zu einer Monographie der Muscaria Schizometopa (exclusive Anthomyidae). Pars I. K. Akad. der Wiss. Wien, *Math.-Nat. Cl. Denkschr.* 56(1):69-189, 11 pls.
1891. Die Zweiflügler des kaiserlichen Museums zu Wien. V. Vorarbeiten zu einer Monographie der Muscaria Schizometopa (exclusive Anthomyidae). Pars III. K. Akad. Wiss. Wien, *Math.-Nat. Cl. Denkschr.* 58:305-446.
1893. Die Zweiflügler des kaiserlichen Museums zu Wien. VI. Vorarbeiten zu einer Monographie der Muscaria Schizometopa (exclusive Anthomyidae). Pars III. K. Akad. Wiss. Wien, *Math.-Nat. Cl. Denkschr.* 60:89-240.
1894. Die Zweiflügler des kaiserlichen Museums zu Wien. VII. Vorarbeiten zu einer Monographie der Muscaria Schizometopa (exclusive Anthomyidae). Pars IV. Pt. 7, 88 pp. Wien. (Also published in K. Akad. der Wiss. Wien, *Math.-Nat. Cl. Denkschr.* 61:537-624, 1895).
- BRÉTHES, J.
1900. *Sarcophaga caridei*, una nueva mosca langosticida. *An. Mus. Nac. Buenos Aires* (3) 6:297-301, 3 figs.
1912. Sobre la *Brachycoma acridiorum* (Weyenb.) (*Nemorea acridiorum* Weyenb.). *An. Mus. Nac. Buenos Aires* 22:441-446.
1916. Descripción de una nueva mosca langosticida. *An. Mus. Nac. Buenos Aires* 28:141-144.

1920. Insectos utiles y daños de Rio Grande do Sul y de la Plata. *An. Soc. Rural Argent.*, Buenos Aires 54:281-290, 307-308.
1928. A proposito de *Masarygus Brèthes* y de *Sarcophaga caridei* Brèthes. *Rev. Soc. Ent. Argent.* 2:73-74.
- CERRUTI, T.
1913. Miasis del oído, la *Sarcophaga nurus*, nueva mosca productora de miasis. *Rev. Med. Rosario* 3(4):199-203, 2 figs.
- CONIL, P. A.
1879. Études sur l'*Acridium parancense* Burm., ses variétés et plusieurs insectes qui le détruisent. *Bol. Acad. Cienc. Córdoba* 3:385-469, 4 pls. (Also published in *Periodico Zoologico, Córdoba* 3:177-254, 4 pls.).
- COQUILLETT, D. W.
1892. The dipterous parasite of *Melanoplus devastator* in California. *Insect Life* 5:22-24.
- 1895a. New genera and species of Tachinidae. *J. New York Ent. Soc.* 3:97-107.
- 1895b. Descriptions of new genera and species, pp. 307-319, in C. W. Johnson, *Diptera of Florida*. *Proc. Acad. Nat. Sci. Philad.* 1895:303-340.
1897. Revision of the Tachinidae of America North of Mexico, a family of parasitic two-winged insects. *U.S. Dept. Agric., Div. Ent. Tech. Ser.* 7:1-156.
1900. Report on a collection of dipterous insects from Porto Rico. *Proc. U.S. Nat. Mus.* 22:249-270.
1902. New cyclorrhaphous Diptera from Mexico and New Mexico. *Canad. Ent.* 34:195-202.
- 1910a. New genera and species of North American Diptera. *Proc. Ent. Soc. Wash.* 12:124-131.
- 1910b. The type-species of the North American genera of Diptera. *Proc. U.S. Nat. Mus.* 37:499-647.
- CROUZEL, I. S.
1944. First stage larva of *Acridiophaga caridei* (Brèthes) (Diptera, Sarcophagidae). *Proc. Ent. Soc. Wash.* 46(9):239-246, 12 figs.
1947. Falta de parasitismo en *Doringia acridiorum* para la langosta. *Cienc. Invest.*, Buenos Aires 3(10):434.
1950. Primer estadio larval de *Doringia acridiorum* (Weyenb.) (Diptera, Sarcophagidae). *Arthropoda*, Buenos Aires 1(2/4):291-297, 8 figs.
- CURRAN, C. H.
1927. Four new American Diptera. *Amer. Mus. Nov.* 275:1-4, 3 figs.
1928. Diptera or two-winged flies. *Sci. Surv. Porto Rico and Virgin Islands*, New York 11:1-118, 39 figs.
1932. Diptera (excl. of Tipulidae and Culicidae) of the Wollleback Galapagos Expedition. *Nyt. Mag. Naturvidensk.* 71:347-366, 1 fig.
1934. The Templeton-Crocker Expedition of the California Academy of Sciences, 1932. N° 12. Diptera. *Proc. Calif. Acad. Sci.* (4) 21:147-172, 4 figs.
1938. New American Diptera. *Amer. Mus. Nov.* 975:1-7.
- CURRAN, C. H. & G. S. WALLEY
1934. Sarcophagidae, pp. 474-491, in C. H. Curran: The Diptera of Kartabo, Bartica District, British Guiana, with description of new species from other British Guiana localities. *Bull. Amer. Mus. Nat. Hist.* 66:287-532, 54 figs.



DODGE, H. R.

1955. Sarcophagid flies parasitic on reptiles (Diptera, Sarcophagidae). *Proc. ent. Soc. Wash.* 57(4):183-187, 11 figs.
1956. Two new sarcophagid flies from Cuba (Diptera). *Mcm. Soc. Cubana Hist. Nat.* 23:97-101, 3 figs.
1963. New Sarcophaginae flies (Diptera, Sarcophagidae). *J. New York Ent. Soc.* 71:238-242, 8 figs.
1964. Sarcophagid flies from Revillagigedo Islands. *Proc. ent. Soc. Wash.* 66(1):47-54, 11 figs.
- 1965a. A new *Tricharaea* Thomson with unusual variation in male chaetotaxy. *Pan-Pacific Ent.* 41(2):90-92.
- 1965b. A review of the genus *Udamopyga* Hall (Diptera, Sarcophagidae). *Canad. Ent.* 97(7):679-684, figs. 1A-I.
- 1965c. Neotropical Sarcophagidae with impressed scutellum. *Ann. Ent. Soc. Amer.* 58(3):252-259, 18 figs.
- 1965d. The Sarcophagidae (Diptera) of the West Indies. I. The Bahama Islands. *Ann. Ent. Soc. Amer.* 58(4):474-497, 57 figs.
- 1965e. The Sarcophagidae (Diptera) of the West Indies. II. Jamaica. *Ann. Ent. Soc. Amer.* 58(4):497-517, 35 figs.
- 1966a. *Sarcophaga utilis* Aldrich and allies (Diptera, Sarcophagidae). *Ent. News* 77(4):85-97, 8 figs.
- 1966b. Some new or little-known Sarcophagidae (Diptera) with a review of the genus *Oxysarcodexia*. *Ann. Ent. Soc. Amer.* 59(4):674-701, figs. 1-6.
1967. New Neotropical Sarcophagidae in the Vienna Museum (Diptera). *Ent. News* 78:123-129, figs. A-F.

ENDERLEIN, G.

- 1928a. Klassifikation der Sarcophagiden. Sarcophagiden Studien I. *Arch. Klass. Phyl. Ent.*, Wien 1:1-56, 7 figs.
- 1928b. *Udamoctis setigena*, eine neue Sarcophaginae aus Paraguay (Dipt.). *Entom. Mitt.* 17(2):129-130.
- 1928c. Sarcophagiden Studien II. *Konowia* 7:147-153, 1 fig.

ENGEL, O.

1931. Die Ausbeute der deutschen Chaco-Expedition 1925/26. Diptera. 28. Sarcophagidae. *Konowia* 10:140-154, pls. 1-6.

FABRICIUS, J. C.

1794. *Entomologia systematica emendata et aucta* 4:472 pp. Hafniae (= Copenhagen).
1805. *Systema antliatorum secundum ordines, genera, species*, 373+30 pp. Brunsvigae (= Brunswick).

GONÇALVES, P. C.

1953. Sobre a presença de *Bercaca haemorrhoidalis* (Fallén, 1816) no Rio de Janeiro (Diptera, Sarcophagidae). *Rev. Brasil. Biol.* 13(4):359-362, 8 figs.

HALL, D. C.


1928. *Sarcophaga pallinervis* and related species in the Americas. *Ann. Ent. Soc. Amer.* 21(2):331-348, pls. 19-22, figs. 1-20.
1929. Corrections to "*Sarcophaga pallinervis* and related species in the Americas". *J. Kansas Ent. Soc.* 2(3):71.
1930. Three new West Indian Sarcophaginae (Dipt.). *Amer. Mus. Nov.* 423:1-4, 3 figs.

1931. New Texas Sarcophaginae (Dipt., Calliphoridae). *Ent. News* 42:280-286, 4 figs.
1932. Biology of *Sarothromyia femoralis* var. *simplex* Aldrich (Diptera: Calliphoridae). *Ann. Ent. Soc. Amer.* 25:641-647, 2 pls.
1933. The Sarcophaginae of Panama (Diptera, Calliphoridae). *Bull. Amer. Mus. Nat. Hist.* 66:251-285, 26 figs.
- 1937a. New Muscoid flies (Diptera) in the United States National Museum. *Proc. U. S. Nat. Mus.* 84:201-216, 8 figs.
- 1937b. *Diptera of Patagonia and South Chile* 7(3):347-384, figs. 61-71 (Fam. Calliphoridae — Sarcophaginae, Calliphorinae).
- 1938a. New genera and species of South American Sarcophagidae (Dipt.). *Arb. morphol. taxon. Ent.*, Berlin-Dahlem 5:253-258, 6 figs.
- 1938b. New South American Sarcophagidae (Diptera). *J. Agric. Univ. Porto Rico* 22:171-175, 1 pl.
- JOHNSTON, T. H. & O. W. TIEGS
1922. Sarcophagid flies in the Australian Museum collection. *Rec. Australian Mus.* 13:175-188, pl. 35, 2 figs.
1921. New and little-known sarcophagid flies from southeastern Queensland. *Proc. Royal Soc. Queensland* 33:46-90, 26 figs.
- LAHILLE, F.
1907. La langosta y sus moscas parasitarias. *An. Min. Agric. Argentina, Sección Zoot., Bact., Vet. y Zool.* 3(4):1-136, 7 pls., 29 figs.
- LOPES, H. S.
1932. Sobre dois paratipos de Townsend do Museu Paulista (Diptera, Sarcophagidae). *Bol. Biol.* 21:45-52, 4 pls.
- 1933a. Sarcophagas do Brasil. *Arch. Esc. Sup. Agric. Med. Vet.* 10:65-71, pls. 1-5.
- 1933b. Sobre algumas especies de *Sarcophaga* do Brasil, com a descrição de cinco especies novas (Dipt., Sarcophagidae). *Rev. Entom.* 3(2):143-158, 5 figs.
- 1933c. A proposito de *Sarcophaga tessclata* (Fabr., 1805). *Rev. Entom.* 3:420.
- 1935a. Algumas especies de *Sarcophaga* do Brazil. *Rev. Entom.* 5:38-46, 12 figs.
- 1935b. Notas sobre Sarcophagidae com descrição de tres especies novas do Brasil (Dipt.). *Rev. Entom.* 5:315-322, 8 figs.
- 1935c. Sobre duas especies de *Sarcophaga* cujas larvas são predadoras (Dipt. Sarcophagidae). *Rev. Entom.* 5:470-479, 16 figs.
- 1936a. Sobre alguns paratipos de Sarcophagidae conservados no Museu Paulista (Dipt.). *Rev. Museu Paulista* 21:839-853, 4 pls.
- 1936b. Sarcophagídeos neotropicos novos ou pouco conhecidos (Dipt.). *Arch. Inst. Biol. Veg.* 3(1):71-90, 28 figs.
- 1938a. *Sarcophaga lutulenta*, nov. nom. (Dipt.). *Rev. Entom.* 8(2):209.
- 1938b. Sobre quatro novas especies de sarcophagídeos do Brasil (Dipt.). *Livro Jubilar Prof. L. Travassos*, Rio de Janeiro: 279-285.
- 1938c. Notas sobre sarcophagídeos neotropicos. Um novo genero e algumas especies novas. *Mém. Inst. Oswaldo Cruz* 33(2):333-348, 6 pls.
- 1938d. Sobre um interessante novo genero de Sarcophagidae que apresenta redução dos esternitos abdominais (Dipt.). *Mem. Inst. Oswaldo Cruz* 33(3):433-435, 1 pl.



- 1938c. Sobre um Rhinophoridae e varios Sarcophagidae da coleção do "Deutsches Entomologisches Institut". *Mem. Inst. Oswaldo Cruz* 33(4):555-565, 4 pls.
- 1939a. Descrição de uma nova especie de *Sarcophaga termitophila* do Brasil (Dipt. Muscidae). *Boll. Lab. Zool. Agric. Portici* 31:63-64, 2 figs.
- 1939b. Sobre alguns sarcophagideos de Misiones, Argentina. *Physis*, B. Aires 17:117-123, figs. 1-2.
- 1939c. Contribuição ao conhecimento do genero *Helleobia* Coquillett (Dipt. Sarcophagidae). *Rev. Entom.* 10:497-517, 51 figs.
1940. Contribuição ao conhecimento do genero *Udamopyga* Hall e de outros sarcophagideos que vivem em moluscos no Brasil (Diptera). *Rev. Entom.* 11(3):924-954, 51 figs.
- 1941a. Sobre o aparelho genital feminino dos Sarcophagidae e sua importancia na classificação (Diptera). *Rev. Brasil. Biol.* 1(2): 215-221, 18 figs.
- 1941b. Sobre alguns sarcophagideos neotropicos da coleção do Museu Britanico. *Arq. Zool. S. Paulo* 2:357-387, 2 pls.
- 1942a. Duas novas especies do genero *Nephoaetopteryx* do Rio de Janeiro (Diptera, Sarcophagidae). *Rev. Chil. Hist. Nat.* 44(1940): 186-189, 6 figs.
- 1942b. *Notochaeta aldrichi* n. sp. parasita de *Oligochaeta* no Brasil (Diptera, Sarcophagidae). *Rev. Brasil. Biol.* 2(3):361-364, 4 figs.
- 1942c. Sobre o genero *Lapoptiloenema* Townsend, com a descrição de uma nova especie (Dipt. Sarcophagidae). *Rev. Entom.* 13(3):296-303, 9 figs.
1943. Contribuição ao conhecimento das larvas de Sarcophagidae, com especial referencia ao esqueleto cefalico. *Mem. Inst. Oswaldo Cruz* 38(3):127-163, 90 figs.
- 1945a. Sobre um novo genero de Sarcophagidae de Surinam (Diptera). *Rev. Brasil. Biol.* 5(2):295-298, 6 figs.
- 1945b. Sobre a presença, no Rio de Janeiro, de *Parasarcophaga ruficornis* (Fabr., 1794), especie oriental, e a possibilidade de seu transporte por avião (Dipt. Sarcophagidae). *Rev. Brasil. Biol.* 5(3):401-403, 11 figs.
- 1945c. Um novo genero e tres novas especies de Sarcophagidae do Brasil. *Rev. Brasil. Biol.* 5(4):453-462, 31 figs.
- 1945d. Sobre dois sarcophagideos neotropicos da coleção do Museu Britanico. *Rev. Entom.* 16:450-456, 13 figs.
- 1946a. Contribuição ao conhecimento das especies do genero *Notochaeta* Aldrich, 1916 (Diptera, Sarcophagidae). *Mem. Inst. Oswaldo Cruz* 42:503-550, 69 figs.
- 1946b. Sarcophagidae do Mexico, capturados pelo Professor A. Dampf (Diptera). *Mem. Inst. Oswaldo Cruz* 44:119-146, 62 figs.
- 1946c. Novos sarcophagideos neotropicos representados na coleção do "Imperial Institute of Entomology" (Diptera, Sarcophagidae). *Rev. Brasil. Biol.* 6(1):117-131, 27 figs.
- 1946d. *Chaetoravintia almeidae*, n. sp., novo Sarcophagidae do Brasil (Dipt.). *Livro Jubilar R. F. d'Almeida*, São Paulo: 227-230.
- 1946e. Contribuição ao conhecimento das especies do genero *Oxysarcodesia* Townsend, 1917 (Diptera — Sarcophagidae). *Bol. Esc. Nac. Vet.*, Rio de Janeiro 1:62-134, 162 figs.

1947. Sobre o genero *Notochaeta* Aldrich, 1916 (Diptera, Sarcophagidae), com a descrição de duas espécies novas do Amazonas. *Rev. Brasil. Biol.* 7:375-382, 13 figs.
1948. Sarcophagidae do Mexico capturados pelo prof. Dampf (2ª nota) (Diptera). *Mem. Inst. Oswaldo Cruz* 45(1947)(3):555-570, 18 figs.
- 1950a. Novas espécies neotropicas de *Notochaeta* Aldrich e *Dexosarcophaga* Townsend (Diptera, Sarcophagidae). *Rev. Brasil. Biol.* 10(3):353-364, 26 figs.
- 1950b. Sobre os generos *Boettcheria* Parker e *Boettcherimima* n. gen. (Diptera, Sarcophagidae). *Mem. Inst. Oswaldo Cruz* 48:687-732, 93 figs.
- 1951a. Sobre o genero *Harpagopyga* Aldrich, 1916 (Diptera, Sarcophagidae). *Rev. Brasil. Biol.* 11(3): 275-288, 43 figs.
- 1951b. Sobre o genero *Lepidodexia* Brauer & Bergenstamm, 1891 (Diptera, Sarcophagidae). *Mem. Inst. Oswaldo Cruz* 49:641-658, 43 figs.
- 1953a. Seis novos Sarcophagidae neotropicos (Diptera). *Rev. Brasil. Biol.* 13(1):41-51, 30 figs.
- 1953b. Considerações sobre os generos de Sarcophagidae (Diptera) propostos por Robineau-Desvoidy em 1830 e 1862. *Arq. Mus. Nac. Rio de Janeiro* 42:265-272.
- 1953c. Sobre *Rafaelia* Townsend e *Sarcodexia* Townsend (Diptera, Sarcophagidae). *Rev. Brasil. Biol.* 13(3):225-233, 19 figs.
1954. Sobre o genero *Xanthobrachycoma* Townsend, 1927 (Diptera, Sarcophagidae). *Rev. Brasil. Biol.* 14(1):65-70, 18 figs.
- 1955a. Contribuição ao conhecimento das espécies do genero *Paraphrissopoda* Townsend (Diptera, Sarcophagidae). *Mem. Inst. Oswaldo Cruz* 52 (1954):83-87, 2 pls.
- 1955b. Contribuição ao conhecimento do genero *Sarcophagula* Wulp. 1887 (Diptera, Sarcophagidae). *Mem. Inst. Oswaldo Cruz* 52 (1954): 587-602, 45 figs.
- 1955c. Catalogo de la fauna cubana. VIII. Catalogo de la familia Sarcophagidae en Cuba. *Circ. Mus. Bibl. Zool. Habana* 1954:1267-1280.
- 1956a. Sobre o genero *Titanogrypa* Townsend, 1917 (Diptera, Sarcophagidae). *Rev. Brasil. Biol.* 16:207-211, 10 figs.
- 1956b. Sobre a genitalia dos Sarcophagidae (Diptera). *Rev. Brasil. Biol.* 16:59-64, 5 figs.
1958. Considerações sobre as espécies de *Peckia* Robineau-Desvoidy e de generos afins (Diptera, Sarcophagidae). *An. Acad. Brasil. Cienc.* 30:211-243, 109 figs.
- 1962a. Contribuição ao conhecimento de *Lepidodexia* Brauer & Bergenstamm, 1891 (Diptera, Sarcophagidae). *Rev. Brasil. Biol.* 21(3):253-256, 4 figs.
- 1962b. Sobre as espécies do genero *Andinoravinia* (Diptera, Sarcophagidae). *Mem. Inst. Oswaldo Cruz* 60(2):165-173, 26 figs.
1966. Sobre *Malacophagomyia* n. gen. (Diptera, Sarcophagidae) cujas larvas vivem em cadaveres de Gastropoda (Mollusca). *Rev. Brasil. Biol.* 26(3):315-321, 18 figs.



LOPES, H. S. & D. O. ALBUQUERQUE

1955. Los insectos de las Islas Juan Fernandez. 22. Caliphoridae et Sarcophagidae (Diptera). *Rev. Chil. Hist. Nat.* 4:113-116, 43 figs.

LOPES, H. S. & W. DOWNS

1951. Contribuição ao conhecimento das espécies do genero *Acanthodotheca* Townsend (Diptera, Sarcophagidae). *Mem. Inst. Oswaldo Cruz* 47(1949):571-603, 59 figs.

LOPES, H. S. & E. VOGELSANG

1953. *Notochaeta bufonivora* n. sp., parasita de *Bufo granulosus* Spix em Venezuela (Diptera, Sarcophagidae). *An. Acad. Brasil. Cienc.* 25(2):139-143, 9 figs.

LUTZ, A.

1910. Notas dipterologicas. III. Especies de *Sarcophaga* de São Paulo. *Mem. Inst. Oswaldo Cruz* 2:58-63.

LYNCH ARRIÁZAGA, E.

1881. In C. Berg: Entomologisches aus dem Indianer-Gebiet der Pampa. IV. Diptera. *Stett. Ent. Zeit.* 46:45-46.

MACQUART, J.

1835. *Histoire naturelle des insectes. Diptères* 2:1-170, pls. 13-24 Paris.

1839. Diptères, pp. 97-119, pl. 4, figs. 2-11, in P. B. Webb & S. Berthelot, eds.: *Histoire naturelle des Iles Canaries* 2(2) Zoologie: Entomologie: 119 pp., 7 pls. Paris, "1836-1844".

1843. Diptères exotiques nouveaux ou peu connus. *Mém. Soc. Roy. des Sci., de l'Agr. et des Arts, Lille* 1842:162-460, 36 pls. (Also separately published, as Vol. 2, Pt. 3, pp. 5-304, 36 pls. Paris. 1843).

1846. *Diptères exotiques nouveaux ou peu connus. (1er. Supplément). Mém. Soc. Roy. Sci., de l'Agric. et des Arts, Lille* (1845) 1844: 133-364, 20 pls. (Also separately published, 1846: pp. 5-238).

1848. *Diptères exotiques nouveaux ou peu connus. (Suite du 2me supplément, i.e., 3e. supplément). Mém. Soc. Roy. des Sci., de l'Agric. et des Arts, Lille* 1847(2):161-237, 7 pls. (Also separately published, 1848:pp. 1-77).

1851. Diptères exotiques nouveaux ou peu connus. (Suite du 4e. supplément). *Mém. Soc. Nat. des Sci., de l'Agric. et des Arts Lille* 1850:134-282 (text), 283-294 (explanation of figs., index), pls. 15-28. (Also separately published, 1861: pp. 161-309 (text), 317-323 (explanation of figs.), 324-336 (combined index of the 2 parts of this supplement).

1855. Diptères exotiques nouveaux ou peu connus. (5e. supplément). *Mém. Soc. Imp. des Sci., de l'Agric. et des Arts, Lille* 1854:25-156. 7 pls. (Also separately published, 1855: pp. 5-136).

MACLOCH, J. R.

1929. Exotic Muscaridae (Diptera). XXVII. *Ann. & Mag. Nat. Hist.* (10) 4:249-257, 6 figs.

1932. New species of *Calliphora* from the Marquesas, with notes on *Sarcophaga taitensis* Schiner. *Bull. Bishop Mus.* 98:13-16, 2 figs.

MASSINI, P. C.

1921. Destruecion de la langosta, lucha biologica. *Prim. Reun. Nac. Soc. Cienc. Nat.*, Buenos Aires (1919):628-638, figs.

MATTOS, W. R. B.

1919. *As sarcophagas de São Paulo* (These, Faculdade de Medicina e Cirurgia de São Paulo), iii + 116 + xii pp., 3 pls. São Paulo.

MEIGEN, W.

1803. Versuch einer neuen Gattungseintheilung der europäischen zweiflügeligen Insekten. *Mag. f. Insektenkunde* 2:259-281.

NEIVA, A. & G. FARIA

1913. Notas sobre um caso de miase humana ocasionada por larvas de *Sarcophaga pyrophila* n. sp. *Mem. Inst. Oswaldo Cruz* 5:16-23.

PANDELLÉ, L.

1896. Etudes sur les Muscides de France. IIe Partie. *Rev. d'Entom.*, Caen 15:1-219.

PARKER, H. L.

1953. Miscellaneous notes on South American dipterous parasites. *Boll. Lab. Ent. Agric. Filippo Silvestri*, Portici 12:46-73.

PARKER, R. R.

1914. Sarcophagidae of New England; males of the genera *Ravinia* and *Boettcheria*. *Proc. Boston Ent. Soc.* 35:1-77, pls. 1-8.
1917. A new *Sarcophaga* from New York. *Canad. Ent.* 49:157-161, fig. 7.
1918. A new species of *Sarcophaga* from California. *J. Ent. Zool.* 10:32-33, fig.

PRADO, A. & F. FONSECA


- 1932a. Algumas especies novas de sarcophagas da cidade de São Paulo. *Rev. Med. Cirurg. Brasil.* 40:3-7, 8 figs.
1932b. Um genero novo e algumas especies de sarcophagas (Diptera, Stephanostomatidae) da cidade de São Paulo. *Mem. Inst. Butantan* 7:161-171, 3 pls.

REINHARD, H. J.

1939. A review of the muscoid genus *Eumacronychia* with key and descriptions of new species (Diptera). *J. New York Ent. Soc.* 74:57-68.
1947. New North American muscoid Diptera. *J. Kansas Ent. Soc.* 20(3):95-116, 19 figs.; (4):117-126, figs. 20-28.
1957. New American muscoid Diptera (Sarcophagidae, Tachinidae). *Ent. News* 68:99-111.
1965a. New American Sarcophagidae. *Ent. News* 74:75-83.
1956b. Review of the miltogrammid genus *Eumacronychia* (Sarcophagidae: Diptera). *Canad. Ent.* 97(4):337-350.

RICHARDS, O. W.

1935. A note on the genus *Pachyophthalmus* B. & B. (Diptera, Tachinidae, Miltogramminae). *Stylops* 4:209-213, 2 figs.
1936. A second note on the genus *Pachyophthalmus* B. & B. (Diptera, Tachinidae). *Proc. Roy. Ent. Soc. London (B)* 5:89-90, fig. 1.



- ROBACK, S. S.
 1952. New species of Sarcophaginae (Diptera, Sarcophagidae). *J. Wash. Acad. Sci.* 42(2):45-49, figs. 1-6.
 1954. The evolution and taxonomy of the Sarcophaginae (Diptera, Sarcophagidae). *Illinois Biol. Monogr.* 23(3-4):1-181, 34 pls.
- ROBINEAU-DESVOIDY, J. B.
 1830. Essai sur les Myodaires. Inst. de France, Cl. des Sci. Math. et Phys., Acad. Roy. des Sci., *Mém. présentées par divers Savans* (ser. 2) 2:1-813.
 1863. *Histoire naturelle des diptères des environs de Paris* 1:xi + 1143 pp. Paris.
- ROHDENDORF, B. B. ..
 1937. *Fauna de l'URSS. Insectes Diptères* 19(1). *Fam. Sarcophagidae*. Inst. Zool., Acad. Sci. de l'URSS 1, 12, 501 pp., 535 figs.
- RONDANI, C.
 1848. Esame di vario specie d'insetti ditteri brasiliani. *Studi Ent.* 1:63-112.
 1850a. Osservazione sopra alquante specie di esapodi del Museo Torinese. *Nuovi Ann. Sci. Nat. Bologna* 3(2):165-197, pl. 4.
 1850b. Dipterorum species aliquae in America Aequatoriali collectae a Cajetano Osculati, observata et distincta novis breviter descriptis. *Nuovi Ann. Sci. Nat. Bologna* 3(2):357-372.
 1856. *Dipterologiae italicae prodromus* 1:1-226. Parma.
 1859. *Dipterologiae italicae prodromus* 3:243 pp. Parma.
 1864. Dipterorum species et genera aliqua revisa et annotata, novis nonnullis descriptis. *Arch. Zool. Anat. di Canestrini* 3:1-99, 5 pls.
 1867. Diptera aliqua in America Meridionalis lecta a Prof. Strobel annis 1866-1867. *Ann. Soc. Nat. Modena* 3:24-40, pl. 4.
- ROSSI, P.
 1790. *Fauna Etrusca* 2:1-348, 10 pls. Livorno.
- SALVIN, R. J.
 1958. Notas biológicas sobre la mosca *Scrvasia* (*Protodexia*) *artegai* (Blanch.) Rob. (Diptera, Sarcophagidae) parasito de la tocura. *Rev. Invest. Agric.* 12(3):299-309, 3 pls.
- SALEN, H. H.
 1935. The Egyptian species of the genus *Sarcophaga* (Diptera, Tachinidae). *Publ. Fac. Med. Egypt* 5:1-61, 19 figs.
- SCHINER, J. R.
 1868. *Reise der österreichischen Fregatte Novara um die Erde*, Zool. Theil, Diptera: vi + 388 pp., 4 pls. Wien.
- SÉGUY, E.
 1926. Sur la *Microcerella rufomaculata* Macq. *Enc. Ent.* (B, II), Diptera 2:184.
 1928. *Études sur les mouches parasites* 1:1-251, 300 figs. Paris.
 1931. Étude sur un nouveau type de diptère parasite, *Pachyophthalmus aurifacies* n. sp. *Zeit. Parasitenk.* 3:780-785, figs. 1-2.
- SILVESTRI, F.
 1903. Contribuzione alla conoscenza dei termitidi o termitofili dell'America Meridionale. *Redia* 1:1-234, pls. 1-6.

SMITH, H. E.


1916. New Tachinidae from North America. *Proc. Ent. Soc. Wash.* 18:94-98.

THOMSON, G.

1869. *Kongliga svenska fregatten Eugénies resa omkring jorden*, Pt. 2 (Zoölogie), 1 (Insekten), 6 (Dipteren):617 pp., 9 pls. Stockholm, "1868".

TOWNSEND, C. H. T.

1891. Notes on North American Tachinidae sens. str. with descriptions of new genera and species. Paper II. *Trans. Amer. Ent. Soc.* 18:349-382.
- 1892a. A dextiid parasite of a longicorn beetle. *J. Inst. Jamaica* 1:105-106.
- 1892b. Description of a *Sarcophaga* bred from *Helix*. *Psyche* 6:220-221.
- 1892c. Catalogue of the described South American species of Calyptratae Muscidae. *Ann. New York Acad. Sci.* 7:1-44.
- 1892d. A sarcophagid parasite of *Cimex americana*. *Canad. Ent.* 24:126-127.
- 1892e. Notes on North American Tachinidae, with descriptions of new genera and species. *Canad. Ent.* 24:64-70, 77-82, 165-172.
- 1892f. Notes on North American Tachinidae sens. str., with descriptions of new genera and species. Paper III. *Trans. Amer. Ent. Soc.* 19:88-132.
- 1893a. A scorpion parasite. *J. Inst. Jamaica* 1:221.
- 1893b. A tachinid reared from cells of a mud-dauber wasp. *Ohio Agric. Sta., Techn. Ser. Bull.* 1:165-166.
1908. The taxonomy of the muscoidean flies including descriptions of new genera and species. *Smiths. Misc. Coll.* 51:1-138.
- 1912a. Descriptions of new genera and species of muscoidean flies from the Andean and Pacific coast regions. *Proc. U. S. Nat. Mus.* 43:301-367.
- 1912b. Foundation of some genera and species of muscoid flies mainly on reproductive and early stage characters. *J. New York Ent. Soc.* 20:107-119.
- 1915a. Muscoidea synonymical notes. *Ins. Insc. Menstruus* 3:115-122.
- 1915b. Proposal of new muscoid genera for old species. *Proc. Biol. Soc. Wash.* 28:19-24.
- 1915c. New Neotropical muscoid flies. *Proc. U. S. Nat. Mus.* 49:405-440.
- 1915d. New Canadian and Alaskan Muscoidea. *Canad. Ent.* 47:385-392.
- 1916a. Elucidations of New England Muscoidea. *Ins. Insc. Menstruus* 4:17-32.
- 1916b. Diagnosis of new genera of muscoid flies founded on old species. *Proc. U. S. Nat. Mus.* 49:617-633.
- 1916c. New genera and species of muscoid flies. *Proc. U.S. Nat. Mus.* 51:299-323.
- 1916d. Designations of types of genera. *Ins. Insc. Menstruus* 4:4-12.
- 1917a. New genera and species of American muscoid Diptera. *Proc. Biol. Soc. Wash.* 30:43-50.
- 1917b. Genera of the dipterous tribe Sarcophagini. *Proc. Biol. Soc. Wash.* 30:189-198.



- 1917c. Second paper on Brazilian Muscoidea collected by Herbert H. Smith. *Bull. Amer. Mus. Nat. Hist.* 37:221-233.
- 1918a. New genera of Amoblinae. *Ins. Insc. Menstruus* 5:157-165.
- 1918b. Some muscoid synonymy with one new genus. *Proc. Ent. Soc. Wash.* 20:19-20.
- 1919a. New muscoid genera, species and synonymy (Diptera). *Ins. Insc. Menstruus* 6:157-182.
- 1919b. New genera and species of muscoid flies. *Proc. U.S. Nat. Mus.* 56:541-592.
- 1927a. Synopse dos generos muscoideos da região humida tropical da America, com generos e especies novas. *Rev. Museu Paulista* 15:205-385, 4 pls.
- 1927b. New Muscoid flies in the collection of the Deutsches Entomologisches Institut, in Berlin. *Ent. Mitt.* 16:277-287.
1928. New Muscoidea from humid tropical South America. *Wien. Ent. Zeit.* 44:143-154.
1929. New species of humid tropical American Muscoidea. *Rev. Chil. Hist. Nat.* 32 (1928):365-382.
- 1931a. Notes on American oestromuscoid types. *Rev. Entom.* 1:65-104.
- 1931b. New genera and species of American oestromuscoid flies. *Rev. Entom.* 1:313-354.
- 1934a. Muscoid notes and descriptions. *Rev. Entom.* 4:110-112.
- 1934b. New Neotropical oestromuscoid flies. *Rev. Entom.* 4:201-212.
- 1935a. New genera, mainly from the Neotropical region. *Rev. Entom.* 5:68-74.
- 1935b. *Manual of Myiology* 2:289 pp., 9 pls. São Paulo.
- 1937a. Two new South American oestromuscarian genera (Diptera). *Rev. Entom.* 7:115-117.
- 1937b. *Manual of Myiology* 5:234 pp. São Paulo.
1938. *Manual of Myiology* 6:242 pp. São Paulo.
- VIMMER, A. & S. S. SOUKUP
1940. Los dípteros de los alrededores de Puno (conclusion). *Bol. Mus. Hist. Nat. "Javier Prado"* 4(4):360-372.
- WALKER, F.
1837. Descriptions of insects collected by Captain P.P. King, R.N., F.R.S., in the survey of the Straits of Magellan. *Diptera. Trans. Linn. Soc. London* 17:331-359.
1849. *List of the specimens of dipterous insects in the collection of the British Museum*, 4:688-1172. London.
1852. *Insecta Saundersiana or characters of undescribed insects in the collections of W. W. Saunders* 4:253-414.
1857. Characters of undescribed Diptera in the collection of W. W. Saunders. *Trans. Ent. Soc. London* (2) 4:119-158, 190-235.
1861. Characters of undescribed Diptera in the collection of W. W. Saunders. *Trans. Amer. Ent. Soc. London* (2) 5:297-334.
- WEYENBERGH, H.
1875. Animales utiles. *Nemorea acridiorum* Weyenb. *An. Agric. Rep. Argentina* 3(9):85-86, 7 figs.
- WIEDEMANN, C. R. W.
1830. *Aussereuropäische zweiflügelige Insekten* 2:xli + 684 pp., 5 pls.

WILLIAMS, R. M.

1958. A study of the summer filth fly populations of the Bermuda Islands. *J. Paras.* 44:339-342.

WILLISTON, S. W.

1896. On the Diptera of St. Vincent (West Indies). *Trans. Ent. Soc. London* 1896 (3):253-446, pls. 8-14.

WULP, F. M. VAN DER

1887. *Sarcophagula*, een nieuw geslacht der Sarcophaginae. *Tijdsch. v. Ent.* 30:173-174, figs. 8-9.

1888-1903. Fam. Muscidae, pp. 2-40 (1888), 41-208 (1890), 209-264 (1891), 265-272 (1895), 273-244 (1896), 345-376 (1897), 377-384 (1898), 385-416 (1899), 417-428 (1900), 429-489 (1903), in F. D. Godman & O. Salvin. *Biologia Centrali-Americana, Zoologia-Insecta-Diptera* 2:489 pp., 11 figs., 13 pls. London.

1892. Diagnosis of new Mexican Muscidae. *Tijdsch. v. Ent.* 35:183-195.

ZETTERSTEDT, J. W.


1844. *Diptera Scandinaviae disposita et descripta* 4:1281-1738. Lund.

INDEX


Abacantha Hall, 1938	103.16
Abapa Dodge, 1965	103.39
abdominalls Dodge, 1965, Tulaeopoda	103.48
abnormalls (Hall), 1937 (Sarcophaga), Pecklamyla	103.38
abnormis Enderlein, 1928, Adiseochaeta	103.38
abrupta Lopes, 1955, Euboetteheria	103.34
Acanthodotheca Townsend, 1918	103.19
acanthoptera (Wulp), 1895 (Sarcophaga), Rafaelia	103.33
Acanthotheca Roback, 1954	103.48
aeurata Hall, 1933 (Sarcophaga)	103.51
Aeridlophaga Townsend, 1917	103.18
aeridlophagoides Lopes & Downs, 1951, Acanthodotheca	103.19
aeridlorum (Weyenbergh), 1875 (Nemorea), Doringia	103.9
acridiorum (Weyenbergh) of Conil, 1879 (Nemorea), Aeridiophaga	103.18
Actenella Enderlein, 1928	103.11
adamsi (Hall), 1928 (Sarcophaga), Chaetoravinla	103.23
addentata (Hall), 1929 (Sarcophaga), Ravinia	103.21
Adiseochaeta Enderlein, 1928	103.38
admixta (Lopes), 1933 (Sarcophaga), Oxysareodexia	103.24
adolenda (Lopes), 1935 (Sarcophaga), Paraphrissopoda	103.36
adunca (Hall), 1933 (Sarcophaga), Dexosarcophaga	103.42
advena Walker, 1852 (Sarcophaga)	103.52
aenigma (Reinhard), 1947 (Sarcophaga), Protodexia	103.18
aequata (Wulp), 1895 (Sarcophaga), Euboetteheria	103.34
affleta (Wulp), 1895 (Sarcophaga), Oxysareodexia	103.24
afflicta (Wulp) of Enderlein, 1928 (Sarcophaga), Euboetteheria	103.35
affinis Lopes, 1936, Nephochaetopteryx	103.28
affinis Lopes, 1948, Sarconelva	103.48
Agriochoeta Enderlein, 1928	103.14
Agriochoetops Townsend, 1934	103.14

103.67

<i>Airypei</i> Dodge, 1965	103.47
<i>Alaccoprosopa</i> Townsend, 1934	103.9
<i>alata</i> (Aldrich), 1916 (Sarcophaga), <i>Titanogrypa</i>	103.48
<i>albicans</i> (Lahille), 1907 (Sarcophaga), <i>Lopesimyia</i>	103.44
<i>albicans</i> (Walker), 1858 (?Dexia), <i>Tricharaca</i>	103.31
<i>albicans</i> (Wiedemann), 1830 (Sarcophaga), <i>Tripanurga</i>	103.46
<i>albida</i> Lopes, 1951, <i>Harpagopyga</i>	103.14
<i>albihirta</i> Dodge, 1965, <i>Harpagopyga</i>	103.14
<i>albisignum</i> Aldrich, 1916 (Sarcophaga)	103.51
<i>alcedo</i> Aldrich, 1916, <i>Acanthodotheca</i>	103.19
<i>aldrichi</i> (Hall), 1937 (Phacosarcodexia), <i>Austrohartigia</i>	103.8
<i>aldrichi</i> Lopes, 1942, <i>Notochaeta</i>	103.11
<i>alcida</i> Lopes, 1951, <i>Pachygraphiops</i>	103.14
<i>alcuaphaga</i> Blanchard, 1942, <i>Protodexia</i>	103.17
<i>Alienancia</i> Townsend, 1935	103.4
<i>almeidai</i> (Lopes), 1938 (Sarcophaga), <i>Dexosarcophaga</i>	103.41
<i>almeidai</i> Lopes, 1946, <i>Chactoravinia</i>	103.22
<i>alpina</i> Townsend, 1912, <i>Euparaphyto</i>	103.8
<i>amarali</i> Prado & Fonseca, 1932, <i>Oxysarcodexia</i>	103.27
<i>amazonica</i> Townsend, 1934, <i>Eunotochacta</i>	103.13
<i>Amblycoryphenes</i> Townsend, 1918	103.18
<i>americanus</i> (Brauer & Bergenstamm), 1891 (<i>Arrenopus</i>), <i>Sco-</i> <i>tainia</i>	103.5
<i>Amesothyrus</i> Enderlein, 1928	103.23
<i>Amobia</i> Robincau-Desvoidy, 1830	103.2
<i>Amobiopsis</i> Townsend, 1915	103.6
<i>amocna</i> (Aldrich), 1916 (Sarcophaga), <i>Paraphrissopoda</i>	103.36
<i>amorosa</i> (Schiner), 1868 (Sarcophaga), <i>Oxysarcodexia</i>	103.24
<i>ampullula</i> (Engel), 1931 (Bercaca), <i>Sarcodexiopsis</i>	103.49
<i>analis</i> Townsend, 1927, <i>Xanthobrachycoma</i>	103.9
<i>Anaravinia</i> Townsend, 1934	103.39
<i>anatina</i> Lopes, 1953, <i>Euboeetcheria</i>	103.34
<i>andina</i> Townsend, 1912, <i>Chloronesia</i>	103.45
<i>Andinoravinia</i> Townsend, 1917	103.22
<i>angolensis</i> (Hall), 1937 (Sarcophaga), <i>Oxysarcodexia</i>	103.24
<i>angrensis</i> (Lopes), 1933 (Sarcophaga), <i>Oxysarcodexia</i>	103.24
<i>angulla</i> (Curran & Walley), 1934 (Sarcophaga), <i>Euboeetcheria</i> ..	103.34
<i>angustifrons</i> (Aldrich), 1916 (Sarcophaga), <i>Acridiophaga</i>	103.18
<i>angustifrons</i> Lopes, 1942, <i>Nephochactopteryx</i>	103.28
<i>angustiventris</i> Curran & Walley, 1934, <i>Harpagopyga</i>	103.14
<i>Anicia</i> Robincau-Desvoidy, 1863	103.4
<i>anitsiana</i> (Enderlein), 1928 (<i>Cricobrachia</i>), <i>Sarcodexia</i>	103.34
<i>annae</i> Dodge, 1965, <i>Discoheilecobia</i>	103.32
<i>Anolisimyia</i> Dodge, 1955	103.11
<i>antennata</i> (Dodge), 1966 (<i>Petriania</i>), <i>Lcpidodexia</i>	103.15
<i>antillarum</i> (Richards), 1935 (<i>Pachyophthalmus</i>), <i>Amobia</i>	103.2
<i>anzia</i> (Walker), 1849 (Sarcophaga), <i>Ravinia</i>	103.21
<i>Apelophyla</i> Hall, 1938	103.24
<i>Aphelomyia</i> Roback, 1954	103.49
<i>apicalis</i> Townsend, 1934, <i>Alaccoprosopa</i>	103.9
<i>apolinari</i> (Lopes), 1938 (Sarcophaga), <i>Udamopyga</i>	103.44
<i>apolinari</i> Lopes, 1951, <i>Lcpidodexia</i>	103.15
<i>Arabiopsis</i> Townsend, 1915	103.5
<i>AArachnidomyia</i> Townsend, 1934	103.42



<i>Arachnidomyia</i> Townsend, 1934	103.42
<i>aragua</i> Dodge, 1966, <i>Chloronesia</i>	103.45
<i>argentea</i> (Townsend), 1912 (<i>Sarcophaga</i>), <i>Argoravinia</i>	103.46
<i>argenteum</i> (Prado & Fonseca), 1932 (<i>Stephanostoma</i>), <i>Townsendimyia</i>	103.9
<i>argentifrons</i> (Townsend), 1891 (<i>Miltogramma</i>), <i>Senotainia</i>	103.5
<i>argentina</i> (Brèthes), 1916 (<i>Sarcophaga</i>), <i>Parasarcophaga</i>	103.50
<i>Argoravinia</i> Townsend, 1917	103.46
<i>argyrocephala</i> (Meigen), 1803 (<i>Tachina</i>), <i>Metopia</i>	103.3
<i>argyrostoma</i> (Robineau-Desvoidy), 1830 (<i>Myophora</i>), <i>Parasarcophaga</i>	103.50
<i>aristata</i> (Aldrich), 1916 (<i>Camptopyga</i>), <i>Xenoppia</i>	103.10
<i>aristata</i> (Aldrich) of Engel, 1931 (<i>Camptopyga</i>), <i>Scopaedisus</i> ..	103.10
<i>arizonica</i> (Townsend), 1919 (<i>Petrosarcophaga</i>), <i>Wohlfahrtiopsis</i> ..	103.51
<i>armilla</i> Reinhard, 1965, <i>Eumaeronychia</i>	103.3
<i>arnaudi</i> Lopes, 1950, <i>Boettcheria</i>	103.42
<i>artegal</i> (Blanchard), 1939 (<i>Opsophyto</i>), <i>Aeridiophaga</i>	103.18
<i>articulata</i> Hall, 1933 (<i>Sarcophaga</i>)	103.52
<i>artigas</i> Dodge, 1966, <i>Oxysarcodexia</i>	103.24
<i>asiliformis</i> Townsend, 1927	103.16
<i>Asilodexia</i> Townsend, 1927	103.16
<i>asinoma</i> Hall, 1938, <i>Euboettcheria</i>	103.34
<i>assidua</i> (Walker) of Aldrich, 1916 (<i>Sarcophaga</i>), <i>Oxysarcodexia</i> ..	103.27
<i>atrata</i> Dodge, 1965, <i>Harpagopyga</i>	103.14
<i>augusta</i> Aldrich, 1925, <i>Notochaeta</i>	103.11
<i>augusta</i> Lopes, 1946, <i>Oxysarcodexia</i>	103.24
<i>Aulacophyto</i> Townsend, 1919	103.7
<i>aura</i> (Hall), 1937 (<i>Sarcophaga</i>), <i>Hybopygia</i>	103.23
<i>aurata</i> (Enderlein), 1928 (<i>Desvoidyella</i>), <i>Sarcophagula</i>	103.29
<i>aurata</i> Townsend, 1927, <i>Discomyiophora</i>	103.13
<i>aurea</i> Townsend, 1918, <i>Ravinioipsis</i>	103.46
<i>aurea</i> Townsend, 1934, <i>Eufletcherimyia</i>	103.13
<i>aureiceps</i> (Macquart), 1855 (<i>Sarcophaga</i>), <i>Oxysarcodexia</i>	103.25
<i>aureopyga</i> (Hall), 1928 (<i>Sarcophaga</i>), <i>Ravinia</i>	103.21
<i>aureostrigata</i> (Enderlein), 1928 (<i>Agriocheata</i>), <i>Gymnocamptops</i> ..	103.14
<i>aureseens</i> (Townsend), 1927 (<i>Helicobiopsis</i>), <i>Helicobia</i>	103.39
<i>auribarbata</i> (Townsend), 1912 (<i>Sarcophaga</i>), <i>Paraphrissopoda</i> ..	103.36
<i>auricauda</i> (Enderlein), 1928 (<i>Dasyproctia</i>), <i>Oxysarcodexia</i>	103.27
<i>aurifacies</i> (Séguy), 1931 (<i>Paehyophthalmus</i>), <i>Amobia</i>	103.2
<i>aurifera</i> Lopes, 1950, <i>Boettcheria</i>	103.42
<i>aurifinis</i> (Walker), 1852 (<i>Sarcophaga</i>), <i>Oxysarcodexia</i>	103.25
<i>aurifrons</i> (Hall), 1937 (<i>Maeronychia</i>), <i>Itamobia</i>	103.6
<i>aurigena</i> (Townsend), 1912 (<i>Sarcophaga</i>), <i>Ravinia</i>	103.21
<i>aurigena</i> (Townsend) of Engel, 1931 (<i>Trioxosarcophaga</i>), <i>Oxysarcodexia</i>	103.25
<i>aurimaculata</i> Vimmer & Soukoup, 1940 (<i>Sarcophaga</i>)	103.52
<i>auromaculata</i> Townsend, 1915, <i>Dolichamobia</i>	103.6
<i>auromaculata</i> (Townsend), 1915 (<i>Punasarcophaga</i>), <i>Ravinia</i>	103.21
<i>auromaculata</i> Townsend, 1919, <i>Aulacophyto</i>	103.7
<i>auromaculata</i> Townsend, 1919, <i>Euparaphyto</i>	103.8
<i>australis</i> (Aldrich), 1916 (<i>Sarcophaga</i>), <i>Oxysarcodexia</i>	103.26
<i>australis</i> Blanchard, 1942, <i>Protodexia</i>	103.17



<i>australis</i> Townsend, 1927, <i>Euboettcheria</i>	103.34
<i>australis</i> (Townsend), 1934 (<i>Euleucomyia</i>), <i>Tricharaea</i>	103.31
<i>Austrohartigia</i> Townsend, 1937	103.8
<i>avuncula</i> (Lopes), 1933 (<i>Sarcophaga</i>), <i>Oxysarcodexia</i>	103.25
<i>azteca</i> Lopes & Downs, 1951, <i>Acanthodotheca</i>	103.19
<i>azurea</i> (Enderlein), 1928 (<i>Robineanella</i>), <i>Chlorosarcophaga</i>	103.45
<i>Bahamiola</i> Dodge, 1965	103.30
<i>bakeri</i> (Aldrich), 1916 (<i>Sarcophaga</i>), <i>Oxysarcodexia</i>	103.25
<i>banksi</i> Dodge, 1966, <i>Emdenimyia</i> (only in key)	103.17
<i>barbata</i> (Thomson), 1869 (<i>Sarcophaga</i>), <i>Parasarcophaga</i>	103.50
<i>barroi</i> Dodge, 1956, <i>Chaetoravinia</i>	103.22
<i>bartica</i> (Curran & Walley), 1934 (<i>Sarcophaga</i>), <i>Udamopyga</i>	103.44
<i>batemani</i> Dodge, 1965, <i>Udamopyga</i>	103.44
<i>baumhoveri</i> Dodge, 1965, <i>Sarcophagrtiopsis</i>	103.30
<i>beeri</i> (Dodge), 1966 (<i>Sarcophaga</i>), <i>Wohlfahrtiopsis</i>	103.51
<i>belfortii</i> (Prado & Fonseca), 1932 (<i>Euravinia</i>), <i>Ravinia</i>	103.21
<i>bellula</i> Dodge, 1965, <i>Abapa</i>	103.39
<i>Bercaea</i> Robineau-Desvoidy, 1863	103.50
<i>bergi</i> (Blanchard), 1939 (<i>Chrysostomia</i>), <i>Paraphrissopoda</i>	103.36
<i>bethae</i> Dodge, 1965, <i>Helicobia</i>	103.40
<i>bicolor</i> Dodge, 1967, <i>Chlorosarcophaga</i>	103.45
<i>bicolor</i> Lopes, 1946, <i>Oxysarcodexia</i>	103.25
<i>bicoloricauda</i> (Enderlein), 1928 (<i>Mesothyrsla</i>), <i>Xanthobrachycoma</i>	103.9
<i>bidentata</i> Dodge, 1966, <i>Dexosarcophaga</i>	103.41
<i>bifrons</i> Walker, 1852 (<i>Sarcophaga</i>)	103.51
<i>bifurcata</i> (Dodge), 1965 (Peckla), <i>Euboettcheria</i>	103.34
<i>blkini</i> Dodge, 1966, <i>Oxysarcodexia</i>	103.25
<i>blinnata</i> Dodge, 1966, <i>Tricharaea</i>	103.31
<i>biplagiata</i> Dodge, 1966, <i>Helicobia</i>	103.40
<i>biseriata</i> (Aldrich), 1916 (<i>Sarcophaga</i>), <i>Sarcodexiopsis</i>	103.49
<i>biseriata</i> Dodge, 1966, <i>Emdenimyia</i>	103.17
<i>biseta</i> (Thomson), 1869 (<i>Miltogramma</i>), <i>Panamaphyto</i>	103.7
<i>bishoppi</i> (Aldrich), 1916 (<i>Sarcophaga</i>), <i>Wohlfahrtiopsis</i>	103.51
<i>bivittata</i> Curran, 1928, <i>Johnsonia</i>	103.16
<i>blakeae</i> (Dodge), 1955 (<i>Anolisimyia</i>), <i>Notochaeta</i>	103.11
<i>blandita</i> (Brèthes), 1920 (<i>Sarcophaga</i>), <i>Mantidophaga</i>	103.18
<i>blodgeti</i> Dodge, 1964, <i>Socorromyia</i>	103.38
<i>Boetia</i> Dodge, 1965	103.43
<i>Boettcheria</i> Parker, 1914	103.42
<i>Boettcherlimina</i> Lopes, 1950	103.17
<i>bogotana</i> (Enderlein), 1928 (<i>Udamoctis</i>), <i>Notochaeta</i>	103.11
<i>boniato</i> Dodge, 1965, <i>Udamopyga</i>	103.41
<i>bonplandi</i> Hall, 1937 (<i>Sarcophaga</i>)	103.53
<i>borgmeieri</i> Lopes, 1939, <i>Helicobia</i>	103.40
<i>borregoensis</i> Reinhard, 1965, <i>Eumacronychia</i>	103.3
<i>bosqui</i> Blanchard, 1939 (<i>Paraphrissopoda</i>)	103.52
<i>boxi</i> Hall, 1938, <i>Acanthodotheca</i>	103.20
<i>brasilliana</i> Townsend, 1929, <i>Anicia</i>	103.4
<i>brasillensis</i> Robineau-Desvoidy, 1830 (<i>Myophora</i>)	103.53
<i>brasillensis</i> (Townsend), 1927 (<i>Pseudosarcophaga</i>), <i>Sarcophaga</i>	
<i>gula</i>	103.29

<i>brasiliensis</i> (Townsend), 1929 (<i>Euselenomyia</i>), <i>Senotainta</i>	103.5
<i>brethesi</i> Blanchard, 1938, <i>Neobellieria</i>	103.51
<i>brevicornis</i> (Wiedemann), 1830 (<i>Sarcophaga</i>), <i>Tricharaea</i>	103.31
<i>brevigaster</i> Lopes, 1945, <i>Geijskesia</i>	103.13
<i>brevirostris</i> Lopes, 1945, <i>Petria</i>	103.16
<i>brevispina</i> (Thomson), 1869 (<i>Sarcophaga</i>), <i>Sarothromyia</i>	103.31
<i>brullei</i> Macquart, 1835 (<i>Phrissopoda</i>)	103.53
<i>buethni</i> Dodge, 1965, <i>Euboettcheria</i>	103.34
<i>bufonivora</i> Lopes & Vogelsang, 1953, <i>Notochaeta</i>	103.11
<i>butantani</i> (Prado & Fonseca), 1932 (<i>Ctenoprosballia</i>), <i>Sarcodexia</i>	103.34
<i>calceata</i> (Dodge), 1967 (<i>Winthemiola</i>), <i>Boettcheria</i>	103.42
<i>calida</i> (Wiedemann), 1830 (<i>Sarcophaga</i>), <i>Nephochaetopteryx</i> ...	103.28
<i>callifornica</i> (Parker), 1918 (<i>Sarcophaga</i>), <i>Kurtomyia</i>	103.20
<i>calliphorina</i> (Enderlein), 1928 (<i>Robineanella</i>), <i>Chlorosarcophaga</i>	103.45
<i>calx</i> (Hall), 1933 (<i>Sarcophaga</i>), <i>Peckiamyia</i>	103.38
<i>camura</i> Hall, 1938 (<i>Sarcophaga</i>)	103.52
<i>Camptopsis</i> Townsend, 1918	103.16
<i>Camptopyga</i> Aldrich, 1916	103.10
<i>cana</i> (Townsend), 1917 (<i>Prosthetocirca</i>), <i>Sarothromylops</i>	103.10
<i>candida</i> (Curran), 1928 (<i>Sarcophagina</i>), <i>Ravinioptis</i>	103.46
<i>cancscens</i> (Thomson), 1869 (<i>Sarcophaga</i>), <i>Tricharaea</i>	103.31
<i>canuta</i> Wulp, 1896, <i>Sarcophagula</i>	103.29
<i>capitata</i> (Aldrich), 1916 (<i>Sarcophaga</i>), <i>Paraphrissopoda</i>	103.36
<i>capitata</i> (Curran), 1928 (<i>Sarcophagina</i>), <i>Sarcophagtiopsis</i>	103.30
<i>caridei</i> (Br��thes), 1906 (<i>Sarcophaga</i>), <i>Acridiophaga</i>	103.18
<i>Carinoclypeus</i> Dodge, 1965	103.49
<i>carolae</i> Dodge, 1965, <i>Tulaepoda</i>	103.48
<i>carvalhoi</i> Lopes, 1946, <i>Oxysarcodexia</i>	103.25
<i>cassidifera</i> Engel, 1931 (<i>Sarcophaga</i>)	103.52
<i>catalina</i> Parker, 1921, <i>Sabinata</i>	103.46
<i>Cataphyto</i> Townsend, 1935	103.30
<i>Catheteronychia</i> Townsend, 1927	103.10
<i>Cattasoma</i> Reinhard, 1947	103.6
<i>cavagnaroi</i> Dodge, 1965, <i>Tolucamyia</i>	103.49
<i>cayennensis</i> Macquart, 1843 (<i>Sarcophaga</i>)	103.53
<i>cessator</i> (Aldrich), 1916 (<i>Sarcophaga</i>), <i>Kurtomyia</i>	103.20
<i>chacoana</i> (Blanchard), 1942 (<i>Oxyhelicobia</i>), <i>Helicobia</i>	103.40
<i>chaetopygialis</i> (Williston), 1896 (<i>Sarcophaga</i>), <i>Oxysarcodexia</i> ...	103.25
<i>Chaetoravinia</i> Townsend, 1917	103.22
<i>chaetosa</i> Blanchard, 1939, <i>Dexosarcophaga</i>	103.41
<i>chaetosa</i> Townsend, 1927, <i>Catheteronychia</i>	103.10
<i>chaetosa</i> Townsend, 1931, <i>Cryptosarcophila</i>	103.30
<i>Chapiniola</i> Dodge, 1965	103.11
<i>chilensis</i> (Enderlein), 1928 (<i>Amesothyrus</i>), <i>Hybopygia</i>	103.23
<i>chilensis</i> (Enderlein) of Engel, 1931 (<i>Amesothyrus</i>), <i>Hypopygia</i>	103.9
<i>chilensis</i> (Hall), 1937 (<i>Mimophytomyia</i>), <i>Alaccoprosopa</i>	103.24
<i>chilensis</i> (Macquart), 1843 (<i>Sarcophaga</i>), <i>Hybopygia</i>	103.52
<i>chirotheca</i> Hall, 1933 (<i>Sarcophaga</i>)	103.45
<i>Chloronesia</i> Townsend, 1912	103.45
<i>Chlorosarcophaga</i> Townsend, 1919	103.10
<i>Chrysagria</i> Townsend, 1935	

<i>chrysella</i> Robineau-Desvoidy, 1830 (Myophora)	103.53
<i>chrysostoma</i> (Wiedemann), 1830 (Sarcophaga), Paraphrissopoda	103.36
<i>Chrysostomyia</i> Townsend, 1931	103.36
<i>chrystomata</i> (Hall), 1933 (Sarcophaga), Paraphrissopoda	103.36
<i>erysura</i> Rondani, 1864 (Sarcophaga)	103.53
<i>elhubutina</i> Blanchard, 1955, Neotephromyiella	103.21
<i>Cicatriella</i> Dodge, 1965	103.20
<i>cimbicis</i> (Townsend), 1892 (Sarcophaga), Boettcheria	103.42
<i>einctus</i> Townsend, 1917, Sarcophromyiops	103.10
<i>cinerascens</i> (Townsend), 1891 (Miltogramma), Senotainia	103.5
<i>cingulata</i> Lopes, 1962, Lepidodexia	103.15
<i>elcumelsa</i> Rondani, 1848 (Sarcophaga)	103.53
<i>clotho</i> Brèthes, 1920 (Sarcophaga)	103.53
<i>clotho</i> (Curran & Walley), 1934 (Sarcophaga), Paraphrissopoda ..	103.36
<i>cochliomyia</i> Townsend, 1919, Chlorosarcophaga	103.45
<i>cognata</i> (Walker), 1852 (Sarcophaga), Notochaeta	103.11
<i>collusor</i> (Curran & Walley), 1934 (Sarcophaga), Euboettcheria ...	103.35
<i>columblana</i> Lopes, 1962, Andinoravina	103.22
<i>comata</i> Aldrich, 1925, Notochaeta	103.12
<i>complicata</i> (Hall), 1937 (Sarcophaga), Hybobygia	103.23
<i>compta</i> (Wiedemann) of Matos, 1919 (Sarcophaga), Oxysarcodexia ..	103.25
<i>comta</i> Wiedemann, 1830, (Sarcophaga)	103.53
<i>conclnata</i> (Williston), 1896 (Sarcophaga), Paraphrissopoda	103.36
<i>conclausa</i> Walker, 1861 (Sarcophaga)	103.53
<i>confundens</i> (Townsend), 1915 (Amobiopsis), Macronychia	103.6
<i>confusa</i> Lopes, 1946, Notochaeta	103.12
<i>confusa</i> Lopes, 1946, Oxysarcodexia	103.25
<i>conjugens</i> (Wulp), 1895 (Sarcophaga), Chaetoravina	103.23
<i>contermina</i> Walker, 1852 (Sarcophaga)	103.53
<i>corolla</i> Dodge, 1965, Oxysarcodexia	103.25
<i>costalis</i> (Townsend), 1927 (Micronotochaeta), Notochaeta	103.12
<i>cotyledonca</i> (Aldrich), 1916 (Sarcophaga), Pattonella	103.38
<i>coval</i> (Dodge), 1965 (Boetia), Ravina	103.21
<i>cralgi</i> (Dodge), 1964 (Peckia), Paraphrissopoda	103.36
<i>crassipalpis</i> (Macquart), 1839 (Sarcophaga), Parasarcophaga	103.50
<i>creameri</i> Dodge, 1965, Carinoclypeus	103.49
<i>Cricobrachia</i> Enderlein, 1928	103.33
<i>crispina</i> (Lopes), 1938 (Sarcophaga), Lipoptilocnema	103.43
<i>erispula</i> (Lopes), 1938 (Sarcophaga), Lipoptilocnema	103.43
<i>crouzelli</i> Blanchard, 1942, Protodexia	103.17
<i>cryptopyga</i> (Lopes), 1956 (Titanogrypa), Alrypel	103.47
<i>Cryptosarcophila</i> Townsend, 1931	103.30
<i>Ctenolioproctia</i> Enderlein, 1928	103.34
<i>Ctenoprosballia</i> Enderlein, 1928	103.34
<i>cubana</i> Lopes, 1940, Udamopyga	103.44
<i>cubana</i> Lopes, 1950, Boettcheria	103.42
<i>cubana</i> Lopes, 1951, Harpagopyga	103.14
<i>eubensis</i> (Dodge), 1965 (Peckia), Euboettcheria	103.35
<i>eubensis</i> Macquart, 1843 (Sarcophaga)	103.53
<i>eubensis</i> Robineau-Desvoidy, 1830, (Myophora)	103.53
<i>Cuculomyia</i> Roback, 1954	103.47
<i>cuernavaca</i> Dodge, 1966, Oxysarcodexia	103.25



<i>culminata</i> (Aldrich), 1916 (Sarcophaga), <i>Oxysarcodexia</i>	103.25
<i>culminiforceps</i> Dodge, 1966, <i>Oxysarcodexia</i>	103.25
<i>cuneata</i> (Townsend), 1935 (Cataphyto), <i>Sarcofahrtiopsis</i>	103.30
<i>curiosa</i> Dodge, 1965, <i>Boetia</i>	103.43
<i>currani</i> (Hall), 1930 (Sarcophaga), <i>Abapa</i>	103.39
<i>currani</i> Lopes, 1951, <i>Lepidodexia</i>	103.15
<i>cyaneiventris</i> Lopes, 1936, <i>Nephochaetopteryx</i>	103.28
<i>cyanciventris</i> Lopes, 1946, <i>Notochaeta</i>	103.12
<i>cyaniforceps</i> (Hall), 1933 (Sarcophaga), <i>Oxysarcodexia</i>	103.25
<i>dampfi</i> (Hali), 1937 (Sarcophaga), <i>Acanthodotheca</i>	103.20
<i>dampfi</i> Lopes, 1946, <i>Chaetoravinia</i>	103.22
<i>dangeloi</i> Blanchard, 1938, <i>Pattonella</i>	103.38
<i>Dasyproctia</i> Enderlein, 1928	103.24
<i>davidsoni</i> (Coquillett), 1892 (Sarcophaga), <i>Arachnidomyia</i>	103.42
<i>deaurata</i> (Blanchard), 1942 (Neosarcodexia), <i>Euboettcheria</i>	103.35
<i>deaurata</i> Blanchard, 1942, <i>Protodexia</i>	103.17
<i>debilis</i> (Wuip), 1895 (Sarcophaga), <i>Helicobia</i>	103.40
<i>decedens</i> Walker, 1857 (Sarcophaga)	103.53
<i>decens</i> Townsend, 1892, <i>Eumacronychia</i>	103.3
<i>decisa</i> (Townsend), 1892 (Mittogramma), <i>Senotainia</i>	103.5
<i>declinata</i> Hali, 1933, (Sarcophaga)	103.52
<i>aelecta</i> Wulp, 1895 (Sarcophaga)	103.52
<i>delpontei</i> Blanchard, 1939, <i>Oxysarcodexia</i>	103.27
<i>denieri</i> Blanchard, 1939, <i>Acanthodotheca</i>	103.19
<i>dentata</i> Dodge, 1966, <i>Boettcheria</i>	103.43
<i>dentifera</i> (Hali), 1937 (Sarcophaga), <i>Euboettcheria</i>	103.35
<i>dereiicta</i> (Walker), 1852 (Sarcophaga), <i>Chaetoravinia</i>	103.22
<i>despecta</i> (Lopes), 1935 (Sarcophaga), <i>Cuculomyia</i>	103.47
<i>despecta</i> (Thomson), 1869 (Sarcophaga), <i>Sarcophagula</i>	103.29
<i>despensa</i> Walker, 1861 (Sarcophaga)	103.53
<i>Desvoidyella</i> Enderlein, 1928	103.29
<i>Dexomyiophora</i> Townsend, 1927	103.32
<i>Dexosarcophaga</i> Townsend, 1917	103.41
<i>diana</i> (Lopes), 1933 (Sarcophaga), <i>Oxysarcodexia</i>	103.25
<i>diembroma</i> Dodge, 1965, <i>Sarcofahrtiopsis</i>	103.30
<i>Dienchaeta</i> Enderlein, 1928	103.39
<i>difficilis</i> Dodge, 1966, <i>Villegasia</i>	103.48
<i>dimidiata</i> (Wiedemann), 1830 (Sarcophaga), <i>Notochaeta</i>	103.12
<i>diminuta</i> (Blanchard), 1942 (Cricobrachia), <i>Sarcodexia</i>	103.34
<i>diponthi</i> Blanchard, 1942, <i>Protodexia</i>	103.18
<i>Discohelicobia</i> Dodge, 1965	103.32
<i>Discomyiophora</i> Townsend, 1927	103.13
<i>dissidia</i> (Parker), 1917 (Sarcophaga), <i>Neobellieria</i>	103.51
<i>dissimilis</i> Dodge, 1965, <i>Harpagopyga</i>	103.14
<i>distincta</i> Lopes, 1947, <i>Notochaeta</i>	103.12
<i>distincta</i> Lopes, 1951, <i>Lepidodexia</i>	103.15
<i>divergens</i> Curran & Walley, 1934, <i>Harpagopyga</i>	103.14
<i>diversa</i> Lopes, 1940, <i>Udamopyga</i>	103.44
<i>diversa</i> Lopes, 1946, <i>Notochaeta</i>	103.12
<i>diversimacuiata</i> Macquart, 1818 (Sarcophaga)	103.53
<i>diversinervis</i> (Wulp), 1895 (Sarcophaga), <i>Notochaeta</i>	103.12
<i>diversipes</i> (Coquillett), 1900 (Sarcophaga), <i>Harpagopyga</i>	103.14



<i>diversipes</i> Coquillett of Aldrich, 1916, Harpagopyga	103.14
<i>diversipes</i> Coquillett of Lopes, 1951, Harpagopyga	103.14
<i>Doliehamobia</i> Townsend, 1915	103.6
<i>dolosa</i> (Dodge), 1966 (<i>Blaesoxipha</i>), <i>Aeridiophaga</i>	103.18
<i>Doringia</i> Weyenbergh, 1875	103.9
<i>dorisae</i> Dodge, 1965, <i>Oxysarcodexia</i>	103.25
<i>downesi</i> Dodge, 1965, Harpagopyga	103.14
<i>downsiana</i> Dodge, 1963 (<i>Sarcophaga</i>)	103.52
<i>duca</i> Hall, 1938, <i>Eumierocerella</i>	103.8
<i>duodecimpunctata</i> Townsend, 1935, <i>Chrysagria</i>	103.10
<i>dupleata</i> Reinhard, 1944, <i>Eumacronychia</i>	102.3
<i>ebi</i> (Hall), 1937 (<i>Sarcophaga</i>), <i>Aeridiophaga</i>	103.19
<i>ecuatorialiana</i> Lopes, 1958, <i>Paraphrissopoda</i>	102.36
<i>ednae</i> Dodge, 1965 (<i>Sarcophaga</i>)	103.53
<i>edwardsi</i> (Hall), 1937 (<i>Phaeosarcodexia</i>), <i>Austrohartigia</i>	103.8
<i>edwardsi</i> (Hall), 1937 (<i>Sarcophaga</i>), <i>Helleobia</i>	103.49
<i>edwardsi</i> Lopes, 1946, <i>Oxysarcodexia</i>	103.26
<i>effrenata</i> (Walker), 1861 (<i>Sarcophaga</i>), <i>Chaetoravinia</i>	103.22
<i>egregia</i> (Hall), 1933 (<i>Sarcophaga</i>), <i>Dexosarcophaga</i>	103.41
<i>elaborata</i> Hall, 1933 (<i>Sarcophaga</i>)	103.52
<i>elegans</i> Aldrich, 1929, <i>Glutoxys</i>	103.32
<i>elegans</i> (Blanchard), 1939 (<i>Sarcophaga</i>), <i>Nephoehaetopteryx</i>	103.28
<i>elegans</i> Coquillett, 1895, <i>Johnsonia</i>	103.16
<i>elegans</i> (Lopes), 1938 (<i>Shannomyiella</i>), <i>Asilodexia</i>	103.16
<i>Eleodimyia</i> Townsend, 1918	102.19
<i>eleutherae</i> (Dodge), 1965 (<i>Sarcophaga</i>), <i>Abapa</i>	103.39
<i>Emblemasoma</i> Aldrich, 1916	103.33
<i>Emdenimyia</i> Lopes, 1946	103.17
<i>enderleini</i> (Engel), 1931 (<i>Adiscochaeta</i>), <i>Paraphrissopoda</i>	103.36
<i>enderleini</i> Townsend, 1913, <i>Dienechaeta</i>	103.39
<i>enderleini</i> (Townsend), 1931 (<i>Ctenoprosballia</i>), <i>Euboettcheria</i> ..	103.35
<i>engeli</i> (Dodge), 1966 (<i>Tricharea</i>), <i>Sarothromyia</i>	103.39
<i>engeli</i> Hall, 1937, <i>Itiophyto</i>	102.8
<i>Engelina</i> Enderlein, 1928	103.22
<i>epimella</i> (Lopes), 1938 (<i>Sarcophaga</i>), <i>Euboettcheria</i>	103.25
<i>errabunda</i> (Wulp), 1895 (<i>Sarcophaga</i>), <i>Chaetoravinia</i>	103.23
<i>erro</i> Aldrich, 1916, <i>Emblemasoma</i>	103.33
<i>erythropysa</i> (Lopes), 1936 (<i>Xenoplia</i>), <i>Doringia</i>	103.9
<i>erythrura</i> (Wulp), 1890 (<i>Miltogramma</i>), <i>Amobia</i>	103.2
<i>Euaraba</i> Townsend, 1915	103.5
<i>Euboettcheria</i> Townsend, 1927	103.34
<i>Eufletcherimyia</i> Townsend, 1934	103.12
<i>Euleucomyia</i> Townsend, 1934	103.31
<i>Eumaeronychia</i> Townsend, 1892	103.2
<i>Eumierocerella</i> Hall, 1938	102.8
<i>Eunotochaeta</i> Townsend, 1934	103.13
<i>Euparaphyto</i> Townsend, 1912	103.8
<i>Euravinia</i> Townsend, 1917	102.21
<i>Euselenomyia</i> Townsend, 1912	102.4
<i>Eusenotainia</i> Townsend, 1915	102.4
<i>excisa</i> (Lopes), 1950 (<i>Dexosarcophaga</i>), <i>Oxyvinia</i>	102.23
<i>excitans</i> (Curran & Walley), 1934 (<i>Sarcophaga</i>), <i>Dexosarcophaga</i>	103.42



<i>expuncta</i> (Lopes), 1935 (Sarcophaga), <i>Peekiamyia</i>	102.38
<i>exuberans</i> Lopes & Downs, 1951, <i>Acanthodotheca</i>	103.20
<i>facialis</i> Townsend, 1927, <i>Dexomyiophora</i>	103.32
<i>fulculata</i> (Pandellé), 1896 (Sarcophaga), <i>Parasarcophaga</i>	103.50
<i>farri</i> Dodge, 1965, <i>Sarcophagtiopsis</i>	103.30
<i>Farrimyia</i> Dodge, 1965	103.41
<i>favorabilis</i> (Lopes), 1935 (Sarcophaga), <i>Oxysarcodexia</i>	103.26
<i>fazi</i> (Hall), 1937 (Sarcophaga), <i>Acridiophaga</i>	103.19
<i>femoralis</i> (Schiner), 1868 (Sarcophila), <i>Sarothromyia</i>	103.30
<i>femoralis</i> Schiner of Engel, 1931, <i>Sarothromyia</i>	103.30
<i>fernaldi</i> Parker, 1914, <i>Boettcheria</i>	103.42
<i>fervens</i> (Wiedemann), 1830 (Dexia), <i>Gymnocamptops</i>	103.14
<i>fervida</i> Robineau-Desvoidy, 1830 (Myophora)	103.53
<i>festinans</i> Reinhard, 1947, <i>Cattasoma</i>	103.6
<i>fiebrigi</i> (Enderlein), 1928 (Aetenella), <i>Notochaeta</i>	103.12
<i>filamenta</i> (Dodge), 1963 (Sarcophaga), <i>Malacophagomyia</i>	103.44
<i>fimbriata</i> (Aldrich), 1916 (Sarcophaga), <i>Sarconeiva</i>	103.43
<i>fissa</i> (Aldrich), 1916 (Sarcophaga), <i>Argoravinia</i>	103.46
<i>flavcola</i> Robineau-Desvoidy, 1830 (Myophora)	103.53
<i>flaviceps</i> Macquart, 1843, <i>Sarcophaga</i>	103.53
<i>flavicornis</i> (Townsend), 1891 (Miltogramma), <i>Senotainia</i>	103.5
<i>flavicostrata</i> (Macquart), 1843 (Sarcophaga), <i>Hybopygia</i>	103.24
<i>flavifrons</i> (Macquart), 1846 (Sarcophaga), <i>Oxysarcodexia</i>	103.26
<i>flavipalpis</i> Lopes, 1936, <i>Nephochaetopteryx</i>	103.28
<i>flavopilosa</i> Lopes, 1951, <i>Harpagopyga</i>	103.14
<i>florenciaei</i> (Prado & Fonseca), 1932 (<i>Ctenoprosballia</i>), <i>Euboet-</i> <i>teheria</i>	103.35
<i>fioridensis</i> (Aldrich), 1916 (Sarcophaga), <i>Ravinia</i>	103.21
<i>fioridensis</i> (Townsend), 1892 (<i>Pachyophthalmus</i>), <i>Amobia</i>	103.2
<i>fluminensis</i> Lopes, 1946, <i>Oxysarcodexia</i>	103.26
<i>fortipes</i> Walker, 1861 (Sarcophaga)	103.53
<i>fraterna</i> Lopes, 1946, <i>Oxysarcodexia</i>	103.27
<i>freirei</i> (Mattos), 1919 (Sarcophaga), <i>Sarcodexia</i>	103.34
<i>fringidea</i> (Curran & Wailey), 1934 (Sarcophaga), <i>Oxysarcodexia</i>	103.26
<i>frontalis</i> Townsend, 1928, <i>Pachygraphia</i>	103.16
<i>fulvicornis</i> (Wulp), 1890 (Miltogramma), <i>Senotainia</i>	103.5
<i>fuivierus</i> Rondani, 1850 (Sarcophaga)	103.53
<i>fuivipes</i> Macquart, 1843 (Sarcophaga)	103.53
<i>fuivittata</i> Walker, 1852, (Sarcophaga)	103.43
<i>fuscipennis</i> Dodge, 1967, <i>Boetia</i>	103.12
<i>fuscianalis</i> (Wulp), 1895 (Sarcophaga), <i>Notochaeta</i>	103.43
<i>fuscipennis</i> Dodge, 1967, <i>Boetia</i>	103.28
<i>fusipennis</i> Lopes, 1941, <i>Nephochaetopteryx</i>	
<i>Gaiapagomyia</i> Bisehof, 1904	103.7
<i>garleppi</i> (Enderlein), 1928 (<i>Udamoetis</i>), <i>Euboetcheheria</i>	103.34
<i>Geljskesia</i> Lopes, 1945	103.13
<i>genalis</i> (Thomson), 1869 (Sarcophaga), <i>Helicobia</i>	103.40
<i>georgina</i> (Wiedemann) of Mattos, 1919 (Sarcophaga), <i>Townsen-</i> <i>dimyia</i>	103.9
<i>Gigantotheca</i> Townsend, 1917	103.47
<i>giovannoli</i> Dodge, 1965, <i>Helicobia</i>	103.40
<i>gladysae</i> Dodge, 1965 (Sarcophaga)	103.52


<i>Glaucosareophaga</i> Townsend, 1917	103.49
<i>globosa</i> Aldrich, 1916, <i>Sthenoyga</i>	103.16
<i>globulosa</i> Lopes, 1946, <i>Dexosareophaga</i>	103.41
<i>globulus</i> (Aldrich), 1916 (<i>Sareophaga</i>), <i>Chaetoravinia</i>	103.23
<i>Glutoxys</i> Aldrich, 1929	103.32
<i>Gonlogramma</i> Townsend, 1931	103.4
<i>grandis</i> Lopes, 1946, <i>Oxysaeodexia</i>	103.26
<i>grata</i> (Lopes), 1953 (<i>Dexosareophaga</i>), <i>Oxyvinia</i>	103.23
<i>grisea</i> Lopes, 1951, <i>Lepidodexia</i>	103.15
<i>griseoflavescens</i> Robineau-Desvoidy, 1830 (<i>Myophora</i>)	103.53
<i>griseescens</i> Townsend, 1927, <i>Gymnocamptops</i>	103.14
<i>guadalupensis</i> (Arnaud), 1963 (<i>Blaesoxipha</i>), <i>Aeridiophaga</i>	103.19
<i>gulanica</i> (Curran & Walley), 1934 (<i>Helicobia</i>), <i>Argoravinia</i> ...	103.46
<i>gulo</i> (Fabrielus), 1805 (<i>Musca</i>), <i>Paraphrissopoda</i>	103.36
<i>guyanensis</i> Lopes, 1946, <i>Araehnidomyia</i>	103.42
<i>Gymnocamptops</i> Townsend, 1927	103.14
<i>haemorrhoidalis</i> (Fallén), 1816 (<i>Musca</i>), <i>Bereaea</i>	103.50
<i>halil</i> (Engel), 1931 (<i>Sareophaga</i>), <i>Townsendimyia</i>	103.9
<i>Hamatomyia</i> Blanchard, 1939	103.19
<i>Harpagogyga</i> Aldrich, 1916	103.13
<i>haydeni</i> Dodge, 1965, <i>Helicobia</i>	103.40
<i>helici</i> (Townsend), 1892 (<i>Sareophaga</i>), <i>Helicobia</i>	103.41
<i>Helicobia</i> Coquillett, 1895	103.39
<i>Helicobiella</i> Blanchard, 1939	103.18
<i>Helicoblopsis</i> Townsend, 1927	103.39
<i>hensellana</i> (Enderlein), 1928 (<i>Mesothyrina</i>), <i>Parasareophaga</i> ...	103.50
<i>herbsti</i> (Townsend), 1931 (<i>Gonlogramma</i>), <i>Opsidia</i>	103.4
<i>Hilarella</i> Rondani, 1856	103.3
<i>Hilarella</i> (Zetterstedt), 1844 (<i>Miltogramma</i>), <i>Hilarella</i>	103.3
<i>hillifera</i> (Aldrich), 1916 (<i>Sareophaga</i>), <i>Paraphrissopoda</i>	103.36
<i>hinei</i> (Aldrich), 1930 (<i>Notochaeta</i>), <i>Dexomyiophora</i>	103.32
<i>hirsuta</i> (Hall), 1933 (<i>Sareophaga</i>), <i>Paraphrissopoda</i>	103.36
<i>hirtipes</i> (Walker), 1852 (<i>Sareophaga</i>), <i>Paraphrissopoda</i>	103.36
<i>hondurana</i> (Townsend), 1934 (<i>Anaravinia</i>), <i>Helicobia</i>	103.40
<i>hondurana</i> (Dodge), 1965 (<i>Peckia</i>), <i>Paraphrissopoda</i>	103.37
<i>hospes</i> Aldrich, 1916, (<i>Sareophaga</i>)	103.52
<i>humboldti</i> Robineau-Desvoidy, 1830 (<i>Myophora</i>)	103.53
<i>humerae</i> (Prado & Fonseca), 1932 (<i>Stephanostoma</i>), <i>Euboettcheria</i>	103.34
<i>hunteri</i> (Hough), 1898 (<i>Sareophaga</i>), <i>Protodexia</i>	103.18
<i>Hybopygia</i> Enderlein, 1928	103.23
<i>hypopygialis</i> Townsend, 1915, <i>Xenoplia</i>	103.10
<i>hypopygialis</i> Lopes, 1950, <i>Boettcherimima</i>	103.17
<i>Hystriocnema</i> Townsend, 1919	103.45
<i>ignota</i> Lopes, 1947, <i>Notochaeta</i>	103.12
<i>iheringi</i> Lopes, 1939, <i>Helicobia</i>	103.40
<i>iheringi</i> (Lopes), 1933 (<i>Sareophaga</i>), <i>Sareoneiva</i>	103.48
<i>illata</i> (Lopes), 1938 (<i>Sareophaga</i>), <i>Oxysaeodexia</i>	103.26
<i>imbecilla</i> Wulp, 1896, <i>Sareophaga</i>	103.29
<i>immanis</i> Walker, 1849 (<i>Sareophaga</i>)	103.53
<i>imperialis</i> Robineau-Desvoidy, 1830, <i>Peckia</i>	103.35
<i>impressa</i> Dodge, 1965, <i>Chapiniola</i>	103.11



<i>impura</i> (Curran & Walley), 1934 (Sarcophaga), Paraphrissopoda	103.36
<i>incerta</i> (Walker), 1852 (Sarcophaga), Paraphrissopoda	103.37
<i>inchoata</i> (Aldrich), 1916 (Sarcophaga), Tulacopoda	103.48
<i>inconismus</i> (Reinhard), 1957 (Phytodes), Neophyto	103.11
<i>indonata</i> Lopes, 1955, Sarcophagula	103.29
<i>ingens</i> Walker, 1849 (Sarcophaga)	103.53
<i>injuneta</i> Walker, 1857 (Sarcophaga)	103.53
<i>innota</i> Walker, 1861 (Sarcophaga)	103.53
<i>inoa</i> (Walker), 1849 (Sarcophaga), Sarothromyiops	103.19
<i>inornata</i> Lopes & Downs, 1951, Acanthodotheca	103.20
<i>insolita</i> Lopes, 1946, Oxysarcodexia	103.26
<i>insularis</i> Lopes, 1946, Arachnidomyia	103.42
<i>insularis</i> Townsend, 1918, Opsophytopsis	103.11
<i>intermedia</i> (Townsend) of Coquillett, 1897 (Sarcotachinella), Xenoplia	103.10
<i>intermedia</i> Lopes, 1962, Andinoravinia	103.22
<i>intermutans</i> Walker, 1861 (Sarcophaga)	103.54
<i>intonsa</i> Aldrich, 1928, Opsidia	103.4
<i>intonsa</i> (Curran & Walley), 1934 (Sarcophaga), Oxysarcodexia	103.26
<i>irwini</i> (Dodge), 1966 (Peckia), Paraphrissopoda	103.37
<i>isla</i> (Curran), 1934 (Sarcophaga), Gigantotheca	103.47
<i>itamobia</i> Townsend, 1927	103.6
<i>itaqua</i> Dodge, 1966, Dexosarcophaga	103.41
<i>itiophyto</i> Hall, 1937	103.8
<i>jamaicensis</i> Dodge, 1965, Sarcofahrtiopsis	103.30
<i>jamaicorum</i> Dodge, 1965, Cicatricia	103.20
<i>Jantia</i> Rohdendorf, 1937	103.50
<i>jejuna</i> Walker, 1849 (Sarcophaga)	103.54
<i>Johnsonia</i> Coquillett, 1895	103.16
<i>juquilana</i> Townsend, 1934, Juquiancia	103.4
<i>Juquiancia</i> Townsend, 1934	103.4
<i>kartabo</i> (Curran & Walley), 1934 (Sarcophaga), Oxysarcodexia	103.24
<i>kesselli</i> (Dodge), 1966 (Sarcophaga), Wohlfahrtiopsis	103.51
<i>keyensis</i> (Dodge), 1965 (Peckia), Paraphrissopoda	103.37
<i>khalili</i> (Salen), 1935 (Sarcophaga), Hystricocnema	103.46
<i>knabi</i> Townsend, 1917, Glaucosarcophaga	103.49
<i>koehleri</i> (Blanchard), 1939 (Parapeltopyga), Lipoptilocnema	103.43
<i>Kurtomyia</i> Roback, 1954	103.20
<i>laginucula</i> (Hall), 1933 (Johnsonia), Helicobia	103.40
<i>lahillei</i> (Blanchard), 1939 (Opsophyto?), Helicobia	103.40
<i>Lahillemyia</i> Blanchard, 1939	103.43
<i>lamanensis</i> (Robineau-Desvoidy), 1830 (Peckia), Paraphrissopoda	103.36
<i>lambens</i> (Wiedemann), 1830 (Sarcophaga), Acridiophaga	103.19
<i>lanei</i> (Lopes), 1933 (Sarcophaga), Acanthodotheca	103.20
<i>lanei</i> Townsend, 1934, Lipoptilocnema	103.43
<i>larvicida</i> (Lopes), 1935 (Sarcophaga), Cuculomyia	103.47
<i>larvivoxax</i> (Lopes), 1935 (Sarcophaga), Sarconeiva	103.48
<i>lateropili</i> Allen, 1926, Metopia	103.4
<i>Lepidodexia</i> Brauer & Bergenstamm, 1891	103.15
<i>Lepyria</i> Roback, 1954	103.48
<i>leucocephala</i> (Rossi), 1790 (Musca), Metopia	103.3

<i>Leucoravinia</i> Townsend, 1928	103.31
<i>lherminieri</i> (Robineau-Desvoidy), 1830 (Myophora), <i>Ravinia</i> ...	103.21
<i>llebermanni</i> Blanchard, 1955, <i>Protodexia</i>	103.18
<i>liguloides</i> (Blanchard), 1939 (<i>Parapeltopyga</i>), <i>Lipoptilocnema</i> ..	103.44
<i>lindneri</i> Townsend, 1931, <i>Neophytodes</i>	103.6
<i>Liopygia</i> Enderlein, 1928	103.50
<i>Liosarcophaga</i> Rohdendorf, 1937	103.50
<i>Lipoptilocnema</i> Townsend, 1934	103.43
<i>lithogaster</i> (Curran & Walley), 1934 (<i>Sarcophaga</i>), <i>Paraphrisso-</i> <i>poda</i>	103.37
<i>litoralis</i> Allen, 1924, <i>Senotainia</i>	103.5
<i>lopesi</i> Dodge, 1966, <i>Oxysarcodexia</i>	103.26
<i>Lopesimyia</i> Blanchard, 1954	103.44
<i>lorosa</i> (Hall), 1937 (<i>Sarcophaga</i>), <i>Acridiophaga</i>	103.19
<i>lubera</i> (Brèthes), 1920 (<i>Sarcophaga</i>), <i>Helicobia</i>	103.40
<i>luculenta</i> (Lopes), 1938 (<i>Sarcophaga</i>), <i>Cuculomyia</i>	103.47
<i>lutzi</i> Lopes, 1958, <i>Paraphrisopoda</i>	103.37
<i>Macronichia</i> , error	103.6
<i>Macronychia</i> Rondani, 1859	103.6
<i>Macronychiinae</i>	103.5
<i>macrophthalma</i> Lopes, 1955, <i>Sarcophagula</i>	103.29
<i>maculata</i> Macquart, 1843 (<i>Phrisopoda</i>)	100.54
<i>maerens</i> Townsend, 1916, <i>Melanophyto</i>	103.32
<i>magellanica</i> Townsend, 1937, <i>Austrohartigia</i>	103.8
<i>magnifica</i> Enderlein, 1928, <i>Pattonella</i>	103.37
<i>major</i> Lopes, 1946, <i>Oxysarcodexia</i>	103.26
<i>malacophaga</i> Lopes, 1940, <i>Udamopyga</i>	103.44
<i>Malacophagomyia</i> Lopes, 1966	103.44
<i>Malacophagula</i> Bequaert, 1925	103.33
<i>mallochi</i> (Hall), 1937 (<i>Mimophytomyia</i>), <i>Alaccoprosopa</i>	103.9
<i>mallogaster</i> (Blanchard), 1939 (<i>Xenagria</i>), <i>Tricharaca</i>	103.31
<i>Mallonotum</i> Aldrich, 1925	103.31
<i>mantidophaga</i> (Blanchard), 1942 (<i>Helicobiella</i>), <i>Mantidophaga</i> ..	103.18
<i>Mantidophaga</i> Townsend, 1919	102.18
<i>margareti</i> (Lahille), 1907 (<i>Sarcophaga</i>), <i>Lipoptilocnema</i>	103.44
<i>marginata</i> (Aldrich), 1916 (<i>Sarcophaga</i>), <i>Acridiophaga</i>	103.19
<i>marina</i> Hall, 1938, <i>Apelophyla</i>	103.24
<i>marstoni</i> Dodge, 1966, <i>Boettcheria</i>	103.43
<i>marstoniana</i> Dodge, 1965, <i>Tolucamyia</i>	103.49
<i>megista</i> (Hall), 1933 (<i>Sarcophaga</i>), <i>Dexosarcophaga</i>	103.41
<i>meinckei</i> Blanchard, 1939, <i>Andinoravinia</i>	102.22
<i>Melanophyto</i> Townsend, 1916	103.32
<i>mentor</i> Curran & Walley, 1934 (<i>Sarcophaga</i>)	103.52
<i>meridiana</i> Townsend, 1912, <i>Metopia</i>	103.3
<i>meridionalis</i> Blanchard, 1942, <i>Oxysarcodexia</i>	103.25
<i>meridionalis</i> (Engel), 1931 (<i>Sarcophaga</i>), <i>Oxysarcodexia</i>	103.26
<i>Merokellymyia</i> Blanchard, 1939	103.44
<i>metallica</i> Townsend, 1928, <i>Sisarsarcophaga</i>	103.13
<i>metamasi</i> Lopes, 1946, <i>Dexosarcophaga</i>	103.41
<i>Metopia</i> Meigen, 1803	103.3
<i>metoploides</i> Allen, 1926, <i>Opsidia</i>	103.4

<i>mexicana</i> Lopes, 1950, <i>Boettcheria</i>	103.43
<i>mexicana</i> Lopes & Downs, 1951, <i>Acanthodotheca</i>	103.19
<i>Microcerella</i> Macquart, 1851	103.7
<i>Microcerella</i> Macquart of Brauer & Bergenstamm, 1891	103.7
<i>Microcerellamyia</i> Hall, 1937	103.7
<i>Microcerellini</i>	103.7
<i>Micronotochaeta</i> Townsend, 1927	103.11
<i>Mieroplugia</i> Townsend, 1915	103.49
<i>miropyga</i> (Wulp), 1895 (<i>Sarcophaga</i>), <i>Notochaeta</i>	103.12
<i>microphygialis</i> Williston, 1896 (<i>Sarcophaga</i>)	103.52
<i>Microsenotainia</i> Townsend, 1916	103.4
<i>milleri</i> (Johnston & Tiegs), 1922 (<i>Sarcophaga</i>), <i>Hybopygia</i>	103.24
<i>Miltogramminae</i>	103.2
<i>Miltorainia</i> Townsend, 1917	103.21
<i>Mimophytomyia</i> Hall, 1937	103.9
<i>minense</i> Lopes & Downs, 1951, <i>Acanthodotheca</i>	103.19
<i>miniatleauda</i> Enderlein, 1928, <i>Ptilocnema</i>	103.39
<i>minima</i> Vimmer & Soukoup, 1940 (<i>Agria</i>)	103.54
<i>minima</i> Vimmer & Soukoup, 1940 (<i>Sarcophila</i>)	103.51
<i>minuta</i> (Hall), 1928 (<i>Sarcophaga</i>), <i>Ravinia</i>	103.21
<i>minuta</i> (Lahille), 1907 (<i>Sarcophaga</i>), <i>Doringia</i>	103.9
<i>minutipenis</i> (Hall), 1937 (<i>Sarcophaga</i>), <i>Peckiamyia</i>	103.38
<i>mirabilis</i> Dodge, 1965, <i>Steatopyga</i>	103.48
<i>misella</i> (Lopes), 1938 (<i>Sarcophaga</i>), <i>Lipoptilocnema</i>	103.44
<i>mitifica</i> Lopes, 1953, <i>Oxysarcodexia</i>	103.26
<i>modesta</i> Lopes, 1946, <i>Oxysarcodexia</i>	103.26
<i>modesta</i> (Wiedemann), 1830 (<i>Sarcophaga</i>), <i>Argoravinia</i>	103.46
<i>modulata</i> Wulp, 1895, <i>Paramintho</i>	103.33
<i>molinai</i> Lopes, 1942, <i>Nephochaetopteryx</i>	103.28
<i>molitor</i> (Curran & Walley), 1934 (<i>Sarcophaga</i>), <i>Oxysarcodexia</i>	103.26
<i>molluscoperda</i> Dodge, 1965, <i>Alrypel</i>	103.47
<i>Mononotochaeta</i> Townsend, 1935	103.13
<i>morionella</i> (Aldrich), 1930 (<i>Sarcophaga</i>), <i>Helicobia</i>	103.40
<i>muehni</i> Blanchard, 1939, <i>Scopaediscus</i>	103.10
<i>mundelli</i> Blanchard, 1939, <i>Boettcheria</i>	103.43
<i>museoides</i> Bigot, 1857 (<i>Sarcophaga</i>)	103.54
<i>myersi</i> (Curran), 1938 (<i>Dexomyiophora</i>), <i>Emdenimyia</i>	103.17
<i>myersi</i> Townsend, 1935, <i>Mononotochaeta</i>	103.13
<i>naides</i> Lopes, 1941, <i>Euboeettcheria</i>	103.35
<i>nana</i> (Enderlein), 1928 (<i>Dasypyga</i>), <i>Parahybopelta</i>	103.8
<i>nana</i> (Wulp), 1892 (<i>Miltogramma</i>), <i>Senotainia</i>	103.5
<i>natiuscula</i> Lopes, 1941, <i>Phagita</i>	103.30
<i>nebulosa</i> (Coquillett), 1902 (<i>Araba</i>), <i>Sphenometopa</i>	103.5
<i>neglecta</i> Lopes, 1946, <i>Helicobia</i>	103.40
<i>nelvai</i> Lopes, 1940, <i>Udamopyga</i>	103.45
<i>nelvai</i> (Mattos), 1919 (<i>Sarcophaga</i>), <i>Oxysarcodexia</i>	103.26
<i>Neobellieria</i> Blanchard, 1939	103.51
<i>Neophyto</i> Townsend, 1908	103.11
<i>Neophytodes</i> Townsend, 1931	103.6
<i>Neopsidia</i> Townsend, 1928	103.4
<i>Neosarcodexia</i> Blanchard, 1942	103.34
<i>Neotephromyiella</i> Blanchard, 1955	103.20
<i>neotropica</i> Bequaert, 1925, <i>Malacophagula</i>	103.33



<i>neotropica</i> le Prado & Fonseca, 1932, Oxysarcodexia	103.25
<i>nephele</i> Lopes, 1941, Paraphrissopoda	103.37
<i>Nephochaetopteryx</i> Townsend, 1934	103.28
<i>neuquenensis</i> (Blanchard), 1942 (Tephromyella), Acridiophaga ..	103.19
<i>nicasia</i> Lopes, 1941, Euboettcheria	103.35
<i>nigribimbo</i> Dodge, 1965, Harpagopyga	103.14
<i>nigricauda</i> (Dodge), 1964 (Peckia), Paraphrissopoda	103.37
<i>nigricornis</i> Allen, 1926, Eumacronychia	103.3
<i>nigricornis</i> (Dodge), 1966 (Petriana), Lepidodexia	103.15
<i>nigropilosa</i> Lopes, 1951, Lepidodexia	103.15
<i>nltens</i> Townsend, 1915, Microplagia	103.49
<i>nobilis</i> (Thomson), 1869 (Sarcophaga), Hybopygia	103.24
<i>nodositas</i> (Blanchard), 1939 (Merokellymyia), Udamopyga	103.45
<i>notata</i> Lopes, 1953, Sarcodexia	103.33
<i>notata</i> Macquart, 1855 (Sarcophaga)	103.54
<i>Notochaeta</i> Aldrich, 1916	103.11
<i>Notochaetini</i>	103.11
<i>Notochaetophyto</i> Hall, 1933	103.39
<i>Notochaetopsis</i> Townsend, 1918	103.19
<i>oberrans</i> (Wulp), 1896 (Sarcophaga), Acridiophaga	103.19
<i>obscura</i> Lopes, 1950, Notochaeta	103.12
<i>obscurior</i> (Lahille), 1907 (Sarcophaga), Acridiophaga	103.19
<i>obscuripes</i> (Hall), 1928 (Sarcophaga), Ravinia	103.22
<i>obsoleta</i> (Wiedemann), 1830 (Sarcophaga), Sarcophagula	103.29
<i>obtusifrons</i> (Thomson), 1869 (Sarcophaga), Oxysarcodexia	103.27
<i>occludua</i> (Fabricius), 1794 (Musca), Sarcophagula	103.29
<i>occlpitalis</i> (Thomson), 1869 (Sarcophaga), Pattonella	103.37
<i>occulta</i> Lopes, 1946, Oxysarcodexia	103.26
<i>ochracea</i> (Aldrich), 1916 (Sarcophaga), Ravinia	103.22
<i>ochrpyga</i> (Wulp), 1895 (Sarcophaga), Oxysarcodexia	103.26
<i>ochristriga</i> (Enderlein), 1928 (Agriochaeta), Gymnocamptops ...	103.15
<i>offecta</i> (Lopes), 1938 (Sarcophaga), Neobellieria	103.51
<i>oglobini</i> Blanchard, 1942, Protodexia	103.18
<i>oglobini</i> (Lopes), 1938 (Sarcophaga), Udamopyga	103.45
<i>oliantaytambensis</i> (Hall), 1928 (Sarcophaga), Ravinia	103.22
<i>omissa</i> Lopes, 1946, Oxysarcodexia	103.26
<i>opifera</i> (Coquillett), 1892 (Sarcophaga), Opsophyto	103.17
<i>oplma</i> (Wiedemann), 1830 (Sarcophaga), Orodexia	103.13
<i>Opsidia</i> Coquillett, 1895	103.4
<i>Opsidiopsis</i> Townsend, 1919	103.4
<i>Opsophyto</i> Townsend, 1915	103.17
<i>Opsophytopsis</i> Townsend, 1918	103.10
<i>orails</i> Rondani, 1868 (Sarcophaga)	103.54
<i>orbitalis</i> (Curran & Walley), 1934 (Sarcophaga), Nephochaetopteryx	103.28
<i>orbitalis</i> Dodge, 1965, Bahamioia	103.30
<i>orbitalis</i> Dodge, 1966, Oxysarcodexia	103.26
<i>ornata</i> (Townsend), 1917 (Amobiopsis), Macronychia	103.6
<i>ornata</i> Townsend, 1927, Itamobia	103.6
<i>ornata</i> Townsend, 1927, Orobrachycoma	103.32
<i>ornata</i> Townsend, 1927, Orodexia	103.13
<i>ornata</i> Townsend, 1927, Orosarcophaga	103.28
<i>ornaticauda</i> (Aldrich), 1923 (Pachyophthalmus), Amobia	103.2



Orobraehycoma Townsend, 1927	103.32
Orodexia Townsend, 1927	103.13
Orosareophaga Townsend, 1927	103.28
otiosa (Williston), 1896 (Sarcophaga), Paraphrissopoda	103.36
Oxyhelicobia Blanchard, 1942	103.39
Oxysareodexia Townsend, 1917	103.24
Oxyvinla Dodge, 1966	103.23
Pachygraphia Brauer & Bergenstamm, 1891	103.16
Pachygraphiops Townsend, 1934	103.13
Pachygraphomyia Hall, 1933	103.46
Pachyophthalmus Brauer & Bergenstamm, 1889	103.2
paeta (Townsend), 1934 (Pachygraphiops), Harpagopyga	103.14
pallidipilosa (Curran & Walley), 1934 (Sarcophaga), Pattonella	103.38
pallincervis (Thomson), 1869 (Sarcophaga), Ravinia	103.21
pallisteri Dodge, 1966, Oxysareodexia	103.26
pallidiventris Townsend, 1934, Nephochaetopteryx	103.28
Panamaphyto Townsend, 1931	103.7
Paradisochaeta Blanchard, 1939	103.36
Parahybopelta Blanchard, 1955	103.8
paralis Lopes, 1938 (Sarcophaga)	103.52
Paramaeronychiinae	103.6
Parametopia Townsend, 1916	103.3
Paramintho Brauer & Bergenstamm, 1891	103.33
Parapeltopyga Blanchard, 1939	103.43
Paraphrissopoda Townsend, 1915	103.35
Parasarcodexia Townsend, 1917	103.42
Parasareophaga Johnston & Tiegs, 1921	103.50
parkeri (Aldrich), 1916 (Sarcophaga), Boettcheria	103.43
parva Lopes, 1946, Notochaeta	103.12
parva Lopes, 1946, Oxysareodexia	103.27
parva (Walker), 1852 (Sarcophaga), Sarcophagula	103.29
parvula (Wiedemann), 1830 (Sarcophaga), Sarcophagula	103.29
parvula (Wiedemann), 1830 (Sarcophaga), Sarcophagula	103.20
paterna Dodge, 1965, Sarcophagula	103.30
patriciae Dodge, 1965, (Sarcophaga)	103.54
Pattonella Enderlein, 1928	103.37
pauciseta Dodge, 1966, Metopia	103.3
pauciseta Lopes & Downs, 1951, Acanthodotheca	103.20
paulistanensis (Mattos), 1919 (Sarcophaga), Oxysareodexia ...	103.27
Peckia Robineau-Desvoidy, 1830	103.35
Peckiamyia Dodge, 1966	103.38
pecorensis Dodge, 1966, Peckiamyia	103.39
peculiaris Dodge, 1966, Promayoa	103.41
pedata (Aldrich), 1916 (Sarcophaga), Abapa	103.39
pedatulla (Dodge), 1965 (Peckia), Abapa	103.39
pedunculata (Hall), 1931 (Sarcophaga), Cueulomyia	103.47
pellita (Enderlein), 1928 (Liopygia), Tricharaea	103.31
peltata (Aldrich), 1916 (Sarcophaga), Oxysareodexia	103.27
penai Dodge, 1965, Tricharaea	103.31
percita (Lopes), 1938 (Sarcophaga), Udampopyga	103.45
percuta (Lopes), 1938 (Sarcophaga), Euboeettcheria	103.35

<i>perlita</i> Lopes, 1953, <i>Euboettcheria</i>	103.35
<i>perneta</i> (Walker), 1861 (Sarcophaga), <i>Oxysarcodexia</i>	103.27
<i>perpendicularis</i> Wulp, 1890, <i>Metopia</i>	103.3
<i>personella</i> Reinhard, 1965, <i>Eumacronychia</i>	103.2
<i>peruana</i> Townsend, 1912, <i>Sarcophagula</i>	103.29
<i>peruviana</i> Lopes, 1950, <i>Boettcheria</i>	103.43
<i>peruviansis</i> (Townsend), 1912 (<i>Euselenomyia</i>), <i>Senotainia</i>	103.5
<i>pervillosa</i> (Aldrich), 1916 (Sarcophaga), <i>Tulacopoda</i>	103.48
<i>Pessoamyia</i> Lopes, 1938	103.33
<i>Petriana</i> Lopes, 1945	103.16
<i>Petrosarcophaga</i> Townsend, 1919	103.51
<i>pexata</i> (Wulp), 1895 (Sarcophaga), <i>Paraphrissopoda</i>	103.37
<i>Phaeosarcodexia</i> Hall, 1937	103.8
<i>Phagita</i> Lopes, 1941	103.30
<i>philippi</i> Rondani, 1861 (Sarcophaga)	103.54
<i>phoenicrus</i> Wiedemann, 1830 (Sarcophaga)	103.54
<i>Phrissopoda</i> Macquart, 1835	103.35
<i>Phytodes</i> Coquillett, 1910	103.11
<i>pillifera</i> Lopes, 1939, <i>Helicobia</i>	103.40
<i>pigmea</i> Rondani, 1850 (Sarcophaga)	103.54
<i>pillipleura</i> Lopes, 1939, <i>Helicobia</i>	103.40
<i>pillventris</i> Dodge, 1966, <i>Oxyvinia</i>	103.23
<i>piophila</i> (Neiva & Gomes), 1913 (Sarcophaga), <i>Sarcodexia</i>	103.34
<i>placida</i> (Aldrich), 1925 (Sarcophaga), <i>Cuculomyia</i>	103.47
<i>plaumanni</i> Lopes, 1946, <i>Oxysarcodexia</i>	103.27
<i>plebeja</i> Lopes, 1946, <i>Oxysarcodexia</i>	103.27
<i>plinthopyga</i> (Wiedemann), 1830 (Sarcophaga), <i>Hystriocnema</i> ...	103.45
<i>plinthopyga</i> (Wiedemann) of Mattos, 1919 (Sarcophaga), <i>Euboettcheria</i>	103.34
<i>plumigera</i> (Wulp), 1895 (Sarcophaga), <i>Notochaeta</i>	103.12
<i>plumigera</i> (Wulp) of Aldrich, 1916 (<i>Notochaeta</i>), <i>Dexomylophora</i>	103.32
<i>plumipes</i> (Robineau-Desvoidy), 1830 (Peckia), <i>Paraphrissopoda</i>	103.37
<i>podeuxaucus</i> (Enderlein), 1928 (<i>Robineanella</i>), <i>Boettcheria</i>	103.43
<i>polita</i> Townsend, 1935, <i>Rupununia</i>	103.4
<i>postilla</i> (Reinhard), 1947 (Sarcophaga), <i>Kurtomyia</i>	103.20
<i>postuncinata</i> (Hall), 1933 (Sarcophaga), <i>Villegasia</i>	103.48
<i>praeceps</i> (Wiedemann), 1830 (Sarcophaga), <i>Peckia</i>	103.35
<i>praevolans</i> (Wulp), 1895 (Sarcophaga), <i>Boettcheria</i>	103.43
<i>pridei</i> (Richards), 1936 (<i>Pachyophthalmus</i>), <i>Amobla</i>	103.2
<i>Probellieria</i> Blanchard, 1942	103.50
<i>proerna</i> Walker, 1849 (Sarcophaga)	103.54
<i>prohlbita</i> (Aldrich), 1916 (Sarcophaga), <i>Acanthodotheca</i>	103.20
<i>prolixa</i> Reinhard, 1939, <i>Eumacronychia</i>	103.3
<i>Promayoa</i> Dodge, 1966	103.41
<i>prosternalis</i> Lopes, 1933, <i>Pessoamyia</i>	103.33
<i>Prosthetocirca</i> Townsend, 1917	103.10
<i>Protodexia</i> Townsend, 1912	103.17
<i>provecta</i> (Lopes), 1939 (Sarcophaga), <i>Udamopyga</i>	103.45
<i>providencia</i> Dodge, 1965, <i>Helicobia</i>	103.40
<i>Pseudosarcophagula</i> Townsend, 1927	103.29
<i>pseudovaria</i> Blanchard, 1939, <i>Hybopygia</i>	103.23
<i>Ptilocnema</i> Enderlein, 1928	103.39
<i>pudica</i> Rondani, 1850 (Sarcophaga)	103.54

puella (Williston), 1896 (Sarcophiloides), Sarcophagula	103.29
Punaphyto Townsend, 1915	103.41
<i>Punasarcophaga</i> Townsend, 1915	103.21
pusilla Bigot, 1857, (Sarcophaga)	103.54
pusiola (Wulp), 1895 (Sarcophaga), Ravinia	103.22
pyrrhopyga (Hall), 1933 (Sarcophaga), Boettcheria	103.43
<i>quadraticcephala</i> (Townsend), 1928 (Leucoravinia), Tricharaea..	103.31
<i>quadriforcps</i> (Curran), 1932 (Sarcophaga), Hystriocnema	103.46
<i>quadrisctosa</i> (Coquillett), 1901 (Helicobia), Chaetoravinia	103.22
quadrivittata (Macquart), 1843 (Sarcophaga), Ravinia	103.22
quaesita (Hall), 1937 (Sarcophaga), Protodexia	103.18
quecetuba Dodge, 1966, Dexosarcophaga	103.42
rabbi Dodge, 1965, Helicobia	103.40
Rafaelia Townsend, 1917	103.33
<i>Raimondia</i> Townsend, 1917	103.15
rapax (Walker), 1849 (Sarcophaga), Helicobia	103.40
Ravinia Robineau-Desvoidy, 1863	103.21
Raviniini	103.21
Raviniopsis Townsend, 1918	103.46
<i>rediviva</i> (Walker), 1849 (Sarcophaga), Ravinia	103.21
<i>reinhardii</i> (Hall), 1928 (Sarcophaga), Chaetoravinia	103.23
resinata (Hall), 1933 (Notochaetophyto), Helicobia	103.41
resona (Lopes), 1935 (Sarcophaga), Pattonella	103.33
rethae Dodge, 1965, Helicobia	103.40
retrocita (Hall), 1933 (Sarcophaga), Paraphrissopoda	103.37
retroversa (Lopes), 1935 (Sarcophaga), Boettcheria	103.43
retusa (Hall), 1933 (Sarcophaga), Dexosarcophaga	103.42
retusa Hall, 1937, Microcerellamyia	103.8
rimosa (Hall), 1938 (Sarcophaga), Acanthodotheca	103.20
riograndensis Lopes, 1946, Oxysarcodexia	103.27
<i>robusta</i> (Aldrich), 1916 (Sarcophaga), Hystriocnema	103.45
rubella Wiedemann, 1830 (Sarcophaga)	103.54
rubrianalis Macquart, 1851 (Sarcophaga)	103.54
rubrigaster Dodge, 1965, Titanogrypa	103.48
rubriventris (Macquart), 1851 (Sarcophaga), Gynnocamptops	103.15
rubriventris Macquart, 1846, Senotainia	103.5
rudicopages Hall, 1933 (Sarcophaga)	103.52
rudis (Aldrich), 1916 (Sarcophaga), Acanthodotheca	103.20
<i>ruficauda</i> (Aldrich), 1928 (Brachicoma), Austrohartigia	103.8
ruficornis (Fabricius), 1794 (Musca), Parasarcophaga	103.50
ruficrura Rondani, 1864 (Sarcophaga)	103.54
rufipalpis Macquart, 1843 (Sarcophaga)	103.54
rufipes Dodge, 1965, Harpagopyga	103.14
<i>rufipes</i> Macquart, 1843, (Sarcophaga)	103.53
rufipes Townsend, 1917, Andinoravinia	103.22
rufiventris Townsend, 1917, Rafaelia	103.33
rufiventris (Wiedemann), 1830 (Sarcophaga), Argoravinia	103.46
<i>rufiventris</i> (Wiedemann) of Aldrich, 1916 (Sarcophaga), Rafaelia	103.33
rufitibia (Wulp), 1895 (Sarcophaga), Sthenopyga	103.16
rufomaculata Macquart, 1851, Microcerella	103.7
Rupununia Reinhard, 1935	103.4

rusca Hall, 1937, Aulacophyto	103.7
rustica Lopes, 1950, Notochaeta	103.12
ruthae (Dodge), 1965 (Farrimyia), Dexosarcophaga	103.42
Sabinata Parker, 1921	103.46
salavini Blanchard, 1955, Parahybopelta	103.8
salobrensis Lopes, 1942, Lipoptilocnema	103.44
sanchezi (Blanchard), 1939 (Helicobiella), Mantidophaga	103.18
sanctijosephi (Engel), 1931 (Sarcophaga), Argoravinia	103.46
sanguisuga (Hall), 1933 (Sarcophaga), Oxysarcodexia	103.27
salutaris Dodge, 1965, Sarcodexiopsis	103.49
Sapromyia Roback, 1954	103.51
sarcinata Lopes, 1953, Oxysarcodexia	103.27
Sarcodexia Townsend, 1892	103.33
Sarcodexiopsis Townsend, 1917	103.49
Sarcophagomyia Hall, 1937	103.32
Sarcophagopsalis Hall, 1933	103.20
Sarcophagobolia Blanchard, 1939	103.28
Sarcophagomyia Townsend, 1892	103.2
Sarcophagomyia Roback, 1954	103.41
Sarcophagina Lopes, 1940	103.48
Sarcophagina Curran, 1928	103.46
sarcophagina Thomson, 1869, Microcerella	103.7
sarcophagina (Townsend), 1927 (Xylocampta), Lepidodexia	103.15
sarcophagina (Wulp), 1890 (Miltogramma), Amobia	103.2
Sarcophaginae	103.7
Sarcophagini	103.31
sarcophagoides (Townsend), 1892 (Sarcophagomyia), Amobia ..	103.2
Sarcophagula Wulp, 1887	103.29
Sarcophagulin	103.29
Sarcophagulus Blanchard, 1939	103.49
Sarcophodexia Townsend, 1918	103.19
Sarcophodexiopsis Blanchard, 1942	103.48
Sarothromyia Brauer & Bergenstamm, 1891	103.30
Sarothromyia Townsend, 1917	103.10
sarracenioides (Aldrich), 1916 (Sarcophaga), Parasarcophaga ...	103.51
satanica Dodge, 1965, Peckia	103.35
Scarabacophaga Townsend, 1918	103.51
scatophagina Thomson, 1869 Tricharaea	103.31
scelesta (Hall), 1931 (Sarcophaga), Dexosarcophaga	103.42
Scopaediscus Blanchard, 1939	103.10
securifera (Villeneuve), 1908 (Sarcophaga), Parasarcophaga	103.50
Senotainia Macquart, 1846	103.4
Senotainia Macquart, 1846	103.4
Senotainia Macquart, 1846	103.4
setifacies Lopes, 1945, Paraphrissopoda	102.37
setiforceps (Hall), 1930 (Sarcophaga), Abapa	103.39
setigena (Enderlein), 1928 (Udamoctis), Udamopyga	103.45
setinervis Lopes, 1939, Helicobia	103.41
setosa Aldrich, 1916, Johnsonia	103.16
Shannomyiella Lopes, 1938	103.16
siccana (Reinhard), 1947 (Sarcophaga), Boettcheria	103.43
similis Lopes, 1946, Boettcheria	103.43



<i>similis</i> (Townsend), 1891 (Miltogramma), Senotainia	103.5
<i>simplex</i> Aldrich, 1916, Sarothromyia	103.31
<i>simplicoides</i> (Lopes), 1933 (Sarcophaga), Oxysarcodexia	103.27
<i>Sisasarcophaga</i> Townsend, 1928	103.13
<i>smagra</i> Hall, 1938, Udamopyga	103.45
<i>smartii</i> Lopes, 1941, Pattonella	103.38
<i>Socorromyia</i> Dodge, 1964	103.33
<i>sp.</i> Dodge, 1965, Harpagopyga	103.14
<i>sp.</i> Dodge, 1966, Aulacophyto	103.7
<i>speciosa</i> (Lopes), 1946 (Fletcherimyia), Speciosia	103.20
<i>Speciosia</i> Roback, 1954	103.20
<i>spectabilis</i> (Aldrich), 1916 (Sarcophaga), Paraphrissopoda	103.37
<i>Sphenometopa</i> Townsend, 1908	103.5
<i>spinigena</i> (Rondani), 1946 (Sarcophaga), Austrohartigia	103.8
<i>spinosa</i> (Hall), 1933 (Pachygraphomyia), Raviniopsis	103.46
<i>spinosa</i> (Hall), 1937 (Phaeosarcodexia), Austrohartigia	103.8
<i>spinosa</i> Lopes, 1948, Emdenimyia	103.17
<i>spitzii</i> (Lopes), 1933 (Sarcophaga), Squamatoides	103.38
<i>Squamata</i> Curran, 1928	103.38
<i>squamata</i> (Walker), 1852 (Tachina), Lepidodexia	103.15
<i>Squamatoides</i> Curran, 1927	103.38
<i>stahli</i> Dodge, 1966, Squamatoides	103.38
<i>stallengi</i> (Lahille), 1907 (Sarcophaga), Lopesimyia	103.44
<i>Steatopyga</i> Dodge, 1965	103.48
<i>steinbachii</i> (Enderlein), 1928 (Tetradiscalis), Dexosarcophaga ...	103.42
<i>steindachneri</i> (Brauer & Bergenstamm), 1891 (Microcerella), Galapagomyia	103.7
<i>stellata</i> (Wulp), 1895 (Sarcophaga), Helicobia	103.41
<i>sternalis</i> Allen, 1926, Eumacronychia	103.3
<i>sternodontes</i> Townsend, 1892, Sarcodexia	103.33
<i>Sthenopyga</i> Aldrich, 1916	103.16
<i>Sthenopygopsis</i> Townsend, 1935	103.41
<i>stigmosa</i> Reinhard, 1965, Eumacronychia	103.3
<i>subandina</i> (Blanchard), 1938 (Doringia), Austrohartigia	103.8
<i>subaurata</i> (Engel), 1931 (Bercaea), Nephochaetopteryx	103.28
<i>subcylindrica</i> Townsend, 1935, Sthenopygopsis	103.41
<i>subducta</i> (Lopes), 1935 (Sarcophaga), Euboettcheria	103.35
<i>subpolita</i> Aldrich, 1916, Notochaeta	103.12
<i>subsericans</i> Walker, 1857 (Sarcophaga)	103.54
<i>sueta</i> (Wulp), 1895 (Sarcophaga), Ravinia	103.22
<i>sugens</i> (Wiedemann), 1830 (Sarcophaga), Sarcophagula	103.29
<i>sulcata</i> (Aldrich), 1916 (Sarcophaga), Zygastropyga	103.47
<i>sulcata</i> (Dodge), 1965 (Peckia), Abapa	103.39
<i>surinamensis</i> Wiedemann, 1830 (Sarcophaga)	103.54
<i>surrubea</i> (Wulp), 1895 (Sarcophaga), Helicobia	103.41
<i>surrubea</i> (Wulp) of Aldrich, 1916 (Sarcophaga), Helicobia	103.40
<i>synthetica</i> Townsend, 1912, Protodexia	103.18
<i>tacituro</i> Roback, 1952, Ravinia	103.22
<i>taitensis</i> (Schiner), 1868 (Sarcophaga), Oxysarcodexia	103.27
<i>taurus</i> Aldrich, 1916 (Sarcophaga)	103.52
<i>teffeensis</i> Townsend, 1927, Xylocamptopsis	103.15
<i>tenta</i> Hall, 1937, Sarcophartimyia	103.33
<i>tenuis</i> Wulp, 1896, Sarcophagula	103.29

<i>tenuiventris</i> (Wulp), 1895 (Sarcophaga), Oxysarcodexia	103.27
<i>Tephromylini</i>	103.17
<i>tergata</i> (Coquillett), 1895 (Araba), Sphenometopa	103.5
<i>terminalis</i> (Wiedemann), 1830 (Sarcophaga), Hybopygia	103.23
<i>termitaria</i> (Lopes), 1939 (Sarcophaga), Dexosarcophaga	103.42
<i>termitophila</i> Silvestri, 1903, Tripanurga	103.46
<i>tesselata</i> (Fabricius), 1805 (Musca), Euboettcheria	103.34
<i>tesselata</i> (Fabricius) of Wiedemann, 1830 (Sarcophaga), Oxysarcodexia	103.25
<i>tesselata</i> (Fabricius) of Mattos, 1919 (Sarcophaga), Ravinia ...	103.21
<i>tesselata</i> (Enderlein), 1928 (Liopygia), Sarcodexia	103.34
<i>tetraptera</i> Brauer & Bergenstamm, 1891, Lepidodexia	103.15
<i>Thomsonia</i> Rohdendorf, 1937	103.50
<i>timida</i> (Aldrich), 1916 (Sarcophaga), Oxysarcodexia	103.27
<i>Titanogrypa</i> Townsend, 1917	103.47
<i>titubata</i> Lopes, 1945, Oxysarcodexia	103.27
<i>Tolucamyia</i> Dodge, 1965	102.49
<i>townsendi</i> Aldrich, 1925, Notochaeta	103.12
<i>townsendi</i> (Engel), 1931 (Sarcophaga), Chaetoravinia	103.23
<i>townsendi</i> (Prado & Fonseca), 1932 (Chrysostoma), Paraphrisopoda	103.36
<i>Townsendimyia</i> Prado & Fonseca, 1932	103.9
<i>transita</i> Townsend, 1917, Dexosarcophaga	103.42
<i>travassosi</i> Lopes, 1938, Nephochaetopteryx	103.28
<i>trejosi</i> Lopes, 1955, Euboettcheria	103.35
<i>Tricharaea</i> Thomson, 1869	103.31
<i>tricos</i> Reinhard, 1965, Eumacronychia	103.3
<i>tridens</i> (Wulp), 1896 (Sarcophaga), Acridiophaga	103.19
<i>tridens</i> Townsend, 1915, Punaphyto	103.41
<i>tridentata</i> (Hall), 1937 (Sarcophaga), Euboettcheria	103.35
<i>trigonomaclulata</i> Macquart, 1843 (Sarcophaga)	103.54
<i>trigonophyni</i> Blanchard, 1939, Sarcophagulopsis	103.50
<i>trilineata</i> (Wulp), 1890 (Miltogramma), Senotainia	103.5
<i>Tripanurga</i> Brauer & Bergenstamm, 1891	103.46
<i>tripartita</i> (Wulp), 1895 (Sarcophaga), Hystriocoenema	103.45
<i>triplasia</i> (Wulp), 1895 (Sarcophaga), Neobeilleria	103.51
<i>trivialis</i> (Wulp), 1895 (Sarcophaga), Oxysarcodexia	103.27
<i>trivittata</i> Macquart, 1843 (Sarcophaga)	103.54
<i>trivittata</i> (Townsend), 1912 (Sarcophaga), Amobia	103.2
<i>trivittata</i> (Townsend), 1927 (Sarcophaga), Chaetoravinia	103.23
<i>trivittatus</i> Curran, 1927, Squamatoides	103.38
<i>Trizosarcophaga</i> Townsend, 1917	103.21
<i>truncata</i> (Schiner), 1868 (Sarcophaga), Hybopygia	103.24
<i>tryporylonis</i> (Townsend), 1893 (Sarcophaga), Amobia ...	103.2
<i>tueumana</i> Blanchard, 1942, Sarcophodexiopsis	103.49
<i>Tulacopoda</i> Townsend, 1917	103.48
<i>turbata</i> Wulp, 1895 (Sarcophaga)	103.52
<i>turbata</i> Wulp of Enderlein, 1928, Dienchaeta	103.39
<i>turkensis</i> Dodge, 1965, Abapa	103.39
<i>Udamoctis</i> Enderlein, 1928	103.11
<i>Udamopyga</i> Hall, 1938	103.44

uncata (Wulp), 1896 (Sarcophaga), Acridiophaga	103.19
uncinata (Hall), 1933 (Sarcophaga), Paraphrissopoda	103.37
unica (Townsend), 1892 (Sarcomacronychia), Amobia	103.2
uniseta Lopes, 1950, Notochaeta	103.12
urceola (Hall), 1933 (Sarcophaga), Paraphrissopoda	103.37
uruhuasi (Townsend), 1917 (Raimondia), Lepidodexia	103.15
vagabunda (Wulp), 1895 (Sarcophaga), Chaetoravinia	103.23
vanvoasti Dodge, 1965 (Sarcophaga)	103.52
varia (Walker), 1837 (Sarcophaga), Hybopygia	103.24
varia (Walker) of Engel, 1931 (Sarcophaga), Hybopygia	103.23
varipes Rondani, 1850 (Sarcophaga)	103.54
vauriae Dodge, 1963, Emdenimyia	103.17
ventricosa (Wulp), 1895 (Sarcophaga), Oxysarcodexia	103.27
venusta (Enderlein), 1928 (Ctenolioproctia), Euboettcheria	103.34
vesica (Hall), 1933 (Sarcophaga), Oxysarcodexia	103.28
villegasi (Dodge), 1966 (Peckia), Paraphrissopoda	103.37
Villegasia Dodge, 1966	103.48
villipes (Wulp), 1895 (Sarcophaga), Sabinata	103.46
villipes (Wulp) of Aldrich, 1916 (Sarcophaga), Glaucosarcophaga	103.49
villipes Dodge, 1965, Harpagopyga	103.14
villosa Lopes, 1945, Oxysarcodexia	103.28
violenta (Walker), 1849 (Sarcophaga), Gigantotheca	103.47
virgata (Wiedemann), 1830 (Dexia), Pachygraphia	103.16
viridis Lopes, 1950, Notochaeta	103.12
vittata Walker, 1837 (Sarcophaga)	103.54
vittata Reinhard, 1965, Opsidia	103.4
volucris (Wulp), 1895 (Sarcophaga), Euboettcheria	103.35
Xanthobrachycoma Townsend, 1927	103.9
xanthophora (Schiner), 1868 (Sarcophaga), Oxyvinia	103.23
xanthophora (Schiner) of Mattos, 1919 (Sarcophaga), Euboettcheria	103.35
xanthophorina Dodge, 1967, Emdenimyia	103.17
xanthopyga (Wulp), 1895 (Sarcophaga), Chaetoravinia	103.23
xanthopyga (Wulp) of Enderlein, 1928 (Engelina), Andinoravinia	103.22
xanthosoma (Aldrich), 1916 (Sarcophaga), Oxysarcodexia	103.28
xanthosoma (Aldrich) of Curran & Walley, 1934 (Sarcophaga), Oxysarcodexia	103.24
Xenagria Blanchard, 1939	103.31
Xenoppia Townsend, 1915	103.10
Xylocampta Townsend, 1927	103.15
Xylocamptopsis Townsend, 1927	103.15
xagneri (Blanchard), 1939 (Sarcophodexia), Acanthodotheca ...	103.20
welchi (Hall), 1930 (Sarcophaga), Sarcodexiopsis	103.49
weyrauchi Lopes, 1954, Xanthobrachycoma	103.9
wicharti (Lopes), 1953 (Dexosarcophaga), Oxyvinia	103.23
wiedemanni (Aldrich), 1916 (Sarcophaga), Paraphrissopoda	103.36
williamsi Curran, 1934 (Sarcophaga)	103.52

<i>Winthemiola</i> Dodge, 1967	103.42
<i>Wohlfahrtiopsis</i> Townsend, 1917	103.51
<i>wygodzinskyi</i> Lopes, 1954, <i>Xanthobrachyeoma</i>	103.9
<i>zayasi</i> Dodge, 1956, <i>Oxysarcodexia</i>	103.28
<i>zetterstedti</i> (Rondani), 1856 (<i>Miltogramma</i>), <i>Hilarella</i>	103.3
<i>Zygastropyga</i> Townsend, 1917	103.47
<i>zygox</i> Hall, 1938, <i>Abacantha</i>	103.16



MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS SOUTH
OF THE UNITED STATES

104

FAMILY TACHINIDAE
(LARVAEVIDAE)J. H. GUIMARÃES¹*Museu de Zoologia
Universidade de São Paulo*

The family Tachinidae is cosmopolitan in distribution. In the Neotropical Region it is represented by an exceedingly large number of species (2,864 species in 944 genera). Because of its parasitic habits, the family has considerable interest and significance in biological control as well as in purely scientific studies of host-parasite adaptations and mechanisms of parasitism. The host preference is varied — most species are parasites of insects, but a few parasitize terrestrial isopods. The main hosts are larvae of Lepidoptera, followed by larvae and adults of Coleoptera, larvae of Hymenoptera, nymphs and adults of Hemiptera, Orthoptera, Mantodea, and other orders of insects.

The adults, ranging in size from 2 to over 20 mm, are usually very strongly bristled. They are frequently found visiting flowers and many feed particularly upon honeydew secreted by homopterous insects or upon plant secretions. Their flight is usually very rapid, and they are commonly seen flying from plant to plant or near the ground in search of a host. For general accounts of biology, habits and immature stages, see Clausen (1940), and Herting (1960). Townsend, in this "Manual of Myiology" (1934-1942), also contributed a large amount of information on the biology and immature stages of many genera. For larval anatomy see Thompson (1963a); for morphology of the genitalia see Herting (1957) and Verbeke (1962, 1963). For descriptions of puparia, the papers of Greene (1921b) and Zucka (1963) may be consulted.

The family Tachinidae is one of the largest, most difficult, and the most challenging of the families of the Oestroidea complex. It is supposed to be of relatively recent origin and still actively radiating, and its members present an extraordinary range of taxonomic characters, creating formidable problems in the definition of groups at all levels. Therefore it is not astonishing that its systematic study became the subject of continuous "overthrows" and that the various authors on this group are far from an agreement. Radically different tribal and generic classifications have been proposed by the various specialists, but the one

¹ Bolsista do Conselho Nacional de Pesquisas, Brasil.

available classification of the Neotropical genera is that of Townsend (l.e.). Because Townsend was an extreme "splitter", he affected drastically the taxonomy of the Neotropical fauna. A recent cataloguing of his life work (Arnaud, 1958), shows that he described 1,491 new generic names (largely Tachinidae), and 1,555 new trivial names. A very high proportion of these new generic names were proposed for Neotropical species, especially from Brazil and Peru. I have decided to follow Townsend's tribal arrangements, with modifications proposed by Sabrosky & Arnaud (1965):

Subfamily RHINOPHORINAE

Melanophorini

Subfamily PHASIINAE

Gymnosomatini
 Trichopodini
 Tarassini
 Phasini
 Cyllindromyini
 Leucostomatini
 Strongygastrini
 Eutherini
 Euseopollopterygini

Subfamily PROSENINAE

Oestrophasini
 Ormini
 Trixodini
 Dexillini
 Trichoprosopini
 Mylophasini
 Melloneurini
 Prosenini
 Theresini

Subfamily TACHININAE

Hypotaenini
 Macromyini
 Tachinini
 Dejeanini
 Jurinini
 Cuphoerini
 Linnaemyini
 Ernestini
 Aphriini
 Lyphini
 Neobracheini
 Germariini
 Campylochetini
 Voriini
 Hesperophasini

Subfamily DEXIINAE

Thelairini
 Dexiini
 Macquartiini
 Icellini
 Zellini
 Trichodurini
 Uramylini
 Urodexilini
 Ebenilini
 Sophiini
 Shannoniellini
 Minthoini
 Leskiini
 Telothyriini

Subfamily GONIINAE

Ophirionini
 Aeemyini
 Blondelliini
 Calodexiini
 Neominthoini
 Exoristini
 Eumasicerini
 Siphonini
 Graphogastrini
 Admontilini
 Oedematocerini
 Erythrocerini
 Gonilini
 Belvosini
 Harrisiini
 Sturmiini
 Masiphylini
 Siphosturmiini
 Careelliini
 Eryelini
 Winthemiini

This catalogue arrangement leaves much to be desired, and the position of genera and species here is merely an attempt to reflect what is believed to be the general consensus of the most authoritative workers at the present time. The work on systematics of the American muscoid flies has been seriously retarded by the impossibility of recognizing many species from the original diagnoses, which were written at a time when many important characters had not yet been discovered. Most of the types of Wulp, Walker, Bigot, Giglio-Tos, and others are still unstudied, and, at the present time, their correct generic placement and status is often a matter of guesswork, leaving much to be cleared up in the future. Except for a few cases, it seems better to recognize most of the species as distinct, until the classification of the entire genus can be thoroughly revised. Judging from the large number of Neotropical species currently being described and the many undescribed ones seen in collections, probably more than 70% are still unnamed.

Many summary citation of new synonymy and new combinations have been made in this catalogue. The primary decisions in these cases were often made by others; for instance, some of them were based on J. M. Aldrich unpublished card files and notes, while others are derived from information passed on to me by Dr. Curtis W. Sabrosky, D. M. Wood, and many others. To simplify what has been a considerable task, I have followed the custom of recent cataloguers in not trying to assign specific credit for all of these taxonomic changes. In doing this, I realize of course that the responsibility for errors in these matters must rest on myself.

Unverified distributional records in the literature are not repeated, and this report is restricted so far as possible to first hand information. For distribution of the Chilean species the papers of Cortés & Hichins (1969), may be consulted.

The preparation of this catalogue was supported by a fellowship from the John Simon Guggenheim Memorial Foundation. This task could not possibly have been brought to the present stage of completeness without full access to the U.S. National Museum collection and its associated card catalogues and library facilities. It was the author's privilege to work for an entire year with his friend Dr. Curtis W. Sabrosky, Laboratory of Systematic Entomology of U.S. Dept. of Agriculture, Washington, D.C., who unfailingly gave his time and the benefit of his experience in Tachinidae systematics, enabling the author to complete a considerable amount of work in a relatively short time. Dr. Sabrosky helped on numerous problems of classification, generic limits, nomenclature and also assisted materially with his personal notes on types in British Museum. Dr. Claude Dupuis, Muséum National d'Histoire Naturelle, Paris, was very generous for critically reading the manuscript on Phasiinae. Finally, I wish to express my gratitude to my friends Dr. George C. Steyskal, Laboratory of Systematic Entomology, U.S. Department of Agriculture, for his valuable assistance on the numerous problems of nomenclature, to Dr. D. M. Wood, Canada Department of Agriculture, Entomology Research Institute, Ottawa, for his generous help in many problems, especially in the sections on Erycinini and Blondellini, and to Dr. Raúl Cortés P., Estación Experimental Agronómica, Universidad de Chile for his generous assistance in numerous problems on Chilean tachinids.

Subfamily RHINOPHORINAE

(Melanophorinae)

Tribe MELANOPHORINI

Genus MELANOPHORA Meigen

MELANOPHORA Meigen, 1803: 279. Type-species, *Musca grossiflacionis* Linnaeus (mon.) = *roralis* (Linnaeus).

RORALIS (Linnaeus), 1758: 597 (*Musca*). Type-locality: "Europe". Distr. — Holarctic; Brazil (immigrant). Ref. — Parker, 1953: 46 (biol. immat. stages).

americana Macquart, 1843: 229 (1843: 72), pl. 8, fig. 4. Type-locality: "Brazil or Chile".

?*brasiliensis* Robineau-Desvoidy, 1863: 1138 (*Illigeria*). Type-locality: "Brazil".

asolis Gundlach, in Pazos, 1914: 1002. *Nomen nudum*.

nitidiventris Curran, 1928a: 41. Type-locality: Jamaica, Cinchona.

Subfamily PHASIINAE

(Gymnosomatinae)

The hosts of genuine representatives of subfamily Phasiinae are adult Hemiptera-Heteroptera. The Strongygastrini have been recorded as parasites of Coleoptera, with few unpublished records on Hemiptera-Heteroptera. Ref. — Dupuis, 1963 (monogr.).

Tribe GYMNOSOMATINI

Ref. — Brooks, 1946 (rev.).

Genus CISTOGASTER Latreille

CISTOGASTER Latreille, 1829: 511. Type-species, *Musca globosa* Fabricius (Blanchard, 1840: 612).

DOMINICA Curran, 1927c: 14. Type-locality: Dominican Republic, Sanchez.

INSULARIS Williston, 1896: 351. Type-locality: St. Vincent, West Indies.

Genus GYMNOSOMA Meigen

Rhodogyne Meigen, 1800: 39. Type-species, *Musca rotundata* Linnaeus (sub. mon.; Hendel, 1908: 66). Suppressed by I.C.Z.N., 1963: 339.

GYMNOSOMA Meigen, 1803: 278. Type-species, *Musca rotundata* Linnaeus (mon.).

FULIGINOSUM Robineau-Desvoidy, 1830: 237. Type-locality: "U.S.A.". Distr. — U.S.A., Mexico.

RUFICORNIS (Wulp), 1892: 187 (*Cistogaster*). Type-locality: Mexico, Tabasco.

Genus GYMNOCLYTIA Brauer & Bergenstamm

GYMNOCLYTIA Brauer & Bergenstamm, 1893: 157 (1893: 69). Type-species, *Cistogaster divisa* Loew (mon.) = *occidua* (Walker).

Procistogaster Townsend, 1934: 208. Type-species, *ferruginea* Townsend (orig. des.) = *immaculata* (Macquart).

FERRUGINOSA (Wulp), 1892: 187 (*Cistogaster*). Type-locality: Mexico, Guerrero and Tabasco.

GRISEONIGRA (Wulp), 1892: 187 (*Cistogaster*). Type-locality: Mexico, Guerrero.

HIRTICOLLIS (Wulp), 1892: 187 (*Cistogaster*). Type-locality: Mexico, Guerrero.

IMMACULATA (Macquart), 1843: 233 (1943: 76) (*Cistogaster*). Type-locality: rero. N. comb.

U.S.A., Carolina. Distr. — U.S.A., Mexico.

ferruginea Townsend, 1934: 208 (*Procistogaster*). Type-locality: "America".

MELANOSOMA (Wulp), 1892: 186 (*Cistogaster*). Type-locality: Mexico, Guerrero.

OCCIDUA (Walker), 1849: 692 (*Gymnosoma*). Type-locality: "Nova Scotia". Distr. — U.S.A., Mexico.

PAULISTA Townsend, 1929: 370. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

PROPINQUA (Wulp), 1892: 187 (*Cistogaster*). Type-locality: Mexico, Guerrero.

SUBPETIOLATA (Wulp), 1892: 186 (*Cistogaster*). Type-locality: Mexico, Guerrero.

VARIEGATA (Wulp), 1892: 187 (*Cistogaster*). Type-locality: Mexico, Guerrero.

Tribe TRICHOPODINI

The hosts of the tribe Trichopodini are adult Hemiptera-Heteroptera.
Ref. — Sabrosky, 1950 (key to genera).

Genus ACAULONA Wulp

- ACAULONA Wulp, 1884: 4. Type-species, *costata* Wulp (mon.). Ref. — Sabrosky, 1950: 369 (taxonomy).
- BRASILIANA Townsend, 1937a: 316. Type-locality: Brazil, São Paulo, Campinas. Distr. — Brazil (São Paulo), Argentina (Bonpland, Misiones, Chaco, Santiago del Estero). Ref. — Mendes, 1938: 216 (hosts).
- COSTATA Wulp, 1888: 4, pl. 3, figs. 1-1a. Type-locality: Mexico, Veracruz, Medellín. Distr. — Mexico (Veracruz, Orizaba, Tabasco, N. Yucatán). Ref. — Williston, 1908: 364 (wing, head).
- ERYTHROPYGA Sabrosky, 1950: 370. Type-locality: Puerto Rico, Guayanilla. Distr. — Puerto Rico (Guayanilla).
- PERUVIANA Townsend, 1913b: 93. Type-locality: (Peru, Plura, Chira Valley, San Jacinto) Neotypes from Peru, Rio Tambo. Distr. — Peru; Puerto Rico (introduced, not established). Ref. — Sabrosky, 1951: 210.

Genus ATRICHOPODA Townsend

- ATRICHOPODA Townsend, 1931a: 322. Type-species, *oviventris* Townsend (orig. des.).
- OVIVENTRIS Townsend, 1931a: 323. Type-locality: Peru, Oxapampa trail at Rio Paucartambo crossing.

Genus BIBIOMIMA Brauer & Bergenstamm

- BIBIOMIMA Brauer & Bergenstamm, 1889: 148 (1889: 80). Type-species, *handlirschi* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.).
- HANDLIRSCHI Brauer & Bergenstamm, 1889: 148 (1889: 80); 1891: 443 (fig.). Type-locality: Brazil, Bahia. Distr. — Panama to Brazil. Ref. — Townsend, 1916g: 22; Aldrich, 1925c: 126 (notes on types).

Genus CYLINDROPHASIA Townsend

- CYLINDROPHASIA Townsend, 1916h: 22. Type-species, *Ocyptera simillima* Fabricius (orig. des.).
- LATERALIS (Walker), 1849: 697 (*Trichopoda*). Type-locality: "Brazil". N. comb.
- OBSCURA (Bigot), 1876: 399 (*Trichopoda*). Type-locality: Argentina, Buenos Aires. N. comb.
- SIMILLIMA (Fabricius), 1805: 313 (*Ocyptera*). Type-locality: "South America". Distr. — Brazil (Amazonia, São Paulo coast, Mato Grosso). Ref. — Sabrosky, 1950: 364, fig. 7.
- TEHUANTEPECA (Townsend), 1908: 130 (*Acaulona*). Type-locality: Mexico, Tehuantepec. Ref. — Sabrosky, 1950: 370.

Genus DALLASIMYIA Blanchard

- DALLASIMYIA Blanchard, 1944: 7. Type-species, *bosqi* Blanchard (orig. des.).
- BOSQI Blanchard, 1944: 7, figs. a-d. Type-locality: Argentina, Buenos Aires, Rivadavia.

Genus EUACAULONA Townsend

- EUACAULONA Townsend, 1908: 131. Type-species, *sumichrasti* Townsend (orig. des.). Ref. — Sabrosky, 1950: 362 (in key), 368.
SUMICHRASTI Townsend, 1908: 131. Type-locality: Mexico, Tehuantepec, Distr. — Mexico (Tehuantepec), Paraguay.

Genus EUTRICHOPODA Townsend

- EUTRICHOPODA Townsend, 1908: 134. Type-species, *nigra* Townsend (orig. des.). Ref. — Sabrosky, 1950: 363 (in key), 366, fig. 3.
ABDOMINALIS Townsend, 1929: 371. Type-locality: Peru, Moyobamba. Distr. — Peru (Rio Negro), Brazil (São Paulo). Ref. — Mendes, 1938: 216 (distribution, hosts).
MELANOPUS (Robineau-Desvoidy), 1830: 284 (*Thichopoda*). Type-locality: "Brazil". Ref. — Townsend, 1931b: 85 (descr. note on type).
NIGRA Townsend, 1908: 134. Type-locality: Mexico, Tehuantepec. Distr. — Mexico (Tehuantepec), Central America, tropical South America.
NITIDIVENTRIS (Wulp), 1892: 183 (*Trichopoda*). Type-locality: Mexico, Atoyac. N. COMB.
PYRRHIOGASTER (Wiedemann), 1830: 272 (*Trichopoda*). Type-locality: "South America". Ref. — Townsend, 1931b: 85 (descr. notes).
TEGULATA (Townsend), 1897a: 29 (*Trichopoda*). Type-locality: Mexico, Veracruz. N. COMB.

Genus EUTRICHOPODOPSIS Blanchard

- EUTRICHOPODOPSIS Blanchard, 1966c: 81. Type-species, *funcbris* Blanchard (orig. des.).
BRUCHI Blanchard, 1966c: 92, pl. 2, fig. 10. Type-locality: Argentina, Córdoba, Villa Cabrera.
FUNEBRIS Blanchard, 1966c: 81, pl. 2, fig. 12. Type-locality: Argentina, Buenos Aires, Capital, Flores.
IMITANS Blanchard, 1966c: 86. Type-locality: Argentina, La Rioja.
NITENS Blanchard, 1966c: 84, pl. 2, fig. 11. Type-locality: "Argentina".
SIMILIS Blanchard, 1966c: 89, pl. 2, fig. 9. Type-locality: Argentina, Buenos Aires, Rivadavia.

Genus FORCIPOPHASIA Townsend

- FORCIPOPHASIA Townsend, 1935d: 216. Type-species, *fusca* Townsend (orig. des.).
FUSCA Townsend, 1935d: 216. Type-locality: Brazil, São Paulo, São Vicente.

Genus HOMOGENIA Wulp

- HOMOGENIA Wulp, 1892: 184. Type-species, *rufipes* Wulp (Brauer & Bergenstamm, 1894: 87; 1895: 623).
Trichopododes Townsend, 1893: 166 (unjustified nom. nov. for *Homogenia* Wulp). Type-species, *Homogenia rufipes* Wulp (aut.).
Euomogenia Townsend, 1908: 132. Type-species, *lacteata* Townsend (orig. des.).
BICOLOR (Bigot), 1876: 396 (*Trichopoda*). Type-locality: Argentina, Buenos Aires. N. COMB.

- DYSDERCI (Townsend), 1937a: 317 (*Euomogenia*). Type-locality: Brazil, São Paulo, Ribeirão Preto. Ref. — Townsend, 1938a: 204 (notes, hosts); Mendes, 1938: 216 (distribution, hosts).
- INCONSTANS (Wiedemann), 1830: 270 (*Trichopoda*). Type-locality: "Brazil".
- LACTEATA (Townsend), 1908: 132 (*Euomogenia*). Type-locality: Mexico, Tabasco, Frontera. Distr. — S. Mexico to S. Brazil and Argentina.
- LATIPENNIS Wulp, 1892: 184. Type-locality: Mexico (several localities). Ref. — Townsend, 1931b: 85 (notes on type).
- NIGROSCUTELLATA Wulp, 1892: 184. Type-locality: Mexico, Guerrero. Ref. — Townsend, 1908: 132 (notes).
- NIGRIPENNIS (Bigot), 1876: 396 (*Trichopoda*). Type-locality: Argentina, Buenos Aires. N. comb.
- RUFIPES Wulp, 1892: 184. Type-locality: "Mexico".

Genus ITAXANTHOMELANA Townsend

- ITAXANTHOMELANA Townsend, 1927a: 214. Type-species, *grandis* Townsend (orig. des.). Ref. — Sabrosky, 1950: 362 (in key), 368.
- GRANDIS Townsend, 1927a: 322. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus MELANOROPHASIA Townsend

- MELANOROPHASIA Townsend, 1934: 205. Type-species, *minuscule* Townsend (orig. des.).
- MINUSCULA Townsend, 1934: 205. Type-locality: Brazil, Pará, Urucurituba, Rio Tapajós. Distr. — Brazil (Pará). Ref. — Sabrosky, 1950: 362 (in key), 367 (nomencl.; genus restr. to lectotype).

Genus PENNAPODA Townsend

- PENNAPODA Townsend, 1897c: 282. Type-species, *Trichopoda phasiana* Townsend (orig. des.).
- PHASIANA (Townsend), 1897c: 282 (*Trichopoda*). Type-locality: Mexico, Veracruz, San Rafael, Rio Nautla. Distr. — South Mexico.

Genus PLATYPHASIA Townsend

- PLATYPHASIA Townsend, 1935d: 216. Type-species, *similis* Townsend (orig. des.).
- SIMILIS Townsend, 1935d: 216. Type-locality: Brazil, São Paulo, São Vicente.

Genus SYRINGOSOMA Townsend

- SYRINGOSOMA Townsend, 1917b: 232. Type-species, *pennipes* Townsend (orig. des.).
- PENNIPES Townsend, 1917b: 233. Type-locality: Brazil, Pará, Ananindeua, Benevides. Distr. — Brazil (Amazonas, Pará).

Genus TAPAJOSIA Townsend

- TAPAJOSIA Townsend, 1934: 205. Type-species, *urucurytuba* Townsend (orig. des.).
- URUCURYTUBA Townsend, 1934: 205. Type-locality: Brazil, Pará, Urucurituba, Rio Tapajós. Distr. — Brazil (Pará, Rio Tapajós).

Genus TRICHOPODA Berthold

TRICHOPODA Berthold, 1827: 508. Type-species, *Thercva plumipes* Fabricius (Coquillett, 1910: 616) (as "Trichopode", vernacular, by Latreille, 1825: 498).

Trichiopoda, emend.

Polistomyia Townsend, 1908: 132. Type-species, *Trichopoda trifasciata* Loew (orig. des.) = *plumipes* (Fabricius). Refs. — Sabrosky, 1950: 362 (in key), 366 (nomencl. notes).

Subgenus TRICHOPODA Berthold

ALIPES Wulp, 1892: 183. Type-locality: Mexico, Guerrero, Omilteme.

APICALIS Wiedemann, 1830: 271. Type-locality: "Brazil". Unrecognized.

CILIATA (Fabricius), 1805: 15 (*Ocyptera*). Type-locality: "South America".

DECISA Walker, 1852: 259. Type-locality: Brazil, Amazon River.

FLAVA Röder, 1885: 343. Type-locality: "Puerto Rico". Ref. — Townsend, 1891: 139 (brief description).

GRADATA Wiedemann, 1830: 275. Type-locality: "Brazil".

HAITENSIS Robineau-Desvoidy, 1830: 285. Type-locality: "Haiti".

INCOGNITA (Blanchard), 1966: 62 (*Trichopodopsis*). Type-locality: Argentina.

INDIVISA Townsend, 1897c: 281 (as *histrio* var.). Type-locality: Mexico, Veracruz, San Rafael.

LANIPES (Fabricius), 1805: 220 (*Thereva*). Type-locality: U.S.A., Carolina. Distr. — U.S.A., Mexico.

tropicalis Townsend, 1897c: 275, 278. Type-locality: Mexico, Veracruz, San Rafael.

LUTEIPENNIS Wiedemann, 1830: 269. Type-locality: "Brazil".

MEXICANA Macquart, 1846: 300 (1846: 172). Type-locality: "Mexico".

NIGRICAUDA Bigot, 1876: 395. Type-locality: "Mexico".

NIGRIPES Wulp, 1892: 184. Type-locality: Mexico, Guerrero, Xucumanatlán and Omilteme.

PICTIPENNIS Bigot, 1876: 398. Type-locality: "South America".

PILIPES (Fabricius), 1805: 220 (*Thercva*). Type-locality: "South America". Ref. — Townsend, 1931b: 84 (notes on type).

Piogaster Pazos, 1914: 1002. *Nomen nudum*.

PENNIPES (Fabricius), 1781: 450 (*Musca*). Type-locality: "North America". Distr. — U.S.A., Mexico. Ref. — Beard, 1940: 624 (biology, immature stages).

SQUAMPES Wulp, 1892: 184. Type-locality: Mexico, Guerrero.

SUBALIPES Townsend, 1897a: 30 (nom. nov. for *subcilipes* Townsend). Type-locality: Dominican Republic, San Domingo.

subcilipes Townsend, 1894: 78 (preocc. Macquart, 1843). Type-locality: Dominican Republic, San Domingo.

SUBCILIPES Macquart, 1843: 243 (1843: 77), pl. 9, fig. 2. Type-locality: "Gulana".

UMBRA Walker, 1849: 698. Type-locality: "Venezuela".

Subgenus GALACTOMYIA Townsend

GALACTOMYIA Townsend, 1908: 135 (as genus). Type-species, *Trichopoda radiata* Loew (orig. des.) = *lanipes* (Fabricius).

- Trichopodopsis* Townsend, 1913: 148, 313. Type-species, *Musca pennipes* Fabricius (orig. des.).
- ARGENTINENSIS (Blanchard), 1966c: 65, pl. 1, fig. 5. (*Trichopodopsis*). Type-locality, Argentina, Córdoba.
- BOSQI (Blanchard), 1966c: 73, pl. 1, fig. 1. (*Trichopodopsis*). Type-locality: Argentina, Buenos Aires, Cañuelas.
- CHRISTENSENI (Blanchard), 1966c: 78, pl. 1, fig. 2. (*Trichopodopsis*). Type-locality: Argentina, José C. Paz.
- GIACOMELLI (Blanchard), 1966c: 75, pl. 2, fig. 7. (*Trichopodopsis*). Type-locality: Argentina, La Rioja.
- LEMBATA (Blanchard), 1966c: 67, pl. 1, fig. 4. (*Trichopodopsis*). Type-locality: Argentina, Buenos Aires, Puente Alsina.
- NIGRIFRONTALIS (Blanchard), 1966c: 70, pl. 1, fig. 3. (*Trichopodopsis*). Type-locality: Argentina, Buenos Aires, Castelar.

Genus URUCURMYIA Townsend

- URUCURMYIA Townsend, 1934: 206. Type-species, *urna* Townsend (orig. des.). Ref. — Sabrosky, 1950: 362 (in key), 367.
- URNA Townsend, 1934: 206. Type-locality: Brazil, Pará, Urucurituba, Rio Tapajós.

Genus XANTHOMELANODES Townsend

- Xanthomelana* Wulp, 1892: 188 (preocc. Waterhouse, 1889). Type-species, *gracilentia* Wulp (Coquillett, 1910: 620).
- XANTHOMELANODES Townsend, 1893: 167 (nom. nov. for *Xanthomelana* Wulp). Type-species, *Xanthomelana gracilentia* Wulp (aut.).
- Erythrophasia* Townsend, 1917c: 127. Type-species, *atripennis* Townsend (orig. des.) = *atripennis* (Say).
- ARCUATUS (Say), 1829: 173 (1859: 360) (*Ocyptera*). Type-locality: U.S.A., Indiana. Distr. — U.S.A., Mexico.
- ATRIFRONS (Wiedemann), 1824: 46 (*Melanophora*). Type-locality: "South America". Ref. — Townsend, 1931b: 85.
- ATRIPENNIS (Say), 1829: 172 (*Phasia*). Type-locality: U.S.A., Indiana. Distr. — U.S.A., also? Mexico. Ref. — Sabrosky, 1950: 363 (key), 365, fig. 10.
- BRASILIENSIS Townsend, 1929: 371. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Ref. — Sabrosky, 1950: 365 (lectotype des.).
- CALIFORNICUS Townsend, 1908: 129. Type-locality: U.S.A., California. Distr. — U.S.A., Mexico.
- DIAPHANUS (Fabricius), 1805: 314 (*Ocyptera*). Type-locality: "South America". Ref. — Townsend, 1931b: 84 (notes on type).
- DORSALIS (Wulp), 1892: 188 (*Xanthomelana*). Type-locality: Mexico, Guerrero and Veracruz. N. comb.
- GRACILENTUS (Wulp), 1892: 189 (*Xanthomelana*). Type-locality: Mexico, Guerrero, Rincon.
- MUTATUS (Wiedemann), 1830: 338 (*Tachina*). Type-locality: "South America".
- RUBICUNDUS (Wulp), 1892: 188 (*Xanthomelana*). Type-locality: Mexico, Tabasco, Teapa. N. comb.
- TRIVITATUS Reinhard, 1955a: 233. Type-locality: Mexico, Nayarit, 15 km N. of Chapalilla.

Genus XANTHOMELANOPSIS Townsend

- XANTHOMELANOPSIS Townsend, 1917c: 126. Type-species, *Xanthomelanodes peruanus* Townsend (orig. des.). Ref. — Sabrosky, 1950: 362 (in key), 367.
- BRASILIENSIS Townsend, 1917b: 233. Type-locality: Brazil, Pará, Ananindeua, Benevides.
- PERUANA (Townsend), 1911: 128 (1912: 302) (*Xanthomelanodes*). Type-locality: Peru, Piura.
- TRIGONALIS (Wulp), 1892: 188 (*Xanthomelana*). Type-locality: Mexico, Guerrero, Xucumanatlán. N. COMB.

Genus XENOPHASIA Townsend

- XENOPHASIA Townsend, 1934: 207. Type-species, *xanthomelanoides* Townsend (orig. des.).
- ARTICULATA (Wulp), 1892:188 (*Xanthomelanodes*). Type-locality: Mexico, Orizaba and Veracruz. Ref. — Sabrosky, 1950: 363.
- XANTHOMELANOIDES Townsend, 1934: 207. Type-locality: Brazil, Pará, Urucurituba, Rio Tapajós.

Tribe TARASSINI

The genus *Tarassus* Aldrich has a doubtful systematic position among the Calyptratae. Aldrich (1934) stated that this genus may represent a new subfamily near *Trichopoda*. Later, Townsend (1935: 92), arguing that it was an aberrant acalyptrate fly established a family for it. According to Dr. Hennig (personal communication) *Tarassus* is an anomalous Tachinidae. Morphologically this genus appears to merit at least tribal status and is left here until its proper position can be established.

Genus TARASSUS Aldrich

- TARASSUS Aldrich, 1933a: 438. Type-species, *shannoni* Aldrich (orig. des.).
- SHANNONI Aldrich, 1933a: 440, figs. 1-2. Type-locality: Brazil, Mato Grosso.

Tribe PHASIINI

The species of Phasiini are parasites of adult Hemiptera-Heteroptera.

Genus ALOPHORELLA Townsend

- ALOPHORELLA Townsend, 1912c: 45. Type-species, *Thercva obesa* Fabricius (orig. des.).
- BRASILIANA Townsend, 1929: 370. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- BRASILIENSIS (Townsend), 1938c: 347 (*Alophoropsis*). Type-locality: Brazil, São Paulo, Tietê and Sorocaba. N. COMB.
- LYSDERCI (Townsend), 1938d: 347 (*Euphorantha*). Type-locality: Brazil, São Paulo, Sorocaba. N. COMB.
- LATFENNIS (Brauer), 1898: 509 (*Hyalomya*). Type-locality: Uruguay, Montevideo. N. COMB.

- METALLICA (Aldrich), 1934: 15 (*Hyalomya*). Type-locality: Chile, Llanquihue, Casa Pangué. N. comb.
- MOERENS (Wulp), 1892: 186 (*Hyalomya*). Type-locality: Mexico, Guerrero, Omitlteme. Ref. — Townsend, 1893: 166 (note). N. comb.
- NIGRENS (Wulp), 1892: 185 (*Hyalomya*). Type-locality: Mexico, Guerrero. N. comb.
- PICEPES (Wulp), 1892: 186 (*Hyalomya*). Type-locality: Mexico, Guerrero, Omitlteme. N. comb.
- subcinerea* (Lynch Arribáizaga) in Brauer, 1898: 509 (*Hyalomya*). *Nomen nudum*.
- VIOLACEIVENTRIS (Brauer), 1898: 509 (*Hyalomya*). Type-locality: Uruguay, Montevideo. N. comb.

Genus ANDROEURYOPS Beneway

- ANDROEURYOPS Beneway, 1961: 44. Type-species, *Hyalomyia ecitonis* Townsend (orig. des.).
- ECTONIS (Townsend), 1897a: 30 (*Hyalomyia*). Type-locality: Mexico, Veracruz, Paso de Telaya, Distr. — Mexico, Panama. Ref. — Rettenmeyer, 1961: 1018 (association with swarm-raids of army ants).

Genus ECTOPHASIOPSIS Townsend

- ECTOPHASIOPSIS Townsend, 1915m: 439. Type-species, *chilensis* Townsend (orig. des.) = *arcuata* (Blgot).
- ARCUATA (Blgot), 1876: 397 (*Trichopoda*). Type-locality: "Chile". Ref. — Aldrich, 1934: 11.
- chilensis* Townsend, 1915m: 440. Type-locality: "Chile".

Genus EPAULOPHASIA Townsend

- EPAULOPHASIA Townsend, 1934: 207. Type-species, *officialis* Townsend (orig. des.).
- OFFICIALIS Townsend, 1934: 207. Type-locality: Brazil, Pará, Urucurituba, Rio Tapajós.

Genus HEYNEOPHASIA Townsend

- HEYNEOPHASIA Townsend, 1934: 208. Type-species, *heynei* Townsend (orig. des.).
- ARGENTICEPS (Wulp), 1892: 185 (*Hyalomyia*). Type-locality: Mexico, Guerrero, Xucumanatlán. N. comb.
- HEBES (Wulp), 1892: 185 (*Hyalomyia*). Type-locality: "Mexico". N. comb.
- HEYNEI Townsend, 1934: 208. Type-locality: Costa Rica, Surrubres. Distr. — Mexico to northern Argentina.
- MUNDA (Wulp), 1892: 185 (*Hyalomyia*). Type-locality: Mexico, Guerrero, Rincón. N. comb.
- OCIIRICEPS (Wulp), 1892: 185 (*Hyalomyia*). Type-locality: Mexico, Guerrero, Rincón. Ref. — Townsend, 1893: 166 (note). N. comb.

Genus HYALOMYA Robineau-Desvoidy

HYALOMYA Robineau-Desvoidy, 1830: 298. Type-species, *Phasia semicinerea* Meigen (Westwood, 1840: 140) = *pusilla* (Meigen).

Hyalomyia, emend.

Allophorellopsis Townsend, 1927a: 209. Type-species, *capitata* Townsend (orig. des.).

ALDRICH Townsend, 1891a: 136. Type-locality: U.S.A., South Dakota. Distr. — Alaska, Canada, U.S.A., Mexico.

ARGENTIFRONS (Brooks), 1945: 675 (*Allophorellopsis*). Type-locality: Canada, British Columbia. Distr. — U.S.A., Canada, Mexico.

CAPITATA (Townsend), 1927a: 284 (*Allophorellopsis*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.

CHILENSIS Macquart, 1851: 189 (1851: 216), pl. 20, fig. 4. Type-locality: "Chile". Distr. — Puerto Rico (introduced, not established), Peru, Uruguay, Argentina, Chile. Ref. — Aldrich, 1934: 14; Berry, 1951: 340 (biol. notes); Cortés, 1963: 249 (notes on type).

CURVIPES Aldrich, 1934: 16, fig. 3. Type-locality: Chile (sev. places).

FREYREISH (Wiedemann), 1830: 263 (*Phasia*). Type-locality: "Brazil". Unrecognized. N. comb.

GLAUCA Aldrich, 1934: 16. Type-locality: Argentina, Río Negro, Bariloche.

MICANS (Wulp), 1883: 14, pl. 1, fig. 12 (*Allophora*). Type-locality: "Argentina". Unrecognized.

PURPURASCENS Townsend, 1891a: 137. Type-locality: U.S.A., Illinois. Distr. — Canada, U.S.A., Mexico.

UMBRIFERA Wulp, 1892: 186 (*Hyalomyia*). Type-locality: Mexico, Guerrero. Ref. — Townsend, 1893: 166 (notes).

UMBROSA Wulp, 1892: 186 (*Hyalomyia*). Type-locality: Mexico, Guerrero, Omilteme.

VILLOSA Wulp, 1892: 185 (*Hyalomyia*). Type-locality: Mexico, Orizaba.

VIOLASCENS Townsend, 1897: 32 (*Hyalomyia*). Type-locality: "Mexico".

Genus PARAPHASIANA Townsend

PARAPHASIANA Townsend, 1940b: 889. Type-species, *dysderci* Townsend (orig. des.).

DYSDERCI Townsend, 1940b: 889. Type-locality: Brazil, São Paulo, Campinas and Sorocaba.

Genus PARAPHORANTHA Townsend

PARAPHORANTHA Townsend, 1915b: 20. Type-species, *Allophora grandis* Coquillett (orig. des.).

AURICAUDATA Brooks, 1945b: 661. Type-locality: U.S.A. Distr. — U.S.A., Mexico.

BRASILIANA Townsend, 1937a: 317. Type-locality: Brazil, São Paulo, Campinas. Ref. — Mendes, 1938: 216 (hosts).

DIMIDIATA Townsend, 1937a: 318. Type-locality: Brazil, São Paulo, Tietê and Campinas. Ref. — Mendes, 1938: 216 (hosts).

PERUVIANA Townsend, 1936c: 489. Type-locality: Peru, Pacasmayo, Jequetepeque.

POLITANA Townsend, 1938c: 347. Type-locality: Brazil, São Paulo, Sorocaba. Ref. — Mendes, 1938: 216 (host).

Genus PHORANTHELLA Brooks

- Phoranthella* Townsend, 1915b: 23. *Nomen nudum* (Opinion 205, I.C.Z.N., 1954: 311).
Phoranthella Townsend, 1936a: 58 (also, 1938b: 68). *Nomen nudum*.
PHORANTHELLA Brooks, 1945b: 672. Type-species, *morrisoni* Brooks (mon.) = *punctigera* (Townsend).
MENDESI Townsend, 1938c: 348. Type-locality: Brazil, São Paulo, Sorocaba and Campinas. Ref. — Mendes, 1938: 216 (host).

Genus TRICHOPIHASIA Townsend

- TRICHOPIHASIA Townsend, 1939c: 447. Type-species, *transita* Townsend (orig. des.).
TRANSITA Townsend, 1939c: 448. Type-locality: Brazil, Mato Grosso, Maracaju.

Genus XANTHOTRICHUS Townsend

- XANTHOTRICHUS Townsend, 1934a: 209. Type-species, *xenos* Townsend (orig. des.).
XENOS Townsend, 1934a: 209. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus NIPHOPHASIA Townsend

- NIPHOPHASIA Townsend, 1937b: 116. Type-species, *ushpayacua* Townsend (mon.).
USHPAYACUA Townsend, 1937b: 117. Type-locality: Peru, Río Ushpayacu.

Tribe CYLINDROMYIINI

The species of Cylindromyini are parasites of adult Hemiptera-Heteroptera.

Genus ARGYROMIMA Brauer & Bergenstamm

- ARGYROMIMA Brauer & Bergenstamm, 1889: 140 (1889: 72). Type-species, *mirabilis* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.).
MIRABILIS Brauer & Bergenstamm, 1889: 140 (1889: 72). Type-locality: "South America". Ref. — Aldrich, 1925c: 122, Townsend, 1931b: 87 (notes on type).

Genus BESKIA Brauer & Bergenstamm

- BESKIA Brauer & Bergenstamm, 1889: 139 (1889: 71). Type-species, *cornuta* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.) =? *aclops* Walker.
Ocypterosipha Townsend, 1894: 79. Type-species, *willistoni* Townsend (orig. des.).
AELOPS (Walker), 1849: 796 (*Tachina*). Type-locality: U.S.A., Georgia. Distr. — U.S.A., Mexico.

? *cornuta* Brauer & Bergenstamm, 1889: 139 (1889: 71), fig. 276.
Type-locality: "Brazil". Ref. — Aldrich, 1925e: 120; Lima, 1935
(host).

? *willstoni* Townsend, 1894: 796 (*Ocypterosipha*). Type-locality:
West Indies, Dominican Republic, St. Domingo.

Genus CATOCYPTERA Townsend

CATOCYPTERA Townsend, 1927a: 215. Type-species, *brasiliiana* Townsend
(orig. des.).

BRASILIANA Townsend, 1927a: 295. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus CLINOGASTER Wulp

CLINOGASTER Wulp, 1892: 189. Type-species, *notabilis* Wulp (mon.).

NOTABILIS Wulp, 1892: 189. Type-locality: Mexico, Guerrero. Ref. —
Townsend, 1931b: 86.

Genus CYLINDROMYIA Meigen

CYLINDROMYIA Meigen, 1803: 279. Type-species, *Musca brassicaria* Fabricius
(mon.).

Aplnocyptera Townsend, 1915k: 94. Type-species, *signata* Townsend (orig.
des.).

Odontocyptera Townsend, 1915g: 233. Type-species, *nana* Townsend (orig.
des.).

Melanocyptera Townsend, 1927a: 215. Type-species, *carinata* Townsend
(orig. des.). N. SYN.

Ocypteryx Townsend, 1931a: 326 (preocc. Leach, 1817: 162). Type-species,
ochrescens Townsend (orig. des.). N. SYN.

Dolichocyptera Townsend, 1931a: 325. Type-species, *pirioni* Townsend
(orig. des.).

Ecatocypteros Townsend, 1935d: 217. Type-species, *ater* Townsend (orig.
des.) = *townsendi* Guimarães. N. SYN.

Ocyptera, authors, not Latreille.

Refs. — Aldrich, 1926e (rev.), Aldrich, 1934: 8 (key to Pata-
gonian spp.).

ALDRICHI Cortés, 1944c: 178. Type-locality: "Chile".

APICALIS (Bigot), 1878: 45 (*Ocyptera*). Type-locality: "Chile". Ref. —
Cortés, 1943: 178.

ATRA (Röder), 1885: 314 (*Ocyptera*). Type-locality: "Puerto Rico". Distr.
— U.S.A., Mexico, Puerto Rico. Ref. — Aldrich, 1926e: 19.

ATRICAUDA Aldrich, 1934: 10. Type-locality: Chile, Llalilal. Ref. — Cortés,
1944c: 179.

BAKERI Aldrich, 1926e: 19, fig. 5. Type-locality: Cuba, Havana.

CARINATA (Townsend), 1927a: 327 (*Melanocyptera*). Type-locality: Brazil,
São Paulo, Itaquaquecetuba. N. COMB.

DECORA Aldrich, 1926e: 21. Type-locality: U.S.A., New Hampshire. Distr.
— U.S.A., Jamaica.

DORSALIS (Wiedemann), 1830: 264 (*Ocyptera*). Type-locality: "Brazil". Ref.
— Aldrich, 1929a: 30.

DOTATAS (Walker), 1849: 694 (*Ocyptera*). Type-locality: West Indies,
Jamaica. Unrecognized.

- MINOR (Röder), 1885: 344 (*Ocyptera*). Type-locality: "Puerto Rico". Ref. — Curran, 1928c: 114.
- NANA (Townsend), 1915g: 233 (*Odontocyptera*). Type-locality: Mexico, Chihuahua, Río Pedras Verdes, Distr. — U.S.A., Mexico.
- NIGRINA (Wulp), 1883: 15 (*Ocyptera*). Type-locality: "Argentina".
- OBSCURA (Bigot), 1878: 44 (*Ocyptera*). Type-locality: "Brazil".
- OECHRESCENS (Townsend), 1931a: 327 (*Ocypteryx*). Type-locality: Paraguay, Santa Trinidad, Distr. — Peru (Lima), Brazil (Guanabara), Paraguay (Santa Trinidad) N. comb.
- PIRIONI (Townsend), 1931a: 326 (*Dolichocyptera*). Type-locality: Chile, San Cristóbal.
- PORTERI (Bréthes), 1925: 208 (*Ocyptera*). Type-locality: Chile, Las Condes.
- PROPUSILLA Sabrosky & Arnaud, 1965: 973. Type-locality: U.S.A., Virginia, Distr. — U.S.A., Mexico.
- SIGNATA (Townsend), 1915h: 94 (*Apinocyptera*). Type-locality: Guatemala, Gualán.
- SIGNATIPENNIS (Wulp), 1892: 187 (*Ocyptera*). Type-locality: "Mexico".
- SIMPLEX (Bigot), 1878: 47 (*Ocyptera*; preocc. Fallén, 1820). Type-locality: Mexico. Unrecognized.
- SOROR (Bigot), 1878: 46 (*Ocyptera*). Type-locality: Mexico. Unrecognized.
- STERNALIS Reinhard, 1955a: 237. Type-locality: Mexico, Nayarit, Ahuacatlán.
- TOWNSENDI Guimarães, nom. nov. for *atra* Townsend. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- ater* Townsend, 1935a: 217 (*Ecatocypteropteryx*; preocc. Röder, 1885). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- UNIFORMIS Aldrich, 1926c: 24, fig. 4. Type-locality: Mexico, D.F., San Jacinto.

Genus GLOSSIDIONOPHORA Bigot

- GLOSSIDIONOPHORA Bigot, 1885c: lv. Type-species, *nigra* Bigot (Townsend, 1916: 7).
- NIGRA Bigot, 1885c: lv. Type-locality: Argentina, Buenos Aires, Distr. — Argentina, Buenos Aires.

Genus HEMYDA Robineau-Desvoidy

- HEMYDA Robineau-Desvoidy, 1830: 226. Type-species, *aurata* Robineau-Desvoidy (mon.).
- Ancylogaster* Bigot, 1884b: lxx. Type-species, *armatus* (Bigot) (mon.). = *aurata* Robineau-Desvoidy.
- AURATA Robineau-Desvoidy, 1830: 226. Type-locality: U.S.A., Philadelphia, Distr. — U.S.A., Mexico. Ref. — Brauer, 1898: 15 (notes on syn.).
- armatus* Bigot, 1884b: lxx (*Ancylogaster*). Type-locality: "Mexico".
- DECUMANA Reinhard, 1958a: 235. Type-locality: Mexico, Veraacruz, Río Blanco.
- ZONULA Reinhard, 1951a: 236. Type-locality: Mexico, Orizaba.

Genus NEOLOPHOSIA Townsend

- NEOLOPHOSIA Townsend, 1939b: 253. Type-species, *shannoni* Townsend (orig. des.).
- SHANNONI Townsend, 1939b: 254. Type-locality: Brazil, Rio de Janeiro, Guanabara.

Genus PENTHOSIA Wulp

PENTHOSIA Wulp, 1892: 189. Type-species, *Scopolia satanica* Bigot (mon.).
SATANICA (Bigot), 1889: 254 (*Scopolia*). Type-locality: "Mexico".

Genus PHASIOCYPTERA Townsend

PHASIOCYPTERA Townsend, 1927a: 215. Type-species, *punctata* Townsend (orig. des.).
PUNCTATA Townsend, 1927a: 346. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus POLISTIOPSIS Townsend

POLISTIOPSIS Townsend, 1915j: 43. Type-species, *mima* Townsend (orig. des.). Ref. — Arnaud, 1966 (rev.).
MIMA Townsend, 1915j: 44. Type-locality: Mexico, Tehuantepec.
WILLIAMSII Arnaud, 1966: 7, figs. 4-6, 12-14, 17-20. Type-locality: Paraguay, Paso Yobay.

Genus TACHINOPHASIA Townsend

TACHINOPHASIA Townsend, 1931a: 323. Type-species, *transita* Townsend (orig. des.).
TRANSITA Townsend, 1931a: 324. Type-locality: Brazil, Santa Catarina, Blumenau.

Tribe LEUCOSTOMATINI

The hosts of Leucostomatini are adult Hemiptera-Heteroptera.

Genus CLAIRVILLIA Robineau-Desvoidy

CLAIRVILLIA Robineau-Desvoidy, 1830: 234. Type-species, *Ocyptera pusilla* Meigen (mon.; misident.) = *biguttata* Meigen.
CUEHALIS Reinhard, 1958a: 235. Type-locality: Mexico, Cuernavaca. Distr. — Mexico (Cuernavaca, Morelos).
FURCATA (Wulp), 1890: 131, pl. 3, fig. 22, 22a (*Labidigaster*). Type-locality: Mexico, Veraacruz, Atoyac. N. COMB.

Genus LEUCOSTOMA Meigen

LEUCOSTOMA Meigen, 1803: 279. Type-species, *Ocyptera simplex* Fallén (sub. mon., Meigen, 1824: 243).
Paradionaea Townsend, 1916e: 631. Type-species, *Leucostoma atra* Townsend (orig. des.) = *simplex* (Fallén).
Neopsalida Townsend, 1916e: 632. Type-species, *Leucostoma neomexicana* Townsend (orig. des.) = *aterrimum* (Villers).
Siphopsalida Townsend, 1915m: 439. Type-species, *meridionalis* Townsend (orig. des.).
ATERRIMUM (Villers), 1789: 548 (*Musca*). Type-locality: Europe. Distr. — U.S.A., Mexico, Puerto Rico, Chile. Ref. — Reinhard, 1965b: 160.
BRASILIANUM (Townsend), 1938a: 204 (*Parapsalida*). Type-locality: Brazil, São Paulo, Campinas.



- GRAVIPES Wulp, 1890: 207. Type-locality: "Mexico". Distr. — Europe; U.S.A., Mexico.
- MERIDIONALE (Townsend), 1915m: 439 (*Siphopsalida*). Type-locality: Peru, San Diego, Casma.
- NIMIRUM Reinhard, 1956b: 163. Type-locality: Cuba, Havana. Distr. — West Indies, Brazil.
- PECCATOR Reinhard, 1956b: 167. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- PERUVIANUM Townsend, 1928c: 159. Type-locality: Peru, Chancal, Esquivel.
- SIMPLEX (Fallén), 1820: 8 (*Ocyptera*). Type-locality: "Sweden". Distr. — Holarctic, Neotropical. Ref. — Reinhard, 1956b: 159.
- VEGETUM Reinhard, 1956b: 163. Type-locality: Peru, Sullana.

Genus VANDERWULFELLA Townsend

- VANDERWULFELLA Townsend, 1919b: 181. Type-species, *Xanthomelana anceps* Wulp (orig. des.).
- ANCEPS (Wulp), 1892: 189 (*Xanthomelana*). Type-locality: Mexico, Veracruz, Atoyac.

Tribe STRONGYGASTRINI

The members of the tribe Strongygastrini are parasites of adult Coleoptera, but few have been reared from Hemiptera-Heteroptera.

Genus HYALOMYODES Townsend

- HYALOMYODES Townsend, 1893: 429. Type-species, *weedi* Townsend (mon.) = *triangulifer* (Loew). Ref. — Brooks, 1942 (key); Thompson, 1954 (biol.).
- ARGENTINENSIS Blanchard, 1942a: 358, fig. 9. Type-locality: Argentina, Buenos Aires.
- BRASILIENSIS Townsend, 1929: 320. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Ref. — Mendes, 1938: 216 (distr., host).
- TRIANGULIFER (Loew), 1863: 319 (Cent. 4, n° 85) (*Hyalomyia*). Type-locality: U.S.A., New York. Distr. — U.S.A., Mexico, Chile.

Tribe EUTHERINI

The species of Eutherini are parasites of adult Pentatomidae. Refs. — Bezzi, 1925 (key), Sabrosky, 1965 (hosts).

Genus EUTHERA Loew

- EUTHERA Loew, 1866: 45 (Cent. 7, n° 85). Type-species, *tentatrix* Loew (mon.).
- BARBIELLINII Bezzi, 1925: 277, fig. 1. Type-locality: Brazil, near São Paulo. Distr. — Brazil (São Paulo, Minas Gerais).
- BICOLOR Coquillett, 1902b: 114. Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico.

Tribe EUSCOPOLIOPTERYGINI

Genus EUSCOPOLIOPTERYX Townsend

EUSCOPOLIOPTERYX Townsend, 1917b: 223. Type-species, *nebulosa* Townsend (orig. des.) = *externa* (Fabricius).

EXTERNA (Fabricius), 1805: 329 (*Dictya*). Type-locality: "South America". — Distr. — Surinam to Mato Grosso. Ref. — Townsend, 1931b: 87 (notes on type).

nebulosa Townsend, 1917b: 224. Type-locality: Brazil, Mato Grosso, Cuiabá River.

Genus SHANNONOMYIELLA Townsend

SHANNONOMYIELLA Townsend, 1939b: 252. Type-species, *ortalidoptera* Townsend (orig. des.).

ORTALIDOPTERA Townsend, 1939b: 253. Type-locality: Brazil, Rio de Janeiro, Guanabara.

UNRECOGNIZED SPECIES OF PHASIINAE

FLAVOCALYPTRATA Bigot, 1889: 255 (*Stercenia*). Type-locality: "Mexico".

PALLIDIVENTRIS Bigot, 1889: 255 (*Stercenia*). Type-locality: "Mexico".

Subfamily PROSENINAE

(Dexiinae, of authors)

Members of the subfamily Proseninae are chiefly parasites of Coleoptera larvae, but some parasitize certain lepidopterous larvae and nocturnal Orthoptera.

Tribe OESTROPHASIINI

(Glaurocarini, in part)

Townsend (1936: 94, 97) erected the tribe Glaurocarini based on *Glaurocara* Thomson from Mauritius including 5 American genera, one from Cape of Good Hope and one from Malaya.

The recent studies of Crosskey (1965) suggest that the genus *Glaurocara* Thomson must be very close phylogenetically to the Ormiini in view of the reproductive habits and the structure of the first instar larvae.

The New World genera assigned to Glaurocarini *sensu* Townsend form a well defined group based on the fundamental similarity in the morphology of the eggs and also of the larvae (cf. Thompson, 1963a: 455, 458, 459). The hosts and biology of this tribe are practically unknown. In the collection of the U.S. National Museum there is a single record of *Cenosoma signiferum* as parasite of *Colaspis pini* (Col., Chrysomelidae, from Louisiana, U.S.A.). In South America, *Euoestrophasia aperta* B. & B. was recorded in Argentina by Parker, *et al.* (1950: 25, fig. 7) as parasite of an adult curculionid of the genus *Listroderes*.

Genus CENOSOMA Wulp

CENOSOMA Wulp, 1890: 44 (key). Type-species, *signiferum* Wulp (sub. mon.; Wulp, 1890: 167).

SIGNIFERUM Wulp, 1890: 167, pl. 4, fig. 13. Type-locality: Mexico, Guerrero, Chilpancingo and Veraacruz, Atoyac. Distr. — Canada, U.S.A., Mexico. Ref. — Thompson, 1963a: 455.

Genus EUOESTROPHASIA Townsend

EUOESTROPHASIA Townsend, 1892f: 133. Type-species, *Oestrophasia aperta* Brauer & Bergenstamm (mon.).

APERTA (Brauer & Bergenstamm), 1889: 146 (1889: 78) (*Oestrophasia*). Type-locality: Brazil, Rio Grande do Sul. Ref. — Parker, *et al.*, 1950: 25 (biol. notes, immature stages).

Genus JAMACARIA Curran

JAMACARIA Curran, 1928a: 42. Type-species, *albofenestrata* Curran (orig. des.).

Jamalcaria, error.

ALBOFENESTRATA Curran, 1928a: 42. Type-locality: Jamaica, Cinchona.

Genus MESEMBRINORMIA Townsend

MESEMBRINORMIA Townsend, 1931a: 321. Type-species, *pertyi* Townsend (orig. des.).

PERTYI Townsend, 1931a: 322. Type-locality: Brazil, Guanabara, Rio de Janeiro (as "Sebastianopolis").

Genus OESTROPHASIA Brauer & Bergenstamm

OESTROPHASIA Brauer & Bergenstamm, 1889: 145 (1889: 77). Type-species, *clausa* Brauer & Bergenstamm (Townsend, 1892f: 133).

CALVA Coquillett, 1902b: 109. Type-locality: U.S.A., Arizona. Distr. — U.S.A., Mexico.

Genus OESTROPHASIANA Townsend

OESTROPHASIANA Townsend, 1931a: 320. Type-species, *Dictya uncani* Fabricius (orig. des.).

UNCANI (Fabricius), 1805: 330 (*Dictya*). Type-locality: "South America".

Tribe ORMIINI

The species of Ormini are parasites of nocturnal Orthoptera.

Genus EUPHASIOPTERYX Townsend

EUPHASIOPTERYX Townsend, 1915b: 23. Type-species, *Phasiopteryx australis* Townsend (orig. des.). Ref. — Sabrosky, 1953 (rev.); Nutting, 1953 (biol.); Tavares, 1964, 1965a, 1965b (Brazilian spp.).

ALDEICHI (Séguy), 1925: 439 (also, 1926c: 4, figs. 4-5) (*Ormia*). Type-locality: French Guiana, La Forestière, upper Maroni River. Ref. — Sabrosky, 1953: 303.

- AUSTRALIS (Townsend), 1911b: 136, 149 (1912a: 352) (*Phasiopteryx*). Type-locality: Peru, Piura. Distr. — Peru, Brazil (Pará, Mato Grosso, Guanabara, São Paulo).
- DEPLETA (Wiedemann), 1830: 289 (*Tachina*). Type-locality: "Brazil". Ref. — Tavares, 1965: 15.
- DOMINICANA (Townsend), 1919a: 548 (*Ormia*). Type-locality: Dominican Republic. Santo Domingo, San Francisco Mts.
- GUIANICA (Curran), 1934c: 496 (*Ormia*). Type-locality: Guyana, Bartica, Kartabo.
- NOCTURNA (Curran), 1934c: 495 (*Ormia*). Type-locality: Guyana, Bartica, Kartabo.
- OCHIRACEA (Bigot), 1889a: 268 (*Pyrrosia*). Type-locality: "Mexico". Distr. — U.S.A., Mexico, Cuba, Puerto Rico, Brazil and Paraguay.
- PELLUCIDA (Séguy), 1925a: 440 (*Ormia*). Type-locality: Unknown. Unrecognized.
- ROSENOI Tavares, 1965a: 22, figs. 30-40. Type-locality: Brazil, Amazonas, Manaus.
- TARSALIS (Séguy), 1925a: 440 (*Ormia*). Type-locality: French Guiana, Kourou River (cf. Séguy, 1927a: 424). Unrecognized.

Genus OPSOPHASIOPTERYX Townsend

- OPSOPHASIOPTERYX Townsend, 1917b: 222. Type-species, *mima* Townsend (orig. des.).
- MIMA Townsend, 1917b: 223. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).

Genus ORMIA Robineau-Desvoidy

- ORMIA Robineau-Desvoidy, 1830: 428. Type-species, *punctata* Robineau-Desvoidy (mon.).
- Phasiopteryx* Brauer & Bergenstamm, 1889: 147 (1889: 79). Type-species, *bilimekii* Brauer & Bergenstamm (Coquillett, 1910: 588).
- Neoptera* Wulp, 1890: 44 (key). Type-species, *rufa* Wulp (sub. mon.; Wulp 1890: 165).
- BILIMEKII (Brauer & Bergenstamm), 1889: 147 (1889: 79) (*Phasiopteryx*). Type-locality: Mexico, Orizaba, Veracruz.
- CARREIRAI Tavares, 1965b: 243, figs. 30-42. Type-locality: Brazil, Rio de Janeiro, Itatiaia.
- CRESPOI Tavares, 1965b: 246. Type-locality: Brazil, Rio de Janeiro, Itaguaí.
- LENTI Tavares, 1965b: 250, figs. 70-82. Type-locality: Brazil, Rio de Janeiro, Itatiaia.
- LENKOI Tavares, 1965b: 241, figs. 17-19. Type-locality: Brazil, Espírito Santo, Conceição da Barra, Vale do Itaúnas, Ribeirão Engano.
- LINEIFRONS Sabrosky, 1953: 175. Type-locality: Puerto Rico, Bosch Finca.
- LOPESI Tavares, 1962: 358, figs. 31-40. Type-locality: Brazil, Rio de Janeiro, Angra dos Reis.
- MENDESI Tavares, 1965b: 239, figs. 5-16. Type-locality: Brazil, Pará, Cachimbo.
- PUNCTATA Robineau-Desvoidy, 1830: 428. Type-locality: "Antilles". Distr. — U.S.A., West Indies.
- RACHOUI Tavares, 1962: 355, figs. 21-30. Type-locality: Brazil, Guanabara, Rio de Janeiro.
- RUFA (Wulp), 1890: 166, pl. 4, figs. 11-12 (*Neoptera*). Type-locality: Mexico, Tabasco, Teapa.

- SABROSKYI Tavares, 1965e: 215. Type-locality: Brazil, Rio de Janeiro, Itatiaia.
billmekii Brauer & Bergenstamm of Tavares, 1962: 349, figs. 1-10 (*Ormia*), misident.
 SERREI Séguy, 1925: 440 (also, 1926: 181). Type-locality: Costa Rica, La Caja. Unrecognized.
 WOLCOTTI Sabrosky, 1953: 174. Type-locality: Puerto Rico, Guánica.
 WYGODZINSKYI Tavares, 1965b: 248, figs. 59-67. Type-locality: Argentina, Tucumán.

Genus ORMIOPHASIA Townsend

- ORMIOPHASIA Townsend, 1919: 164. Type-species *busckii* Townsend (orig. des.).
Pseudormia Séguy, 1926c: 5 (as *Pendormia*, p. 9, error); (also, 1927b: 262). Type-species, *inflata* Séguy (orig. des., as gen. n., sp. n.).
Pseudoneoptera Séguy, 1926b: 19, 20 (in key). Type-species, *morardi* Séguy (mon.).
Plagiotormia Séguy, 1926b: 19, 20 (as *Plagiotormia*, p. 20). Type-species, *obscura* Séguy (mon.).
Plagiotormia, error.
 Ref. — Sabrosky, 1963; Tavares, 1964.
 BUSCKII Townsend, 1919b: 165. Type-locality: Panama, Cabima.
 CAUSEYI Tavares, 1964: 42, figs. 13, 25. Type-locality: Brazil, Pará, Serra do Cachimbo.
 COSTALIMAI Tavares, 1964: 44, figs. 25-36. Type-locality: Brazil, Pará, Serra do Cachimbo.
 CRUZI Tavares, 1964: 47, figs. 37-47. Type-locality: Brazil, Guanabara, Rio de Janeiro.
 INFLATA (Séguy), 1927b: 262 (*Pseudormia*). Type-locality: French Guiana, Kourou River.
 LANEI Tavares, 1964: 39, figs. 1-12. Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia.
 MORARDI (Séguy), 1926b: 19 (*Pseudoneoptera*). Type-locality: "French Guiana". Ref. — Townsend, 1931b: 82.
 OBSCURA (Séguy), 1926b: 19 (*Plagiotormia*). Type-locality: Argentina, upper Paraná River, near San Ignacio.
 TRAVASSOSI Tavares, 1964: 44, figs. 48-60. Type-locality: Brazil, Pará, Serra do Cachimbo.

Tribe TRIODINI

Genus TRIODES Coquillett

- TRIODES Coquillett, 1902:201. Type-species, *obesus* Coquillett (orig. des.).
 OBESUS Coquillett, 1902:202. Type-locality: Mexico, Chihuahua, Sierra Madre, Rio Piedras Verdes. Distr. — U.S.A., Mexico.

Tribe DEXILLINI

(Dexillini of authors)

The species of the tribe Dexillini are parasites of certain scarabaeid larvae.

Genus ACRONACANTHA Wulp

- ACRONACANTHA Wulp, 1891:212 (key). Type-species, *nubilipennis* Wulp (sub. mon.; Wulp, 1891:243).
NUBILIPENNIS Wulp, 1891:243, pl. 5, fig. 2. Type-locality: Costa Rica, Volcán de Irazú.

Genus ANGIORRHINA Brauer & Bergenstamm

- ANGIORRHINA Brauer & Bergenstamm, 1889:163 (1889:95). Type-species, *Tachina crudelis* Wiedemann (mon.).
CRUDELIS (Wiedemann), 1830:300 (*Tachina*). Type-locality: "West Indies".

Genus CERATOMETOPA Townsend

- CERATOMETOPA Townsend, 1931a:240. Type-species, *cornifera* Townsend (orig. des.).
CORNIFERA Townsend, 1931a:341. Type-locality: Argentina, Mendoza.

Genus EUTRINOIDES Walton

- EUTRINOIDES Walton, 1913b:50. Type-species, *jonesii* Walton (orig. des.).
JONESII Walton, 1913b:50, pl. 3, figs. a-f. Type-locality: Puerto Rico, Añasco.

Genus IRENGIA Townsend

- IRENGIA Townsend, 1935:217. Type-species, *guianensis* Townsend (orig. des.).
GUIANENSIS Townsend, 1935d:217. Type-locality: Guyana, upper Ireng River, Pacaraima Mts. Distr. — Venezuela, Guyana, Brazil.
LATIVENTRIS (Curran), 1934:497 (*Microphthalma*). Type-locality: "Guyana".

Genus MACROMETOPA Brauer & Bergenstamm


- MACROMETOPA Brauer & Bergenstamm, 1889:117 (1889:49). Type-species, *mericana* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.) = *calogaster* (Bigot).
CALOGASTER (Bigot), 1889a:226 (*Microphthalma*). Type-locality: Mexico, Orizaba. Ref. — Aldrich, 1924a:211 (note on type).
mericana Brauer & Bergenstamm, 1889:117 (1889:49). Type-locality: "Mexico".

Genus MEGAPARIA Wulp

- MEGAPARIA Wulp, 1891:212 (key). Type-species, *venosa* Wulp (sub. mon., Wulp, 1891:240).
VENOSA Wulp, 1891:240, pl. 5, fig. 9-9a. Type-locality: Mexico, Durango, Ciudad.

Genus MEGAPROSOPUS Macquart

- MEGAPROSOPUS Macquart, 1843:240 (1843:83). Type-species, *rufiventris* Macquart (mon.). Ref. — Aldrich, 1924a:211 (notes).



ANDINUS Townsend, 1912a:365. Type-locality: "Peru".
RUFIVENTRIS Macquart, 1843:241 (1843:84), pl. 10, fig. 1. Type-locality:
"Mexico".

Genus MICROPHTHALMA Macquart

- MICROPHTHALMA Macquart, 1843:241 (1843:84). Type-species, *nigra* Macquart (orig. des.) = *disjuncta* (Wiedemann).
Microphthalmia (error) Adams in Williston, 1908:376.
PERUA Townsend, 1912a:364. Type-species, *cuzcana* Townsend (orig. des.).
Ref. — Aldrich, 1926a (rev.); Curran, 1934e (key to American spp.).
ASCITA Reinhard, 1953b:89. Type-locality: Mexico, Morelos, Acatilpa.
CUZCANA (Townsend), 1912a:365 (*Perua*). Type-locality: Peru, Cuzco.
CROUZELI Blanchard, 1966b:192, fig. 7. Type-locality: "Argentina".
DIFFERENS (Wulp), 1890:88 (*Trixa*). Type-locality: Mexico, Guerrero, Omilteme.
DISJUNCTA (Wiedemann), 1824:45 (*Tachina*, as *disjuncta*). Type-locality: "North America". Distr. — U.S.A., Mexico, Guatemala.
distincta (error), Aldrich, 1926a: 1, 5.
OBSOLETA (Wulp), 1890:87 (*Trixa*). Type-locality: Mexico, Guerrero. Distr. — U.S.A., Mexico, Guatemala.
sordida Giglio-Tos, 1893:3. Type-locality: "Mexico".
nigra Macquart of Aldrich, 1926a:3 (*Microphthalma*), misident.
FEDALIS Reinhard, 1953b:90. Type-locality: Mexico, Nuevo León, Chapinque.
VIBRISATA (Wulp), 1891:244, pl. 5, fig. 13 (*Dexiosoma*). Type-locality: Mexico, Tabasco, Teapa. Distr. — Mexico, Costa Rica, Guyana.
VIRENS Aldrich, 1926a:6. Type-locality: Peru, Jaén, Huascaray Ridge.

Genus MORPHODEXIA Townsend

- MORPHODEXIA Townsend, 1931a:342. Type-species, *microphthalmoides* Townsend (orig. des.) = *barrosi* (Brèthes).
BARROSI (Brèthes), 1919:42 (*Camaronia*). Type-locality: Chile, Aconcagua, Río Blanco. Ref. — Cortés, 1967b:13.
microphthalmoides Townsend, 1931a:343. Type-locality: Chile, Valparaíso, Perales.
CLAUSA Aldrich, 1934:149. Type-locality: Argentina, Río Negro, Puerto Blest.
FACIALIS (Aldrich), 1923b:23 (*Selenomyia*). Type-locality: Chile, Valparaíso, Perales.
NIGRA Aldrich, 1934:149. Type-locality: Chile, Ancud.
PALPALIS Aldrich, 1934:150. Type-locality: Chile, Angol, Concepción and Victoria.
SUBAENEAE Aldrich, 1934:149 (as *nigra* var.). Type-locality: Argentina, Río Negro, Lake Gutiérrez.

Genus PROMEGAFARIA Townsend

- PROMEGAFARIA Townsend, 1931a:339. Type-species, *petiolata* Townsend (orig. des.).
PETIOLATA Townsend, 1931a:340. Type-locality: Peru, Chosica Canyon.

Genus PYRRHODEXIA Townsend

- PYRRHODEXIA Townsend, 1931a:341. Type-locality: *Dexia pyrrhoprocta* Wiedemann (orig. des.).
PYRRHIOPROCTA (Wiedemann), 1830:381 (*Dexia*). Type-locality: "Brazil".

Tribe TRICHOPROSOPINI

Genus DAETALEUS Aldrich

- DAETALEUS Aldrich, 1928b:13. Type-species, *purpureus* Aldrich (orig. des.).
Zuanalia Curran, 1934d:457. Type-species, *azurea* Curran (orig. des.).
AZUREA (Curran), 1934d:468, figs. 147, 177 (*Zuanalia*). Type-locality: Panama, Canal Zone, Barro Colorado I. (Cf. Arnaud, 1963a:132).
PURPUREUS Aldrich, 1928b:13, fig. 1. Type-locality: "Brazil".

Genus EUOESTROPSIS Townsend

- Oestropsis* Townsend, 1912a:355 (preocc. Brauer, 1868). Type-species, *viridis* Townsend (orig. des.).
EUOESTROPSIS Townsend, 1913:133 (nom. nov. for *Oestropsis* Townsend). Type-species, *Oestropsis viridis* Townsend (aut.).
VIRIDIS (Townsend), 1912a:356 (*Oestropsis*). Type-locality: Peru, Huarochiri, Tamboraque.

Genus GNADOCHAETA Macquart

- GNADOCHAETA Macquart, 1851:200 (1851:377). Type-species, *caerulea* Macquart (orig. des.).
CAERULEA Macquart, 1851:201 (1851:228), pl. 21, fig. 7 (as *coerulea*). Type-locality: Brazil, Minas Gerais.

Genus PROTRICHOPROSOPIS Blanchard

- PROTRICHOPROSOPIS Blanchard, 1966b:198. Type-species, *chaetosus* Blanchard (orig. des.).
CHAETOSUS Blanchard, 1966b:199, fig. 9. Type-locality: Argentina, Rio Negro, Bariloche, Puente Limay.

Genus PSECACERA Bigot

- PSECACERA Bigot, 1880: liii (1880: 69). Type-species, *chiliensis* Bigot (mon.). Ref. — Aldrich, 1934: 151 (key to males).
ATRIVENTRIS Aldrich, 1934: 154. Type-locality: Chile, Llanquihue, Ensenada. Distr. — Chile (Malleco, Llanquihue).
CHILIENSIS Bigot, 1880: liii (1880: 70). Type-locality: "Chile". Distr. — Chile (Angol).
plena Aldrich, 1928b: 23 (*Selenomyia*). Type-locality: Chile, Angol.
LATIFORCEPS Aldrich, 1934:155, fig. 20c. Type-locality: Argentina, Rio Negro, Lake Gutiérrez.
ROBUSTA Aldrich, 1934:154, fig. 20b. Type-locality: Chile, Concepción, San Rosendo.
TIBIALIS Aldrich, 1934:154, fig. 20d. Type-locality: Chile, Llanquihue, Casa Pangué.

VIRENS (Aldrich), 1928b:22 (*Selenomyia*). Type-locality: Chile, Valparaiso, Perales.

Genus STUARDOMYIA Cortés

STUARDOMYIA Cortés, 1945c:157. Type-species, *crassiset* Cortés (orig. des.).

CRASSISETA Cortés, 1945c:158. Type-locality: Chile, Malleco, Angol.

Genus TRICHOCERONIA Cortés

TRICHOCERONIA Cortés, 1945c:150. Type-species, *thermitana* Cortés (orig. des.).

LATIFRONS (Aldrich), 1934:20 (*Trichoprosopus*). Type-locality: "South Patagonia".

THERMITANA Cortés, 1945c:151. Type-locality: Chile, Cautin, Curacautin, Terinas de Río Blanco.

Genus TRICHOPROSOPUS Macquart

TRICHOPROSOPUS Macquart, 1843:227 (1843:70). Type-species, *durvillei* Macquart (orig. des.).

Trichoprosopa Cortés, 1963:249, error.

DURVILLE Macquart, 1843:228 (1843:71), pl. 8, fig. 2. Type-locality: Chile, Concepción. Ref. — Cortés, 1945c:149; 1963:249.

Genus TRIXODORSIS Townsend

TRIXODORSIS Townsend, 1933:527. Type-species, *facialis* Townsend (orig. des.).

FACIALIS Townsend, 1933:527. Type-locality: "Chile".

Tribe MYIOPHASIINI

The species of the tribe Myiophasiini are parasites of weevil grubs.

Genus EULOEWIOPSIS Townsend

EULOEWIOPSIS Townsend, 1917b:222. Type-species, *setosa* Townsend (orig. des.) = *anthracina* (Wiedemann).

Xystotirix Townsend, 1927a:214. Type-species, *Tachina anthracina* Wiedemann (orig. des.).

ANTHRACINA (Wiedemann), 1830:324 (*Tachina*). Type-locality: "Brazil".
setosa Townsend, 1917:222. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).

Genus GONZALEZODORIA Cortés

GONZALEZODORIA Cortés, 1967b:18. Type-species, *gonioides* Cortés (orig. des.).

GONIOIDES Cortés, 1967b:19, figs. 1-3. Type-locality: Chile, Coquimbo, Bosque Fray Jorge, 15 km SW of Pachingo.

Genus METAMYIOPHASIA Blanchard

- METAMYIOPHASIA Blanchard, 1966b:183. Type-species, *nigricauda* Blanchard (orig. des.).
NIGRICAUDA Blanchard, 1966b:184, fig. 4. Type-locality: Argentina, Río Negro, Bariloche.

Genus MYIOPHASIA Brauer & Bergenstamm

- MYIOPHASIA Brauer & Bergenstamm, 1891:362 (1891:58). Type-species, *Tachina aenea* Wiedemann (mon.) = *australis* Townsend.
Pseudoclista Brauer & Bergenstamm, 1893:192 (1893:104). Type-species, *atra* Brauer & Bergenstamm (orig. des.).
Ref. — Townsend, 1915 (key); Aldrich, 1934:167 (Patagonian spp.).
ANTENNALIS Aldrich, 1934:166. Type-locality: Chile, Llanquihue, Puerto Blest.
ATRA (Brauer & Bergenstamm), 1893:192 (1893:104) (*Pseudoclista*). Type-locality: "Brazil".
AUSTRALIS Townsend, 1916c:11 (nom. nov. for *aenea* Wiedemann). Type-locality: Uruguay, Montevideo.
aenea Wiedemann, 1830:298 (*Tachina*, preocc. Meigen, 1824). Type-locality: Uruguay, Montevideo.
DIFFICILIS Aldrich, 1934:166. Type-locality: Argentina, Río Negro, Lake Gutiérrez.
GLOBOSA (Townsend), 1892d:129 (*Lococia*). Type-locality: U.S.A., Florida. Distr. — U.S.A., Trinidad.
MADERA (Townsend), 1915e:110, 112 (*Ennyomma*; as *robusta* ssp.). Type-locality: Mexico, Chihuahua, San Pedro de Madera.
MUSCAEFORMIS (Wulp), 1890:207, pl. 4, fig. 18 (*Clista*). Type-locality: Mexico, Veracruz, near Medellín. Unrecognized.
MADRENSIS (Townsend), 1915e:109, 113 (*Eulocia*). Type-locality: "Mexico".
NEOMEXICANA (Townsend), 1915e:110, 112 (*Ennyomma*, as *robusta* ssp.). Type-locality: U.S.A., New Mexico. Distr. — U.S.A., Mexico.
OCHREICORNIS (Townsend), 1916h:15 (*Eulocia*). Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).
POLLINOSA Curran, 1926:104. Type-locality: Jamaica, Cinchona.
SIERRICOLA (Townsend), 1917:110, 112 (*Ennyomma*, as *clistoides* ssp.). Type-locality: Mexico, Chihuahua, San Pedro de Madera.

Genus MYIOPHASIOMIMA Blanchard

- MYIOPHASIOMIMA Blanchard, 1966b:186. Type-species, *lloydi* Blanchard (orig. des.).
LLOYDI Blanchard, 1966b:186, fig. 5. Type-locality: Argentina, Río Negro, Bariloche.

Genus PLESIODEXILLA Blanchard

- Plesiodesilla* Blanchard in Crouzel, 1965:84. *Nomen nudum*.
PLESIODEXILLA Blanchard, 1966b:192. Type-species, *crouzeli* Blanchard (orig. des.).
CROUZELI Blanchard, 1966b:192, fig. 7. Type-locality: "Argentina".

Genus PROTONOTODYTES Blanchard

- PROTONOTODYTES Blanchard, 1966b:196. Type-species, *ventripyga* Blanchard (orig. des.).
VENTRIPYGA Blanchard, 1966b:196, fig. 8. Type-locality: Argentina, Rio Negro, Bariloche.

Tribe MELISONEURINI

Genus AGLUMMYIA Townsend

- AGLUMMYIA Townsend, 1912a:354. Type-species, *percinerea* Townsend (orig. des.).
PERCINEREA Townsend, 1912a:355. Type-locality: Peru, Plura.
ssp. FLAYIDA Townsend, 1915d:65. Type-locality: Peru, Chosica.

Genus MICROCHAETINA Wulp

- MICROCHAETINA Wulp, 1891:212 (key). Type-species, *cinerea* Wulp (sub. mon.; Wulp, 1891:240).
Almugmyia Townsend, 1911b:136 (1912a:354). Type-species, *arida* Townsend (orig. des.).
ARIDA (Townsend), 1911b:136, 148 (1912a:354) (*Almugmyia*). Type-locality: Peru, Piura.
CINEREA Wulp, 1891:241, pl. 5, fig. 10-10a. Type-locality: Mexico, Sinaloa, Presidio de Mazatlán.
major Townsend, 1911a:154 (*Almugmyia*). *Nomen nudum*.
VALIDA (Townsend), 1892e:167 (*Rhinophora*). Type-locality: U.S.A., New Mexico, Distr. — U.S.A., Mexico.

Genus REINHARDIANA Arnaud

- Hypenomyia* Townsend, 1919a:545 (preocc. Grimschaw, 1901). Type-species, *petiolata* Townsend (orig. des.).
REINHARDIANA Arnaud, 1952:58 (nom. nov. for *Hypenomyia*). Type-species, *petiolata* Townsend (aut.).
PETIOLATA (Townsend), 1919a:545 (*Hypenomyia*). Type-locality: U.S.A., Arizona, Distr. — U.S.A., Mexico.

Tribe PROSENINI

Members of the tribe Prosenini are parasites of white grubs, elaterid grubs and other coleopterous larvae.

Genus CALLOTROXIS Aldrich

- CALLOTROXIS Aldrich, 1929b:7. Type-species, *edwardsi* Aldrich (orig. des.).
EDWARDSI Aldrich, 1929b:8. Type-locality: Chile, Malleco, Angol.

Genus CHAETOCALLIRHOE Townsend

- CHAETOCALLIRHOE Townsend, 1935d:217. Type-species, *grandis* Townsend (orig. des.).
GRANDIS Townsend, 1935d:218. Type-locality: Haiti, Kenscoff.

Genus CORDILLERODEXIA Townsend

CORDILLERODEXIA Townsend, 1927a:226. Type-species, *orientalis* Townsend (orig. des.).

COLOMBIANA Townsend, 1929:366. Type-locality: "Colombia".

ORIENTALIS Townsend, 1927a:300. Type-locality: Peru, Esealera Mt.

Genus DOLICHOCODIA Townsend

DOLICHOCODIA Townsend, 1908:59. Type-species, *Myiocera bivittata* Coquillett (orig. des.).

ERRATILIS Reinhard, 1958c:278. Type-locality: Mexico, Teotihuacán.

Genus DOLICHODINERA Townsend

DOLICHODINERA Townsend, 1935d:218. Type-species, *divaricata* Townsend (orig. des.).

DIVARICATA Townsend, 1935d:218. Type-locality: Guyana, Mount Roraima.

Genus ECHINODEXIA Brauer & Bergenstamm

ECHINODEXIA Brauer & Bergenstamm, 1893:174 (1893:86). Type-species, *Hystriosphona pseudohystriicia* Brauer & Bergenstamm (mon.).

CUBENSIS Malloch, 1932:203. Type-locality: Cuba, Sierra Maestra.

PSEUDOHYSTRICIA (Brauer & Bergenstamm), 1889:120 (1889:52) (*Hystriosphona*). Type-locality: Mexico, Tacubaya. Ref. — Aldrich, 1934:213 (notes).

Genus EUDEXIA Brauer & Bergenstamm

EUDEXIA Brauer & Bergenstamm, 1889:120 (1889:52). Type-species, *goliath* Brauer & Bergenstamm (mon.) = *formidabilis* (Bigot).

DREISEACHI Reinhard, 1956a:127. Type-locality: Mexico, Michoacán, Tzitzutzán.

FORMIDABILIS (Bigot), 1889:264 (*Rhamphina*). Type-locality: "Mexico".

goliath Brauer & Bergenstamm, 1889:128 (1889:52). Type-locality: "Venezuela". Ref. — Aldrich, 1924a:213 (notes).

memorina Schiner in Brauer & Bergenstamm, 1891:417 (1891:113). *Nomen nudum*.

OBSCURA (Bigot), 1889:264 (*Prosenia*). Type-locality: "Mexico".

Genus EXODEXIA Townsend

EXODEXIA Townsend, 1927a:227. Type-species, *uruhuasi* Townsend (orig. des.).

URUHUASI Townsend, 1927a:310. Type-locality: Peru, Uruhuasi.

Genus GIGAMYIOPSIS Reinhard

GIGAMYIOPSIS Reinhard, 1964c:5. Type-species, *funbris* Reinhard (orig. des.).

FUNEBRIS Reinhard, 1964c:6. Type-locality: Mexico, Morelos, Acatlipa.

Genus HYSTRICHODEXIA Röder

- HYSTRICHODEXIA Röder, 1886:266 (also 1892:11). Type-species, *armata* Röder (mon.).
- ARMATA Röder, 1886:266 (also 1892:12, figs. 3, 3a, 3b). Type-locality: Ecuador, Cerro del Altar.
- BREVICORNIS (Macquart), 1851:203 (1851:230) (*Prosenia*). Type-locality: Brazil, Bahia.
- ECHINATA Wulp, 1891:220. Type-locality: Costa Rica, Caché.
- INSOLITA (Walker), 1852:318 (*Dexia*). Type-locality: "Brazil".
- MELLEA Giglio-Tos, 1893:2. Type-locality: Mexico, Oaxaca.
- PUEYRREDONI Brèthes, 1918b:115. Type-locality: Argentina, Córdoba.

Genus HYSTRYSYPHONA Bigot

- HYSTRYSYPHONA Bigot, 1859:309. Type-species, *nigra* Bigot (mon.). *Hystriisiphona*, emend.
- NIGRA Bigot, 1859:309. Type-locality: "Mexico".

Genus JURINODEXIA Townsend

- JURINODEXIA Townsend, 1915b:22. Type-species, *Hystriisiphona bicolor* Giglio-Tos (orig. des.).
- BICOLOR (Giglio-Tos), 1893:1 (*Hystriisiphona*). Type-locality: "Mexico".

Genus LEPTODEXIA Townsend

- LEPTODEXIA Townsend, 1919a:550. Type-species, *gracilis* Townsend (orig. des.).
- Shermanalia Curran, 1934d:460. Type-species, *pretiosa* Curran (orig. des.). N. SYN.
- GRACILIS Townsend, 1919a:550. Type-locality: Peru, Jaén, Huascaray Ridge.
- PRETIOSA (Curran), 1934d:470, figs. 176, 185 (*Shermanalia*). Type-locality: Panama, Canal Zone, Fort Sherman. N. COMB.

Genus MICROCHAETOGYNE Townsend

- MICROCHAETOGYNE Townsend, 1931a:344. Type-species, *Prosenia melacna* Wulp (orig. des.).
- MELACNA (Wulp), 1891:217 (*Prosenia*). Type-locality: Mexico, Guerrero.

Genus MOCHLOSOMA Brauer & Bergenstamm

- MOCHLOSOMA Brauer & Bergenstamm, 1889:126 (1889:58). Type-species, *validum* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.).
- TROCHILODEXIA Townsend, 1915b:22. Type-species, *Mochlosoma anale* Giglio-Tos (orig. des.).
- Ref. — Reinhard, 1958c (rev.).
- ADUSTUM Reinhard, 1958c:107. Type-locality: Mexico, Chihuahua, Mound Valley.
- ANALE Giglio-Tos, 1893:8. Type-locality: "Mexico". Ref. — Reinhard, 1958:107.
- DEMISSUM Reinhard, 1958c:102. Type-locality: Mexico, Rio Frio.
- DUPLARE Reinhard, 1958c:100. Type-locality: Mexico, Cuernavaca. Distr. — U.S.A., Mexico.

- FURTUM Reinhard, 1958c:108. Type-locality: Mexleo, Oaxaca.
 LACERTOSUM (Wulp), 1891:215, pl. 5, fig. 1, 1a (*Prosenia*). Type-locality: "Mexleo". Ref. — Reinhard, 1958c:110.
 LAUDATUM Reinhard, 1958c:106. Type-locality: México, Río Frio.
 MEXICANUM (Maequart), 1851:204 (1851:231), pl. 21, fig. 12 (*Prosenia*). Type-locality: Mexico, Guerrero, Durango, Ciudad. Ref. — Reinhard, 1958c:108.
tesselans Wulp, 1891:216. Type-locality: "Mexleo".
 OPIPARE Reinhard, 1958c:105. Type-locality: U.S.A., Arizona. Distr. — U.S.A., Mexico.
 RUFIPES Coquillett, 1902a:202. Type-locality: Mexico, Chihuahua, Río Piedras Verdes, Sierra Madre.
 RUSSULUM Reinhard, 1958c:105. Type-locality: Mexico, D.F. Mixquile.
 SABROSKYI Reinhard, 1958c:109. Type-locality: Mexleo, Michoacán.
 SARCINALE Reinhard, 1958c:103. Type-locality: Mexico, Chihuahua, Sierra Madre Valley.
 SERICEUM Giglio-Tos, 1893:1. Type-locality: "Mexleo".
 SEROTINUM Reinhard, 1958c:101. Type-locality: Mexico, Cuernavaca.
 VALIDUM Brauer & Bergenstamm, 1889:126 (1889:58, 100). Type-locality: U.S.A., Pennsylvania. Distr. — U.S.A., Mexico.

Genus MYIOMIMA Brauer & Bergenstamm

- MYIOMIMA Brauer & Bergenstamm, 1889:119 (1889:51). Type-species, *sarcophagina* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.).
 APPENDICULATA (Blot), 1889:269 (*Phorostoma*). Type-locality: "Mexleo". Unrecognized.
 SARCOPHAGINA Brauer & Bergenstamm, 1889:119 (1889:51). Type-locality: "Brazil" (as Central America, error). Ref. — Aldrich, 1926c:124.

Genus MYIOSCOTIPTERA Giglio-Tos

- MYIOSCOTIPTERA Giglio-Tos, 1893:2. Type-species, *cincta* Giglio-Tos (orig. des.).
 CINCTA Giglio-Ts, 1893:2. Type-locality: "Mexleo".

Genus NEOMYOSTOMA Townsend

- NEOMYOSTOMA Townsend, 1935d:218. Type-species, *ptilodexioides* Townsend (orig. des.).
 PTILODEXIOIDES Townsend, 1935d:218. Type-locality: Brazil, São Paulo, São Vicente.

Genus NIMIOGLOSSA Reinhard

- NIMIOGLOSSA Reinhard, 1945:35. Type-species, *ravida* Reinhard (orig. des.).
 RAVIDA Reinhard, 1945:36. Type-locality: U.S.A., N. Mexico. Distr. — U.S.A., Mexico.

Genus OBERONOMYIA Reinhard

- OBERONOMYIA Reinhard, 1964:1. Type-species, *palpalis* Reinhard (orig. des.).
 PALPALIS Reinhard, 1964:2. Type-locality: Mexico, Veraacruz, Jalapa.

Genus PIRIONIMYIA Townsend

PIRIONIMYIA Townsend, 1931a:343. Type-species, *paradoxa* Townsend (orig. des.).

PARADOXA Townsend, 1931a:344. Type-locality: Chile, Valparaíso, Perales.

Genus PLATYRRHINODEXIA Townsend

PLATYRRHINODEXIA Townsend, 1927a:228. Type-species, *punctulata* Townsend (orig. des.).

MOYOBAMBENSIS Townsend, 1929:366. Type-locality: Peru, Moyobamba.

FUNCTULATA Townsend, 1927a:349. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus PROPHOROSTOMA Townsend

PROPHOROSTOMA Townsend, 1927a:227. Type-species, *pulchrum* Townsend (orig. des.; as *pulcra*).

FULCHRUM Townsend, 1927a:352. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus PRORHYNCHOPS Brauer & Bergenstamm

PRORHYNCHOPS Brauer & Bergenstamm, 1891:364 (1891:60). Type-species, *bilimeki* Brauer & Bergenstamm (mon.).

BILIMEKI Brauer & Bergenstamm, 1891:364 (1891:60). Type-locality: Mexico, Mazatlán, Sinaloa. Ref. — Aldrich, 1924b:212.

ERRANS Curran, 1927c:13, fig. 5. Type-locality: Puerto Rico, Manatí.

Genus PROSENOIDES Brauer & Bergenstamm

PROSENOIDES Brauer & Bergenstamm, 1891:370 (1891:60). Type-species, *papilio* Brauer & Bergenstamm (mon.) = *curvirostris* (Bigot). *Neoprosena* Townsend, 1927a:221. Type-species, *haustellata* Townsend (orig. des.).

Ref. — Reinhard, 1954 (rev.).

ASSIMILIS Reinhard, 1954:412. Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico.

curvirostris authors, not Bigot.

CURVIROSTRIS (Bigot), 1889:264 (*Prosenia*). Type-locality: "Brazil". Distr. — Mexico, Central America to Ecuador, Peru, Brazil. Ref. — Thompson, 1963b:525.

papilio Brauer & Bergenstamm, 1891:370 (1891:16). Type-locality: "Brazil".

DIACRITA Reinhard, 1954:411. Type-locality: Guatemala, San Cristóbal, Alta Vera Paz.

FLAVIPES Coquillett, 1895b:314. Type-locality: U.S.A. (Georgia, Florida). Distr. — U.S.A., Cuba.

GRANDIS Reinhard, 1954:413. Type-locality: U.S.A., Arizona. Distr. — U.S.A., Mexico (Chihuahua).


HAUSTELLATA (Townsend), 1927a:335 (*Neoprosena*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.

ISODOMOS Reinhard, 1954:413. Type-locality: Argentina, Tucumán.

TRILINEATA Reinhard, 1954:412. Type-locality: Argentina, Tucumán. Distr. — Paraguay (Villarica), Argentina (Tucumán).

Genus PTILODEXIA Brauer & Bergenstamm

- PTILODEXIA Brauer & Bergenstamm, 1889:119 (1889:51). Type-species, *carolinensis* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.).
- Clinoneura* Brauer & Bergenstamm, 1889:119 (1889:51). Type-species, *Dexia rubriventris* Macquart (mon.).
- Rhamphiniina*, authors, not Bigot.
- Rhynchiodexia*, authors, not Elgot.
- Rhynchodexia*, emend.
- Ref. — Curran, 1930 (key).
- ANGULATA (Wulp), 1891:233 (*Rhynchodexia*). Type-locality: Mexico, Guerrero. N. comb.
- ANTIIRACINA (Bigot), 1889:265 (*Rhamphiniina*). Type-locality: "Mexico".
- ARGENTINA (Bigot), 1889:265 (*Rhamphiniina*). Type-locality: Argentina, Buenos Aires. Unrecognized. N. comb.
- CINGULIPES Blanchard, 1966:202, fig. 10. Type-locality: Argentina, Buenos Aires, Castelar.
- cingulipes* Blanchard in Crouzel, 1965:83. *Nomen nudum*.
- CONJUNCTA (Wulp), 1891:228 (*Rhynchodexia*). Type-locality: Mexico, Guerrero, and Durango. N. comb.
- CONSTRICANS (Wulp), 1891:221 (*Hystriodexia*). Type-locality: Mexico, Guerrero, Omilteme. N. comb.
- DISCOLOR (Wulp), 1891:227 (*Rhynchodexia*). Type-locality: Mexico, Guerrero and Veracruz. N. comb.
- FRATERNALIS (Wulp), 1891:229 (*Rhynchodexia*). Type-locality: Mexico, Guerrero, Morcós, Tabasco. Ref. — Thompson, 1963:517.
- FUMIPENNIS (Bigot), 1889:270 (*Dexiosoma*). Type-locality: "Mexico".
- IMITATRIX (Wulp), 1891:236 (*Rhynchodexia*). Type-locality: Mexico, Guerrero and Tabasco. N. comb.
- LATERALIS (Walker), 1837:352 (*Sarcophaga*). Type-locality: Brazil, São Paulo. Ref. — Aldrich, 1930:14. N. comb.
- MACROPTERA (Wulp), 1891:232 (*Rhynchodexia*). Type-locality: Mexico, Guerrero. N. comb.
- MAJOR (Bigot), 1889:265 (*Rhamphiniina*). Type-locality: U.S.A., Washington. Distr. — U.S.A., Mexico. N. comb.
- MUSCARIA (Walker), 1852:308 (*Dexia*). Type-locality: "Brazil". Ref. — Austen, 1907:343. N. comb.
- OMISSA (Wulp), 1891:235 (*Rhynchodexia*). Type-locality: Mexico (sev. places). N. comb.
- PLANIFRONS (Wulp), 1891:234, pl. 5, fig. 7 (*Rhynchodexia*). Type-locality: Mexico, Durango, Ciudad. N. comb.
- PRAEUSTA (Wulp), 1891:234 (*Rhynchodexia*). Type-locality: Mexico, Durango, Ciudad. N. comb.
- PUNCTIPENNIS (Wulp), 1891:233 (*Rhynchodexia*). Type-locality: Mexico (sev. places). N. comb.
- RUBRICAUDA (Bigot), 1889:265 (*Rhamphiniina*). Type-locality: "Cuba". N. comb.
- RUBRICORNIS (Wulp), 1891:230 (*Rhynchodexia*). Type-locality: Mexico (sev. places). N. comb.
- RUBRIVENTRIS (Macquart), 1846:316 (1846:188), pl. 20, fig. 10 (*Dexia*). Type-locality: "Mexico". N. comb.
- RUFIANALIS (Wulp), 1891:231 (*Rhynchodexia*). Type-locality: Mexico (sev. places). N. comb.
- RUTILANS (Wulp), 1891:227 (*Rhynchodexia*). Type-locality: Mexico, Guerrero, Omilteme. N. comb.



- SCUTELLATA (Wulp), 1891:230 (*Rhynchodexia*). Type-locality: Mexico, Durango, Ciudad. N. comb.
- SIMPLEX (Bigot), 1889:266 (*Myocera*). Type-locality: "Mexico". N. comb.
- SIMULANS (Wulp), 1891:229 (*Rhynchodexia*). Type-locality: Mexico, northern Sonora. N. comb.
- SORORIA (Williston), 1896:360, pl. 11, fig. 98 (*Rhynchodexia*). Type-locality: West Indies, St. Vincent. Ref. — Curran, 1934. N. comb.
- SPINOSA (Bigot), 1889:266 (*Rhynchodexia*). Type-locality: "Haiti". Unrecognized. N. comb.
- STRENUA (Robineau-Desvoidy), 1830:315 (*Zelia*). Type-locality: Haiti, Port-au-Prince. N. comb.
- STRIATA (Wulp), 1891:234, pl. 5, fig. 6-6a (*Rhynchodexia*). Type-locality: Panama, Volcán de Chiriquí. N. comb.
- STRIGILATA (Wulp), 1891:232 (*Rhynchodexia*). Type-locality: Mexico, Durango, Ciudad. N. comb.
- TINCTICORNIS (Bigot), 1889:266 (*Rhynchodexia*). Type-locality: Mexico, Paso de Macho; Costa Rica. N. comb.
- TINCTIPENNIS (Curran), 1934c:499 (*Rhynchodexia*). Type-locality: Guyana, Kaletur. N. comb.
- VARIPEX (Wulp), 1891:231 (*Rhynchodexia*). Type-locality: Mexico, Guerrero, Omilteme. N. comb.
- VITTIGERA Curran, 1934c:500. Type-locality: Guyana, Bartica, Kartabo. N. comb.
- YPSILOFORMIS Blanchard, 1966b:205, fig. 11. Type-locality: Argentina, North of Riestra.

Genus PUNAMYOCERA Townsend

- PUNAMYOCERA Townsend, 1919a:549. Type-species, *oroyensis* Townsend (orig. des.).
- OROYENSIS Townsend, 1919a:549. Type-locality: Peru, Oroya.

Genus RHAMPHININA Bigot

- RHAMPHININA Bigot, 1885a:xi. Type-species, *dubia* Bigot (mon.) = *pica* (Fabricius).
- Paramyocera* Townsend, 1915m:405. Type-species, *discaulis* Townsend (orig. des.).
- DISCALIS (Townsend), 1915m:406 (*Paramyocera*). Type-locality: Peru, Lima, Rimac River.
- PICA (Fabricius), 1805:293 (*Musca*). Type-locality: "South America".
- dubia* Bigot, 1885a:xi. Type-locality: "Mexico".

Genus SARCOCALIRRHOE Townsend

- SARCOCALIRRHOE Townsend, 1928b:145. Type-species, *zuercheri* Townsend (orig. des.).
- TRIVITTATA (Curran), 1925e:7 (*Myiomima*). Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).
- ZUERCHERI Townsend, 1928b:146. Type-locality: Paraguay, Santa Trinidad.

Genus SCOTIPTERA Macquart

- SCOTIPTERA Macquart, 1835:215. Type-species, *Dexia melaleuca* Wiedemann (Macquart, 1843:240; 1843:83) = *venatoria* (Fabricius).

- CYANEA Giglio-Tos, 1893:2. Type-species, Mexico, Michoacán, Anganguco (as "Angang").
- GAGATEA (Robineau-Desvoidy), 1830:318 (*Sophia*). Type-locality: "Brazil". Unrecognized.
- PELLUCIDA (Robineau-Desvoidy), 1830:318 (*Sophia*). Type-locality: "Brazil". Unrecognized.
- ROBUSTA Curran, 1925a:262. Type-locality: Colombia, Magdalena, Sierra San Lorenzo, Hacienda Vista Nieve.
- VARIPENNIS Wulp, 1891:224. Type-locality: Mexico, Veracruz. N. comb.
- VENATORIA (Fabricius), 1805:285 (*Musca*). Type-locality: "South America".
- melaleuca* Wiedemann, 1830:369 (*Dexia*). Type-locality: Brazil, Rio de Janeiro.
- punctata* Robineau-Desvoidy, 1830:318 (*Sophia*). Type-locality: "Brazil".
- vittata* Guérin-Meneville in Duperrey [1831], also (1838:297), pl. 21, figs. 5-5a (*Sophia*). Type-locality: Brazil, Santa Catarina, Florianópolis.

Genus SITELLITERGUS Reinhard

- SITELLITERGUS Reinhard, 1964c:3. Type-species, *acmulus* Reinhard (orig. des.).
- AEMULUS Reinhard, 1964c:4. Type-locality: Mexico, Morcos, Cuernavaca.
- SIMOLUS Reinhard, 1964c:5. Type-locality: Mexico, Veracruz, Jalapa.

Genus STURMIODENIA Townsend

- STURMIODENIA Townsend, 1919:549. Type-species, *rubescens* Townsend (orig. des.).
- RUBESCENS Townsend, 1919a:550. Type-locality: Peru, Yahuar Mayo, Inambari River.

Genus SUMICHRASTIA Townsend

- SUMICHRASTIA Townsend, 1916b:178. Type-species, *Hystrichodexia aurea* Giglio-Tos (orig. des.).
- AUREA (Giglio-Tos), 1893:2 (*Hystrichodexia*). Type-locality: "Mexico".

Genus TESSERACEPHALUS Reinhard

- TESSERACEPHALUS Reinhard, 1955c:123. Type-species, *lenis* Reinhard (orig. des.).
- LENIS Reinhard, 1955c:124. Type-locality: Mexico, Morelos, Acatlipa.

Genus TROPIDODENIA Townsend

- TROPIDODENIA Townsend, 1915c:66. Type-species, *lutzi* Townsend (orig. des.).
- CORACINA (Wulp), 1891:221 (*Hystrichodexia*). Type-locality: Mexico, Guerrero.
- LUTZI Townsend, 1915e:67. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus TROPIDOPSIOMORPHIA Townsend

TROPIDOPSIOMORPHIA Townsend, 1927a:226. Type-species, *tropica* Townsend (orig. des.).

TROPICA Townsend, 1927a:362. Type-locality: Brazil, São Paulo, Alto da Serra (= Paranaipicaba).

Tribe THERESIINI

The hosts of Theresiini are wood-boring coleopterous larvae, and some lepidopterous stalk-borers. The genus *Paratheresia* parasitizes various species of *Diatraea* (Lep.).

Genus BATHYDEXIA Wulp

BATHYDEXIA Wulp, 1891:222. Type-species, *Phorostoma appendiculata* Bigot (Coquillett, 1910:513; misident.) = *wulphi* Townsend.

ALBOLINEATA Wulp, 1891:223. Type-locality: Costa Rica, San José, Rio Sucho

WULPHI Townsend, 1913a:347. Type-locality: Guatemala, San Gerónimo. *appendiculata* Bigot of Coquillett, 1910:513, misident.

Genus BLEPHAROPODA Rondani

BLEPHAROPODA Rondani, 1850:362. Type-species, *pilittarsis* Rondani (mon.).

PILITARSIS Rondani, 1850:362. Type-locality: Ecuador, Rio Napo.

Genus CAMARONA Wulp

CAMARONA Wulp, 1891:241. Type-species, *xanthogastra* Wulp (Brauer & Bergenstamm, 1893:94).

CAERULEONIGRA Wulp, 1891:242. Type-locality: Mexico, Guerrero, Omilteme.

XANTHOGASTRA Wulp, 1891:241, pl. 5, figs. 11-11a. Type-locality: Mexico, Guerrero, Omilteme. Ref. — Townsend, 1931b:103 (note on type).

Genus CHAETOGYNE Brauer & Bergenstamm

CHAETOGYNE Brauer & Bergenstamm, 1889:125 (1889:57). Type-species, *Stomoxys vexans* Wiedemann (mon.).

ANALIS Curran, 1937a:334. Type-locality: Brazil, São Paulo, Campos do Jordão.

VEXANS (Wiedemann), 1830:248 (*Stomoxys*). Type-locality: Brazil, São Paulo. Ref. — Aldrich, 1925c:116; Townsend, 1931b:102-3.

Genus CHAETOTHERESIA Townsend

CHAETOTHERESIA Townsend, 1931a:346. Type-species, *Musca crassa* Wiedemann (orig. des.).

CRASSA (Wiedemann), 1830:387 (*Musca*). Type-locality: "Brazil".

brasiliana Brauer & Bergenstamm, 1891:364 (1891:60) (*Myiomima*). Type-locality: "Brazil". Ref. — Townsend, 1931b:102.

Genus CHARAPOZELIA Townsend

CHARAPOZELIA Townsend, 1927a:227. Type-species, *fulviventris* Townsend (orig. des.).

FULVIVENTRIS Townsend, 1927a:297. Type-locality: Peru, Charape River.

Genus EUCHAETOGYNE Townsend

EUCHAETOGYNE Townsend, 1908:59. Type-species, *Hystrihodexia roederi* Williston (orig. des.).

ROEDERI (Williston), 1893:77 (*Hystrihodexia*). Type-locality: U.S.A., Arizona. Distr. — U.S.A., Mexico.

Genus CYRTOSOMA Townsend

Cyrtosoma Brauer & Bergenstamm, 1893:132 (1894:44) (preocc. Curtis, 1829:30). Type-species, *rufum* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.).

OCYRTOSOMA Townsend, 1912c:48 (nom. nov. for *Cyrtosoma* Brauer & Bergenstamm). Type-species, *Cyrtosoma rufum* Brauer & Bergenstamm (aut.).

RUFUM (Brauer & Bergenstamm), 1893:132 (1893:44) (*Cyrtosoma*). Type-locality: Mexico, Guerrero, Omilteme.

Genus PACHYMYIA Macquart

PACHYMYIA Macquart, 1843:272 (1843:115). Type-species, *Stomoxys vexans* Wiedemann (mon.; misident.) = *macquartii* Townsend.

MACQUARTII Townsend, 1916:11. Type-locality: Brazil, São Paulo. Ref. — Townsend, 1931b:102.

vexans Wiedemann of Macquart, 1843:272 (1843:115) (*Stomoxys*), misident.

Genus PARATHERESIA Townsend

PARATHERESIA Townsend, 1915c:65. Type-species, *signifera* Townsend (orig. des.) = *claripalpis* (Wulp).

BATHYTERESIA Townsend, 1928:146. Type-species, *bassleri* Townsend (orig. des.) = *claripalpis* (Wulp).

PARABILLAEA Blanchard, 1937:44. Type-species, *rhynchophorae* Blanchard (orig. des.) = *brasiliensis* Townsend.

Theresia, authors not Robineau-Desvoidy, in part.

Ref. — Van Emden, 1949 (nomenclature).

CLARIPALPIS (Wulp), 1896:280 (*Sarcophaga*). Type-locality: Mexico, Guerrero, Chilpancingo. Ref. — Thompson, 1960:217 (larva); Parker, 1953:49 (imm. stages).

signifera Townsend, 1915d:66. Type-locality: Peru, Rio Chira, near Sullana.

diatraeae Brèthes, 1927:165 (*Sarcophaga*). Type-locality: Argentina, Tucumán.

- ? *bassleri* Townsend, 1929:367. (*Bathitheresia*). Type-locality: Peru, Ushpayacu River.
- rufiventris* Townsend, 1929:367. Type-locality: Peru, Ushpayacu River.
- ? *pernambucensis* Townsend, 1939d:548. Type-locality: Brazil, Pernambuco, Tapera.
- ? *argentaurea* Townsend, 1939d:548. Type-locality: Brazil, Mato Grosso, Maracaju.
- BRASILIENSIS Townsend, 1917b:222. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).
- rhynchophorae* Blanchard, 1937:44, fig. 5 (*Parabillaea*). Type-locality: Argentina, Misiones, San Ignacio.

Genus SARCOPROSENA Townsend

- SARCOPROSENA Townsend, 1927a:228. Type-species, *triangulifera* Townsend (orig. des.).
- RUFIVENTRIS Townsend, 1929:367. Type-locality: Peru, Ushpayacu River.
- TRIANGULIFERA Townsend, 1927a:356. Type-locality: Peru, Yahuar Mayo.

Genus SCHISTOSTEPHANA Townsend

- SCHISTOSTEPHANA Townsend, 1919a:551. Type-species, *aurifrons* Townsend (orig. des.).
- AURIFRONS Townsend, 1919a:552. Type-locality: Peru, Charape River.

Genus THERESIA Robineau-Desvoidy

- THERESIA Robineau-Desvoidy, 1830:325. Type-species, *tandree* Robineau-Desvoidy (mon.) = *rutilans* (Fabricius).
- RECTA Aldrich, 1934:107. Type-locality: Chile, Peulla.

Genus TYREOMMA Brauer & Bergenstamm

- TYREOMMA Brauer & Bergenstamm, 1891:381 (1891:77). Type-species, *muscinum* Wulp (sub. mon.; Wulp, 1896:293).
- Eutheresiops* Townsend, 1917a:48. Type-species, *trixoides* Townsend (orig. des.).
- Pterinopterna* Townsend, 1919a:553. Type-species, *ciliata* Townsend (orig. des.).
- CILIATUM (Townsend), 1919a:553 (*Pterinopterna*). Type-locality: Peru, Jaén, Huascaray.
- MUSCINUM Wulp, 1896:293, pl. 7, figs. 13-13a. Type-locality: Mexico, Guerrero, Omilteme.
- trixoides* Townsend, 1917a:49 (*Eutheresiopsis*). Type-locality: Mexico, Chihuahua, Mound Valley.

Genus XANTHOTHERESIA Townsend

- XANTHOTHERESIA Townsend, 1931a:345. Type-species, *bicolor* Townsend (orig. des.).
- BICOLOR Townsend, 1931a:346. Type-locality: Colombia, "tierra templada".

UNRECOGNIZED SPECIES OF PROSENIINAE

- LONGIPALPIS Wulp, 1883:30, pl. 2, figs. 1, 2 (*Prosenia*). Type-locality: "Argentina".
- MELANAEA Wulp, 1896:295, pl. 7, figs. 15-15a (*Calliphora*). Type-locality: Mexico, Guerrero, Amula.
- RAVA Wulp, 1891:237 (*Myocera*). Type-locality: Mexico, Durango and Guerrero.
- SARCOPHAGINA Wulp, 1883:31, pl. 2, figs. 5,6 (*Prosenia*). Type-locality: "Argentina".

Subfamily TACHININAE

(Larvaevorinae)

The members of the subfamily Tachininae parasitize many kinds of heterocerous caterpillars, especially armyworms, cutworms and stalk borers such as *Diatraea*.

Tribe HYPOTACHININI


Genus HYPOTACHINA Brauer & Bergenstamm

- HYPOTACHINA Brauer & Bergenstamm, 1891:351 (1891:47). Type-species, *disparata* Brauer & Bergenstamm (mon.) = *chrysophora* (Wiedemann).
- Phyllolabella* Townsend, 1919b:168. Type-species, *robusta* Townsend (orig. des.) = *chrysophora* (Wiedemann).
- CHIRYSOPHORA (Wiedemann), 1830:316 (*Tachina*). Type-locality: "Brazil". Ref. — Aldrich, 1925d:469; Thompson, 1963a:429.
- disparata* Brauer & Bergenstamm, 1891:351 (1891:47). Type-locality: "Brazil".
- robusta* Townsend, 1919b:168 (*Phyllolabella*). Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).

Tribe MACROMYINI

Genus BOMBYLIOMYIA Brauer & Bergenstamm

- BOMBYLIOMYIA Brauer & Bergenstamm, 1889:131 (1889:63). Type-species, *Hystricia flavipalpis* Macquart (mon.).
- Eublepharipeza* Townsend, 1914:46. Type-species, *hystrix* Townsend (orig. des.).
- ALBICEPS (Wulp), 1888:12, pl. 1, fig. 6 (*Hystricia*). Type-locality: Costa Rica, Rio Sueio.
- CONCOLOR Engel, 1920:320, figs. W,X. Type-locality: Brazil, Rio Grande do Sul.
- ochracea* Townsend, 1929:331 (*Eublepharipeza*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- FLAVIPALPIS (Macquart), 1846:275 (1846:147), pl. 12, fig. 10 (*Hystricia*). Type-locality: "Brazil".
- hystrix* Townsend, 1914:47 (*Eublepharipeza*). Type-locality: Peru, Charape.
- FLAVITARSIS (Macquart), 1846:276 (1846:148), pl. 13, fig. 9 (*Hystricia*). Type-locality: "Colombia".
- GABANA (Townsend), 1914:47 (*Eublepharipeza*; as *hystrix* ssp.). Type-locality: Peru, San Gabán Canyon, Casahuiri.



- NIGRA (Townsend), 1914:47 (*Eubtepharipeza*). Type-locality: Peru, Matucana, and Verrugas Canyon.
 PATULA (Walker), 1849:712 (*Tachina*). Type-locality: unknown (probably American, cf. Austen, 1907:328). N. comb.
 PURPUREA Thompson, 1963a:425. Type-locality: Trinidad.
 ROROR (Williston), 1886:298 (*Hystricia*). Type-locality: U.S.A., Arizona. Distr. — Canada, U.S.A., Mexico.
flavipalpis, authors, not Macquart.

Genus BOMBYLIOPSIS Townsend

- BOMBYLIOPSIS Townsend, 1915b:23. Type-species, *Tachina abrupta* Wiedemann (orig. des.).
Hystricia, authors, not Macquart.
 Ref. — Curran, 1942 (key as *Hystricia*).
 ABRUITA (Wiedemann), 1830:293 (*Tachina*). Type-locality: "North America". Distr. — U.S.A., Mexico.
 ARGENTINENSIS Blanchard, 1941:342, fig. 1. Type-locality: Argentina, Tucumán, Rodeo de la Vaca.
 MICANS (Wulp), 1888:16, pl. 1, fig. 12 (*Hystricia*). Type-locality: Costa Rica, Rio Sucho, Volcan de Irazú.
 KUFOHRTA (Engel), 1920:318 (*Bombyliomyia*; as *micans* var.). Type-locality: Bolivia, Cuesta de Chlunticara. N. comb.
 TESTACEA (Macquart), 1843:201 (1843:44), pl. 4, fig. 4 (*Hystricia*). Type-locality: "Mexico". Unrecognized.

Genus CALOHYSTRICIA Townsend

- CALOHYSTRICIA Townsend, 1931a:350. Type-species, *Hystricia velutina* Wulp (orig. des.).
 ALBOSIGNATA (Wulp), 1892:190 (*Hystricia*). Type-locality: "Mexico". N. comb.
 GERSTCHII Curran, 1942:84. Type-locality: Panama, Chiriquí, El Volcán.
 VELUTINA (Wulp), 1888:15, pl. 1, fig. 10 (*Hystricia*). Type-locality: Costa Rica, Rio Sucho.

Genus CHLOROHYSTRICIA Townsend

- CHLOROHYSTRICIA Townsend, 1927a:244. Type-species, *purpurea* Townsend (orig. des.) = *reinwardtii* (Wiedemann).
 CYANEIVENTRIS (Wulp), 1884:cxcel (*Hystricia*). Type-locality: Mexico, Guanajuato. Distr. — U.S.A., Mexico.
 REINWARDTII (Wiedemann), 1830:315 (*Tachina*). Type-locality: "Brazil".
 Ref. — Townsend, 1913b:104, 169 (notes on type).
purpurea Townsend, 1927a:297. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus DIOTREPES Reinhard

- DIOTREPES Reinhard, 1964c:7. Type-species, *formosus* Reinhard (orig. des.).
 FORMOSUS Reinhard, 1964c:8. Type-locality: Mexico, Morelos, Cuernavaca.

Genus ENGELOMYIA Townsend

- Bombyliomyia*, subg. *Hystriciella* Engel, 1929:311 (preocc. Townsend, 1915). Type-species, *Hystricia tarsata* Schiner (orig. des.; misident.) = *alpina* Townsend.

ENGELONYIA Townsend, 1913b:349 (nom. nov. for *Hystrieciella* Engel). Type-species, *Hystrieciella tarsata* Schiner (misident.) = *alpina* Townsend.
 ALPINA Townsend, 1931a:350. Type-locality: Bolivia, Cuesta de Cillunticara.
tarsata Schiner of Engel, 1920:34 (*Hystrieciella*), misident.
 CUESTAE Engel, 1920:327. Type-locality: Bolivia, Cuesta de Cillunticara.
 PUMIPENNIS Engel, 1920:327. Type-locality: Bolivia, Cuesta de Cillunticara.
 ORNATIPENNIS Engel, 1920:328, fig. B2. Type-locality: Bolivia, Cuesta de Cillunticara.
 TARSATA (Schiner), 1868:333 (*Hystrieciella*). Type-locality: "South America".

Genus EUHYSTRICIA Townsend

EUHYSTRICIA Townsend, 1914:84. Type-species, *nigra* Townsend (orig. des.).
 CUSSILIHIS Reinhard, 1953:91. Type-locality: Mexico, Rio Frio.
 NIGRA Townsend, 1914:84. Type-locality: Peru, San Gabán Canyon, Casahuiri.

Genus HYSTRICIA Macquart

HYSTRICIA Macquart, 1843:200 (1843:43). Type-species, *amoena* Macquart (Coquillett, 1910:555).
 Ref. — Curran, 1942 (key).

According to Aldrich's notes this genus has very few species; most of those referred to it belong to *Jurinia* in the usual sense, not of Townsend, based on the specimens of *gagatea* now in Paris.

ALBIMANA Curran, 1942:81. Type-locality: Ecuador, Bolivar, Hacienda Talahua.
 AMOENA Macquart, 1843:201 (1843:44) (as *amaena*, error). Type-locality: "Mexico".
 BROWNI Curran, 1942:81. Type-locality: Ecuador, Tunguragua Volcán, Minza Rldge.
 CALIGINOSA (Walker), 1852:268 (*Tachina*). Type-locality: "Brazil".
 CONDOR Curran, 1942:80. Type-locality: Ecuador, Bolivar, Hacienda Talahua.
 FLAVITIBIA Curran, 1942:82. Type-locality: Ecuador, Bolivar, Hacienda Talahua.
 HUMERALIS Curran, 1942:81. Type-locality: Ecuador, Bolívar, Hacienda Talahua.
 LAXA Curran, 1942:83. Type-locality: Panama, Chiriqui, El Volcán.
 NIGROTEBIATA Curran, 1942:82. Type-locality: Ecuador, Tungarahua, Volcán Minza Rldge.
 NIGROSCUTATA Rondani, 1863:18. Type-locality: "Colombia".
 OBESA Engel, 1920:306, fig. N. Type-locality: "Venezuela, Peru, Mexico".
 PALPINA Rondani, 1850a:362. Type-locality: Ecuador, Rio Napo.
 EUFIPES Macquart, 1851:145 (1851:172), pl. 15, fig. 11. Type-locality: "Brazil". Unrecognized.
 EUFOHISTA Curran, 1942:83. Type-locality: Ecuador, Bolívar, Hacienda Talahua.
 VARGAS Curran, 1942:83. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
 VULTUR Curran, 1942:82. Type-locality: Panama, Chiriqui, El Volcán.

Genus HYSTRICIOPSIS Townsend

- HYSTRICIOPSIS Townsend, 1914:85. Type-species, *obscura* Townsend (orig. des.).
 COPULATA (Wiedemann), 1830:295 (*Tachina*). Type-locality: "Brazil". Ref. — Aldrich, 1927c:10 (notes).
 OBSCURA Townsend, 1914:86. Type-locality: Peru, Matucana.
 TESTACEIVENTRIS (Wulp), 1892:190 (*Hystricla*). Type-locality: "Mexico".
 N. COMB.

Genus LASIONA Wulp

- LASIONA Wulp, 1890:43 (key). Type-species, *multisetosa* Wulp (sub. mon.; Wulp, 1890:127).
 MULTISETOSA Wulp, 1890:128, pl. 3, figs. 20-20a. Type-locality: Costa Rica, Rio Suelo. Ref. — Townsend, 1913b:157 (notes on type).

Genus LASIOPALPUS Macquart

- LASIOPALPUS Macquart, 1847:79 (1847:63). Type-species, *flavitaris* Macquart (mon.).
 FLAVITARIS Macquart, 1847:80 (1847:64), pl. 3, fig. 1. Type-locality: "South America" (as unknown). Ref. — Aldrich, 1925e:118.

Genus MACROMYA Robineau-Desvoidy

- MACROMYA Robineau-Desvoidy, 1830:322. Type-species, *depressa* Robineau-Desvoidy (Townsend, 1916c:7).
Gymnostylia Macquart, 1835:216. Type-species, *Macromya depressa* Robineau-Desvoidy (Townsend, 1916c:7).
Tropidopsis Brauer & Bergenstamm, 1889:132 (1889:64). Type-species, *Tachina pyrrhaspis* Wiedemann (mon.).
Xanthohystricla Townsend, 1913a:348. Type-species, *Tachina amisias* Walker (orig. des.) = *pyrrhaspis* (Wiedemann).
 Ref. — Reinhard, 1968 (rev.).
 ANTHEMON (Walker), 1849:733 (*Tachina*). Type-locality: "Brazil".
 CONNECTANS (Townsend), 1912a:312 (*Tropidopsis*). Type-locality: Peru, Jaén, Charape River.
 CINISCUOLA Reinhard, 1968:1287. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
 CROCATA Reinhard, 1968:1283. Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico, San Salvador, Brazil.
 DEPRESSA Robineau-Desvoidy, 1830:322. Type-locality: "Brazil". Ref. — Reinhard, 1968:1284.
dorsalis Wulp, 1888:17, pl. 1, fig. 13 (*Hystricla*). Type-locality: Guatemala (San Gerónimo); Costa Rica (Caché), Panama (Volcán de Chiriquí).
 FASCIATA Macquart, 1848:212 (1848:52), pl. 6, fig. 3. Type-locality: "Brazil". Unrecognized.
 LUCENS Reinhard, 1968:1285. Type-locality: Ecuador, Rio Mapota.
 PYRRHASPIS (Wiedemann), 1830:307 (*Tachina*). Type-locality: "Brazil". Ref. — Aldrich, 1925c:125 (notes on types); Curran, 1939:2; Reinhard, 1968:1285.
analís Robineau-Desvoidy, 1830:322. Type-locality: "Brazil".

analis Macquart, 1843:245 (1843:88) (*Gymnostylia*). Type-locality: "Guiana".

amisias Walker, 1849:734 (*Tachina*). Type-locality: Unknown.

pachecoi Curran, 1939a:2. Type-locality: Guatemala, Guatemala City.

bassleri Curran, 1939a:2. Type-locality: Peru, Achinamiza.

Genus PHOBETROMYIA Reinhard

PHOBETROMYIA Reinhard, 1964c:9. Type-species, *dumalis* Reinhard (orig. des.).

DUMALIS Reinhard, 1964c:10. Type-locality: Mexico, Rio Frio.

Genus PSEUDOBOMBYLIOMYIA Townsend

PSEUDOBOMBYLIOMYIA Townsend, 1931a:347. Type-species, *linellii* Townsend (orig. des.).

LINELLII Townsend, 1931a:348. Type-locality: "Venezuela".

Tribe TACHININI

(Larvaevorini, Echinomyini)

Genus FABRICIELLA Bezzi

Fabricia Latreille, 1829:510 (preocc. Blainville, 1828). Type-species, *Musca fera* Linnaeus (mon.; msident.) = *ferox* (Panzer).

Fabricia Robineau-Desvoidy, 1830:42 (preocc. Blainville, 1828). Type-species, *Musca ferox* Panzer (mon.).

FABRICIELLA Bezzi, 1906:49 (nom. nov. for *Fabricia* Robineau-Desvoidy). Type-species, *Musca ferox* Panzer (aut.).

ACUMINATA Tothill, 1924:260. Type-locality: U.S.A., Colorado. Distr. — U.S.A., Mexico.

FLORUM (Walker), 1849:722 (*Tachina*). Type-locality: Canada, Ontario. Distr. — Canada, U.S.A., Mexico (Chihuahua).

Genus PELETERIA Robineau-Desvoidy

PELETERIA Robineau-Desvoidy, 1830:39. Type-species, *abdominalis* Robineau-Desvoidy (Coquillett, 1910:586).

Peletieria, emend.

Sphyromyia Bigot, 1884c:cviii. Type-species, *malleola* Bigot (mon.).

Peleteropsis Townsend, 1916e:630. Type-species, *Echinomyia flaviventris* Wulp (orig. des.).

Oxydosphyria Townsend, 1926:40. Type-species, *infernalis* Townsend (orig. des.) = *iterans* (Walker).

Aphriosphyria Townsend, 1927a:238. Type-species, *communis* Townsend (orig. des.) = *robusta* (Wiedemann).

Cuphoceropsis Townsend, 1935d:220. Type-species, *facialis* Townsend (orig. des.).

Aphriosphyriopsis Blanchard, 1943b:134. Type-species, *nemochaetoides* Blanchard (orig. des.). N. SYN.

Cuphoceromyia Blanchard, 1943b:135. Type-species, *aldrichi* Blanchard (orig. des.). N. SYN.

- Prosteatosoma* Blanchard, 1943b:150. Type-species, *lineata* Blanchard (orig. des.). N. SYN.
 Ref. — Curran, 1925c (key); Guimarães, 1962 (Brazilian spp.).
- ACLISTA Reinhard, 1956c:109. Type-locality: Mexico, Mt. Popocatepetl, 3000 m. Distr. — U.S.A., Mexico.
- ALBUQUERQUEI Guimarães, 1962:491, figs. 38-47. Type-locality: Brazil, São Paulo, São José dos Campos.
- ANIALIS (Macquart), 1846:272 (1846:144), pl. 12, fig. 3 (*Echinomyia*). Type-locality: "New Granada" (Colombia), (as "Mexico", error).
- ANDINA (Blanchard), 1943b:138, fig. 24 (*Cuphoceropsis*). Type-locality: Argentina, Jujuy.
- APICALIS (Walker), 1852:275 (*Tachina*). Type-locality: "Colombia".
- APICATA Curran, 1925c:244. Type-locality: Mexico, Chihuahua, Sierra Madre.
- BLANCHARDI Guimarães, nom. nov. for *aldrichi* Blanchard. Type-locality: Argentina, Santa Cruz, Rio Marta, Lago Tar.
aldrichi Blanchard, 1943b:136 (*Cuphocromyia*; preocc. Curran, 1925:240). Type-locality: Argentina, Santa Cruz, Rio Marta, Lago Tar.
- CARNATA Reinhard, 1953b:92. Type-locality: Mexico, Michoacán, Zamora.
- CINERASCENS (Bigot), 1889a:256 (*Echinomyia*). Type-locality: "Mexico". Ref. — Curran, 1925c:244, figs. 23, 24.
- COMPASCUA (Wulp), 1892:192 (*Echinomyia*). Type-locality: Mexico, Guerrero. Distr. — U.S.A., Mexico.
- CONVEXA Curran, 1925c:251, fig. 32. Type-locality: U.S.A., New Mexico. Distr. — U.S.A., Mexico.
- CORA (Bigot), 1888a:81 (*Echinomyia*). Type-locality: "Mexico". Unrecognized. Ref. — Brauer, 1898:5 (note on type).
- FILIPALPIS (Rondani), 1863:15 (*Echinomyia*). Type-locality: Chile, Santiago. Ref. — Aldrich, 1934:120.
andicola Bigot, 1888b:86 (*Fabricia*). Type-locality: "Chile".
inca Curran, 1925c:247. Type-locality: Peru, Matucana.
- FLAVIVENTRIS (Wulp), 1888:32, pl. 2, fig. 11 (*Echinomyia*). Type-locality: Mexico, Durango, Ciudad.
- FUSCISQUAMA Curran, 1925c:237. Type-locality: Colombia, La Cumbre.
- GENEROSA (Wulp), 1892:192 (*Echinomyia*). Type-locality: Mexico, N. COMB.
- GIACOMELLI (Blanchard), 1943b:125, fig. 21 (*Aphriophyria*). Type-locality: Argentina, La Rioja.
- GRIOTI (Blanchard), 1943b:127, fig. 22 (*Aphriophyria*). Type-locality: Argentina, Santa Fé, Gálvez.
- INCONTESTA Curran, 1926b:171 (nom. nov. for *neglecta* Curran). Type-locality: U.S.A., New Mexico. Distr. — U.S.A., Mexico.
neglecta Curran, 1925:243 (preocc. Townsend, 1897). Type-locality: U.S.A., New Mexico.
- ITERANS (Walker), 1949:727 (*Tachina*). Type-locality: Canada, Nova Scotia. Distr. — Canada, U.S.A., Mexico.
- LALANDII Robineau-Desvoidy, 1830:40. Type-locality: "Brazil". Unrecognized.
- LATIFASCIATA (Blanchard), 1943b:130 (*Aphriophyria*). Type-locality: Argentina, Posadas, Posadas.
- LINEATA (Blanchard), 1943b:150 (*Prosteatosoma*). Type-locality: Argentina, Santa Cruz, Rio Marta, Lago Tar. N. COMB.
- MACROCERA (Bigot), 1888a:81 (*Echinomyia*). Type-locality: "Mexico". Unrecognized. Ref. — Brauer, 1898:6 (notes on type).
- MALLEOLA (Bigot), 1884c:clx (*Sphyromyia*). Type-locality: U.S.A., California. Distr. — U.S.A., Mexico.
- MEXICANA Curran, 1925c:256, fig. 39. Type-locality: Mexico, Hidalgo, Real del Monte.

- MULTISPINOSA Thompson, 1963a:413. Type-locality: Trinidad, Santa Cruz Valley.
- NEMOCHARTOIDES Blanchard, 1943b:134, fig. 23 (*Aphriosphyriopsis*). Type-locality: Argentina, Valle Túnel. N. comb.
- OBSOLETA Curran, 1925c:251, fig. 31. Type-locality: U.S.A., New Mexico. Distr. — U.S.A. to Costa Rica.
- PYGMIAEA (Macquart), 1851:143 (1851:170), pl. 15, fig. 10 (*Echinomyia*). Type-locality: "Chile". Distr. — Brazil (Pernambuco to São Paulo), Paraguay, Argentina, Chile (Coquimbo to Malleco). Ref. — Cortés, 1963:248 (notes on type).
- sordida* Aldrich, 1934:122. Type-locality: Chile, Angol.
- facialis* Townsend, 1935d:220 (*Cuphoceropsis*). Type-locality: Brazil, Pernambuco.
- filipalpis* Rondani of Curran, 1925c:257 (*Peleteria*), misident.
- ROBUSTA (Wiedemann), 1830:290 (*Tachina*). Type-locality: Uruguay, Montevideo. Distr. — Neotropical. Ref. — Guimarães, 1962.
- marmorata* Townsend, 1915h:185 (as *robusta* ssp.). Type-locality: Peru, Chosilca.
- ? *texensis* Curran, 1925c:246. Type-locality: U.S.A., Texas.
- communis* Townsend, 1927a:287 (*Aphriosphyria*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- RUBRIFRONS (Blot), 1888a:80 (*Echinomyia*). Type-locality: "Mexico". Unrecognized.
- SEABRAI Guimarães, 1962:492, figs. 48-55. Type-locality: Brazil, São Paulo, Campos do Jordão.
- SEMIURFA (Blanchard), 1943b:132 (*Aphriosphyria*). Type-locality: Argentina, Salta.
- SETOSA Curran, 1925c:252, fig. 34. Type-locality: Mexico, Chihuahua, Mound Valley.
- SIMILIS (Walker), 1852:269 (*Tachina*) (preocc. Walker, 1852:266). Type-locality: "South America". Unrecognized. Ref. — Austen, 1907:344.
- SUBANDINA (Blanchard), 1943b:140 (*Cuphoceropsis*). Type-locality: Argentina, Rioja. N. comb.
- TEGULATA (Townsend), 1916e:630 (*Peleteropsis*). Type-locality: "Mexico".
- TOWNSENDI Curran, 1925c:252, fig. 33. Type-locality: Mexico, Chihuahua, Mound Valley. Distr. — U.S.A., Mexico.
- TRINITATIS Thompson, 1963a:415, fig. 92. Type-locality: Trinidad, Holy Reservoir.

Genus RUIZIELLA Cortés

- RUIZIELLA Cortés, 1951:254. Type-species, *frontosa* Cortés (orig. des.).
- FRONTOSA Cortés, 1951:255. Type-locality: Chile, Santiago, Punta de Damas.
- LUCTUOSA Cortés, 1951:257. Type-locality: Chile, Chillán, Termas de Chillán.

Genus STEATOSOMA Aldrich

- STEATOSOMA Aldrich, 1934:112. Type-species, *rufiventris* Aldrich (orig. des.).
- NIQRIVENTRIS Aldrich, 1934:115. Type-locality: Argentina, Tierra del Fuego, Lake Yuvín. Ref. — Cortés, 1963:246 (notes on type).
- RUFIVENTRIS Aldrich, 1934:112, fig. 14. Type-locality: Argentina, Tierra del Fuego, Río Grande, Estancia Viamonte. Distr. — South Chile, Argentina (Tierra del Fuego). Ref. — Cortés, 1963:246 (notes on type).

Tribe DEJEANIINI

The members of the tribe Dejeanini parasitize a great variety of lepidopterous larvae such as cutworms, armyworms, etc.

Genus ADEJEANIA Townsend

- ADEJEANIA Townsend, 1913c:103, 104. Type-species, *Tachina armata* Wiedemann (orig. des.).
- Echinotachina* Townsend, 1913c:105. Type-species, *Tachina corpulenta* Wiedemann (orig. des.).
- Trichodejeania* Townsend, 1913c:105. Type-species, *Dejeania vexatrix* Osten Sacken (orig. des.).
- Dejeania*, authors, not Robineau-Desvoidy.
Ref. — Curran, 1947 (rev.); Guimarães, 1966b (rev. of Brazilian spp.).
- ANALIS (Macquart), 1843:191 (1843:34), pl. 3, fig. 3 (*Dejeania*). Type-locality: "Colombia" (as Mexico, error). Ref. — Guimarães, 1966b:189.
- ANDINA (Townsend), 1912a:333 (errata, 1915a:28) (*Dejeania*). Type-locality: Peru, Ollachea, Canyon of San Gabán River.
- ANDUZEI Curran, 1947:54. Type-locality: Venezuela, D.F., Junquito (cf. Arnaud, 1963:105).
- ARMATA (Wiedemann), 1830:287 (*Tachina*). Type-locality: "Cuba". Ref. — Curran, 1947:58.
- AUREA (Giglio-Tos), 1893:3 (*Dejeania*). Type-locality: Mexico, Solco.
- BICAUDATA Curran, 1947:61, figs. 21-22. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
brasiliensis Townsend, 1929:382 (*Trichodejeania*, preocc. Robineau-Desvoidy, 1830:33). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- BIORNATA Curran, 1947:59, figs. 16-17. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- BRASILIENSIS (Robineau-Desvoidy), 1830:33 (*Dejeania*). Type-locality: Brazil, Paraná, Guaratuba. Ref. — Townsend, 1931:161 (notes on synonymy).
- BREVIHIRTATA Curran, 1947:56, fig. 5. Type-locality: "Venezuela".
- BREVIROSTRIS Curran, 1947:56. Type-locality: Mexico, Tlaltenpautla.
- BROWNI Curran, 1947:54. Type-locality: Ecuador, Runtun. Distr. — Colombia, Ecuador.
- CONCLUSA Curran, 1947:61. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- CORPULENTA (Wiedemann), 1830:280 (*Tachina*). Type-locality: "Mexico".
- GRANDIS Guimarães, 1966b:174, figs. 27-28. Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia.
- HONESTA (Rondani), 1850:260 (*Dejeania*). Type-locality: Ecuador, Rio Napo. Ref. — Curran, 1947:58.
- LOPESI Guimarães, 1966b:175, figs. 29-30. Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil (Rio de Janeiro, São Paulo, Paraná, Rio Grande do Sul).
- MAGALHAESI Guimarães, 1966b:183, figs. 55, 56. Type-locality: Brazil, São Paulo, Campos do Jordão.
- MARGINALIS Curran, 1947:60, figs. 27, 32. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- NIGROTHORACICA (Vimmer & Soukup), 1940:219 (*Dejeania*). Type-locality: Peru, Puno. Unrecognized.

- PALPALIS* Curran, 1947:53, fig. 15. Type-locality: Panama, Chiriquí, El Volcán.
- PELLUCENS* Guimarães, 1966b:168, figs. 11, 13. Type-locality: Brazil, Rio de Janeiro, Teresópolis.
- RUFIPALPIS* (Macquart), 1843:192 (1843:35), pl. 3, fig. 1 (*Dejeania*). Type-locality: "Mexico".
- SABROSKYI* Guimarães, 1966b:173, figs. 25, 26. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil (Rio de Janeiro to Santa Catarina).
- SAETIGERA* Guimarães, 1966b:178, figs. 40, 41. Type-locality: Brazil, Rio de Janeiro, Itatiaia.
- SANCTIPauli* Guimarães, 1966b:185, figs. 52, 57. Type-locality: Brazil, São Paulo, Campos do Jordão.
- SPINIVENTRIS* Guimarães, 1966b:181, figs. 47, 48. Type-locality: Brazil, Santa Catarina, Corupá.
- SPINOSA* Guimarães, 1966b:177, figs. 31, 35. Type-locality: Brazil, São Paulo, Barueri.
- THOMPSONI* Guimarães, 1966b:175, figs. 32, 33. Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia.
- TOWNSENDI* Curran, 1947:60, figs. 25, 26. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- TRIDENS* Curran, 1947:61, figs. 30, 31. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- UNIFORMIS* Curran, 1947:58, figs. 28, 29. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- VERRUGANA* (Townsend), 1914:172 (*Trichodejeania*). Type-locality: Peru, Verrugas Canyon.
- VEXATRIX* (Osten Sacken), 1877:343 (*Dejeania*). Type-locality: U.S.A., Colorado.
- rufipalpis*, authors, not Macquart.
- WYGODZINSKYI* Guimarães, 1966b:178, figs. 36, 37. Type-locality: Brazil, São Paulo, Campos do Jordão, Umuarama.
- XANTHOPILOSA* Guimarães, 1966b:182, figs. 49, 50. Type-locality: Brazil, Rio de Janeiro, Teresópolis.
- YPSILON* Curran, 1947:57. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Rio de Janeiro to Santa Catarina).

Genus *AMICROTRICHOMMA* Townsend

- AMICROTRICHOMMA* Townsend, 1927a:245. Type-species, *orbitalis* Townsend (orig. des.).
- ADA* Curran, 1947:83, figs. 71, 73, 74. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- ORBITALIS* Townsend, 1927a:285. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus *ARCHYTAS* Jaennicke


- ARCHYTAS* Jaennicke, 1867:392 (1868:84). Type-species, *bicolor* Jaennicke (mon.) = *diaphanus* (Fabricius).
- Nemochaeta* Wulp, 1885:38. Type-species, *disstimilis* Wulp (orig. des.).
- Tachinodes* Brauer & Bergenstamm, 1889:133 (1889:65). Type-species, *Musca hystrix* Fabricius (mon.; misident.) = *metallica* (Robineau-Desvoidy).

- Parafabricia* Brauer & Bergenstamm, 1894:76 (1895:612). Type-species, *Tachina bicolor* Wiedemann (mon.; misident) = *perplexa* Townsend.
- Eufabricia* Townsend, 1908:111. Type-species, *flavicans* Townsend (orig. des.) = *diaphanus* (Fabricius).
- Pseudoarchytas* Townsend, 1915h:185. Type-species, *marmoratus* Townsend (orig. des.).
- Neoarchytas* Townsend, 1915m:430. Type-species, *inambaricus* Townsend (orig. des.).
- Makasinocera* Townsend, 1915m:431. Type-species, *unguis* Townsend (orig. des.).
- Pseudoarchytopsis* Townsend, 1927a:252. Type-species, *brasiliensis* Townsend (orig. des.). = *incertus* (Macquart).
- Proarchytas* Townsend, 1931a:351. Type-species, *Tachina daemon* Wiedemann (orig. des.).
- Makasinoceros* Townsend, 1935d:219. Type-species, *fulviventris* Townsend (orig. des.). = *shannoni* Guimarães.
- Itarchytas* Blanchard, 1940:225. Type-species, *pseudodaemon* Blanchard (orig. des.).
- Archynemochaeta* Blanchard, 1941:345. Type-species, *frenguelli* Blanchard (orig. des.).
- Proarchytoides* Blanchard, 1941:365. Type-species, *glacomelli* Blanchard (orig. des.).
- Archytodejeania* Blanchard, 1941:348. Type-species, *bruchii* Blanchard (orig. des.). N. SYN.
- Ref. — Curran, 1928b (rev.), Cortés, 1944a (Chilean spp.): Guimarães, 1960, 1961a, 1961b, 1963a, 1963b (Brazilian spp.).
- ABERRANS (Giglio-Tos), 1893:2 (*Nemochaeta*). Type-locality: Mexico, Metztitlán. Unrecognized.
- ALBICEPS (Walker), 1860:295 (*Echlinomyia*). Type-locality: "Brazil". Ref. — Austen, 1907:333.
- ANALIS (Fabricius), 1805:311 (*Tachina*). Type-locality: "South America". Unrecognized. Ref. — Townsend, 1931:157 (note on type).
- ANALIS (Macquart), 1843:196 (1843:39), pl. 3, fig. 8 (*Jurinea*). Type-locality: "Brazil". (This species is a secondary homonym of the preceding one; as both are unrecognized I prefer not to rename it at present). Unrecognized.
- ANGRENSIS Guimarães, 1963b:161, figs. 25, 31. Type-locality: Brazil, Rio de Janeiro, Angra dos Reis.
- APICIFER (Walker), 1849:718 (*Tachina*). Type-locality: "North America". Distr. — U.S.A., Mexico, Tropical South America. Ref. — Guimarães, 1961:262.
- vulgaris* Curran, 1928:204, 276. Type-locality: U.S.A., Kansas.
- analís* authors, in part, not fabricius.
- basilfinus* authors, not Fabricius.
- ARAUJOI Guimarães, 1963d:336, figs. 35, 41. Type-locality: Brazil, Pará, Utinga.
- ARNAUDI Guimarães, 1963b:158, figs. 9-16. Type-locality: Brazil, Mato Grosso, Fazenda Murtinho.
- ATERRIMUS (Robineau-Desvoidy), 1830:35 (*Jurinea*). Type-locality: U.S.A., Carolina. Distr. — U.S.A., Mexico.
- AUSTRALIS (Macquart), 1855:117 (1854:97) (*Jurinea*). Type-locality: "America" (cf. Brauer, 1898:500) (as "New Holland", error). Unrecognized.

- AURIFRONS (Townsend), 1917b:228 (*Ncoarchytas*; as *inambaricus* ssp.).
Type-locality: "Brazil". Ref. — Thompson, 1963a:371. N. comb.
- BASIFULVUS (Walker), 1849:725 (*Tachina*). Type-locality: Jamaica, West Indies. Distr. — Neotropical.
- andicula* Townsend, 1915h:186. Type-locality: Peru, Chosica. N. syn.
- antillicolla* Curran, 1927c:2, fig. 2. Type-locality: Porto Rico, Albonito. N. syn.
- BENNETTI Thompson, 1963a:366, figs. 39, 42, 44, 46. Type-locality: Trinidad, Simla, Santa Cruz.
- BIEZANKOI Guimarães, 1961b:390, figs. 97-101. Type-locality: Brazil, Rio Grande do Sul, Pelotas.
- BRUCHI (Blanehard), 1941:360 (*Parafabricia*). Type-locality: Argentina, Buenos Aires. N. comb.
- BRUCHI (Blanchard), 1941:348, figs. 5, 6 (*Archytodejeania*). Type-locality: Argentina, Salta. N. comb. (This species is a secondary homonym of the preceding, but will not be renamed because their identities require clarification).
- CARONIENSIS Thompson, 1963a:380, figs. 49, 51, 53. Type-locality: Trinidad, Brazil Village.
- CARRERA Guimarães, 1961b:389, figs. 92-95. Type-locality: Brazil, Goiás, Anápolis.
- CIRPHIS Curran, 1927a:497. Type-locality: Mexico, Sinaloa, Los Moctis. Distr. — West Indies, Tropical South America.
- cirphisac* Curran in Swezey, 1926:226. *Nomen nudum*.
- CRUCIUS (Giglio-Tos), 1893:2 (*Nemochacta*). Type-locality: Mexico, Tuxpango. Unrecognized.
- DAEMON (Wiedemann), 1830:292 (*Tachina*). Type-locality: "Brazil".
- DAMIPPUS (Walker), 1849:719 (*Tachina*). Type-locality: Mexico (as unknown). Ref. — Austen, 1907:337.
- DIAPHANUS (Fabricius), 1805:349 (*Tachina*). Type-locality: "South America". Ref. — Guimarães, 1960:116.
- immaculata* Macquart, 1851:142 (1851:169), pl. 15, fig. 8 (*Echinomyia*). Type-locality: Brazil, Minas Gerais.
- pumila* Macquart, 1851:143 (1851:170) (*Echinomyia*). Type-locality: Brazil, Minas Gerais.
- bicolor* Jaennicke, 1867:392 (1868:84), pl. 44, fig. 8. Type-locality: "Venezuela".
- flavicans* Townsend, 1908:112 (*Eufabricia*). Type-locality: "Brazil".
- DISSIMILIS (Wulp), 1888:39, pl. 2, fig. 18 (*Nemochacta*). Type-locality: Costa Rica, Caché.
- DISSIMILIDES Thompson, 1963a:369, figs. 55, 58, 67. Type-locality: Trinidad, Maracas Valley.
- DIVISUS (Walker), 1852:270 (*Tachina*). Type-locality: Brazil, Pará. Distr. — Neotropical.
- incasanus* Townsend, 1912a:331. Type-locality: Peru, Piura. N. syn.
- DUBIUS (Giglio-Tos), 1893:2 (*Nemochacta*). Type-locality: "Mexico". Unrecognized.
- LUCKE Guimarães, 1961b:378, figs. 50-56. Type-locality: Brazil, Amazonas. Distr. — Brazil (Amazonas, São Paulo).
- DUX Curran, 1928b:279, figs. 35-36. Type-locality: "Ecuador".
- FLAVIFACIES (Macquart), 1851:144 (1851:171) (*Jurinia*). Type-locality: "Brazil" N. comb.
- FLAVIFRONS (Jaennicke), 1867:39 (1868:82) (*Jurinia*). Type-locality: "Mexico".

- FRENGUELLI (Blanchard), 1941:345, figs. 2-3 (*Archynemochacta*). Type-locality: Argentina, Salta.
- FRONTALIS (Wulp), 1892:194 (*Nemochacta*). Type-locality: Mexico, Atoyac. Unrecognized.
- FULVIVENTRIS (Robineau-Desvoidy), 1830:37 (*Jurinia*). Type-locality: Brazil, Rio Grande do Sul. Unrecognized.
- GIACOMELLI (Blanchard), 1941:365, figs. 13-15 (*Proarchytoides*). Type-locality: Argentina, La Rioja.
- GONCALVESI Guimarães, 1963b:159, figs. 17-24. Type-locality: Brazil, Goiás, Anápolis.
- HIEMALIS Thompson, 1963a:368, figs. 47, 50, 57. Type-locality: Trinidad, Santa Cruz Valley.
- INAMBARICUS (Townsend), 1915m:431 (*Neoarchytas*). Type-locality: Peru, junction of Yahuar Mayo and Inambari Rivers.
- INCERTUS (Giglio-Tos), 1893:2 (*Nemochacta*). Type-locality: Mexico, Oaxaca. Unrecognized. (This species is a secondary homonym of the following, but will not be renamed because its identity requires clarification).
- INCERTUS (Macquart), 1851:152 (1851:179) (*Gonia*). Type-locality: Argentina, Corrientes. Distr. — Neotropical. Ref. — Sabrosky, 1955:78; Guimarães, 1961a.
- pilliventris* Wulp, 1883:22 (*Echinomyia*). Type-locality: "Argentina".
- brasiliensis* Townsend, 1927a:354 (*Pseudoarchytopsis*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- pilliventris* of authors, not Wulp (in part).
- INFUMATUS (Bigot), 1887:cxxi (*Fabricia*). Type-locality: "Mexico". Unrecognized. Ref. — Brauer, 1898:495 (notes on type).
- INFUSCATUS (Wulp), 1892:194 (*Nemochacta*). Type-locality: Mexico, Teapa. Unrecognized.
- INNOVATUS (Walker), 1860:296 (*Jurinia*). Type-locality: "Mexico". Ref. Austen. 1907:336.
- INTRITUS (Walker), 1860:297 (*Nemoraea*). Type-locality: Mexico. Unrecognized.
- JURINOIDES (Giglio-Tos), 1893:2 (*Nemochacta*). Type-locality: Mexico, Oaxaca. Unrecognized.
- LANEI Guimarães, 1961b:386, figs. 80-85. Type-locality: Brazil, São Paulo, Barueri. Distr. — Colombia, Brazil (São Paulo).
- LATERALIS (Macquart), 1843:199 (1843:42), pl. 3, fig. 10 (*Jurinia*). Type-locality: "Mexico".
- apicalis* Jaenicke, 1867:390 (1868:82) (*Jurinia*). Type-locality: "Mexico".
- gonioides* Bigot, 1887:cxi (*Jurinia*). Type-locality: "Mexico".
- dispar* Wulp, 1890:34, pl. 2, fig. 14a (*Echinomyia*). Type-locality: Mexico, Northern Sonora.
- congruens* Wulp, 1892:192 (*Jurinia*). Type-locality: "Mexico".
- LENKOL Guimarães, 1961b:387, figs. 86, 91. Type-locality: Brazil, São Paulo, Ilha Sêca.
- LESCHENALDI (Robineau-Desvoidy), 1830:40 (*Peleteria*). Type-locality: "Surinam". Ref. — Brauer, 1898:497. Unrecognized.
- LOBULATUS Curran, 1928b:279, figs. 34-34a. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).
- LOPEZI Guimarães, 1961b:383, figs. 68-73. Type-locality: Brazil, Mato Grosso, Bodoquena. Distr. — Brasil (Mato Grosso to Santa Catarina).

- MARMORATUS (Townsend), 1915h:186 (*Pseudoarchytas*). Type-locality: Peru, Chosica. Distr. — U.S.A., Neotropical. Ref. — Sabrosky, 1955:78, figs. 1-3.
- incertus* N. American authors, not Macquart.
- piliventris*, authors, not Wulp (in part).
- METALLICUS Robineau-Desvoidy, 1830:35 (*Jurinia*). Type-locality: U.S.A., Carolina. Distr. — U.S.A., Mexico.
- hystrix* authors, not Fabricius.
- pilosa* Drury of Curran, 1928:206.
- MISIONENSIS (Blanchard), 1941:363 (*Parafabricia*). Type-locality: Argentina, Misiones.
- NEITILUCUS (Wulp), 1892:191 (*Jurinia*). Type-locality: "Mexico". Unrecognized.
- NIGRIVENTRIS (Wulp), 1882:81 (*Jurinia*). Type-locality: "Chile".
- NIGROCALYPTRATUS (Maequart), 1846:273 (1846:145), pl. 21, fig. 6 (*Jurinia*). Type-locality: Brazil, Minas Gerais, Unrecognized.
- NIGROCALYPTRATUS (Wulp), 1888:33 (*Echinomyia*). Type-locality: "Mexico". Unrecognized. (This species is a secondary homonym of the preceding; as both are unrecognized I prefer not to rename it at present).
- NITIDUS (Wulp), 1892:194 (*Nemochaeta*). Type-locality: "Mexico". Ref. — Brauer & Bergenstamm, 1891:131 (referred to *Tachinodes*).
- PALLIATUS Curran, 1928b:250, fig. 19. Type-locality: Costa Rica, Carillo.
- PEARSONI Guimarães, 1963d:337, figs. 42-48. Type-locality: Brazil, Pará.
- PERUANUS Curran, 1928b:249, figs. 17-18. Type-locality: Peru, Oroya. Distr. — Peru, Bolivia.
- PERNOX (Giglio-Tos), 1893:2 (*Nemochaeta*). Type-locality: Mexico, Orizaba. Unrecognized.
- PERPLEXA (Townsend), 1931a:354 (*Parafabricia*; nom. nov. for *bicolor* Wiedemann of Brauer & Bergenstamm). Type-locality: "Brazil".
- bicolor* Wiedemann of Brauer & Bergenstamm, 1894:76 (misident., according to Townsend, 1931a:354).
- PIARCONENSIS Thompson, 1963a:387, figs. 54, 59, 62. Type-locality: Trinidad, Piarco.
- PILIFRONS (Schiner), 1868:331 (*Echinomyia*). Type-locality: "Chile". Ref. — Sabrosky, 1955:83 (syn. note).
- nudigaena* Brauer, 1898:500 (*Jurinia*). Type-locality: "Chile and Montevideo".
- pollinosus* Curran, 1928b:251, fig. 20. Type-locality: "Chile".
- scutellatus* authors, not Macquart.
- PILOSUS (Walker), 1852:266 (*Tachina*). Type-locality: "South America". Unrecognized.
- PLANGENS Curran, 1928b:255, fig. 27. Type-locality: "Ecuador". Distr. — Ecuador, Guatemala.
- PRODUCTUS Curran, 1928b:276. Type-locality: Peru, Chosica.
- PRUDENS Curran, 1928b:275, fig. 28a. Type-locality: Mexico, Yucatán.
- PSEUDODAEEMON (Blanchard), 1940:225, fig. 1 (*Itarchytas*). Type-locality: Argentina, Córdoba, Cruz Chica. Distr. — Brazil (Goiás to São Paulo), Argentina.
- PURSEGLOVEI Thompson, 1963a:364, figs. 38, 41, 45. Type-locality: Trinidad, Santa Cruz Valley.
- RUSSATUS Reinhard, 1962b:170. Type-locality: Mexico, Cuernavaca, Morelos.
- SABROSKYI Guimarães, 1963b:162, figs. 32-37. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Goiás to Santa Catarina).



- SANCTAECRUCIS* Thompson, 1963a:386, figs. 60, 63, 66. Type-locality: Trinidad, Santa Cruz Valley.
- SCUTELLATUS* (Macquart), 1843:198 (1843:41) (*Jurinia*). Type-locality: "Chile, Colombia, Bogotá". Ref. — Cortés, 1963:246 (notes on types).
- infirmus* (Walker), 1849:719 (*Tachina*). Type-locality: "Chile".
- andana* Robineau-Desvoidy, 1863:657 (*Jurinia*). Type-locality: "Chile".
- ignobilis* Rondani, 1863:15 (*Echinomyia*). Type-locality: "Chile".
- chilensis* Curran, 1928b:222. Type-locality: "Chile".
- SEABRAI* Guimarães, 1961b:376, figs. 45-49. Type-locality: Brazil, Guanabara, Rio de Janeiro.
- FEMINIGER* (Wiedemann), 1830:296 (*Tachina*). Type-locality: "Brazil". Ref. — Guimarães, 1963b:155, figs. 1-8.
- SETIFACIES* Curran, 1928b:250. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).
- SHANNONI* Guimarães, 1960:120, figs. 21-32. Type-locality: Brazil, Mato Grosso, Maracaju. Distr. — Brazil (Mato Grosso, Pernambuco, Rio de Janeiro).
- fulvicentris* Townsend, 1935:219 (*Makasinocerops*; preocc. Robineau-Desvoidy, 1830). Type-locality: Brazil, Pernambuco, Tapera. N. SYN.
- SIBILLANS* Curran, 1928b:225, fig. 16. Type-locality: Guyana, Kartabo and Peru, Puerto Bermúdez. Ref. — Curran, 1934c:519 (redescr.).
- SMARAGDINUS* (Macquart), 1843:196 (1843:39), pl. 3, fig. 6 (*Jurinia*). Type-locality: "Brazil". Unrecognized.
- TOWNSENDI* Curran, 1928b:219, fig. 9. Type-locality: Mexico, Veracruz, San Rafael.
- TRANSLUCENS* (Macquart), 1846:273 (1846:145) (*Jurinia*). Type-locality: Brazil, Minas Gerais.
- TRAVASSOSI* Guimarães, 1961a:173, figs. 29-35. Type-locality: Brazil, Rio de Janeiro, Angra dos Reis.
- TRINITATIS* Thompson, 1963a:374, figs. 48, 52, 56. Type-locality: Trinidad, St. Augustine.
- UNGUIS* (Townsend), 1915m:432 (*Makasinocera*). Type-locality: Peru, Santa Eulalia.
- VARICORNIS* Curran, 1928b:251, fig. 21. Type-locality: Mexico, Veracruz, San Rafael.
- VERNALIS* Curran, 1928b:280, fig. 37. Type-locality: Guyana, Bartica. Distr. — Colombia, Guyana, Brazil (Rio de Janeiro to Santa Catarina).
- VEXOR* Curran, 1928b:221, fig. 10. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).
- WAGNERI* (Blanchard), 1941:351, figs. 7-8 (*Archytodesjeania*). Type-locality: Argentina, Santiago del Estero. N. COMB.
- WILLISTONI* Curran, 1925c:9. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães). Distr. — Brazil (Mato Grosso, Goiás, Guanabara). Ref. — Guimarães, 1963d:335.
- ZIKANI* Guimarães, 1961b:375, figs. 41-44. Type-locality: Brazil, Minas Gerais, Passa Quatro.

Genus DEJEANIOPS Townsend

- DEJEANIOPS Townsend, 1913e:104. Type-species, *ollachea* Townsend (orig. des.). Ref. — Curran, 1947:86.
 BECKERI Engel, 1920:293, fig. E. Type-locality: "Venezuela". Distr. — Venezuela, Ecuador, San Gabán Canyon.
 FALLACIOSUS Engel, 1920:292. Type-locality: Unknown.
 OLLACHEUS Townsend, 1913e:105. Type-locality: Peru, Ollachea.

Genus DUMERILLIA Robineau-Desvoidy

- DUMERILLIA Robineau-Desvoidy, 1830:38. Type-species, *rubida* Robineau-Desvoidy (mon.).
Tuberculocera Townsend, 1927a:346 (as *Tuberulocera* by misprint, corrected in errata). Type-species, *ochracea* Townsend (orig. des.). = *rubida* (Robineau-Desvoidy).
 RUBIDA Robineau-Desvoidy, 1830:39. Type-locality: "Brazil". Distr. — Brazil (São Paulo to Rio Grande do Sul).
furiosa Walker, 1857:194 (*Echinomyia*). Type-locality: Brazil, Rio Grande do Sul.
ochracea Townsend, 1927a:363 (*Tuberculocera*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus EUDEJEANIA Townsend

- EUDEJEANIA Townsend, 1912a:334. Type-species, *subalpina* Townsend (orig. des.).
Bombylojeania Townsend, 1913a:352. Type-species, *Dejeania canescens* Macquart (orig. des.).
Eudejeaniops Blanchard, 1941:353. Type-species, *pseudopyrrhopoda* Blanchard (orig. des.).
 Ref. — Sabrosky, 1941 (rev.); Curran, 1941:3 (rev.).
 ALBIPILA Curran, 1941:3. Type-locality: Ecuador, Tungurahua Volcán, Minza Chica.
 ALDRICHI Sabrosky, 1947:146. Type-locality: Colombia, Bogotá. Distr. — Ecuador, Colombia.
 ANDEANA Sabrosky, 1947:152. Type-locality: Bolivia, Cuesta de Cillunticara. Distr. — Venezuela, Bolivia.
 ARGYROPA (Schiner), 1868:337 (*Dejeania*). Type-locality: "South America". Distr. — Colombia, Ecuador. Ref. — Curran, 1941:4.
 ATRATA (Guérin-Ménéville), 1844:549 (*Dejeania*). Type-locality: Colombia, Bogotá. Unrecognized.
 BIRABENI (Blanchard), 1941:355, figs. 9-10 (*Eudejeaniops*). Type-locality: Argentina, Salta, Lumbreras. Ref. — Sabrosky, 1947:145.
 BROWNI Curran, 1941:5. Type-locality: Ecuador, Tungurahua Volcán, Minza Chica.
 CANESCENS (Macquart), 1846:271 (1846:143), pl. 12, fig. 1 (*Dejeania*). Type-locality: "Colombia" (as "Nouvelle Grenade"). Ref. — Curran, 1941:3.
 FEMORALIS Curran, 1941:4. Type-locality: Panama, Chiriquí, El Volcán.
 HUASCARAYANA Townsend, 1914:171. Type-locality: Peru, Huascaray.
 MELANAX (Walker), 1849:700 (*Tachina*). Type-locality: "Venezuela". Distr. — Venezuela, Colombia, Ecuador. Ref. — Sabrosky, 1947:155.

- podleera* Rondani, 1864:17, pl. 5, fig. 14 (*Dejeania*). Type-locality: "South America".
- MEXICANA (Robineau-Desvoidy), 1863:652 (*Dejeania*). Type-locality: "Mexico". Unrecognized. Ref. — Sabrosky, 1947:155.
- NIGRA Townsend, 1912a:335. Type-locality: Peru, Río Rímac Valley, Matucana.
- alpina* Townsend, 1913:106. Type-locality: Peru, Pachacayo Valley.
- NUDITIBIA Sabrosky, 1947:152. Type-locality: Colombia, Cundinamarca, Usaquén. Distr. — Venezuela, Colombia, Ecuador.
- PACHECOI Curran, 1941:5. Type-locality: Guatemala, Guatemala City.
- PALLIDA (Robineau-Desvoidy), 1863:653 (*Dejeania*). Type-locality: "Mexico". Unrecognized.
- PALLIPES (Macquart), 1843:191 (1843:34), pl. 2, fig. 9 (*Dejeania*). Type-locality: Colombia, Bogotá. Ref. — Sabrosky, 1947:149.
- PILOSA Curran, 1941:3. Type-locality: Bolivia, Mairi.
- PSEUDOPYRRHOPODA (Blanchard), 1941:353 (*Eudejeaniops*). Type-locality: Argentina, Jujuy. Ref. — Sabrosky, 1947:151.
- PUNENSIS Townsend, 1913e:105. Type-locality: Peru, Puno, Pachacayo Valley. Ref. — Sabrosky, 1947:144.
- PYRRHOPODA Engel, 1920:281, 287 (as *pallipes* var.). Type-locality: Peru, Madre de Dios and Urubamba.
- SUBALPINA Townsend, 1912:334 (as *pallida* var.). Type-locality: Peru, San Gabán River.

Genus EUJURINIODES Townsend

- EUJURINIODES Townsend, 1935d:219. Type-species, *eva* Townsend (orig. des.).
- ASSIMILIS (Wulp), 1892:192 (*Jurinia*). Type-locality: Mexico (sev. places). N. COMB.
- EVA Townsend, 1935d:219. Type-locality: "Trinidad".

Genus EULASIOPALPUS Townsend

- EULASIOPALPUS Townsend, 1913e:105. Type-species, *Lasiopalpus albipes* Townsend (orig. des.). Ref. — Curran, 1947:46 (key).
- ALDIPES (Townsend), 1913e:105 (*Lasiopalpus*). Type-locality: Peru, Huascaray.
- CORPULENTUS Townsend, 1914:173. Type-locality: Peru, Verrugas Canyon.
- GERTSCHII Curran, 1947:47. Type-locality: Panama, Chiriquí, El Volcán.
- NIVEUS Townsend, 1914:175. Type-locality: Peru, Verrugas Canyon.
- OBSCURUS Townsend, 1914:174. Type-locality: Peru, Matucana.
- TATEI Curran, 1947:47. Type-locality: Brazil, Roraima, Roraima Mts.
- TYPICUS Curran, 1947:46. Type-locality: Colombia, Call Distr., West Cordillera.
- SUBALPINUS Townsend, 1912a:335. Type-locality: Peru, Canyon of San Gabán River, Watkins camp.
- VITTATUS Curran, 1947:46, figs. 1-4. Type-locality: Ecuador, Bolívar, Hacienda Talahua.

Genus EXOPALPUS Macquart

- EXOPALPUS Macquart, 1851:149 (1851:176). Type-species, *bicolor* Macquart (orig. des.).

- Arthrochaeta* Brauer & Bergenstamm, 1889:134 (1889:66). Type-species, *demoticooides* Brauer & Bergenstamm (orig. des., as gen. n. sp. n.).
- Microtrichomma* Giglio-Tos, 1893:1. Type-species, *Nemoraea forreri* Wulp (Coquillett, 1910:570). N. SYN.
- Eurythiopsis* Townsend, 1914:30. Type-species, *ochracea* Townsend (orig. des.). N. SYN.
- Microtrichommodes* Townsend, 1927:245. Type-species, *elegans* Townsend (orig. des.). N. SYN.
- ANDINA (Townsend), 1914:169 (*Jurinella*). Type-locality: "Peru".
- BICOLOR Maequart, 1851:150 (1851:177), pl. 16, fig. 1. Type-locality: "Colombia".
- DEMOTICOIDES (Brauer & Bergenstamm), 1889:134 (1889:66) (*Arthrochaeta*). Type-locality: "Venezuela" (as "Colombia", error). Ref. — Aldrich, 1925:130; Townsend, 1931:162 (notes).
- ELEGANS (Townsend), 1927a:330 (*Microtrichommodes*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. N. COMB.
- FORRERI (Wulp), 1894:49, pl. 2, figs. 21-21a (*Nemoraea*). Type-locality: Mexico, Veracruz, Atoyac. Ref. — Townsend, 1931:159. N. COMB.
- INFUSCATA (Wulp), 1888:12 (*Hystriola*). Type-locality: Mexico, near Veraeruz. N. COMB.
- INTERMEDIA (Wulp), 1890:50 (*Nemoraea*). Type-locality: Mexico, Guerrero and Mexico City. N. COMB.
- MAJOR (Curran), 1925e:8 (*Jurinella*). Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães). N. COMB.
- MINOR (Curran), 1925e:7 (*Jurinella*). Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães). N. COMB.
- NOTATA (Bigot), 1888a:82 (*Echinomyia*). Type-locality: "Mexico". Unrecognized. N. COMB.
- OCHRACEA (Townsend), 1914:31 (*Eurythiopsis*). Type-locality: Peru, San Gabán Canyon. Ref. — Curran, 1947:86. N. COMB.
- SMITHI (Wulp), 1891:50 (*Nemoraea*). Type-locality: Mexico, Veraeruz. Distr. — U.S.A., Mexico.

Genus FABRICIOPSIS Townsend

- FABRICIOPSIS Townsend, 1914:83. Type-species, *hystrix* Townsend (orig. des.).
- ARGENTINENSIS Blanchard, 1941:357, fig. 11. Type-locality: Argentina, Misiones.
- HYSTRIX Townsend, 1914:83. Type-locality: Peru, San Gabán, Casahuiri.

Genus HEGESINUS Reinhard

- HEGESINUS Reinhard, 1964e:11. Type-species, *griphus* Reinhard (orig. des.).
- GRIPHUS Reinhard, 1964c:11. Type-locality: Mexico, Morelos, Topoztlán.

Genus HYSTRICIELLA Townsend

- HYSTRICIELLA Townsend, 1915k:95. Type-species, *aurifrons* Townsend (orig. des.) = *pilosa* (Drury).
- PILOSA (Drury), 1770:109, pl. 45, fig. 7 (*Musca*). Type-locality: "West Indies, Jamaica".

hystriz Fabricius, 1775:777 (*Musca*). Type-locality: "America".
exilis Townsend, 1892b:146 (*Pseudohystricia*). Type-locality: "Jamaica".
aurifrons Townsend, 1915k:95. Type-locality: "Jamaica".

Genus JURINELLA Brauer & Bergenstamm

- JURINELLA Brauer & Bergenstamm, 1889:132 (1889:64). Type-species,
Jurinea caeruleonigra Macquart (mon.).
Pseudohystricia Brauer & Bergenstamm, 1889:132 (1889:64). Type-species,
Hystricia ambigua Macquart (mon.).
? *Eufurinia* Townsend, 1908:113. Type-species, *Hystricia pollinosa* Wulp
(misident.) = *jicaltepecki* Townsend.
? *Neofurinia* Townsend, 1914:48, 81. Type-species, *abscondens* Townsend
(orig. des.).
? *Eufurinella* Townsend, 1914:86. Type-species, *abdominalis* Townsend
(orig. des.).
? *Tachinosoma* Townsend, 1927a:246. Type-species, *corpulentum* Townsend
(orig. des.).
? *Saundersiopsimima* Townsend, 1927a:246. Type-species, *spinosa* Townsend
(orig. des.).
? *Eufuriniopsis* Townsend, 1931a:351. Type-species, *Hystricia pollinosa*
Wulp (orig. des.).
? *Gigantachinosoma* Townsend, 1932b:106. Type-species, *gigantum* Town-
send (orig. des.).
? *Tachinosomopsis* Blanchard, 1941:370. Type-species, *jufuyensis* Blanchard
(orig. des.).
Ref. — Curran, 1947:63 (key).
ABDOMINALIS (Townsend), 1914:87 (*Eufurinella*). Type-locality: Peru, San
Gabán Canyon, Uruhuasi.
ABSCONDENS (Townsend), 1914:81 (*Neofurinia*). Type-locality: Peru, Santa
Eulalia.
AJAX Curran, 1947:81, figs. 67, 69. Type-locality: Brazil, Santa Catarina,
Nova Teutônia.
AMBIGUA (Macquart), 1851:145 (1851:172) (*Hystricia*). Type-locality: Mexico,
Guerrero. Ref. — Curran, 1947:66.
spinigera Wulp, 1892:192 (*Jurinia*). Type-locality: Mexico, Guerrero.
ANAX Curran, 1947:74, fig. 50. Type-locality: Brazil, Santa Catarina, Nova
Teutônia.
ANDICOLA Townsend, 1914:169. Type-locality: Peru, Chosica, Santa Eulalia
and Verrugas Canyon.
APICATA Curran, 1947:77, figs. 56, 60. Type-locality: Peru, upper River
Tapiche.
ARIEL Curran, 1947:80, fig. 64. Type-locality: Brazil, Santa Catarina, Nova
Teutônia.
BELLA Curran, 1947:71, fig. 46. Type-locality: Brazil, Santa Catarina, Nova
Teutônia.
BICOLOR (Wiedemann), 1830:282 (*Tachina*). Type-locality: "Brazil".
CIRCULARIS Curran, 1947:69, fig. 41. Type-locality: Brazil, Santa Catarina,
Nova Teutônia.
CAERULEONIGRA (Macquart), 1846:274 (1846:146), pl. 12, fig. 8 (as *cocculi-*
nigra, index, pg. 361) (*Jurinia*). Type-locality: "Colombia" (as
"Nouvelle Grenade"). Unrecognized. Ref. — Curran, 1947:66, fig. 33.
CONNOTA Curran, 1947:76, fig. 57. Type-locality: Brazil, Santa Catarina,
Nova Teutônia.

- CORPULENTA (Townsend), 1927a:359 (*Tachinosoma*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. Ref. — Curran, 1947:69, figs. 39, 43.
- CROSSI (Blanchard), 1942:361, fig. 10 (*Pseudohystricia*). Type-locality: Argentina, Tucumán.
- DEBITRIX (Walker), 1860:296 (*Jurinia*). Type-locality: "Mexico".
- EGLÉ Curran, 1947:78. Type-locality: "Mexico".
- EPILEUCA (Walker), 1849:716 (*Tachina*). Type-locality: "Jamalca". N. COMB.
- FEMINEA Curran, 1947:73. Type-locality: Brazil, Rio Grande do Sul, Santa Cruz.
- FERRUGINEA Townsend, 1929:382. Type-locality: Brazil, São Paulo, São Paulo, Cantareira.
- FUSCICORNIS Curran, 1925e:8. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães). Ref. — Curran, 1947:74, fig. 70.
- fusicornis* Curran, 1947:64 (error).
- GERTSCHII Curran, 1947:75, figs. 52, 53, 55. Type-locality: Panama, Chiriquí, El Volcán.
- GIGANTEA (Townsend), 1932b:106 (*Gigantachinosoma*). Type-locality: Brazil, Minas Gerais, Guaxupé.
- HUNTINGTONI Curran, 1947:82. Type-locality: Colombia, between Quereima and Buenaventura.
- JICALTEPECIA (Townsend), 1931a:354 (*Eufurinia*; nom. nov. for *pollinosa* Wulp of Townsend). Type-locality: Mexico, San Rafael, Jicaltepec, Veracruz.
- pollinosa* Wulp of Townsend, 1908:113 (*Eufurinia*), misident.
- JUJUYENSIS (Blanchard), 1941:370 (*Tachinosomopsis*). Type-locality: Argentina, Jujuy.
- KOEHLERI (Blanchard), 1941:368, fig. 16 (*Pseudohystricia*). Type-locality: Argentina, Salta, Metán.
- LATA Curran, 1947:68. Type-locality: Ecuador, Upano River, Sucre.
- LUTZI Curran, 1947:66, figs. 35-37. Type-locality: U.S.A., New Mexico. Distr. — U.S.A., Mexico.
- MACULATA Vimmer & Soukup, 1940:361 (as sp. I, "*maculata*"). Type-locality: Peru, Puno. Unrecognized.
- MAJOR Curran, 1925e:8. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães). Ref. — Curran, 1947:70, fig. 72.
- MEXICANA Curran, 1947:66, fig. 34. Type-locality: Mexico, Tlaxiapa.
- MILLERI Curran, 1947:67. Type-locality: Panama, Chiriquí, El Volcán.
- MINUTA Curran, 1947:72, figs. 45, 49. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- NIGRIVENTRIS Vimmer & Soukup, 1940:361 (? *Jurinella*). Type-locality: Peru, Puno. Unrecognized.
- NIVEISQUAMMA Engel, 1920:328, fig. B (*Bombyliomyia*). Type-locality: Bolivia, Cuesta de Cilluncara.
- OBESA (Wiedemann), 1830:282 (*Tachina*). Type-locality: "Brazil". Unrecognized.
- PALPALIS Curran, 1947:82. Type-locality: Mexico, Nayarit, Compostela.
- PANAMENA Curran, 1947:70. Type-locality: Panama, Canal Zone, Barro Colorado.
- POLLINOSA (Wulp), 1888:14, pl. 1, fig. 8 (*Hystricia*). Type-locality: Guatemala, San Gerónimo and Costa Rica (Río Sucio and Caché). Ref. — Curran, 1947:73, fig. 44.
- PROCTERI Curran, 1947:77, fig. 65. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- PRODUCTA Curran, 1947:72, figs. 47-48. Type-locality: Brazil, Santa Catarina, Nova Teutônia.

- PROFUSA* Curran, 1947:79. Type-locality: Guatemala, Guatemala City.
REDUCTA Curran, 1947:79, figs. 63, 68. Type-locality: Brazil, Rio de Janeiro.
RUFIVENTRIS Vimmer & Soukup, 1940:361 (as sp. II, "*rufiventris*"). Type-locality: Peru, Puno. Unrecognized.
SALLA Curran, 1947:71, fig. 42. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
SCHWARZI Curran, 1947:78, fig. 59. Type-locality: Colombia, West Cordillera, Cali Distr.
SPINOSA (Townsend), 1927a:356 (*Saundersi*pmima). Type-locality: Brazil, São Paulo, Itaquaquecetuba. Ref. — Curran, 1947:69, fig. 40.
TIHORACICA Curran, 1925c:9. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).
VAGA Curran, 1947:75, figs. 51, 54. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
VARGAS Curran, 1947:68, fig. 38. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
VARIANS Curran, 1947:76, figs. 58, 61. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
ZETEKI Curran, 1947:80. Type-locality: Panama, Canal Zone, Corozal.

Genus JURINIOPSIS Townsend

- JURINIOPSIS* Townsend, 1916d:73. Type-species, *floridensis* Townsend (orig. des.). Ref. — Sabrosky, 1969 (revision).
ADUSTA (Wulp), 1890:28, pl. 2, fig. 7 (*Jurinia*). Type-locality: Mexico, Durango, Ventanas.
AURIFRONS Brooks, 1949:21. Type-locality: U.S.A., New Mexico. Distr. — U.S.A., Mexico.
metallica authors, in part, not Robineau-Desvoidy.
FLORIDENSIS Townsend, 1916e:73. Type-locality: U.S.A., Florida.
basalis Walker, 1849:713 (*Tachina*; preocc. Walker, 1837). Type-locality: "Jamaica".
adusta, authors, in part, not Wulp.
INSULARIS Curran, 1960:6. Type-locality: Cuba, Havana, San Antonio, Baños (cf. Arnaud, 1963a:119).
LAMPURIS Reinhard, 1953b:93. Type-locality: Mexico, Morelos. Distr. — U.S.A., Mexico.
NITIDULA (Wulp), 1892:191 (*Jurinia*). Type-locality: Mexico, Guerrero (sev. places).
NIDITIVENTRIS (Curran), 1928b:205, figs. 2-3 (*Jurinia*). Type-locality: Mexico, Veracruz, San Rafael.
PERUANA Curran, 1960:4. Type-locality: Peru, Chosica, Verrugas Canyon.

Genus LINDIGIA Townsend

- LINDIGIA* Townsend, 1931a:352. Type-species, *Hystericia plagiata* Schiner (orig. des.).
 Ref. — Curran, 1947:84 (key).
BROWNI Curran, 1947:84, figs. 75, 78, 83. Type-locality: Ecuador, Bolivar, Hacienda Talahua.
PLAGIATA (Schiner), 1868:332 (*Hystericia*). Type-locality: "South America".
VARICOLOR Curran, 1947:84, figs. 77, 79, 82. Type-locality: Ecuador, Bolivar, Hacienda Talahua.
VIERECKI Curran, 1947:85. Type-locality: Colombia, Vista Nieve.

Genus MACROJURINIA Townsend

MACROJURINIA Townsend, 1916h:20. Type-species, *Jurinia brasiliensis* Robineau-Desvoidy (orig. des.).

BRASILIENSIS (Robineau-Desvoidy), 1830:35 (*Jurinia*). Type-locality: "Misiones" (= Brazil, Rio Grande do Sul, western part).

Genus PALPOTACHINA Townsend

PALPOTACHINA Townsend, 1915g:229. Type-species, *similis* Townsend (orig. des.).

SIMILIS Townsend, 1915g:230. Type-locality: Mexico, Chihuahua, Sierra Madre Mts., Rio Piedras Verdes Canyon.

Genus PARADEJEANIA Brauer & Bergenstamm

Jurinia, subg. PARADEJEANIA Brauer & Bergenstamm, 1893:147, 184 (1893:59, 96). Type-species, *Dejeania rutiloides* Jaenicke (Coquillett, 1910:584). Ref. — Arnaud, 1951 (rev.).

COLOMBIENSIS Arnaud, 1951:326, fig. 1. Type-locality: Colombia, Boyaca, Páramo, Pomeca River.

RUTILOIDES (Jaenicke), 1867:394 (1868:86) (*Dejeania*). Type-locality: "Mexico". Distr. — U.S.A., to Central America.

Genus PARAJURINIA Townsend

PARAJURINIA Townsend, 1928b:153. Type-species, *Tachina obesa* Townsend (orig. des.).

OBESA Townsend, 1928b:153. Type-locality: Brazil, Espírito Santo.

Genus PARARCHYTAS Brauer & Bergenstamm

PARARCHYTAS Brauer & Bergenstamm, 1894:76 (1895:612). Type-species, *Tachina decisa* Walker (mon.).

DECISUS (Walker), 1849:715 (*Tachina*). Type-locality: Canada, Ontario. Distr. — Canada, U.S.A., Mexico.

punctata Wulp, 1892:191 (*Jurinia*). Type-locality: "Mexico". N. SYN.

Genus PARECHINOTACHINA Townsend

PARECHINOTACHINA Townsend, 1931a:353. Type-species, *Dejeania plumitarsis* Wulp (orig. des.).

PLUMITARSIS (Wulp), 1886:xxxii (*Dejeania*). Type-locality: "Guatemala".

Genus PROTODEJEANIA Townsend

PROTODEJEANIA Townsend, 1915b:21. Type-species, *Dejeania hystricosa* Williston (orig. des.). Ref. — Curran, 1947 (key).

DICHROMA (Wulp), 1888:27, pl. 2, figs. 5-5a (*Jurinia*). Type-locality: Mexico, Durango. N. COMB.

DOWNSI Curran, 1947:50. Type-locality: Mexico, Mexico, D. F.

HYSTRICOSA (Williston), 1886:297 (*Dejeania*). Type-locality: U.S.A., New Mexico. Distr. — U.S.A., Mexico.

- MAJOR Curran, 1947:49, fig. 8. Type-locality: Guatemala, Guatemala City.
 MONTANA (Wulp), 1891:190 (*Dejeania*). Type-locality: Mexico, Guerrero.
 Ref. — Curran, 1947:48.
 PACHECOI Curran, 1947:48, fig. 9. Type-locality: Guatemala, Guatemala City.

UNPLACED SPECIES OF DEJEANIINI

- ATRATA Wulp, 1888:8, pl. 1, fig. 2 (*Dejeania*). Type-locality: Costa Rica, Volcan de Irazú.
 BADVENTRIS Wulp, 1888:28, pl. 2, fig. 6 (*Jurinia*). Type-locality: "Costa Rica".
 EYTHIRINA Bigot, 1888a:79 (*Hystrix*). Type-locality: Brazil, Bahia.
 TRIANGULI Walker, 1849:706 (*Tachina*). Type-locality: "Venezuela".

Tribe JURINIINI

Members of the tribe Jurinini parasitize larvae of Lepidoptera.

Genus ANEPALPUS Townsend

- ANEPALPUS Townsend, 1931a:444. Type-species, *hystrix* Townsend (orig. des.).
 HYSTRIX Townsend, 1931a:445. Type-locality: Peru, near Oxapampa, Río Tingo de Chontabamba.

Genus ARCHYTOEPALPUS Townsend

- ARCHYTOEPALPUS Townsend, 1927a:255. Type-species, *rufiventris* Townsend (orig. des.).
 RUFIVENTRIS Townsend, 1927a:268. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus AUSTENIOPS Townsend

- AUSTENIOPS Townsend, 1915h:96. Type-species, *Saundersia truncaticornis* Wulp (orig. des.).
 TRUNCATICORNIS (Wulp), 1888:26, pl. 2, fig. 4 (*Saundersia*). Type-locality: Panama, Chiriquí Volcano.

Genus BISCHOFIMYIA Townsend

- BISCHOFIMYIA Townsend, 1927a:250. Type-species, *atra* Townsend (orig. des.).
 ATRA Townsend, 1927a:290. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus CAMPOSIANA Townsend

- CAMPOSIANA Townsend, 1915k:95. Type-species, *emarginata* Townsend (orig. des.).
 EMARGINATA Townsend, 1915k:96. Type-locality: Ecuador, Quito.

Genus CHILOEPALPUS Townsend

- CHILOEPALPUS Townsend, 1927a:281. Type-species, *aureifacies* Townsend (orig. des.) = *callipyga* (Bigot). Ref. — Aldrich, 1934:122.

- AUREUS (Aldrich), 1926b:25 (*Cuphocera*). Type-locality: Chile, Angol. Distr. — Chile (Angol, Santiago).
 CALLIPYGUS (Bigot), 1857a:299 (*Jurinia*). Type-locality: Chile, Angol.
aurifacies Townsend, 1927a:281. Type-locality: Chile, Concepción.

Genus CHROMOEPALPUS Townsend

- CHROMOEPALPUS Townsend, 1914:158. Type-species, *uruhuasi* Townsend (orig. des.).
 URUHUASI Townsend, 1914:159. Type-locality: Peru, Uruhuasi bridge.

Genus CORPULENTOEALPUS Townsend

- CORPULENTOEALPUS Townsend, 1927a:249. Type-species, *rufus* Townsend (orig. des.).
 RUFUS Townsend, 1927a:300. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus CORPULENTOSOMA Townsend

- CORPULENTOSOMA Townsend, 1914:31. Type-species, *cornutum* Townsend (orig. des.).
 CORNUTUM Townsend, 1914:42. Type-locality: Peru, San Gabán Canyon, Uruhuasi.

Genus CRYPTOPALPUS Rondani

- CRYPTOPALPUS Rondani, 1850:169. Type-species, *Micropalpus ornatus* Macquart (Coquillett, 1910:528).
Saundersia Schiner, 1868:333. Type-species, *Micropalpus ornatus* Macquart (orig. des.).
Euquadratosoma Townsend, 1915m:75. Type-species, *rubrum* Townsend (orig. des.).
 AEQUABILIS (Walker), 1849:704 (*Tachina*). Type-locality: "Venezuela". Unrecognized.
 DISCALIS (Brèthes), 1909:95 (*Saundersia*). Type-locality: Argentina, Mendoza. Unrecognized.
 DIVERSUS (Walker), 1849:703 (*Tachina*). Type-locality: "Venezuela".
hystrix (Schiner) in Brauer & Bergenstamm, 1891:409 (1891:105) (*Saundersia*; preocc. Rondani, 1865). *Nomen nudum*.
 MARGINALIS (Brèthes), 1909:95 (*Saundersia*). Type-locality: Argentina, Mendoza. Unrecognized.
 NIGRIVENTRIS (Macquart), 1843:201 (1843:44), pl. 4, fig. 3 (*Hystrioides*). Type-locality: "Colombia".
hystrix Rondani, 1865:18. Type-locality: Colombia, Bogotá.
 ORNATUS (Macquart), 1843:204 (1843:47), pl. 4, fig. 6 (*Micropalpus*). Type-locality: "Mexico". Ref. — Townsend, 1931:167 (descr. notes).
pictipennis (Macquart) in Brauer & Bergenstamm, 1891:409 (1891:105) (*Saundersia*). *Nomen nudum*.
 RUBER (Townsend), 1915m:75 (*Euquadratosoma*). Type-locality: Peru, Uruhuasi.

- rubripes* (Schiner) in Brauer & Bergenstamm, 1891:409 (1891:105) (*Saundersia*). Colombia. *Nomen nudum*.
RUFIVENTRIS (Macquart), 1846:279 (1846:151), pl. 13, fig. 3 (*Micropalpus*). Type-locality: "Colombia" (as "Nouvelle Grenade").
SEMIATER (Schiner), 1868:334 (*Saundersia*). Type-locality: "South America".
TRANSIENS (Walker), 1849:706 (*Tachina*). Type-locality: Ecuador, Quito. Unrecognized.
TRANSVERSUS (Walker), 1852:274 (*Tachina*). Type-locality: "Brazil". Unrecognized.

Genus DIAPHANOMYIA Townsend

- DIAPHANOMYIA* Townsend, 1917b:229. Type-species, *aurea* Townsend (orig. des.).
Xanthozonella Townsend, 1931b:164. Type-species, *brasiliensis* Townsend (orig. des.).
AUREA Townsend, 1917:229. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).
AURIFACIES (Robineau-Desvoidy), 1830:38 (*Jurinta*). Type-locality: "Brazil".
brasiliensis Townsend, 1927a:365 (*Xanthozonella*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
diaphana Brauer & Bergenstamm, 1891:348 (1891:80) (*Paragymnomma*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
ludens Walker, 1860:295 (*Echinomyia*). Type-locality: "Brazil".

Genus ECHINOPYRRHOSIA Townsend

- ECHINOPYRRHOSIA* Townsend, 1914:90. Type-species, *alpina* Townsend (orig. des.). Ref. — Curran, 1941:1 (key).
ALPINA Townsend, 1914:91. Type-locality: Peru, Oroya, Pachacayo. Distr. — Peru, Bolivia.
ATYPICA Townsend, 1914:92. Type-locality: Peru, Pachacayo.
BROWNI Curran, 1941:2. Type-locality: Ecuador, Minza Chica, Volcano Tungurahua.
MELANICA Townsend, 1914:92. Type-locality: Peru, Pachacayo.
NOTATA (Walker), 1852:267 (*Tachina*). Type-locality: "Colombia".
PICTIPENNIS Curran, 1941:1. Type-locality: Ecuador, Tungurahua Volcán, Minza Chica.
TROPHOCYON Aldrich, 1928b:18. Type-locality: Ecuador, Tungurahua Mt.
VARIA (Walker), 1852:286 (*Tachina*). Type-locality: "Colombia".

Genus ECHINOPYRRHOSIOPS Townsend

- ECHINOPYRRHOSIOPS* Townsend, 1931a:443. Type-species, *Saundersia decorata* Townsend (orig. des.).
DECORATUS (Townsend), 1931a:443 (*Saundersia*). Type-locality: Bolivia, Cuesta de Chlunticara.

Genus EDWYNIA Aldrich

- Reedia* Aldrich, 1928b:17 (preocc. Ashmead, 1904). Type-species, *robusta* Aldrich (orig. des.).

- EDWYNIA Aldrich, 1930b:26 (nom. nov. for *Reedia*). Type-species, *Reedia robusta* Aldrich (aut.).
- ROBUSTA (Aldrich), 1928b:18 (*Reedia*). Type-locality: Chile, Concepción. Distr. — Chile, Argentina (Mendoza).

Genus EMPHEREMYIA Bischof

- EMIPHEREMYIA Bischof, 1904:87. Type-species, *atra* Bischof (Townsend, 1916:6).
- ATRA Bischof, 1904:88. Type-locality: Brazil, Rio Grande do Sul.
- LEOPOLDIENSIS (Brauer), 1897:369 (1897:41) (*Micropalpus*). Type-locality: Brazil, Rio Grande do Sul, São Leopoldo. Ref. — Bischof, 1904:88 (as *leopoldensis*, error).

Genus EMPHEREMYIOPS Townsend

- EMIPHEREMYIOPS Townsend, 1927a:254. Type-species, *discahis* Townsend (orig. des.).
- DISCALIS Townsend, 1927a:303. Type-locality: Peru, São Paulo, Itaquaquecetuba.

Genus EPALPELLUS Townsend

- EPALPELLUS Townsend, 1914:89. Type-species, *corpulentus* Townsend (orig. des.).
- CORPULENTUS Townsend, 1914:89. Type-locality: Peru, San Gabán Canyon, Uruhuasi.

Genus EPALPODES Townsend

- EPALPODES Townsend, 1912a:330. Type-species, *equatorialis* Townsend (orig. des.).
- ALBOLINEATUS (Macquart), 1854:119 (1854:99) (*Micropalpus*). Type-locality: "Colombia". N. COMB.
- CHILLANENSIS Cortés, 1951:258. Type-locality: Chile, Suble, Chillán, Termas de Chillán.
- EQUATORIALIS (Macquart), 1854:19 (1854:99) (*Micropalpus*). Type-locality: "Colombia". N. COMB.
- VITTATUS (Walker), 1852:273 (*Tachina*). Type-locality: "Colombia".
- EQUATORIALIS Townsend, 1912:330. Type-locality: Ecuador. (This species is a secondary homonym of the preceding one, but I prefer not to rename it at present).
- ssp. *rimacensis* Townsend, 1915h:184. Type-locality: "Peru".

Genus EPALPUS Rondani

- EPALPUS Rondani, 1850:169. Type-species, *Micropalpus rufipennis* Macquart (Coquillett, 1910:538).
- Eusignosoma* Townsend, 1914:123. Type-species, *aurcum* Townsend (orig. des.).
- Argentocpalpus* Townsend, 1919:178. Type-species, *Epalpus niveus* Townsend (orig. des.).

- AFFINIS (Schiner), 1868:336 (*Saundersia*). Type-locality: "South America". Unrecognized.
- ALBOMACULATUS (Jaennicke), 1867:368 (1868:80) (*Micropalpus*). Type-locality: "Mexico". Distr. — U.S.A., Mexico, Guatemala.
- ALLIGANS (Walker), 1849:713 (*Tachina*). Type-locality: "Venezuela". Unrecognized. N. comb.
- ALTERNUS (Walker), 1849:701 (*Tachina*). Type-locality: "Venezuela". Unrecognized. N. comb.
- AUREOLATUS (Townsend), 1914:125 (*Eusignosoma*). Type-locality: Peru, San Gabán Canyon, Casahuari.
- AUREUS (Giglio-Tos), 1893:3 (*Saundersia*). Type-locality: "Mexico". Unrecognized.
- AUREUS (Townsend), 1914:124 (*Eusignosoma*). Type-locality: Peru, San Gabán Canyon, Uruhuasi. (This species is a secondary homonym of the preceding one, but I prefer not to rename, at present).
- AURIFERUS (Walker), 1849:702 (*Tachina*). Type-locality: "Venezuela". Unrecognized.
- BOLIVIANUS Bischof, 1904:90. Type-locality: Bolivia (Corolco), Peru (*Callanga*) and Venezuela.
- BRUNNEIPENNIS Bischof, 1904:93. Type-locality: Peru, Callanga.
- CALLANGANUS Bischof, 1904:92. Type-locality: Peru, Callanga.
- CANUS (Wulp), 1888:25, pl. 2, fig. 8 (*Saundersia*). Type-locality: "Costa Rica".
- CONEANGUINEUS (Wulp), 1892:191 (*Saundersia*). Type-locality: "Mexico".
- CONTRARIUS (Walker), 1849:716 (*Jurinia*). Type-locality: "Mexico". N. comb.
- DENUDATUS Bischof, 1904:92. Type-locality: "Venezuela".
- FEMORATUS (Wulp), 1892:191 (*Saundersia*). Type-locality: Mexico, Guerrero. N. comb.
- FLAVICANS (Macquart), 1846:279 (1846:151), pl. 13, fig. 4 (*Micropalpus*). Type-locality: "Colombia". N. comb.
- FLAVIPES Vimmer & Soukup, 1940:220, 372. Type-locality: Peru, Ollachea, Puno. Unrecognized.
- FURCANIPENNIS Bischof, 1904:92. Type-locality: Bolivia (*Cillunticara*) and Brazil (Rio Grande do Sul).
- IMITATOR (Townsend), 1929:381 (*Argentoepalpus*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- JAENNICKI (Giglio-Tos), 1894:22 (*Saundersia*; nom. nov. for *rufipes* Jaennicke). Type-locality: "Panama". Unrecognized.
- rufipes* Jaennicke, 1867:387 (1867:79) (*Micropalpus*; preocc. Macquart, 1851). Type-locality: "Panama".
- LATICORNIS (Wulp), 1888:20, pl. 1, fig. 15-15a (*Saundersia*). Type-locality: Costa Rica, Rio Sucio. N. comb.
- LATIVITTUS (Walker), 1852:269 (*Tachina*). Type-locality: "Colombia" (as unknown) (cf. Austen, 1907:327).
- LEUCOMELANUS (Walker), 1849:714 (*Tachina*). Locality: unknown. N. comb.
- LINDIGI Bischof, 1904:90. Type-locality: "Venezuela".
- LINEATUS Townsend, 1914:136. Type-locality: Peru, Matucana.
- MACULUS (Macquart), 1843:203 (1843:45), pl. 5, fig. 2 (*Micropalpus*). Type-locality: "South America".
- melica* (Macquart) in Brauer, 1898:498 (*Hystriola*). Locality: "Nova Granada". *Nomen nudum*.
- MONTIVAGUS (Wulp), 1892:190 (*Saundersia*). Type-locality: Mexico, Guerrero. N. comb.
- NATTERERI Bischof, 1904:91. Type-locality: Brazil, Rio Grande do Sul, Minas Gerais and Espírito Santo.

- NIGER (Townsend), 1914:125 (*Eusignosoma*). Type-locality: Peru, San Gabán Canyon, Uruhuasi.
- NITIDUS (Macquart), 1851:147 (1851:174), pl. 15, fig. 4 (*Micropalpus*). Type-locality: "South America".
- NIVUS Townsend, 1914:136. Type-locality: Peru, Uruhuasi Canyon.
- OCHIRACEUS (Townsend), 1929:381 (*Argentoepalpus*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- PALLIATARSIS Rondani, 1850b:170. Type-locality: "Venezuela".
- PERUVIANUS (Macquart), 1848:205 (1848:45), pl. 5, fig. 2 (*Micropalpus*). Type-locality: Ecuador, Quito.
- PICEUS (Giglio-Tos), 1893:3 (*Saundersia*). Type-locality: "Mexico". N. COMB.
- PICTUS (Schiner), 1868:335 (*Saundersia*). Type-locality: "South America".
- PORTERI Brèthes, 1918:50. Type-locality: Chile, La Ligua. Ref. — Cortés, 1963:250 (notes on type).
- ROSTRATUS Rondani, 1868:25, pl. 4, figs. 1-2. Type-locality: Argentina, Mendoza.
- RUFIPENNIS (Macquart), 1846:280 (1846:152), pl. 13, fig. 5 (*Micropalpus*). Type-locality: "Colombia" (as "Nouvelle Grenade"). Ref. — Townsend, 1931b:166.
- RUFITIBIA (Wulp), 1888:24 (*Saundersia*). Type-locality: Mexico, Orizaba. N. COMB.
- RUFIVENTRIS (Macquart), 1843:198 (1843:41), pl. 3, fig. 9 (*Jurinia*). Type-locality: Colombia, Bogotá.
- SEMIATER Bischof, 1904:93. Type-locality: Peru, Callanga.
- SEMIFLAVUS Bischof, 1904:91. Type-locality: Peru (Callanga) and Bolivia (Songo).
- TARSALIS (Schiner), 1868:334 (*Saundersia*). Type-locality: "South America".
- TESTACEUS (Wulp), 1888:24, pl. 2, fig. 1-1a (*Saundersia*). Type-locality: "Costa Rica". N. COMB.
- UNICOLOR (Wulp), 1888:23 (*Saundersia*). Type-locality: Mexico, Orizaba. N. COMB.

Genus ERYTHROEPALPUS Townsend


- ERYTHROEPALPUS Townsend, 1931a:438. Type-species, *aurantiacus* Townsend (orig. des.).
- AURANTIACUS Townsend, 1931a:439. Type-locality: "Mexico".

Genus EUBISCHOFIMYIA Townsend

- EUBISCHOFIMYIA Townsend, 1927a:249. Type-species, *analis* Townsend (orig. des.).
- ANALIS Townsend, 1927a:305. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus EUCORPULENTOSOMA Townsend

- EUCORPULENTOSOMA Townsend, 1914:87. Type-species, *simile* Townsend (orig. des.).
- SIMILE Townsend, 1914:88. Type-locality: Peru, San Gabán Canyon, Uruhuasi.



Genus EUEMPHEREMYIA Townsend

- EUEMPHEREMYIA Townsend, 1927a:249. Type-species, *paulensis* Townsend (orig. des.). Ref. — Guimarães, 1963c (rev.).
- ALBUQUERQUEI Guimarães, 1963c:344, figs. 7-11. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- ELYOWALDI Guimarães, 1963c:348, figs. 22-29. Type-locality: Brazil, Rio de Janeiro, Teresópolis.
- NEMO (Curran), 1947:100, fig. 124 (*Euhuascaraya*). Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- PAULENSIS Townsend, 1927a:307. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus EUFPALPODES Townsend

- EUFPALPODES Townsend, 1915m:429. Type-species, *arcuatus* Townsend (orig. des.).
- ARCUATUS Townsend, 1915m:429. Type-locality: Peru, Uruhuasi.

Genus EUEPALPUS Townsend

- EUEPALPUS Townsend, 1908:115. Type-species, *flavicauda* Townsend (orig. des.).
- Xanthozonopsis* Townsend, 1916g:314. Type-species, *vestita* Townsend (orig. des.). N. SYN.
- FLAVICAUDA Townsend, 1908:115. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).
- VESTITUS (Townsend), 1916g:314 (*Xanthozonopsis*). Type-locality: Paraguay, Sapucay. N. COMB.

Genus EUFABRICIOPSIS Townsend

- EUFABRICIOPSIS Townsend, 1915b:22. Type-species, *Gymnomma quadrisetosa* Coquillett (orig. des.).
- QUADRISETOSA (Coquillett), 1902:120 (*Gymnomma*). Type-locality: Mexico, Chihuahua, Rio Piedras Verdes.

Genus EUHUASCARAYA Townsend

- EUHUASCARAYA Townsend, 1927a:255. Type-species, *atra* Townsend (orig. des.). Ref. — Curran, 1947:99 (key).
- ATRA Townsend, 1927a:307. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- MEDIA Curran, 1947:100, fig. 122. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- OBSCURA Curran, 1947:99, figs. 121, 123. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- SIESTA Curran, 1947:100, figs. 119, 120. Type-locality: Brazil, Santa Catarina, Nova Teutônia.

Genus EUMELANEPALPUS Townsend

- EUMELANEPALPUS Townsend, 1915m:74. Type-species, *ruber* Townsend (orig. des.).



RUBER Townsend, 1915m:74. Type-locality: Peru, San Gabán Canyon, Uruhuasi.

Genus EUSAUNDERSIOPS Townsend

EUSAUNDERSIOPS Townsend, 1915m:76. Type-species, *notata* Townsend (orig. des.) = *inortata* (Sehlnr).

INORTATUS (Sehlnr), 1868:335 (*Saundersia*). Type-locality: "South America".

notata Townsend, 1915m:76. Type-locality: Peru, Jaén, Huasearay.

Genus GIGANTOEPALPUS Townsend

GIGANTOEPALPUS Townsend, 1913a:443. Type-species, *heros* Townsend (orig. des.).

HEROS Townsend, 1931a:443. Type-locality: "Venezuela".

Genus GYMNOGMA Wulp

GYMNOGMA Wulp, 1888:38. Type-species, *nitidiventris* Wulp (orig. des.).

DIAPHANOIDES Curran, 1925e:10. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).

NOVUM Giglio-Tos, 1893:158. Type-locality: "Mexico".

NITIDIVENTRIS Wulp, 1888:38, pl. 5, fig. 17. Type-locality: Mexico, Durango. Ref. — Townsend, 1931b:167 (note on type).

Genus HOMOSAUNDERSIA Townsend

HOMOSAUNDERSIA Townsend, 1931a:439. Type-species, *Saundersia rufa* Sehlnr (orig. des.).

Rufa (Sehlnr), 1868:335 (*Saundersia*). Type-locality: "South America".

Genus HOMOSAUNDERSIOPS Townsend

HOMOSAUNDERSIOPS Townsend, 1931a:445. Type-species, *haenschi* Townsend (orig. des.).

HAENSCHI Townsend, 1931a:446. Type-locality: Ecuador, Baños.

Genus HUASCARAYOPSIS Townsend

HUASCARAYOPSIS Townsend, 1927a:254. Type-species, *paulensis* Townsend (orig. des.).

PAULENSIS Townsend, 1927a:317. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus ITASAUNDERSIA Townsend

ITASAUNDERSIA Townsend, 1927a:254. Type-species, *robusta* Townsend (orig. des.).

ROBUSTA Townsend, 1927a:321. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus JURINIA Robineau-Desvoidy

- JURINIA Robineau-Desvoidy, 1830:34. Type-species, *gagatea* Robineau-Desvoidy (Coquillett, 1910:556).
Proepalpus Townsends, 1927a:255. Type-species, *paulensis* Townsend (orig. des.).
 BARBATA Bigot, 1887:cxl. Type-locality: "Mexico".
 FELIGINIPENNIS Bigot, 1888a:79. Type-locality: Uruguay, Montevideo. Unrecognized.
 GAGATEA Robineau-Desvoidy, 1830:36. Type-locality: "Brazil".
 HYALIPENNIS (Macquart), 1835:80 (*Echinomyia*). Type-locality: "Brazil". Unrecognized.
 LATICORNIS Macquart, 1846:274 (1846:146). Type-locality: "Colombia".
 NIGRIVENTRIS Robineau-Desvoidy, 1863:659. Type-locality: Chile, Argentina.
 OLIVAUREA Townsend, 1914:159. Type-locality: Peru, Huascaray Ridge.
 PAULENSIS (Townsend), 1927a:350 (*Proepalpus*). Type-locality: Brasil, São Paulo, Itaquaquecetuba.
 EUFIPALPIS Macquart, 1843:197 (1843:40). Type-locality: "Guyane". Unrecognized.
 SURINAMENSIS Macquart, 1843:197 (1843:40), pl. 4, fig. 1. Type-locality: "Surinam".
 VERSICOLOR Robineau-Desvoidy, 1863:662. Type-locality: "Mexico".

Genus JURINIOSOMA Townsend

- JURINIOSOMA Townsend, 1927a:255. Type-species, *gagateum* Townsend (orig. des.).
 GAGATEUM Townsend, 1927a:323. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus LINDIGEPALPUS Townsend

- LINDIGEPALPUS Townsend, 1931a:444. Type-species, *Hystriola testacea* Macquart (orig. des.; preocc. Macquart, 1843) = *townsendi* Guimarães.
 TOWNSENDI Guimarães, nom. nov. for *testaceus* Macquart, 1846. Type-locality: "Colombia".
testaceus (Macquart), 1846:276 (1846:148) (*Hystriola*; preocc. Macquart, 1843). Type-locality: "Colombia".

Genus MELANEPALPELLUS Townsend

- MELANEPALPELLUS Townsend, 1927a:250. Type-species, *corpulentus* Townsend (orig. des.).
 CORPULENTUS Townsend, 1927a:327. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus MELANEPALPUS Townsend

- MELANEPALPUS Townsend, 1914:154. Type-species, *albipes* Townsend (orig. des.).
 ALBIPES Townsend, 1914:155. Type-locality: Peru, Huascaray Ridge.
 FELVUS Townsend, 1914:155. Type-locality: Peru, San Gabán River, Uruhuasi.

Genus NEOGYMNOMMA Townsend

- NEOGYMNOMMA Townsend, 1915e:69. Type-species, *rufa* Townsend (orig. des.).
RUFFA Townsend, 1915e:70. Type-locality: Peru, Huariaca, Río Hualliaga Canyon.

Genus OCHROEPALPUS Townsend

- OCHROEPALPUS Townsend, 1927a:255. Type-species, *ochraceus* Townsend (orig. des.).
CITRINUS Blanchard, 1941:373. Type-locality: Argentina, Misiones.
OCHIRACEUS Townsend, 1927a:337. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus OESTROHYSTRICIA Townsend

- OESTROHYSTRICIA Townsend, 1912:332. Type-species, *subalpina* Townsend (orig. des.).
SUBALPINA Townsend, 1912:333. Type-locality: Peru, San Gabán River.

Genus OPSOEMPHERIA Townsend

- OPSOEMPHERIA Townsend, 1927a:249. Type-species, *atra* Townsend (orig. des.).
ATRA Townsend, 1927a:340. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus OXYEPALPUS Townsend

- OXYEPALPUS Townsend, 1927a:253. Type-species, *Saundersia flavoscutellata* Bischof (orig. des.).
FLAVOSCUTELLATUS (Bischof), 1904:90 (*Epalpus*). Type-locality: "Brazil".
brasilensis Townsend, 1927a:343. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus PAREPALPUS Coquillett

- PAREPALPUS Coquillett, 1902:120. Type-species, *flavidus* Coquillett (mon.).
Oxapampoepalpus Townsend, 1931a:441. Type-species, *auroanalis* Townsend (orig. des.).
Ref. — Curran, 1947:101 (key).
AUROANALIS (Townsend), 1931a:441 (*Oxapampoepalpus*). Type-locality: Peru, Oxapampa.
CONSTANS (Walker), 1849:705 (*Tachina*). Type-locality: "Venezuela". N. COMB.
nigripilosus Wuip, 1858:23 (*Saundersia*). Type-locality: Mexico, Guerrero. N. SYN.
DISCORS (Wuip), 1852:193 (*Gymnomma*). Type-locality: Mexico, Guerrero.
LABEOSUS Reinhard, 1957:102. Type-locality: Mexico, Cuernavaca.
SIMILIS Townsend, 1914:134. Type-locality: Peru, San Gabán Canyon, Uruhuasi.

Genus PICTOEPALPUS Townsend

- PICTOEPALPUS Townsend, 1915c:71. Type-species, *clarus* Townsend (orig. des.).
CLARUS Townsend, 1915c:72. Type-locality: Peru, Tamboraque and Matucana.

Genus PSEUDOXANTHOZONA Townsend

- PSEUDOXANTHOZONA Townsend, 1931a:442. Type-species, *denudata* Townsend (orig. des.).
DENUDATA Townsend, 1931a:442. Type-locality: "Venezuela".

Genus PSEUDOXANTHOZONELLA Townsend

- PSEUDOXANTHOZONELLA Townsend, 1931a:437. Type-species, *similis* Townsend (orig. des.).
SIMILIS Townsend, 1931a:437. Type-locality: Ecuador, Balzapampa.

Genus QUADRATOSOMA Townsend

- QUADRATOSOMA Townsend, 1914:153. Type-species, *rufum* Townsend (orig. des.).
RUFUM Townsend, 1914:154. Type-locality: Peru, Uruhuasi.

Genus RHACHOEPALPUS Townsend

- RHACHOEPALPUS Townsend, 1908:114. Type-species, *Saundersia testacea* Wulp (Coquillett, 1910:599).
Rhachoealpodes Townsend, 1935d:220. Type-species, *quatuornotatus* Townsend (orig. des.).
Ref. — Curran, 1947:86 (key).
ANDINUS Townsend, 1914:142. Type-locality: Peru, Huascaray.
ARGENTUS Townsend, 1914:153. Type-locality: Peru, Verrugas Canyon and Matucana.
BEATUS Curran, 1947:89, fig. 92. Type-locality: Ecuador, River Mapóto.
BORNATUS Curran, 1947:88, fig. 91. Type-locality: Panama, Chiriquí, El Volcán.
BLANDUS Curran, 1947:90, fig. 93. Type-locality: Colombia, Vista Nieve.
CINEREUS Townsend, 1914:144. Type-locality: Peru, Matucana.
ETHELUS Curran, 1947:90. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
FLAVITARSIS (Macquart), 1843:204 (1843:47), pl. 5, fig. 1 (*Micropalpus*). Type-locality: "Mexico". N. comb.
IMMACULATUS (Macquart), 1846:276 (1846:148), pl. 12, fig. 9 (*Hystricia*). Type-locality: "Colombia".
METALLICUS Curran, 1947:87, figs. 84, 88, 89. Type-locality: Ecuador, Bolívar, Hacienda Talahua.
NITIDUS Townsend, 1914:144. Type-locality: Peru, Matucana.
NOTATUS Curran, 1947:87, figs. 85, 87. Type-locality: Ecuador, Bolívar, Hacienda Talahua.
NOVA Curran, 1947:89, fig. 95. Type-locality: Colombia, Vista Nieve.
OCHRIPEX (Wulp), 1888:19, pl. 1, fig. 14-14a (*Saundersia*). Type-locality: Costa Rica (sev. places). N. comb.

- OLIVACEUS Townsend, 1908:114. Type-locality: Mexico, Chihuahua, Rio Piedras Verdes.
- PULVERULENTUS (Schiner), 1868:335 (*Saundersia*). Type-locality: "South America". Ref. — Curran, 1947:88.
- palliceps* Bigot, 1888a:94 (*Cryptopalpus*). Type-locality: "Colombia".
- QUATUORNOTATUS (Townsend), 1935d:220 (*Rhachocarpodes*). Type-locality: Venezuela, Silla de Caracas.
- TRIFORMIS (Walker), 1852:267 (*Tachina*). Type-locality: "Colombia".
- TUCUMANUS Blanchard, 1941:375, figs. 17, 18 (as *Rachocarpus*). Type-locality: Argentina, Tucumán. Ref. — Curran, 1947:89.

Genus RHACHOSAUNDERSIA Townsend

- RHACHOSAUNDERSIA Townsend, 1931a:439. Type-species, *boliviana* Townsend (orig. des.).
- BOLIVIANA Townsend, 1931a:440. Type-locality: "Bolivia".

Genus SAUNDERSIOPS Townsend

- SAUNDERSIOPS Townsend, 1914:138. Type-species, *confluens* Townsend (orig. des.).
- Signoecarpus* Townsend, 1931:446. Type-species, *spinosus* Townsend (orig. des.).
- Ref. — Curran, 1947:94 (key).
- BROWNAE Curran, 1947:95, fig. 118. Type-locality: Ecuador, Bolivar, Hacienda Talahua.
- CAYENSIS Townsend, 1914:141 (as *cruciata* ssp.). Type-locality: Peru, Huancayo.
- COLOMBIENSIS Curran, 1947:97, fig. 114. Type-locality: Colombia, Vista Nieve.
- CONFLUENS Townsend, 1914:140. Type-locality: Peru, San Gabán Canyon.
- CRUCIATUS Townsend, 1914:140. Type-locality: Peru, Matucana.
- DISCALIS (Townsend), 1914:134 (*Epalpus*). Type-locality: Peru, Huascaray.
- HARPEZUS Reinhard, 1952a:8. Type-locality: Mexico, Rio Frio.
- METALLICUS Curran, 1947:98, fig. 115. Type-locality: Panama, Chiriqui, El Volcán.
- NIGRICORNIS (Townsend), 1914:135 (*Epalpus*). Type-locality: "Peru".
- OCULATUS Curran, 1947:94, figs. 101, 102, 105, 107. Type-locality: Ecuador, Bolivar, Hacienda Talahua.
- PACHECOI Curran, 1947:95. Type-locality: Guatemala, Guatemala City.
- SCHWARZI Curran, 1947:96, figs. 109, 112. Type-locality: Colombia, between Queremal and Buenaventura.
- SIESTUS Curran, 1947:97, fig. 113. Type-locality: Colombia, Popayán.
- SIMILLIMUS Townsend, 1914:141. Type-locality: Peru, Matucana.
- ssp. *punensis* Townsend, 1914:142. Type-locality: Peru, Paehacayo.
- SPINOSUS (Townsend), 1931a:446 (*Signoecarpus*). Type-locality: Peru, Oxapampa.
- TATEI Curran, 1947:96, fig. 108. Type-locality: Brazil, Roraima, Mt. Arabipu.
- MACULIVENTRIS (Brèthes), 1909:99 (*Trichophora*). Type-locality: Argentina, Mendoza, Valle del Tupungato. Unrecognized.

Genus SIGNOSOMA Townsend

- SIGNOSOMA Townsend, 1914:93. Type-species, *impressum* Townsend (orig. des.).
- IMPRESSUM Townsend, 1914:94. Type-locality: Peru, Uruhuasi.

Genus SIGNOSOMOPSIS Townsend

- SIGNOSOMOPSIS Townsend, 1914:95. Type-species, *argentea* Townsend (orig. des.).
 ARGENTEA Townsend, 1914:96. Type-locality: Peru, Uruhuasi.
 BRONIS Curran, 1929:505. Type-locality: Peru, Ollachea and Uruhuasi.
 TOWNSENDI Curran, 1929:504. Type-locality: Peru, Matucana.

Genus TRICHOEPALPUS Townsend

- TRICHOEPALPUS Townsend, 1914:156. Type-species, *emarginatus* Townsend (orig. des.).
 EMARGINATUS Townsend, 1914:156. Type-locality: Ecuador, Manchi.

Genus TRICHOPHORA Macquart

- TRICHOPHORA Macquart, 1847:78 (1847:62). Type-species, *nigra* Macquart (orig. des.).
Siphoniomyia Blgot, 1885a:xii. Type-species, *melas* Blgot (mon.).
Paragymnomma Brauer & Bergenstamm, 1891:384 (1891:80). Type-species, *hystrix* Brauer & Bergenstamm (Coquillett, 1910:584) = *nigra* Macquart.
Gabanimyia Townsend, 1914:44. Type-species, *polita* Townsend (orig. des.) = *nigra* Macquart.
Huascaraya Townsend, 1914:82. Type-species, *tegulata* Townsend (orig. des.).
Parepalpodes Townsend, 1915m:428. Type-species, *rimacensis* Townsend (orig. des.).
 ALBOCALYPTRATA Blgot, 1888a:82. Type-locality: Peru, Quito and Ecuador. Unrecognized.
 ANALIS Schlner, 1868:330. Type-locality: "Venezuela".
 CONVEXINERVIS Wulp, 1892:193. Type-locality: Mexico, Guerrero.
 MELAS (Blgot), 1885:xii (*Siphoniomyia*). Type-locality: "Mexico". Ref. — Townsend, 1931:166.
mitis Schlner in Brauer & Bergenstamm, 1891:406 (1891:102). *Nomen nudum*.
 NIGRA Macquart, 1843:79 (1847:63), pl. 3, fig. 7. Type-locality: "Brazil".
hystrix Brauer & Bergenstamm, 1891:384 (1891:80) (*Paragymnoma*). Type-locality: "Brazil".
polita Townsend, 1914:45 (*Gabanimyia*). Type-locality: Peru, San Gabán Canyon, Casahuari.
hystricosa Townsend, 1914:46 (*Gabanimyia*). Type-locality: Peru, Charape River.
 RIMACENSIS (Townsend), 1915m:429 (*Parepalpodes*). Type-locality: Peru, Matucana.
 RUFINA Wulp, 1888:36, pl. 2, fig. 16. Type-locality: Guatemala and Costa Rica.
 TEGULATA (Townsend), 1914:82 (*Huascaraya*). Type-locality: Peru, Huascaray.

Genus TRICHOSAUNDERSIA Townsend

- TRICHOSAUNDERSIA Townsend, 1914:137. Type-species, *lineata* Townsend (orig. des.). Ref. — Curran, 1947:92 (key).

- DORSOPUNCTATA (Macquart), 1843:207 (1843:47), pl. 4, fig. 5 (*Micropalpus*).
Type-locality: Colombia, Bogotá.
- LINEATA Townsend, 1914:138. Type-locality: Peru, San Gabán Canyon, Uruhuasi.
- NORA Curran, 1947:93, fig. 106. Type-locality: Guatemala, Guatemala City.
- RUBRIPILA (Rondani), 1850b:169 (*Epalpus*). Type-locality: "Venezuela".
- RUFOPILOSA (Wulp), 1888:22, pl. 1, fig. 18 (*Saundersia*). Type-locality: "Guatemala and Costa Rica".

Genus URUHUASIA Townsend

- URUHUASIA Townsend, 1914:126. Type-species, *cruciata* Townsend (orig. des.).
- CRUCIATA Townsend, 1914:127. Type-locality: Peru, San Gabán Canyon, Uruhuasi.
- DELTA Townsend, 1914:128. Type-locality: Peru, San Gabán Canyon, Uruhuasi.

Genus URUHUASIOPSIS Townsend

- URUHUASIOPSIS Townsend, 1915n:70. Type-species, *analis* Townsend (orig. des.).
- ANALIS Townsend, 1915n:71. Type-locality: Peru, Huariaca.

Genus VERTEPALPUS Curran

- VERTEPALPUS Curran, 1947:91. Type-species, *verdans* Curran (orig. des.).
- VERDANS Curran, 1947:92, figs. 96, 97, 100. Type-locality: Ecuador, Bolivar, Haelenda Talahua.

Genus VIBRISSEOEPALPUS Townsend

- VIBRISSEOEPALPUS Townsend, 1915n:73. Type-species, *flavipes* Townsend (orig. des.).
- FLAVIPES Townsend, 1915n:74. Type-locality: Peru, Pachacayo.

Genus XANTHOEPALPODES Townsend

- XANTHOEPALPODES Townsend, 1913a:437. Type-species, *bischofi* Townsend (orig. des.).
- BISCHOFI Townsend, 1913a:438. Type-locality: "Bolivia".

Genus XANTHOEPALPUS Townsend

- XANTHOEPALPUS Townsend, 1914:157. Type-species, *Saundersia bipartita* Wulp (orig. des.) = *bicolor* (Williston).
- BICOLOR (Williston), 1886:304 (*Saundersia*). Type-locality: U.S.A., New Mexico. Distr. — U.S.A., Mexico.
- bipartitus* Wulp, 1888:25, pl. 2-2a (*Saundersia*). Type-locality: Mexico, Durango, Ciudad.
- GABANUS Townsend, 1914:157. Type-locality: Peru, San Gabán Canyon, Casahuari.

- LAETUS (Wulp), 1892:191 (*Saundersia*). Type-locality: Mexico, Guerrero.
N. COMB.
SEMIFLAVUS (Bischof), 1904:91 (*Epalpus*). Type-locality: "Peru and
Bolivia".

Genus XANTHOZONA Townsend

- XANTHOZONA Townsend, 1908:116. Type-species, *Tachina melanopyga*
Wiedemann (orig. des.).
MELANOPYGA (Wiedemann), 1830:292 (*Tachina*). Type-locality: "Surinam".
Ref. — Travassos & Carrera, 1941 (biology).
SCUTELLARIS (Robineau-Desvoidy), 1830:36 (*Jurinia*). Type-locality: Brazil.
Paraná, Guaratuba. Ref. — Townsend, 1931:165.

Genus ZONOEPALPUS Townsend

- ZONOEPALPUS Townsend, 1927a:240 (errata). Type-species, *brasilensis*
Townsend (orig. des.) = *testaceus* (Robineau-Desvoidy).
ARGENTINENSIS Blanchard, 1941:378, fig. 19. Type-locality: Argentina,
Misiones.
TESTACEUS (Robineau-Desvoidy), 1830:38 (*Jurinia*). Type-locality: Brazil,
Rio Grande do Sul. Ref. — Townsend, 1931b:165.
brasilensis Townsend, 1927a:369. Type-locality: Brazil, São Paulo,
Itaquaquecetuba.

Tribe CUPHOCERINI

The hosts of Cuphocerini are lepidopterous larvae, chiefly cutworms
and armyworms.

Genus ABEPALPUS Townsend

- ABEPALPUS Townsend, 1931a:449. Type-species, *archytoides* Townsend (orig.
des.).
ARCHYTODES Townsend, 1931a:449. Type-locality: Brazil, São Paulo.

Genus ACROCERONIA Cortés

- ACROCERONIA Cortés, 1951b:251. Type-species, *elquiensis* Cortés (orig. des.).
ELQUIENSIS Cortés, 1951b:252. Type-locality: Chile, Coquimbo, Elqui,
Gualliguaica.

Genus ACUPHOCEROPSIS Blanchard

- ACUPHOCEROPSIS Blanchard, 1943b:123. Type-species, *nigricornis* Blanchard
(orig. des.).
NIGRICORNIS Blanchard, 1943b:123, fig. 20. Type-locality: Argentina, Salta.

Genus AGICUPHOCERA Townsend

- AGICUPHOCERA Townsend, 1915m:430. Type-species, *nigra* Townsend (orig.
des.).
NIGRA Townsend, 1915m:430. Type-locality: Peru, Chosica.

Genus ANDINOMYIA Townsend

- ANDINOMYIA Townsend, 1912a:329. Type-species, *cruciata* Townsend (orig. des.).
- Chaetoandinomyia* Vimmer & Soukup, 1940:218. Type-species, *townsendi* Vimmer & Soukup (mon.) (as *towscndi*, misprint). N. syn.
- COMPLANATA (Vimmer & Soukup), 1940:218 (*Chaetoandinomyia*; as *towscndi* var.). Type-locality: Peru, Puno. N. comb.
- CRUCIATA Townsend, 1912a:329. Type-locality: Peru, Tirapata.
- NIGRA Vimmer & Soukup, 1940:216. Type-locality: Peru, Puno, Ollachea and Cuzco. Unrecognized.
- RUFOMACULATA Vimmer & Soukup, 1940:216. Type-locality: Peru, Ollachea, Puno and Cuzco. Unrecognized.
- TOWNSENDI (Vimmer & Soukup), 1940:218 (*Chaetoandinomyia*; as *towscndi*, misprint). Type-locality: Peru, Ollachea, Puno and Cuzco. N. comb.

Genus BESKIOCEPHALA Townsend

- BESKIOCEPHALA Townsend, 1916h:17. Type-species, *flava* Townsend (orig. des.).
- FLAVA Townsend, 1916h:17. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães). Distr. — Brazil (Mato Grosso, Goiás, Guanabara, São Paulo). Ref. — Guimarães, 1963:62.

Genus CATAJURINIA Townsend

- CATAJURINIA Townsend, 1927a:245. Type-species, *angusta* Townsend (orig. des.).
- ANGUSTA Townsend, 1927a:294 (as *angusta*, misprint). Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus COMOPS Aldrich

- COMOPS Aldrich, 1934:40. Type-locality, *ruficornis* Aldrich (orig. des.).
- RUFICORNIS Aldrich, 1934:40. Type-locality: Argentina, Rio Negro, Lake Nahuel Huapi.
- penai* Cortés, 1967:16 (*Enchomyia*). Type-locality: Chile, Hacienda Llapel, Coquimbo, Illapel, 2.500-2.800m. N. syn. (R. Cortés, in litt.).

Genus COPECRYPTA Townsend

- COPECRYPTA Townsend, 1908:109. Type-species, *Schineria ruficauda* Wulp (mon.).
- NITENS (Wiedemann), 1830:294 (*Tachina*). Type-locality: "Brazil". Ref. — Guimarães, 1963:44.
- saginata* Walker, 1861:298 (*Eurigaster*). Type-locality: "Mexico". N. syn.
- devia* Townsend, 1917b:227 (as *ruficauda* ssp.). Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).
- andina* Townsend, 1919a:591. Type-locality: Peru, Verrugas Canyon.
- RUFICAUDA (Wulp), 1867:146, pl. 4, fig. 17 (*Schineria*). Type-locality: "U.S.A." Distr. — U.S.A., Mexico. Ref. — Guimarães, 1963:47.
- nitidifrons* Wulp, 1888:37 (*Trichophora*). Type-locality: "Mexico".

? *trisetosa* Wulp, 1888:36, pl. 2, fig. 16-16a (*Trichophora*). Type-locality: Costa Rica, Caché.

nitens of authors, not Wiedemann.

Genus CYANOGYMNOMMA Townsend

CYANOGYMNOMMA Townsend, 1927a:250 (errata). Type-species, *coerulea* Townsend (orig. des.).

Cianogymnomma, idem, pg. 50 (misprint).

COERULEA Townsend, 1927a:301. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus CYANOPSIS Townsend

CYANOPSIS Townsend, 1917b:228. Type-species, *costalis* Townsend (orig. des.). Ref. — Guimarães, 1963:50; 1964:178.

COSTALISMAI Guimarães, 1964:178, figs. 14, 23. Type-locality: Brazil, Goiás, Campinas.

COSTALIS Townsend, 1917b:228. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).

Genus DEOPALPUS Townsend

DEOPALPUS Townsend, 1908:110. Type-species, *hirsutus* Townsend (orig. des.).

Spanipalpus Townsend, 1908:110. Type-species, *Trichophora miscelli* Coquillett (orig. des.).

Procyanopsis Townsend, 1934:209. Type-species, *pictipennis* Townsend (orig. des.).

Cuphocera, authors, not Macquart.

Cyphocera, error or emend.

Ref. — Guimarães, 1963:64.

ALBIMACULA (Wiedemann), 1830:328 (*Tachina*). Type-locality: "South America". Unrecognized.

ALDRICHI (Townsend), 1931b:169 (*Spanipalpus*). Type-locality: "Brazil". *macrocera*, authors, not Wiedemann.

AUSTRALIS (Townsend), 1928c:164 (*Spanipalpus*). Type-locality: Chile, Punta Arenas.

BUCCATUS (Reinhard), 1934c:59 (*Cuphocera*). Type-locality: Cuba, Havana.

CONTIGUUS (Reinhard), 1934c:61 (*Cuphocera*). Type-locality: U.S.A., California. Distr. — U.S.A., Mexico.

DECURATUS (Rondani), 1850a:361 (*Cuphocera*). Type-locality: "America". Unrecognized.

FUCATUS (Wulp), 1892:193 (*Trichophora*). Type-locality: "Mexico".

HIRBUTUS Townsend, 1908:110. Type-locality: Mexico, Chihuahua, Sierra Madre, Meadow Valley. Distr. — Canada, U.S.A., Mexico.

MACROCERUS (Wiedemann), 1830:290 (*Tachina*). Type-locality: "Brazil". Ref. — Aldrich, 1929:29. N. comb.

OCIRICORNIS (Bigot), 1888a:95 (*Epalpus*). Type-locality: "Chile". N. comb. (R. Cortés, in litt.).

PICTIPENNIS (Townsend), 1934:210 (*Procyanopsis*). Type-locality: Brazil, Pará, Urucurituba. Ref. — Guimarães, 1963:63, figs. 106, 113.

PRETIOSUS (Curran), 1934c:521 (*Cuphocera*). Type-locality: Guyana, Bartica, Kartabo.

- PRUINOSUS (Rondani), 1863:16 (*Cyphocera*). Type-locality: "Chile". Unrecognized.
- FULCHRICEPS (Aldrich), 1934:128 (*Cuphocera*). Type-locality: Argentina, Río Negro, Bariloche. Unrecognized.
- KATZBURGI (Jaenicke), 1867:386 (1868:78) (*Demoticus*). Type-locality: "Chile". Unrecognized.
- REINHARDI Guimarães, 1963:70, figs. 114, 121. Type-locality: Brazil, São Paulo, Pôrto Cabral, R. Paraná.
- TRINITATIS Thompson, 1963a:396, figs. 77, 82. Type-locality: "Trinidad".

Genus DOLICHOSTOMA Townsend

- DOLICHOSTOMA Townsend, 1912a:325. Type-species, *alpina* Townsend (orig. des.).
- Erigonopsis* Townsend, 1912a:326. Type-species, *arequipae* Townsend (orig. des.).
- Epidolichostoma* Townsend, 1927a:238. Type-species, *andinum* Townsend (orig. des.).
- Ref. — Aldrich, 1934:37.
- ALPINUM Townsend, 1912a:325. Type-locality: Peru, Tirapata.
- ANDINUM (Townsend), 1927a:304 (*Epidolichostoma*). Type-locality: Peru, Oroya.
- AREQUIPAE (Townsend), 1912a:326 (*Erigonopsis*). Type-locality: Peru, Arequipa.
- NIGRICAUDUM Blanchard, 1963:178, fig. 16. Type-locality: Argentina, Tucumán.
- PUNTARENENSIS (Townsend), 1928c:163 (*Erigonopsis*). Type-locality: Chile, Punta Arenas. Distr. — South Chile, South Argentina.

Genus ERICUPHOCERA Townsend

- ERICUPHOCERA Townsend, 1927a:240. Type-species, *andina* Townsend (orig. des.).
- ANDINA Townsend, 1927a:304. Type-locality: Peru, Matucana.

Genus EUHELIOPROSOPA Reinhard

- EUHELIOPROSOPA Reinhard, 1964b:123. Type-species, *pacilis* Reinhard (orig. des.).
- PACILIS Reinhard, 1964b:124. Type-locality: Chile, 50 km south of La Serena.

Genus EUTRICHOPHORA Townsend

- EUTRICHOPHORA Townsend, 1915h:183. Type-species, *puncensis* Townsend (orig. des.).
- PUNENSIS Townsend, 1915h:184. Type-locality: Peru, Pachacayo.

Genus FORMICOMYIA Townsend

- FORMICOMYIA Townsend, 1916h:17. Type-species, *ovata* Townsend (orig. des.).
- OVATA Townsend, 1916h:18. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).

Genus GANOPROCTUS Aldrich

- GANOPROCTUS Aldrich, 1934:36. Type-species, *argentifer* Aldrich (orig. des.).
ARGENTIFER Aldrich, 1934:36. Type-locality: Chile, Lake Correntoso.
LONGICORNIS Aldrich, 1934:37. Type-locality: Chile, Lonquimay, Alto Biobío.

Genus GYMNO-MMOPSIS Townsend

- GYMNO-MMOPSIS Townsend, 1927a:255. Type-species, *gagatea* Townsend (orig. des.).
Metagymnopsis Blanchard, 1943b:145. Type-species, *missionensis* Blanchard (orig. des.) N. SYN.
Gymnocuphocera Blanchard, 1943b:143. Type-species, *cordubensis* Blanchard (orig. des.) N. SYN.
CORDUBENSIS (Blanchard), 1943b:143, fig. 25 (*Gymnocuphocera*). Type-locality: Argentina, Córdoba, Los Cocos. N. COMB.
GAGATEA Townsend, 1927a:312. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
HAYWARDI (Blanchard), 1943b:148 (*Metagymnopsis*). Type-locality: Argentina, Misiones.
MISSIONENSIS (Blanchard), 1943b:146, fig. 26 (*Metagymnopsis*). Type-locality: Argentina, Misiones. N. COMB.

Genus HELIOPROSOPA Townsend

- HELIOPROSOPA Townsend, 1927a:239. Type-species, *facialis* Townsend (orig. des.).
Ref. — Reinhard, 1964 (key).
AURIFODINA Reinhard, 1964b:119. Type-locality: Colombia, Valle, 6 miles west of Cali.
ELECTILIS Reinhard, 1964b:120. Type-locality: Mexico, Colima.
FACIALIS Townsend, 1927a:314. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
FINITA Reinhard, 1964b:121. Type-locality: Chile, Coquimbo, 5 miles northwest of Laguna Dam. Distr. — Mexico, Colombia (Cundinamarca).
LICIATA Reinhard, 1964b:123. Type-locality: Mexico, Hidalgo, 7 miles northeast of Jacala. Distr. — Mexico, Colombia.
MACROCERA Reinhard, 1964b:120. Type-locality: "Colombia".
VELADA Reinhard, 1964b:121. Type-locality: Colombia, Cundinamarca, Fusagasugá.

Genus ITACUPHOCERA Townsend

- ITACUPHOCERA Townsend, 1927a:238. Type-species, *ocellaris* Townsend (orig. des.).
Ref. — Guimarães, 1964.
BORGMEIERI Guimarães 1964:188, figs. 47, 52. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
CARRERAÍ Guimarães, 1964:185, figs. 40, 46. Type-locality: Brazil, São Paulo, Campos do Jordão.
OCELLARIS Townsend, 1927a:320. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus MICROGYMNOMMA Townsend

- MICROGYMNOMMA Townsend, 1916h:18. Type-species, *orbitalis* Townsend (orig. des.).
- ORBITALIS Townsend, 1916h:18. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).
- PAULENSIS Townsend, 1929:380. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus NEOCUPHOCERA Townsend

- NEOCUPHOCERA Townsend, 1927a:239. Type-species, *nepos* Townsend (orig. des.).
- Myocuphocera* Townsend, 1931b:168. Type-species, *Tachina macrocera* Wiedemann (mon., ? misident.).
Ref. — Guimarães, 1963a:57.
- NEPOS Townsend, 1927a:333. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Tropical South America.
aureifacies Guimarães, 1963a:60, figs. 69, 70. Type-locality: Brazil (São Paulo, Mato Grosso, Golás). N. SYN.
? *macrocera* Wiedemann of Townsend.
- ORBITALIS (Aldrich), 1929a:26 (*Copecrypta*). Type-locality: Ecuador, Posorja. N. COMB.
nepos of Guimarães, nec Townsend.

Genus NEOSSARROMYIA Townsend

- NEOSSARROMYIA Townsend, 1927a:237. Type-species, *neotropica* Townsend (orig. des.).
- Antillicolla* Curran, 1927e:1. Type-species, *aureiceps* Curran (orig. des). N. SYN.
- AURICEPS (Curran), 1927e:1, fig. 1 (*Antillicolla*). Type-locality: Puerto Rico, Adjuntas. N. COMB.
- NEOTROPICA Townsend, 1927a:335. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- TRINITAS Thompson, 1963a:422. Type-locality: "Trinidad".

Genus PALPOLINNAEMYIA Townsend

- PALPOLINNAEMYIA Townsend, 1927a:238. Type-species, *perorbitalis* Townsend (orig. des.).
- PERORBITALIS Townsend, 1927a:344. Type-locality: Peru, Matucana.

Genus PERIOPTICHOCHAETA Townsend

- PERIOPTICHOCHAETA Townsend, 1927a:245. Type-species, *pendula* Townsend (orig. des.).
- PENDULA Townsend, 1927a:345. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus PHOSOCEPHALA Townsend

- PHOSOCEPHALA Townsend, 1908:69. Type-species, *metallica* Townsend (orig. des.).

Phosococephala Townsend, 1936b:285, emend.

METALLICA Townsend, 1908:72. Type-locality: Costa Rica, Tucurrique.

Genus PROSPANIFALPUS Townsend

PROSPANIFALPUS Townsend, 1931a:448. Type-species, *peruvianus* Townsend (orig. des.).

PERUVIANUS Townsend, 1931a:449. Type-locality: Peru, Choslea Canyon.

Genus PUNAMYIA Townsend

PUNAMYIA Townsend, 1915m:428. Type-species, *translionalis* Townsend (orig. des.).

TRANSITIONALIS Townsend, 1915m:428. Type-locality: Peru, Oroya.

Genus PYRRHOTACHINA Townsend

PYRRHOTACHINA Townsend, 1931a:447. Type-species, *proboscidea* Townsend (orig. des.).

PROBOSCIDEA Townsend, 1931a:448. Type-locality: Argentina, Mendoza.

Genus SOROCHEMYIA Townsend

SOROCHEMYIA Townsend, 1915f:45. Type-species, *oroya* Townsend (orig. des.).

OROYA Townsend, 1915f:45. Type-locality: Peru, Oroya.

Genus TALAROCERA Williston

TALAROCERA Williston, 1888:152. Type-species, *smithii* Williston (orig. des.)
= *nigripennis* (Wiedemann).

Talacrocer Townsend, 1936b:285, error.

Ref. — Bezzi, 1923:652.

NIGRIPENNIS (Wiedemann), 1830:286 (*Tachina*). Type-locality: Brazil. Distr.
— Brazil (Mato Grosso, Minas Gerais), Paraguay.

smithii Williston, 1888:152. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).

Genus TRICHOPIHOROPSIS Townsend

TRICHOPIHOROPSIS Townsend, 1914:42. Type-species, *puna* Townsend (orig. des.).

NITENS Townsend, 1914:44. Type-locality: Peru, Oroya.

PUNA Townsend, 1914:43. Type-locality: Peru, Oroya.

Genus VIBRISSOMYIA Townsend

VIBRISSOMYIA Townsend, 1912a:327. Type-species, *lineata* Townsend (orig. des.).

Ref. — Cortés, 1945c:29.

BICOLOR Townsend, 1912a:328. Type-locality: Peru, Tlapata.

ERYTHROSTOMA (Bigot), 1888a:95 (*Epalpus*). Type-locality: "Chile". Unrecognized.

- LINEOLATA (Bigot), 1888a:94 (*Epalpus*). Type-locality: "Chile". Distr. — Chile, Peru (Tirapata) and Argentina (Santa Cruz).
lineata Townsend, 1912a:328. Type-locality: Peru, Tirapata.
albilineata Blanchard, 1943b:152, fig. 27. Type-locality: Argentina, Santa Cruz, Valle Tunel.
 NOTATA Cortés, 1967b:14. Type-locality: Chile, Santiago, Cajón del Rio Maipo, El Yeso.
 OROYENSIS Townsend, 1914:175 (as *lineata* ssp.). Type-locality: Peru, Oroya.
 PULLATA Cortés, 1951:260. Type-locality: Chile, Santiago, El Canelo.

Tribe LINNAEMYINI

Members of the tribe Linnaemyini parasitize various species of lepidopterous larvae.

Genus BONNETIA Robineau-Desvoidy

- BONNETIA Robineau-Desvoidy, 1830:55. Type-species, *oecanthis* Robineau-Desvoidy (Townsend, 1916:6) = *comta* (Fallén).
 COMTA (Fallén), 1810:277 (*Tachina*). Type-locality: "Sweden". Distr. — Canada, U.S.A., Mexico. Palearctic.

Genus CHRYSOTACHINA Brauer & Bergenstamm

- CHRYSOTACHINA Brauer & Bergenstamm, 1889:161 (1889:93). Type-species, *Tachina reinicardtii* Wiedemann (mon., misident.). = *braueri* Townsend.
Eugymnochaeta Townsend, 1912a:314. Type-species, *Gymnochaeta alcedo* Loew (orig. des.).
Paragymnochaeta Townsend, 1915b:21. Type-species, *Gymnochaeta equatorialis* Townsend (orig. des.). Ref. — Curran, 1939b:1 (key), Aldrich, 1926d (key).
Chrysoerigone Townsend, 1927a:244. Type-species, *ornata* Townsend (orig. des.). N. SYN.
 ALCEDO (Loew), 1869:36 (Cent. 8, n° 61) (*Gymnochaeta*). Type-locality: U.S.A., Massachusetts. Distr. — U.S.A., Mexico.
 BRAUERI Townsend, 1931a:452. Type-locality: "Brazil".
reinicardtii authors, not Wiedemann.
 EQUATORIALIS (Townsend), 1912a:314 (*Eugymnochaeta*). Type-locality: Peru, Piura. Distr. — Peru, Brazil.
 ORNATA (Townsend), 1927a:298 (*Chrysoerigone*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. N. COMB.
 PANAMENSIS Curran, 1939b:2. Type-locality: Panama, Canal Zone, Barro Colorado I.
 PERUVIANA Curran, 1939b:590. Type-locality: Peru, Chosica.
 PURPUREA Curran, 1939b:2. Type-locality: "Venezuela". Distr. — Guatemala, Panama (cf. Austen, 1907:335).
 RUFICORNIS (Walker), 1852:304 (*Tachina*). Type-locality: "South America". Unrecognized.
 SUBCYANEA (Wulp), 1890:58 (*Mystacelia*). Type-locality: Mexico, Guerrero, Sierra, de las Aguas Escondidas. N. COMB.
 SUBVIRIDIS (Wulp), 1892:194 (*Gymnochaeta*). Type-locality: "Mexico". Unrecognized.



- TATEI Curran, 1939b:1. Type-locality: Brazil, Roraima Mt., Roraima, Philipp camp.
 TOWNSENDI Curran, 1939b:2. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).
 WILLISTONI Curran, 1939b:2. Type-locality: Colombia, Cauca Valley, Quebrada Seca. Distr. — Colombia, West Indies, Brazil (Mato Grosso).

Genus ELEUTHROMYIA Reinhard

- ELEUTHROMYIA Reinhard, 1964c:14. Type-species, *inuitata* Reinhard (orig. des.).
 INUITATA Reinhard, 1964c:15. Type-locality: Mexico, Veraacruz, Orizaba.

Genus GYMNOCHIAETOPSIS Townsend

- GYMNOCHIAETOPSIS Townsend, 1914:15. Type-species, *analís* Townsend (orig. des.).
 ANALIS Townsend, 1914:29. Type-locality: Peru, Rio Chira Valley, Cañada de Samán.
 FULVICAUDA (Walton), 1914:93 (*Linnaemyia*). Type-locality: "Puerto Rico".

Genus MERICINA Curran

- MERICINA Curran, 1927c:6. Type-species, *ruficauda* Curran (orig. des.).
 RUFICAUDA Curran, 1927c:6. Type-locality: Puerto Rico, Arecibo.

Genus NEOERIGONE Townsend

- NEOERIGONE Townsend, 1919a:590. Type-species, *cinerea* Townsend (orig. des.).
 CINEREA Townsend, 1919a:591. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).

Genus NEOTRAFOIOPSIS Townsend

- NEOTRAFOIOPSIS Townsend, 1931a:451. Type-species, *andina* Townsend (orig. des.).
 ANDINA Townsend, 1931a:452. Type-locality: Peru, Oxapampa and Uruhuasi.

Genus PARABRACHYCOMA Blanchard

- PARABRACHYCOMA Blanchard, 1940:229. Type-species, *ruficauda* Blanchard (orig. des.).
 RUFICAUDA Blanchard, 1940:229, fig. 2a-f. Type-locality: Argentina, Salta. Distr. — northern Argentina.

Genus PYRRHOERNESTIA Townsend

- PYRRHOERNESTIA Townsend, 1931a:450. Type-species, *petiolata* Townsend (orig. des.).
 PETIOLATA Townsend, 1931a:450. Type-locality: Bolivia, Cuesta de Cillunticara.

Genus TRAFOLA Brauer & Bergenstamm

- TRAFOLA Brauer & Bergenstamm, 1893:142 (1893:54). Type-species, *monticola* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.).
- NEOTRAFOLA Townsend, 1912a:313. Type-species, *incarum* Townsend (orig. des.).
- CHARAPEMYIA Townsend, 1919a:589. Type-species, *callida* Townsend (orig. des.). = *incarum* (Townsend).
- INCARUM (Townsend), 1912a:314 (*Neotrafola*). Type-locality: Peru, Cuzco. Ref. — Aldrich, 1928:144 (notes).
- callida* Townsend, 1919a:590 (*Charapemyia*). Type-locality: Peru, Jaén, Charape River.
- RUFIPALPIS (Bigot), 1889:256 (*Exorista*). Type-locality: "Mexico". Distr. — U.S.A., El Salvador.
- hispidula* Wulp, 1890:65 (*Exorista*). Type-locality: Mexico, Orizaba and Guerrero.
- SETULOSA (Wulp), 1890:58 (*Mystacella*). Type-locality: Mexico, Guerrero. N. comb.
- TRINITATIS (Thompson), 1963a:463, fig. 130 (*Neotrafola*). Type-locality: "Trinidad".

UNPLACED SPECIES OF LINNAEMYIINI

- ANGUSTIFRONS Wulp, 1892:193 (*Micropalpus*). Type-locality: Mexico, Tabasco.
- COMPACTA Curran, 1926a:113 (*Linnaemyia*). Type-locality: "Jamaica".
- NIGRIFRONS Bigot, 1889:263 (*Micropalpus*). Type-locality: "Mexico".

Tribe ERNESTIINI

The hosts of Ernestiini are lepidopterous larvae, principally those of the family Noctuidae.

Genus ECUADORANA Townsend

- ECUADORANA Townsend, 1912a:324. Type-species, *bicolor* Townsend (orig. des.).
- BICOLOR Townsend, 1912a:324. Type-locality: Ecuador, Manchi.

Genus EUCHEIROPHAGA James

- EUCHEIROPHAGA James, 1945:328. Type-species, *lugubris* James (orig. des.).
- LUGUBRIS James, 1945:329, fig. 1. Type-locality: Mexico, Morelos.

Genus ERNESTIOPSIS Townsend

- ERNESTIOPSIS Townsend, 1931a:454. Type-species, *crigonopsidis* Townsend (orig. des.).
- ERIGONOPSIDIS Townsend, 1931a:454. Type-locality: Chile, Valparaiso, Perales.

Genus EXOERNESTIA Townsend

- EXOERNESTIA Townsend, 1927a:244. Type-species, *uruhuasi* Townsend (orig. des.).

LLUYI Townsend, 1929:380. Type-locality: Peru, Lluy.
URUHUASI Townsend, 1927a:310. Type-locality: Peru, Uruhuasi.

Genus FASSLOMYIA Townsend

FASSLOMYIA Townsend, 1931a:452. Type-species, *fantastica* Townsend (orig. des.).
FANTASTICA Townsend, 1931a:453. Type-locality: Bolivia, Cuesta de Cillunticara.

Genus MESEMBRERIGONE Townsend

MESEMBRERIGONE Townsend, 1931a:457. Type-species, *alpina* Townsend (orig. des.).
ALPINA Townsend, 1931a:458. Type-locality: Bolivia, Cuesta de Cillunticara.

Genus OPTICOPTERYX Townsend

OPTICOPTERYX Townsend, 1931a:455. Type-species, *alpina* Townsend (orig. des.).
ALPINA Townsend, 1931a:456. Type-locality: Bolivia, Cuesta de Cillunticara.

Genus RHAMPHOPTERYX Townsend

RHAMPHOPTERYX Townsend, 1931a:456. Type-species, *retrorsa* Townsend (orig. des.).
PETRORSA Townsend, 1931a:457. Type-locality: Peru, near Huancabamba.

Genus TELODYTES Aldrich

TELODYTES Aldrich, 1934:50. Type-species, *analis* Aldrich (orig. des.).
ANALIS Aldrich, 1934:50. Type-locality: Argentina, Rio Negro, Lake Gutlérrez.

Genus XANTHOPHYTO Townsend

XANTHOPHYTO Townsend, 1916e:627. Type-species, *Nemoraca labis* Coquillett (orig. des.).
ERYTHROPYGA (Wulp), 1882:83 (*Nemoraca*). Type-locality: "Chile". Distr. — Chile (Santiago, Angol).
VERSICOLOR (Wulp), 1890:130 (*Macquartia*). Type-locality: Mexico, Guerrero. N. COMB.

Genus XYLOCAMPTOMIMA Townsend

XYLOCAMPTOMIMA Townsend, 1927a:243. Type-species, *oculata* Townsend (orig. des.) = *usta* (Wiedemann).
USTA (Wiedemann), 1830:313 (*Tachina*). Type-locality: "Brazil".
oculata Townsend, 1927a:366. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Tribe APHRIINI

The hosts of Aphriini are lepidopterous larvae including some wood and stalk-borers.

Genus MICRONYCHOPS Townsend

MICRONYCHOPS Townsend, 1915m:423. Type-species, *aureescens* Townsend (orig. des.).

AURESCENS Townsend, 1915m:424. Type-locality: Peru, Huancayo.

Genus NEAPHRIA Townsend

NEAPHRIA Townsend, 1914:12. Type-species, *dezina* Townsend (orig. des.).

DEXINA Townsend, 1914:13. Type-locality: Peru, Chosica.

Genus NEOSOLIERIA Townsend

NEOSOLIERIA Townsend, 1927a:211. Type-species, *nasuta* Townsend (orig. des.).

Parcipromus Reinhard, 1958b:227. Type-species, *silus* Reinhard (orig. des.).

NASUTA Townsend, 1927a:336. Type-locality: Peru, Lima, San Cristóbal Mt.

SILA (Reinhard), 1958b:228 (*Parcipromus*). Type-locality: Mexico, Baja California, Agua Verde Bay. Distr. — U.S.A., Mexico.

Tribe LYPHINI

The species of the tribe Lyphini parasitize lepidopterous larvae. *Lypha scitificies* (West.) is an important parasite of the spruce budworm *Choristoneura fumiferana* (Clem.), in the Nearctic Region.

Genus ARCTOSOMA Aldrich

ARCTOSOMA Aldrich, 1934:18. Type-species, *nigripalpis* Aldrich (orig. des.).

NIGRIPALPIS Aldrich, 1934:19. Type-locality: Argentina, Rio Negro, Lake Nahuel Huapi.

Genus EMBIOMYIA Aldrich

EMBIOMYIA Aldrich, 1934:29. Type-species, *australis* Aldrich (orig. des.).

AUSTRALIS Aldrich, 1934:30. Type-locality: Argentina, Rio Negro, Lake Nahuel Huapi. Distr. — Argentina, Chile.

Genus LYPHA Robineau-Desvoidy

LYPHA Robineau-Desvoidy, 1830:141. Type-species, *Tachina dubia* Fallén (Robineau-Desvoidy, 1863:196).

Ref. — Aldrich, 1934:51 (key to Chilean spp.).

ANGOLENSIS Aldrich, 1934:58. Type-locality: Chile, Angol.

CHAETOSA Aldrich, 1934:59. Type-locality: Argentina, Rio Negro, Lake Nahuel Huapi.

CORAX Aldrich, 1934:61. Type-locality: Argentina, Rio Negro, Bariloche.

- EDWARDSI Aldrich, 1934:53, fig. 8. Type-locality: Argentina, Río Negro, Lake Correntoso.
- LITURATA Aldrich, 1934:59. Type-locality: Argentina, Río Negro, Bariloche.
- LONGICORNIS Aldrich, 1934:62. Type-locality: Chile, Llanquihue, Casa Pangue.
- NOCTIFER Aldrich, 1943:62. Type-locality: Argentina, Río Negro, Lake Nahuel Huapi.
- ORBITALIS Aldrich, 1934:60. Type-locality: Chile, Llanquihue, Ensenada.
- ORNATA Aldrich, 1934:56. Type-locality: Argentina, Tierra del Fuego, Estancia Viamonte.
- TRIANGULIFERA (Jacobs), 1900:107 (*Hystriola*). Type-locality: Chile, Canal del Beagle, Harberton Harbour. Ref. — Cortés, 1968c.
- TRUNCATA Aldrich, 1934:55, fig. 9. Type-locality: Argentina, Tierra del Fuego, Lake Yuvín.

Genus PROLESKIA Townsend

- PROLESKIA Townsend, 1927a:211. Type-species, *hirta* Townsend (orig. des.).
- HIRTA Townsend, 1927a:350. Type-locality: Brazil, São Paulo, São Paulo Cantareira.

Tribe NEOBRACHELINI

Members of the tribe Neobracheilini have been recorded as parasites of Hemiptera, Pentatomidae.

Genus NEOBRACHELIA Townsend

- NEOBRACHELIA Townsend, 1931a:458. Type-species, *charapemyioides* Townsend (orig. des.).
- Xenopyxis Townsend, 1940b:889. Type-species, *mirabilis* Townsend (orig. des.). N. SYN.
- Xenophyxis, error.
- CHARAPEMYIOIDES Townsend, 1931a:459. Type-locality: Peru, Oxapampa.
- EDESSAE (Townsend), 1942d:438 (as *Xenophyxis*). Type-locality: Uruguay, Montevideo. N. COMB. Ref. — Parker, 1953:52 (immature stages).
- GRANDIS (Townsend), 1940b:891 (*Xenopyxis*). Type-locality: Brazil, Mato Grosso, Maracaju. N. COMB.
- MIRABILIS (Townsend), 1940b:890 (*Xenopyxis*). Type-locality: Brazil, Mato Grosso, Maracaju. N. COMB.

Tribe GERMARIINI

The species of Germariini parasitize lepidopterous larvae; a few species have been bred from crickets, beetles, centipedes, and earwigs (Dermaptera).

Genus ACTINOPLAGIA Blanchard

- ACTINOPLAGIA Blanchard, 1940:234. Type-species, *koehleri* Blanchard (orig. des.).
- KOEHLERI Blanchard, 1940:234, fig. 3. Type-locality: Argentina, Arrecifes. Ref. — Parker, 1953:50 (immature stages).

Genus AMBLYCHAETA Aldrich

AMBLYCHAETA Aldrich, 1934:67. Type-species, *picticornis* Aldrich (orig. des.).
PICTICORNIS Aldrich, 1934:68, fig. 10. Type-locality: Argentina, Rio Negro,
Lake Correntoso.

Genus CHAETODEMOTICUS Brauer & Bergenstamm

CHAETODEMOTICUS Brauer & Bergenstamm, 1891:385 (1891:81). Type-species,
Demoticus chilensis Schiner (orig. des.).
CHILENSIS (Schiner), 1868:324 (*Demoticus*). Type-locality: "Chile". Distr.
— Chile, Santiago. Ref. — Cortés, 1945e:24.

Genus DELOBLEPHARIS Aldrich

DELOBLEPHARIS Aldrich, 1934:74. Type-species, *nigra* Aldrich (orig. des.).
NIGRA Aldrich, 1934:74. Type-locality: Chile, Llanquihue, Casa Pangué.

Genus EXORISTOPSIS Townsend

EXORISTOPSIS Townsend, 1915m:426. Type-species, *setifera* Townsend (orig.
des.).
SETIFERA Townsend, 1915m:427. Type-locality: Peru, Chosica.
VERTICALIS (Reinhard), 1935a:161 (*Exoristoides*). Type-locality: "Mexico".

Genus JALAPAMYIA Reinhard

JALAPAMYIA Reinhard, 1964e:12. Type-species, *callitris* Reinhard (orig. des.).
CALLITRIS Reinhard, 1964e:13. Type-locality: Mexico, Veraacruz, Jalapa.

Genus HELICLYDELLOPS Townsend

Heliolydella Townsend, 1934a:210 (preocc. Townsend, 1927). Type-species,
homconychioides Townsend (orig. des.).
HEMOLYDELLOPS Townsend, 1934a:405 (n. name for *Heliolydella* Townsend).
Type-species, *homconychioides* Townsend (aut.).
HOMEONYCHIOIDES (Townsend), 1934a:210 (*Heliolydella*). Type-locality:
Brazil, São Paulo, Itaquaquecetuba.

Genus HELIOPLAGIA Townsend

HELIOPLAGIA Townsend, 1934a:211. Type-species, *amazonica* Townsend (orig.
des.).
AMAZONICA Townsend, 1934a:211. Type-locality: Brazil, Amazonas, Rio
Tapajós, Urucurituba.
URIEHI (Aldrich), 1932:25 (*Exoristoides*). Type-locality: "Trinidad".

Genus LYDINA Robineau-Desvoidy

LYDINA Robineau-Desvoidy, 1830:124. Type-species, *nitida* Robineau-Des-
voidy (Robineau-Desvoidy, 1863:111) = *acnea* (Meigen).
IMMISTA Reinhard, 1955c:128. Type-locality: Mexico, Popocatepetl.

Genus MELLACHNUS Aldrich

- MELLACHNUS Aldrich, 1934:41. Type-species, *velutinus* Aldrich (orig. des.).
VELUTINUS Aldrich, 1934:41. Type-locality: Argentina, Rio Negro, Lake Nahuel Huapi.

Genus OLLACHERYPHE Townsend

- OLLACHERYPHE Townsend, 1927a:256. Type-species, *facialis* Townsend (orig. des.).
Aeglops Aldrich, 1934:47. Type-species, *acnea* Aldrich (orig. des.) Ref. — Cortés, 1945c:30.
AENEAE (Aldrich), 1934:47 (*Aeglops*). Type-locality: Argentina, Lake Correntoso.
FACIALIS Townsend, 1927a:339. Type-locality: Peru, Ollachea. Distr. — Peru, south Chile, south Argentina. Ref. — Cortés, 1945c:30.

Genus PARKERELLA Townsend

- PARKERELLA Townsend, 1942c:149. Type-species, *parva* Townsend (orig. des.).
PARVA Townsend, 1942c:149. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus PROLYPHA Townsend

- PROLYPHA Townsend, 1934a:210. Type-species, *palmarum* Townsend (orig. des.).
PALMARUM Townsend, 1934a:210. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus TARPESSITA Reinhard

- TARPESSITA Reinhard, 1967:95. Type-species, *fulgens* Reinhard (orig. des.).
FULGENS Reinhard, 1967:96. Type-locality: Colombia, Cundinamarca, Fusagasugá.

Genus TROCHILCHAETA Townsend

- TROCHILCHAETA Townsend, 1940b:891. Type-species, *transcendens* Townsend (orig. des.).
TRANSCENDENS Townsend, 1940b:892. Type-locality: Brazil, Mato Grosso, Maracaju.

Genus VISAYALYDINA Townsend

- VISAYALYDINA Townsend, 1926:31. Type-species, *sierricola* Townsend (orig. des.).
SIERRICOLA Townsend, 1926:32. Type-locality: Mexico, Chihuahua, near San José de Barbicora, Las Visayas.

Genus XANTHOPELTA Aldrich

- XANTHOPELTA Aldrich, 1934:48. Type-species, *scutellaris* Aldrich (orig. des.).
SCUTELLARIS Aldrich, 1934:49. Type-locality: Argentina, Rio Negro, Bariloche.

Tribe CAMPYLOCHETINI

The hosts of the species of Campylochetini are lepidopterous larvae.

Genus CHAETOPHLEPSIS Townsend

- CHAETOPHLEPSIS Townsend, 1915m:422. Type-species, *tarsalis* Townsend (orig. des.).
Ref. — Reinhard, 1952b (rev.).
ANCISA Reinhard, 1952b:17. Type-locality: "Mexico". Distr. — Central America and West Indies.
LIPERNIS Reinhard, 1952b:20. Type-locality: Brazil, Mato Grosso.
TARSALIS Townsend, 1915m:423. Type-locality: Peru, San Rafael, Casma.
Ref. — Reinhard, 1952b:16.

Genus HOMOHYPCHAETA Townsend

- HOMOHYPCHAETA Townsend, 1927a:255. Type-species, *reclinata* Townsend (orig. des.).
RECLINATA Townsend, 1927a:316. Type-locality: Peru, Charape River.
UCAYALI (Townsend), 1929:379 (*Chaetophlepsis*). Type-locality: Peru, Ucayali River, Yarina-cocha.

Genus HYPOCHAETOPSIS Townsend

- HYPOCHAETOPSIS Townsend, 1915m:422. Type-species, *chaetosa* Townsend (orig. des.).
CHAETOSA Townsend, 1915m:422. Type-locality: Peru, Rio Huallaga Canyon, Huariaca.

Genus NEOCAMPYLOCHAETA Townsend

- NEOCAMPYLOCHAETA Townsend, 1927a:255. Type-species, *genalis* Townsend (orig. des.).
GENALIS Townsend, 1927a:332. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Tribe VORIINI

Members of the tribe Voriini parasitize many kinds of lepidopterous larvae.

Ref. — D'Aguiar, 1957 (morphology).

Genus ALPINOPLAGIA Townsend

- ALPINOPLAGIA Townsend, 1931a:475. Type-species, *boliviana* Townsend (orig. des.).
BOLIVIANA Townsend, 1931a:476. Type-locality: Bolivia, Cuesta de Cillunticara.

Genus ATEOGLUTUS Aldrich

- ATEOGLUTUS Aldrich, 1934:24. Type-species, *ruficornis* Aldrich (orig. des.).
CHILENSIS (Brèthes), 1920b:42 (*Phorichaeta*). Type-locality: Chile, Santiago,

- Río Blanco. Distr. — Chile (Santiago) and Argentina. Ref. — Cortés, 1963:251 (notes on types).
- NITENS Aldrich, 1934:26. Type-locality: Argentina, Río Negro, Lake Nahuel Huapi.
- RUFICORNIS Aldrich, 1934:25. Type-locality: Argentina, Río Negro, Lake Nahuel Huapi. Distr. — Argentina (Lake Nahuel Huapi), Chile (Angol, Valparaiso).

Genus *ATHRYCIA* Robineau-Desvoidy

- ATHRYCIA Robineau-Desvoidy, 1830:111. Type-species, *erythrocerus* Robineau-Desvoidy (Robineau-Desvoidy, 1863:830).
- CINEREA (Coquillett), 1895d:101 (*Paraplagia*). Type-locality: U.S.A., Illinois. Distr. — U.S.A., Mexico.
- spinosula*, authors, not Bigot.

Genus *CATALINOVORIA* Townsend

- CATALINOVORIA Townsend, 1926:37. Type-species, *cauta* Townsend (orig. des.).
- CAUTA Townsend, 1926:38. Type-locality: U.S.A., Arizona. Distr. — Canada, U.S.A., Mexico.

Genus *CHAETOPLAGIA* Coquillett

- CHAETOPLAGIA Coquillett, 1895d:98. Type-species, *atripennis* Coquillett (orig. des.).
- ATRIPENNIS Coquillett, 1895d:98. Type-locality: U.S.A. Distr. — U.S.A., Mexico.

Genus *COWANIA* Reinhard

- COWANIA Reinhard, 1952a:3. Type-species, *icheeleri* Reinhard (orig. des.).
- WHEELERI Reinhard, 1952a:3. Type-locality: Mexico, Zacatecas, Fresnillo.

Genus *ERGOLABUS* Reinhard

- ERGOLABUS Reinhard, 1964c:16. Type-species, *bonasus* Reinhard (orig. des.).
- BONASUS Reinhard, 1964c:16. Type-locality: Mexico, Morelos, Cuernavaca.

Genus *EUCYRTOPHLOEBA* Townsend

- EUCYRTOPHLOEBA Townsend, 1916f:316. Type-species, *rhois* Townsend (orig. des.).
- HORREDA (Giglio-Tos), 1893:6 (*Cyrtophloeoba*). Type-locality: "Mexico".
- RHOIS Townsend, 1916f:317. Type-locality: Mexico, west Chihuahua, Meadow Valley.

Genus *HELIAEA* Curran

- HELIAEA Curran, 1934c:492 (key). Type-species, *mirabilis* Curran (mon.).
- MIRABILIS Curran, 1934c:517. Type-locality: Venezuela, Guanoco.

Genus ITAVORIA Townsend

ITAVORIA Townsend, 1931a:474. Type-species, *aureescens* Townsend (orig. des.).

AURESCENS Townsend, 1931a:475. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus MELETERUS Aldrich

MELETERUS Aldrich, 1926c:20. Type-species, *montanus* Aldrich (orig. des.).

MONTANUS Aldrich, 1926c:20. Type-locality: Mexico, Chihuahua, Sierra Madre Mts. Distr. — U.S.A., Mexico (Chihuahua).

Genus MICROPLAGIA Townsend

MICROPLAGIA Townsend, 1915:437. Type-species, *nitens* Townsend (orig. des.).

NITENS Townsend, 1915:437. Type-locality: Peru, Lima.

Genus MINTHOPLAGIA Townsend

MINTHOPLAGIA Townsend, 1915h:92. Type-species, *rafaeli* Townsend (orig. des.).

RAFAELI Townsend, 1915h:92. Type-locality: Mexico, Veraacruz, San Rafael. Ref. — Thompson, 1961:27 (Immature stages).

SETIFRONS (Wulp), 1890:101 (*Plagia*). Type-locality: Mexico, Guerrero. N. COMB.

Genus MYIOCHAETA Cortés

MYIOCHAETA Cortés, 1967b:24. Type-species, *marnefi* Cortés (orig. des.).

MARNEFI Cortés, 1967b:25, fig. 5. Type-locality: Chile, Santiago, Rincónada de Malpu, Quebrada de La Plata.

Genus NEOPAEDARIUM Blanchard

NEOPAEDARIUM Blanchard, 1943b:155. Type-species, *subauratum* Blanchard (orig. des.).

SUBAURATUM Blanchard, 1943b:155, fig. 28. Type-locality: Argentina, Salta.

Genus NEPHOPLAGIA Townsend

NEPHOPLAGIA Townsend, 1919:171. Type-species, *arcuata* Townsend (orig. des.).

ARCUATA Townsend, 1919:172. Type-locality: Peru, Jaén, Huascaray.

Genus NEOCHAETOPLAGIA Blanchard

NEOCHAETOPLAGIA Blanchard, 1963:173. Type-species, *pastranai* Blanchard (orig. des.).

PASTRANAI Blanchard, 1963:173, fig. 14. Type-locality: Argentina, Buenos Aires.

Genus NEOCYRTOPHOEBIA Vimmer & Soukup

NEOCYRTOPHOEBIA Vimmer & Soukup, 1940:363. Type-species, *heyrovskyi* Vimmer & Soukup (mon.).

HEYROVSKYI Vimmer & Soukup, 1940:363. Type-locality: Peru, Puno.

Genus OPSOPHAGUS Aldrich

- OPSOPHAGUS Aldrich, 1926c:15. Type-species, *ornatus* Aldrich (orig. des.).
 Ref. — Caltagirone, 1966 (key).
 CORTESI Caltagirone, 1966:64, figs. 1,2. Type-locality: Chile, Talca, Gualleco.
 ORNATUS Aldrich, 1926c:15. Type-locality: Peru, Jaén, Huascaray.
 NIGRIPALPIS Aldrich, 1926c:16. Type-locality: Chile, Valparaiso, Perales.

Genus PACHYNOCERA Townsend

- PACHYNOCERA Townsend, 1919a:585. Type-species, *petiolata* Townsend (orig. des.).
 PETICLATA Townsend, 1919a:585. Type-locality: Peru, Jaén, Charape River.

Genus PAEDARIUM Aldrich

- PAEDARIUM Aldrich, 1926c:22. Type-species, *parvum* Aldrich (orig. des.).
Vorallia Curran, 1934d:448. Type-species, *Voria neotropica* Curran (orig. des.).
 NEOTROPICUM (Curran), 1926a:109 (*Voria*). Type-locality: Jamaica, West Indies, Mona and Hope Gardens.
 PARVUM Aldrich, 1926c:22. Type-locality: Jamaica, West Indies, Montego Bay.
 PUNCTIPENNIS (Walker), 1857:208 (? *Sarcophaga*). Type-locality: "Colombia".

Genus PERISCEPSIA Gistel

- Scopolia* Robineau-Desvoidy, 1830:268 (preocc. Hübner, 1825 or earlier).
 Type-species, *Musca carbonaria* Panzer (Zetterstedt, 1844:1239).
 PERISCEPSIA Gistel, 1848:lx (nom. nov. for *Scopolia* Robineau-Desvoidy).
 Type-species, *Musca carbonaria* Panzer (aut.).
Wagneria authors, in part, not Robineau-Desvoidy.
 Ref. — Reinhard, 1955 (rev. as *Wagneria*, sensu lat.).
 LAEVIGATA (Wulp), 1890:205, pl. 4, fig. 17-17a (*Rhinophora*). Type-locality: Mexico, Veracruz, Atoyac. Distr. — Nearctic, Mexico, Guatemala.
 NIGRA (Bigot), 1857:814, pl. 20, fig. 8-8a (French edition), (Spanish edition, p. 342) (*Scopolia*). Type-locality: "Cuba". Unrecognized.

Genus PLAGIOMIMA Brauer & Bergenstamm

- PLAGIOMIMA Brauer & Bergenstamm, 1891:384 (1891:80). Type-species, *disparata* Brauer & Bergenstamm (mon.).
 FACETA Reinhard, 1957:105. Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico.
 INCOGNITA (Wulp), 1890:103 (*Plagia*). Type-locality: Mexico, Guerrero.
 RIGIDIROSTRIS (Wulp), 1890:102 (*Plagia*). Type-locality: "Mexico". N. comb.

Genus PSEUDORRHINACTIA Thompson

- PSEUDORRHINACTIA Thompson, 1968:120. Type-species, *rubricornis* Thompson (orig. des.).
 RUBRICORNIS Thompson, 1968:120. Type-locality: Trinidad, Brazil (village).

Genus VORIA Robineau-Desvoidy

- VORIA Robineau-Desvoidy, 1830:195. Type-species, *latifrons* Robineau-Desvoidy (mon.) = *ruralis* (Fallén).
Plagia Meigen, 1838:201. Type-species, *Tachina verticalis* Meigen (Rondani, 1856:69) = *ruralis* Fallén.
Ref. — Aldrich, 1926; d'Agullar, 1957:259; Thompson, 1961:28.
DICTA (Giglio-Tos), 1893:5 (*Plagia*). Type-locality: "Mexico". Unrecognized.
PARVA (Johnson), 1920:436 (*Plagia*). Type-locality: Jamaica, Liguanea Plain, near Kingston.
rufitorax (Pazos), 1914:1002 (*Plagia*). *Nomen nudum*.
RURALIS (Fallén), 1810:265 (*Tachina*). Type-locality: "Sweden". Distr. — Europe, U.S.A., Mexico, South America.
americana Wulp, 1890:102, pl. 3, fig. 19 (*Plagia*). Type-locality: Mexico, several places.
? *mexicana* Giglio-Tos, 1893:5 (*Plagia*). Type-locality: "Mexico".
brasiliiana Townsend, 1929:380. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
brasiliensis Townsend, error.
? *ayerzai* Blanchard, 1943:157, fig. 29. Type-locality: "Argentina".
SIGNATA (Walker), 1861:298 (*Eurigaster*). Type-locality: "Mexico". Unrecognized. Ref. — Austen, 1907:337.

Genus XENOPLAGIA Townsend

- XENOPLAGIA Townsend, 1914:13. Type-species, *setosa* Townsend (orig. des.).
SETOSA Townsend, 1914:14. Type-locality: Peru, Samán, Río Chira.

Tribe HESPEROPHASIINI

This tribe proposed by Townsend, was originally placed in the Phasiinae. The study of anatomy of male genitalia of some representatives of this tribe shows a close relationship with the Voriini.

Ref. — Curran, 1927b (key to American spp.).

Genus HESPEROPHASIA Townsend

- HESPEROPHASIA Townsend, 1915g:220. Type-species, *setosa* Townsend (orig. des.).
AENESCEENS Curran, 1927b:302. Type-locality: Mexico, Chihuahua, Sierra Madre.
SETOSA Townsend, 1915g:221. Type-locality: U.S.A., New Mexico. Distr. — U.S.A., Mexico.

Subfamily DEXIINAE

The members of the subfamily Dexiinae have a great variety of hosts. The main hosts are lepidopterous larvae, some coleopterous larvae and other insects.

Tribe THELAIRINI

The species of tribe Thelairini are parasites of various lepidopterous larvae, chiefly Noctuidae and Arctiidae.

Genus ARRHINACTIA Townsend

- ARRHINACTIA Townsend, 1927a:258. Type-species, *cylindrica* Townsend (orig. des.).
Archynactia, misprint.
Minthomima Townsend, 1927a:380. Type-species, *chaetosa* Townsend (orig. des.). N. SYN.
CHAETOSA (Townsend), 1927a:381 (*Minthomima*). Type-locality: Brazil, Amazonas, Manaus. N. COMB.
CYLINDRICA Townsend, 1927a:288. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus GYMNOPALPUS Townsend

- GYMNOPALPUS Townsend, 1919b:172. Type-species, *setipennis* Townsend (orig. des.).
SETIPENNIS Townsend, 1919b:172. Type-locality: Guatemala, Los Amates and unknown locality. Ref. — Thompson, 1963:508.

Genus NEPHOCHAETONA Townsend

- NEPHOCHAETONA Townsend, 1919b:174. Type-species, *mima* Townsend (orig. des.).
MIMA Townsend, 1919b:174. Type-locality: Peru, Jaen, Huascaray.

Genus PARODOMYIOPS Townsend

- PARODOMYIOPS Townsend, 1935d:221. Type-species, *thelairodops* Townsend (orig. des.).
THELAIRODOPS Townsend, 1935d:221. Type-locality: Guyana, Kanaku Mts. Ref. — Thompson, 1963a:505.

Genus POLYGASTER Wulp

- POLYGASTER Wulp, 1890:43 (key). Type-species, *egregia* Wulp (sub. mon.; Wulp, 1890:139).
Polygastropsis Townsend, 1919b:173. Type-species, *Polygaster brasiliensis* Townsend (orig. des.).
Actinomintho Townsend, 1928b:144. Type-species, *ivu* Townsend (orig. des.).
BRASILIENSIS Townsend, 1917b:224. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães). Ref. — Arnaud, 1963a:125 (notes).
EGREGIA Wulp, 1890:139, pl. 4, figs. 3-3a. Type-locality: Mexico, Tabasco, Teapa.
IVU (Townsend), 1928b:145 (*Actinomintho*). Type-locality: Paraguay, Ivu.

Genus RHOMBOTHYRIA Wulp

- RHOMBOTHYRIA Wulp, 1891:213 (key). Type-species, *flavicosta* Wulp (sub. mon.; Wulp, 1891:259).
FLAVICOSTA Wulp, 1891:259, pl. 6, fig. 13. Type-locality: Mexico, Guerrero, Venta de Zopilote. Ref. — Townsend, 1931b:89.

Genus SPATHIDEXIA Townsersd

- SPATHIDEXIA Townsend, 1912b:110. Type-species, *clemonsi* Townsend (orig. des.).
- Minthohoughia* Townsend, 1919a:581. Type-species, *cylindrica* Townsend (orig. des.). N. SYN.
- Stenaulacodoria* Townsend, 1928e:161. Type-species, *spatulata* Townsend (orig. des.).
Ref. — Arnaud, 1960 (rev.).
- ANTILLENIS Arnaud, 1960:6, figs. 2, 11, 15, 27. Type-locality: Puerto Rico, Mayaguez. Distr. — Puerto Rico and Cuba.
- BRASILIENSIS Arnaud, 1960:10, figs. 5, 9, 18, 19. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- EINEREMCOLLIS (Wulp), 1891:255 (*Thelairodes*). Type-locality: Mexico, Guerrero. N. COMB.
- EYLINDRICA (Townsend), 1919a:581 (*Minthohoughia*). Type-locality: Peru, Lima. N. COMB.
- DUNNINGH (Coquillett), 1895e:54 (*Thryptocera*). Type-locality: U.S.A., Illinois. Distr. — U.S.A., Jamaica, Puerto Rico.
- NEXA Reinhard, 1953b:94. Type-locality: Mexico, Mexico City, D F. Distr. — U.S.A., Mexico.
- NIVEOMARGINATA (Wulp), 1890:200 (*Anisia*). Type-locality: Mexico, Guerrero. N. BOMB.
- SPATULATA (Townsend), 1928e:162 (*Stenaulacodoria*). Type-locality: Peru, Cafete. Ref. — Thompson, 1963a:497.

Genus THELAIRA Robineau-Desvoidy

- THELAIRA Robineau-Desvoidy, 1830:214. Type-species, *abdominalis* Robineau-Desvoidy (Townsend, 1916:9) = *nigripes* (Fabrieius).
- AMERICANA Brooks, 1945:88. Type-locality: Canada, Quebec. Distr. — Canada, U.S.A., Mexico.

Genus THELAIRODES Wulp

- THELAIRODES Wulp, 1891:213 (key), 254. Type-species, *Homoderia vittigera* Bigot (Brauer & Bergenstamm, 1891:377; 1891:73).
- Parodomysia* Townsend, 1917b:225. Type-species, *paradoxica* Townsend (orig. des.).
Ref. — Thompson, 1963a:502.
- LAVINIA Curran, 1934e:503. Type-locality: Guyana, Bartica, Penal Settlement.
- PALLIDUS Wulp, 1891:255, pl. 6, fig. 10, 10a. Type-locality: Mexico, Guerrero, Venta de Zopilote.
- PARADOXIEUS (Townsend), 1917b:226 (*Parodomysia*). Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).
- SPINOSUS (Bigot), 1889:268 (*Homoderia*). Type-locality: "Mexico". Ref. — Thompson, 1963a:501.
- VITTIGERUS (Bigot), 1889:267 (*Homoderia*). Type-locality: "Mexico". Ref. — Thompson, 1963a:501.

Genus THELAIROMIMA Townsend

- THELAIROMIMA Townsend, 1935d:222. Type-species, *plutipennis* Townsend (orig. des.).
PLUTIPENNIS Townsend, 1935d:222. Type-locality: Brazil, São Paulo, São Vicente.

Genus XANTHODEXIA Wulp

- XANTHODEXIA Wulp, 1891:213 (key). Type-species, *Tachina sericea* Wiedemann (sub. mon.; Wulp, 1891:256; misident.) = *semipicta* (Walker).
Minthodexia Brauer & Bergenstamm, 1891:371 (1891:67). Type-species, *gravipes* Brauer & Bergenstamm (sub. des.; Brauer & Bergenstamm, 1893:131) = *sericea* (Wiedemann).
SEMIPICTA (Walker), 1852:316 (*Dexia*). Type-locality: Brazil, Amazonas.
suffusa Walker, 1852:317 (*Dexia*). Type-locality: unknown.
extrema Walker, 1857:203 (*Dexia*). Type-locality: Brazil, Amazonas.
sericea Wiedemann of Wulp.
SERICEA (Wiedemann), 1830:136 (*Tachina*). Type-locality: "Brazil".
gravipes Brauer & Bergenstamm, 1891:376 (1891:72) (*Minthodexia*). Type-locality: "Venezuela".

Genus ZONALIA Curran

- ZONALIA Curran, 1934d:429. Type-species, *nitens* Curran (orig. des.).
NITENS Curran, 1934d:464, figs. 188, 195. Type-locality: Panama, Canal Zone, Corozal (cf. Arnaud, 1963a:132).

Tribe DEXIINI

Genus ITAMINTHO Townsend

- ITAMINTHO Townsend, 1931a:329. Type-species, *erro* Townsend (orig. des.).
ERRO Townsend, 1931a:330. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus MASTIGIOMYIA Reinhard

- MASTIGIOMYIA Reinhard, 1964c:17. Type-species, *delusa* Reinhard (orig. des.).
DELUSA Reinhard, 1964c:18. Type-locality: Mexico, Cuernavaca.

Genus MICROAPORIA Townsend

- MICROAPORIA Townsend, 1919a:560. Type-species, *elegans* Townsend (orig. des.).
ELEGANS Townsend, 1919a:561. Type-locality: Peru, Cuzco.

Genus PARAZELIA Townsend

- PARAZELIA Townsend, 1919a:556. Type-species, *pulchra* Townsend (orig. des.).
PULCHRA Townsend, 1919a:556. Type-locality: Mexico, Cuernavaca.

Genus TRINITODEXIA Townsend

TRINITODEXIA Townsend, 1935d:222. Type-species, *trichops* Townsend (orig. des.).

TRICHOPS Townsend, 1935d:222. Type-locality: "Trinidad".

UNPLACED SPECIES OF DEXIINI

PICTIGASTER Bigot, 1889:261 (*Ceromasia*). Type-locality: "Mexico".

Tribe MACQUARTIINI

The species of Macquartiini have been reared from a variety of hosts, including chrysomelid beetles and several lepidopterous and sawflies.

Genus ALEXOGLOBLINIA Cortés

ALEXOGLOBLINIA Cortés, 1945b:256. Type-species, *Metopomuscopteryx shannoni* Aldrich (orig. des.).

SHANNONI (Aldrich), 1934:46 (*Metopomuscopteryx*). Type-locality: Argentina, Río Negro, Lake Nahuel Huapi.

Genus CHILOCLISTA Townsend

CHILOCLISTA Townsend, 1931a:334. Type-species, *bicolor* Townsend (orig. des.).

BICOLOR Townsend, 1931a:334. Type-locality: Chile, Tanumé.

Genus DASYUROMYIA Bigot

DASYUROMYIA Bigot, 1885c:liv. Type-species, *penicillata* Bigot (mon.) = *inornata* (Walker).

Selenomyia Brauer & Bergenstamm, 1891:361 (1891:57). Type-species, *brevicornis* Brauer & Bergenstamm (mon.) = *sarcophagidea* (Bigot).

Mesembriophyto Townsend, 1916g:301. Type-species, *magellana* Townsend (orig. des.).

Ref. — Aldrich, 1934:157 (key to species).

APERTA Aldrich, 1934:161. Type-locality: Argentina, Río Negro, Lake Nahuel Huapi.

INORNATA (Walker), 1837:349 (*Tachina*). Type-locality: Chile, Cape Gregory. Ref. — Cortés, 1963:244.

penicillata Bigot, 1885c:lv. Type-locality: "Chile".

magellana Townsend, 1916g:301 (*Mesembriophyto*). Type-locality: Chile, Straits of Magellan.

LLOYDI Blanchard, 1947b:258, figs. a-g. Type-locality: Argentina, Río Negro, El Bolsón.

NERVOSA (Walker), 1837:349 (*Tachina*). Type-locality: Chile, Port Famine, Str. of Magellan. Ref. — Cortés, 1963:243 (note on type).

NIGRICEPS Aldrich, 1934:162. Type-locality: Chile, Llanquihue, Casa Pangué.

SARCOPHAGIDEA (Bigot), 1888b:28, pl. 3, fig. 8 (*Hyadesimyia*). Type-locality: Chile, Cape Horn.

- brevicornis* Brauer & Bergenstamm, 1891:361 (1891:57) (*Selenomyia*).
Type-locality: "Chile".
- STERNALIS Aldrich, 1934:160. Type-locality: Chile, Malleco, Angol.
- TARSALIS Aldrich, 1934:160. Type-locality: Argentina, Rio Negro, Lake
Gutiérrez. Ref. — Cortés, 1963:245.

Genus DISCHOTRICHIA Cortés

- DISCHOTRICHIA Cortés, 1944d:54. Type-species, *caellobata* Cortés (orig. des.).
- CAELLOBATA Cortés, 1944d:56. Type-locality: Chile, Valparaíso, Marga-Marga.

Genus EUTHYPROSOPIELLA Blanchard

- EUTHYPROSOPIELLA Blanchard, 1963:186. Type-species, *mendocina* Blanchard
(orig. des.).
- MENDOCINA Blanchard, 1963:186, fig. 19. Type-locality: Argentina, Mendoza,
Uspallata.

Genus GANOPLEURON Aldrich

- GANOPLEURON Aldrich, 1934:118. Type-species, *divergens* Aldrich (orig. des.).
- DIVERGENS Aldrich, 1934:119. Type-locality: Chile, Chiloé I, Castro.

Genus GONATORRHINA Röder

- GONATORRHINA Röder, 1886:264 (also, 1892:10). Type-species, *paramonensis*
Röder (mon.).
- PARAMONENSIS Röder, 1886:265 (also, 1892:10), figs. 2, 2a, 2b. Type-locality:
Colombia, Páramo, 3600 m.

Genus HYOSOMA Aldrich

- HYOSOMA Aldrich, 1934:139. Type-species, *limbisquama* Aldrich (orig. des.).
- LIMBISQUAMUM Aldrich, 1934:140, figs. 16-17. Type-locality: Argentina,
Puerto Blest.

Genus LAFUENTEMYIA Marnef

- LAFUENTEMYIA Marnef, 1965:243. Type-species, *yanezi* Marnef (orig. des.).
- YANEZI Marnef, 1965:246, figs. 1-2. Type-locality: Chile, Valparaíso, Parque
Nacional del Lago Peñuelas.

Genus MATUCANIA Townsend

- MATUCANIA Townsend, 1919a:568. Type-species, *mellisquama* Townsend
(orig. des.).
- MELLISQUAMA Townsend, 1919a:568. Type-locality: Peru, Matucana.

Genus MUSCOPTERYX Townsend

- MUSCOPTERYX Townsend, 1892:170. Type-species, *chaetosula* Townsend
(orig. des.).

- Psammopia* Townsend, 1915b:20. Type-species, *Brachycoma pulvrea* Coquillett (orig. des.) = *chaetosula* Townsend.
CHAETOSULA Townsend, 1892:171. Type-locality: Mexico, Chihuahua. Distr. — U.S.A., Mexico.
PETENTIS Reinhard, 1958e:280. Type-locality: U.S.A., Arizona. Distr. — U.S.A., Mexico.

Genus NOTODYTES Aldrich

- NOTODYTES* Aldrich, 1934:163. Type-species, *variabilis* Aldrich (orig. des.). Ref. — Aldrich, 1934:163 (key to spp.).
AUREA Aldrich, 1934:165. Type-locality: Chile, Llanquihue, Ensenada.
MAJOR Aldrich, 1934:165. Type-locality: Chile, Malleco, Angol.
VARIABILIS Aldrich, 1934:164. Type-locality: Argentina, Rio Negro, Bariloche.

Genus NOTOMANES Aldrich

- NOTOMANES* Aldrich, 1934:93. Type-species, *Tachina maura* Walker (orig. des.) = *basalis* (Walker).
BASALIS (Walker), 1837:351 (*Tachina*). Type-locality: Chile, Strait of Magellan, Port Famine (Aldrich, 1934, adopted *maura*, but Austen, 1907:330, was the first reviser).
maura Walker, 1837:352 (*Tachina*). Type-locality: Chile, Strait of Magellan, Port Famine.

Genus PELYCOPS Aldrich

- PELYCOPS* Aldrich, 1934:168. Type-species, *darwinii* Aldrich (orig. des.).
DARWINI Aldrich, 1934:169. Type-locality: Chile, Port Famine, Tierra del Fuego. Ref. — Verbeke, 1962:98, pl. V, fig. 6 (genitalia).

Genus PIRIONA Aldrich

- PIRIONA* Aldrich, 1928b:24. Type-species, *fasciculata* Aldrich (orig. des.).
FASCICULATA Aldrich, 1928b:24, fig. 2. Type-locality: Chile, Marga-Marga. Distr. — Argentina (Bariloche), Chile.

Genus PROCLEONICE Townsend

- PROCLEONICE* Townsend, 1935d:223. Type-species, *prolixa* Townsend (orig. des.).
PROLIXA Townsend, 1935d:223. Type-locality: "Trinidad".

Genus PROSOPOCHAETA Macquart

- PROSOPOCHAETA* Macquart, 1851:83 (1851:209). Type-species, *nitidiventris* Macquart (orig. des.).
Punaclista Townsend, 1915m:406. Type-species, *setosa* Townsend (orig. des.).
ANOMALA Aldrich, 1934:118. Type-locality: Argentina, Rio Negro, Lake Nahuel Huapi, Jones Estancia. Distr. — Chile (Concepción, Angol), Argentina.
FIDELIS (Reinhard), 1967:98 (*Punaclista*). Type-locality: Peru, Lima. N. COMB.

NITIDIVENTRIS Macquart, 1851:184 (1851:211), pl. 19, fig. 5. Type-locality: "Chile".

SETOSA (Townsend), 1915m:407 (*Punaclista*). Type-locality: Peru, Oroya.

Genus PSEUDOMORINIA Wulp

PSEUDOMORINIA Wulp, 1891:213 (key). Type-species, *picitipennis* Wulp (sub. mon.; Wulp, 1891:259).

PICITIPENNIS Wulp, 1891:260, pl. 6, fig. 14-14a. Type-locality: Mexico, Guerrero, Xucumanatlán.

Genus RHINOMACQUARTIA Brauer & Bergenstamm

RHINOMACQUARTIA Brauer & Bergenstamm, 1891:380 (1891:76). Type-species, *chaetophora* Brauer & Bergenstamm (mon.).

CHAETOPHORA Brauer & Bergenstamm, 1891:381 (1891:76). Type-locality: Brazil, São Paulo, Varnhagen (formerly Ypanema). Ref. — Aldrich, 1927c:7 (note).

Genus SCHLINGERMYIA Cortés

SCHLINGERMYIA Cortés, 1967b:20. Type-species, *venusta* Cortés (orig. des.).

VENUSTA Cortés, 1967b:22, fig. 4. Type-locality: Chile, Valparaíso, Margamarga, Los Perales.

Genus SETOLESTES Aldrich

SETOLESTES Aldrich, 1934:142. Type-species, *genalis* Aldrich (orig. des.).

GENALIS Aldrich, 1934:142. Type-locality: Chile, Chiloé I, Castro.

Genus TRICHODISCHIA Bigot

TRICHODISCHIA Bigot, 1885c:xliv. Type-species, *soror* Bigot (Townsend, 1916:9). Ref. — Cortés, 1969:97.

CAERULEA Bigot, 1885c:xliv. Type-locality: Argentina, Buenos Aires. Distr. — Argentina (Buenos Aires), Chile (Coquimbo).

SOROR Bigot, 1885c:xlvi. Type-locality: Argentina, Buenos Aires. Distr. — Argentina (Buenos Aires), Chile (Santiago).

caerulea Bigot of Cortés, 1944:49.

UNRECOGNIZED SPECIES OF MACQUARTIINI

QUADRIMACULATA Macquart, 1855:200 (1855:227) (*Gymnostylia*). Type-locality: "South America". Ref. — Crosskey, 1971:271.

SOLITARIA Aldrich, 1934:168 (*Mylophasia*). Type-locality: Chile, Malleco, Angol.

Tribe ICELIINI

Genus ERVIA Robineau-Desvoidy

ERVIA Robineau-Desvoidy, 1820:225. Type-species, *Ocyptera triquetra* Olivier (mon.).

- ANGULATA (Wulp), 1890:136 (*Myobia*). Type-locality: Mexico, Guerrero, Chilpancingo. N. COMB.
 TRIQUETRA (Olivier), 1812:423 (*Ocyptera*). Type-locality: U.S.A., Carolina. Distr. — U.S.A., Central America and West Indies. Ref. — Thompson, 1963:446.

Genus ERVIOPSIS Townsend

- ERVIPSIS Townsend, 1934:211. Type-species, *aurata* Townsend (orig. des.).
 AURATA Townsend, 1934a:211. Type-locality: Brazil, Pará, Boa Vista, Tapajós River.

Genus ICELIA Robineau-Desvoidy

- ICELIA Robineau-Desvoidy, 1830:224. Type-species, *flavescens* Robineau-Desvoidy (Townsend, 1916c:7).
 BRASILIENSIS Robineau-Desvoidy, 1830:224. Type-locality: "Brazil".
 FLAVESCENS Robineau-Desvoidy, 1830:224. Type-locality: Brazil, Minas Gerais.

Tribe ZELIINI

The members of the tribe Zeliini have been recorded as parasites of passalid, elaterid and scarabaeid larvae.

Genus DIAUGHIA Perty

- DIAUGHIA Perty, 1833:187. Type-species, *angusta* Perty (mon.).
 ANGUSTA Perty, 1833:187, pl. 37, fig. 9. Type-locality: Brazil, Minas Gerais.

Genus HUASCARODENIA Townsend

- HUASCARODENIA Townsend, 1919b:176. Type-species, *pulchra* Townsend (orig. des.).
 PULCHRA Townsend, 1919b:176. Type-locality: Peru, Jaen, Huascaray.

Genus METADEXIA Coquillett

- METADEXIA Coquillett, 1899:20. Type-species, *tricolor* Coquillett (orig. des., as gen. n., sp. n.).
 TRICOLOR Coquillett, 1899:220. Type-locality: U.S.A., Louisiana. Distr. — U.S.A., Mexico.

Genus OPHIRODEXIA Townsend

- OPHIRODEXIA Townsend, 1911b:134 (also, 1912a:307). Type-species, *pulchra* Townsend (orig. des., as gen n., sp. n.).
 PULCHRA Townsend, 1911b:134 (also, 1912a:308). Type-locality: Peru, Piura.

Genus OPSOZELIA Townsend

- OPSOZELIA Townsend, 1919a:575. Type-species, *discahis* Townsend (orig. des.)
 = *lateralis* (Fabricius).

LATERALIS (Fabricius), 1805:286 (*Musca*). Type-locality: "South America".
Ref. — Townsend, 1931b:96.
discaelis Townsend, 1919a:557. Type-locality: Guyana, Bartica, Kartabo.

Genus TAPERAMYIA Townsend

TAPFRAMYIA Townsend, 1935d:223. Type-species, *pickeli* Townsend (orig. des.).
PICKELI Townsend, 1935d:223. Type-locality: Brazil, Pernambuco, Tapera.

Genus TROMODESIANA Townsend

TROMODESIANA Townsend, 1931a:339. Type-species, *Dexia thomae* Wiedemann (orig. des.).
THOMAE (Wiedemann), 1830:379 (*Dexia*). Type-locality: West Indies, St. Thomas I.

Genus TROMODESIOPSIS Townsend

TROMODESIOPSIS Townsend, 1927a:221. Type-species, *Tromodesia haemorrhoidalis* Bigot (orig. des.) = *atrifrons* (Wiedemann).
ATRIFRONS (Wiedemann), 1830:403 (*Musca*). Type-locality: Unknown. Ref. — Aldrich, 1929:4 (as *Zelia*).
haemorrhoidalis Bigot, 1889:267 (*Tromodesia*; as *hoemorrhoidalis*, error). Type-locality: "Mexico".

Genus USHPAYACUA Townsend

USHPAYACUA Townsend, 1928a:145. Type-species, *ureophila* Townsend (orig. des.).
UREOPHILA Townsend, 1928b:145. Type-locality: Peru, Ushpayacu River.

Genus YAHUARMAYOIA Townsend

YAHUARMAYOIA Townsend, 1927a:228. Type-species, *analís* Townsend (orig. des.) = *phacoptera* (Wiedemann).
PHAEOPTERA (Wiedemann), 1830:370 (*Dexia*). Type-locality: "Brazil". Ref. — Aldrich, 1929a:2.
analís Townsend, 1927a:367. Type-locality: Peru, Yahuarmayo.

Genus ZELIA Robineau-Desvoidy

ZELIA Robineau-Desvoidy, 1830:314. Type-species, *rostrata* Robineau-Desvoidy (Coquillett, 1910:621) = *vertebrata* (Say).
Leptoda Wulp, 1885:196. Type-species, *Dexia gracilis* Wiedemann (Wulp, 1891:250) = *vertebrata* (Say).
Melaleuca Wulp, 1891:213 (key). Type-species, *spectabilis* Wulp (sub. mon.; Wulp, 1891:247).
Ref. — Aldrich, 1925a:23 (notes), 1929:1 (notes).
LIMBATA (Wiedemann), 1830:371 (*Dexia*). Type-locality: "Brazil". Ref. — Aldrich, 1929a:5.
PERUVIANA (Brèthes), 1920:31 (*Melaleuca*). Type-locality: Peru, Lima. Unrecognized.



- PICTA (Bigot), 1889:265 (*Rhamphina*). Type-locality: "Cuba". Unrecognized. Ref. — Aldrich, 1929a:2.
- PLUMOSA (Wiedemann), 1830:370 (*Dexia*). Type-locality: "Brazil". Ref. — Aldrich, 1929:6.
- POTENS (Wiedemann), 1830:368 (*Dexia*). Type-locality: "Brazil". Ref. — Aldrich, 1929a:2.
- SEMIURFA (Wulp), 1891:250 (*Leptoda*). Type-locality: Mexico, N. Yucatán, Temax.
- SPECTABILIS (Wulp), 1891:248, pl. 6, figs. 3-3a (*Melaleuca*). Type-locality: Mexico, Guerrero, Xucumanatlán.
- VERTEBRATA (Say), 1829:176 (1859:366) (*Dexia*). Type-locality: U.S.A., Indiana. Distr. — U.S.A., Mexico.
- gracilis* Wiedemann, 1830:373 (*Dexia*). Type-locality: Unknown.

Tribe TRICHODURINI

Genus TRICHODURA Macquart

- TRICHODURA Macquart, 1843:248 (1843:91). Type-species, *Musca anceps* Fabricius (orig. des.).
- Trichoduropsis* Townsend, 1919b:169. Type-species, *Trichodura recta* (orig. des.).
- ANCEPS (Fabricius), 1805:296 (*Musca*). Type-locality: "South America".
- DORSALIS (Walker), 1852:308 (*Dexia*). Type-locality: "South America".
- GUIANENSIS (Townsend), 1919:553 (*Trichoduropsis*). Type-locality: Guyana, Bartica.
- LINEATA Townsend, 1934:390. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- vittata* Curran, 1937a:333. Type-locality: Brazil, São Paulo, Campos do Jordão. N. SYN.
- RECTA Schiner, 1868:320, pl. 4, fig. 1, a-d. Type-locality: "South America".
- VIDUA Schiner, 1868:321. Type-locality: "Brazil".

Tribe URAMYINI

Members of the tribe Uramyini parasitize lepidopterous larvae of various families.

Genus GYMNAFORIA Townsend

- GYMNAFORIA Townsend, 1919b:170. Type-species, *Gymnostylia fasciata* Macquart (orig. des.).
- FASCIATA (Macquart), 1848:212 (1848:52), pl. 6, fig. 3 (*Gymnostylia*). Type-locality: "Brazil".

Genus ORTHAPORIA Townsend

- ORTHAPORIA Townsend, 1919b:167. Type-species, *similis* Townsend (orig. des.) = *longa* (Walker).
- LONGA (Walker), 1852:311 (*Dexia*). Type-locality: "South America".
- similis* Townsend, 1916b:167. Type-locality: Brazil, Rio de Janeiro.

Genus PARAPORIA Townsend

- Aporia* Macquart, 1846:296 (1846:168) (preocc. Hübner, 1819). Type-species, *quadrimaculata* Macquart (mon.).
- Neaporia* Townsend, 1908:67 (nom. nov. for *Aporia* Macquart, but preocc. Gorham, 1897). Type-species, *Aporia quadrimaculata* Macquart (aut.).
- PARAPORIA Townsend, 1912c:48 (nom. nov. for *Neaporia* Townsend). Type-species, *Aporia quadrimaculata* Macquart (aut.).
- ACUMINATA (Wulp), 1890:130 (*Macquartia*). Type-locality: Mexico, Guerrero.
- ELEGANS (Giglio-Tos), 1893:3 (*Aporia*). Type-locality: Mexico, Tuxpango.
- QUADRIMACULATA (Macquart), 1846:297 (1846:169), pl. 15, fig. 7 (*Aporia*). Type-locality: "Colombia". Ref. — Aldrich, 1927c:26. Distr. — U.S.A., Mexico, Neotropical.

Genus PSEUDEUANTHA Townsend

- PSEUDEUANTHA Townsend, 1915m:416. Type-species, *linellii* Townsend (orig. des.) = *indita* (Walker). Ref. — Aldrich, 1921:87 (key).
- ATRIFRONS (Bigot), 1889:259 (*Macquartia*). Type-locality: "Mexico". Unrecognized.
- INDITA (Walker), 1860:306 (? *Lydella*). Type-locality: "Mexico". Distr. — U.S.A., Mexico, El Salvador.
- caloptera* Bigot, 1889:263 (*Tricoliga*). Type-locality: "Mexico".
- linellii* Townsend, 1915m:416. Type-locality: Mexico, Tehuantepec.
- OCTOMACULATA Townsend, 1919a:560. Type-locality: Peru, Huadquina.
- UMBRATILIS Reinhard, 1935a:164. Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico.
- VENUSTA Wulp, 1890:130 (*Macquartia*). Type-locality: Mexico, Guerrero. Unrecognized.

Genus URAMYA Robineau-Desvoidy

- URAMYA Robineau-Desvoidy, 1830:215. Type-species, *producta* Robineau-Desvoidy (mon.).
- Oxydextia* Bigot, 1885b:xxxiii. Type-species, *acuminata* Bigot (mon.).
- Uramya*, emend.
Ref. — Aldrich, 1921:85 (key).
- ACUMINATA (Bigot), 1885b:xxxiii (*Oxydextia*). Type-locality: "Mexico".
- ALDRICHI Reinhard, 1935a:163. Type-locality: Mexico, Oaxaca.
- BREVICAUDA Curran, 1934c:503. Type-locality: Venezuela, Guanoco.
- HARIOLA Reinhard, 1961:206. Type-locality: "Colombia".
- NITENS (Schiner), 1868:320 (*Aporia*). Type-locality: "South America". Unrecognized.
- NUBILIS (Townsend), 1929:367 (*Pseudeuanta*). Type-locality: Brazil, São Paulo, São Paulo, Cantareira.
- PRODUCTA Robineau-Desvoidy, 1830:216. Type-locality: "Brazil".
- sermyla* (Walker), 1849:850 (*Dextia*). Type-locality: "Brazil".
- SETIVENTRIS (Wulp), 1890:129, pl. 3, figs. 21-21a (*Macquartia*). Type-locality: Mexico, Orizaba and Guerrero. N. COMB.
- TRINITATIS (Thompson), 1963a:352, fig. 30 (*Uromacquartia*). Type-locality: Trinidad, Mt. Lambert. N. COMB.



Genus URAPORIA Townsend

URAPORIA Townsend, 1919b:170. Type-species, *Aporia caudata* Schiner (orig. des.).

CAUDATA (Schiner), 1868:320 (*Aporia*). Type-locality: "South America".

Tribe URODEXIINI

The tribe Urodexiini contains parasites of beetle larvae.

Genus BINGHAMIMYIA Townsend

BINGHAMIMYIA Townsend, 1919b:175. Type-species, *reclinata* Townsend (orig. des.).

RECLINATA Townsend, 1919b:175. Type-locality: Peru, Santa Ana.

Genus CALOTHELAIRA Townsend

CALOTHELAIRA Townsend, 1927a:218. Type-species, *Myobia flavicornis* Wulp (orig. des.).

FLAVICORNIS (Wulp), 1890:133, pl. 4, fig. 1 (*Myobia*). Type-locality: Mexico, Tabasco, Teapa. Ref. — Townsend, 1931b:94.

STRIGOSA (Wulp), 1890:147 (*Hypostena*). Type-locality: Mexico, Tabasco, Teapa. N. comb.

Genus CHAETONA Wulp

CHAETONA Wulp, 1891:253. Type-species, *Dexia longiseta* Wiedemann (Coquillett, 1910:521).

CONGRUA Wulp, 1891:253. Type-locality: Mexico, Guerrero, Xucumanatlán.

CRUENTA Giglio-Tos, 1893:3. Type-locality: "Mexico".

ICTERICA (Wiedemann), 1830:321 (*Tachina*). Type-locality: "Brazil". Ref. — Aldrich, 1927c:18.

citrina Bigot, 1889:262 (*Viviania*). Type-locality: "Mexico".

LONGISETA (Wiedemann), 1830:381 (*Dexia*). Type-locality: "Brazil".

PILISETA (Wulp), 1890:110 (*Masicera*). Type-locality: Mexico, Yucatán.

TUCHUCHEENSIS Thompson, 1968:87. Type-locality: Trinidad, El Tucuche.

Genus CHAETONALIA Curran

CHAETONALIA Curran, 1934d:431. Type-species, *lateralis* Curran (orig. des.).

LATERALIS Curran, 1934d:465, figs. 186, 190. Type-locality: Panama, Canal Zone, Barro Colorado I. (cf. Arnaud, 1963a:111).

Genus CHOLOMYIA Bigot

CHOLOMYIA Bigot, 1884a:xxxvii. Type-species, *inaequipes* Bigot (mon.).

ACROMIODEXIA Townsend, 1931a:335. Type-species, *Musca acromion* Wiedemann (orig. des.).

ACROMION (Wiedemann), 1824:47 (*Musca*). Type-locality: "South America". Ref. — Parker, 1953:54 (immature stages).

longipes Fabricius, 1805:398 (*Musca*; preocc. Scopoli, 1763). Type-locality: "Brazil".



CALCEATA (Wulp), 1891:258 (*Calodexia*). Type-locality: Mexico, Veracruz, Tabasco and Guerrero.
 FILIPES (Walker), 1857:202 (*Dexia*). Type-locality: Brazil, Amazon.
 INAEQUIPES Bigot, 1884a:xxxvii. Type-locality: "Mexico". Distr. — Canada, U.S.A., Mexico, Central America.
basalis Giglio-Tos, 1893a:3 (*Thelairodes*). Type-locality: "Mexico".
nigriceps Williston, 1908:353, figs. 16-17. Type-locality: Unknown.
longipes, authors, not Fabricius.

Genus ERYTHROMELANA Townsend

ERYTHROMELANA Townsend, 1919b:174. Type-species, *jaena* Townsend (orig. des.).
 JAENA Townsend, 1919b:175. Type-locality: Peru, Jaen, Huascaray.

Genus EUCALODEXIA Townsend

EUCALODEXIA Townsend, 1908:64. Type-species, *Homodexia flavipes* Bigot (orig. des.).
 FLAVIPES (Bigot), 1889:268 (*Homodexia*). Type-locality: "Mexico".

Genus ICTERICODEXIA Townsend

ICTERICODEXIA Townsend, 1934a:391. Type-species, *aristata* Townsend (orig. des.).
 ARISTATA Townsend, 1934:391. Type-locality: Brazil, Pará, Boa Vista, Tapajós River.

Genus ISCHYROPHAGA Townsend

ISCHYROPHAGA Townsend, 1915b:23. Type-species, *Thelairodes ischryri* Coquillett (orig. des.).
 ISCHYRI (Coquillett), 1905:362 (*Thelairodes*). Type-locality: Cuba, Santiago de Las Vegas.

Genus LEPTOMACQUARTIA Townsend

LEPTOMACQUARTIA Townsend, 1919a:568. Type-species, *planifrons* Townsend (orig. des.).
 PLANIFRONS Townsend, 1919a:569. Type-locality: Peru, Matucana.

Genus MEDINOPHYTO Townsend

MEDINOPHYTO Townsend, 1927a:219. Type-species, *gracilis* Townsend (orig. des.) = *dilecta* (Wiedemann).
Callesthes Aldrich, 1928b:11. Type-species, *histrio* Aldrich (orig. des.).
 DILECTA (Wiedemann), 1830:419 (*Musca*). Type-locality: "Brazil". Ref. — Aldrich, 1929a:19.
gracilis Townsend, 1927a:326. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
 HISTRIO (Aldrich), 1928b:12 (*Callesthes*). Type-locality: Ecuador, Tena. Ref. — Townsend, 1931b:93.



Genus MICROCHAETONA Townsend

- MICROCHAETONA Townsend, 1919a:565. Type-species, *gracilis* Townsend (orig. des.).
- Epiphyllophila* Townsend, 1927a:234. Type-species, *yahuarmayana* Townsend (orig. des.) = *gracilis* Townsend. N. SYN.
- Microchaetonops* Townsend, 1934a:391. Type-species, *medinops* Townsend (orig. des.). N. SYN.
- DOLICHOTARSIS (Curran), 1934c:504 (*Phillophilopsis*). Type-locality: Guyana, Bartica, Kartabo.
- GRACILIS Townsend, 1919a:565. Type-locality: Peru, Inambari River, Yahuarmayo.
- yahuarmayana* Townsend, 1927a:304 (*Epiphyllophila*). Type-locality: Peru, Yahuarmayo River and Casahuari. N. SYN.
- MEDINOPS (Townsend), 1934a:392 (*Microchaetonops*). Type-locality: Brazil, Pará, Tapajós River, Boa Vista. N. COMB.

Genus MINTHODEXIOPSIS Townsend

- MINTHODEXIOPSIS Townsend, 1927a:221. Type-species, *Minthodexia flavicornis* Brauer & Bergenstamm (orig. des.).
- ELEGANS Reinhard, 1964a:37. Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico (Veracruz).
- FLAVICORNIS (Brauer & Bergenstamm), 1891:376 (1891:72) (*Minthodexia*). Type-locality: "Venezuela".

Genus MONOESTROGASTER Townsend

- MONOESTROGASTER Townsend, 1939c:449. Type-species, *mattoensis* Townsend (orig. des.).
- MATTOENSIS Townsend, 1939c:450. Type-locality: Brazil, Mato Grosso, Maracaju.

Genus NEOPHYLLOPHILA Townsend

- NEOPHYLLOPHILA Townsend, 1927a:235. Type-species, *neotropica* Townsend (orig. des.).
- Urophyllophila* Townsend, 1927a:235. Type-species, *caudata* Townsend (orig. des.). N. SYN.
- Maracajuia* Townsend, 1939c:448. Type-species, *caudata* Townsend (1939) (orig. des.) = *caudata* (Townsend, 1927). N. SYN.
- CAUDATA (Townsend), 1927a:363 (*Urophyllophila*). Type-locality: Brazil, São Paulo, Itaquacuctuba. N. COMB.
- caudata* Townsend, 1939c:449 (*Maracajuia*). Type-locality: Brazil, Mato Grosso, Maracaju. N. SYN.
- NEOTROPICA Townsend, 1927a:334. Type-locality: Brazil, São Paulo, Itaquacuctuba.

Genus OMMALESKIA Townsend

- OMMALESKIA Townsend, 1917b:227. Type-species, *fumosa* Townsend (orig. des.).
- FUMOSA Townsend, 1917b:227. Type-locality: Brazil, Rio de Janeiro.



Genus OPSOLESKIA Townsend

OPSOLESKIA Townsend, 1919a:565. Type-species, *flava* Townsend (orig. des.).
FLAVA Townsend, 1919a:566. Type-locality: Guyana, Bartica, Kartabo.
MAJOR Thompson, 1968:89. Type-locality: Trinidad, St. Augustine.

Genus PHYLLOPHILOPSIS Townsend

Phyllophila Townsend, 1915b:21 (preocc. Guénnée, 1852). Type-species,
Chaetona nitens Coquillett (orig. des.).
PHYLLOPHILOPSIS Townsend, 1915o:78 (mon. nov. for *Phyllophila* Townsend).
Type-species, *Chaetona nitens* Coquillett (aut.).
Phillophilopsis, error.
TENUIFRONS Curran, 1934c:505. Type-locality: Guyana, Bartica, Kartabo.

Genus PSEUDCHAETONA Townsend

PSEUDCHAETONA Townsend, 1919a:563. Type-species, *polita* Townsend (orig. des.).
POLITA Townsend, 1919a:563. Type-locality: Panama (Trinidad River) and Peru (Yahuarmayo).

Genus PSEUDODEXIA Brauer & Bergenstamm

PSEUDODEXIA Brauer & Bergenstamm, 1891:678 (1891:74). Type-species,
Dexia eques Wiedemann (mon.).
Ref. — Aldrich, 1929a:28.
ALBIFACIES (Bigot), 1889:268 (*Oplisa*). Type-locality: "Mexico".
EQUES (Wiedemann), 1830:378 (*Dexia*). Type-locality: Brazil, Bahia. Ref.
— Aldrich, 1929a:28, fig. 3.
LONGICORNIS (Bigot), 1889:267 (*Homodexia*). Type-locality: "Mexico".
PALLIDICORNIS (Bigot), 1889:270 (*Anthracomyia*). Type-locality: "Mexico".
Ref. — Brauer, 1897:357, 360 (notes on Bigot's types).

Genus SCHWARZALIA Curran

SCHWARZALIA Curran, 1934d:431. Type-species, *luteipennis* Curran (orig. des.).
LUTEIPENNIS Curran, 1934d:466, figs. 154, 180. Type-locality: Panama, Canal Zone, Corozal (cf. Arnaud, 1963:127).

Genus SQUAMOMEDINA Townsend

SQUAMOMEDINA Townsend, 1934a:392. Type-species, *squamata* Townsend (orig. des.).
SQUAMATA Townsend, 1934a:392. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus STENODEXIA Wulp

STENODEXIA Wulp, 1891:212. Type-species, *albicincta* Wulp (sub. mon.; Wulp, 1891:246).
ALBICINCTA Wulp, 1891:246, pl. 5, figs. 15-15a. Type-locality: Mexico, sev. places.
FOXII Johnson, 1919:438. Type-locality: Jamaica, Port Antonio.



Genus UROCHAETONA Townsend

UROCHAETONA Townsend, 1919a:562. Type-species, *longipes* Townsend (orig. des.).

LONGIPES Townsend, 1919a:562. Type-locality: Peru, Yahuar Mayo, Río Inambari.

Genus UROHYPOMYIA Townsend

UROHYPOMYIA Townsend, 1939:450. Type-species, *anomala* Townsend (orig. des.).

ANOMALA Townsend, 1939:450. Type-locality: Brazil, Mato Grosso, Maracaju.

Genus XANTHOCHAETONA Townsend

XANTHOCHAETONA Townsend, 1934a:392. Type-species, *similis* Townsend (orig. des.).

SIMILIS Townsend, 1934a:393. Type-locality: Brazil, Pará, Boa Vista, Tapajós River.

Genus XANTHOPHYLLOPHILA Townsend

XANTHOPHYLLOPHILA Townsend, 1927a:234. Type-species, *gracilis* Townsend (orig. des.).

GRACILIS Townsend, 1927a:365. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

UNPLACED SPECIES OF URODESMINI

LONGITARSIS Wulp, 1895:261 (*Morinia*). Type-locality: Mexico, Guerrero.

Tribe EBENIINI

Genus COMYOPS Wulp

COMYOPS Wulp, 1891:213 (key), 262. Type-species, *nigripennis* Wulp (Brauer & Bergenstamm, 1893:95).

NIGRIPENNIS Wulp, 1891:262, pl. 6, figs. 15a-15b. Type-locality: Mexico, Tabasco, Teapa. Distr. — Mexico, Trinidad. Ref. — Thompson, 1963a:472.

STRIATICOLLIS Wulp, 1891:262. Type-locality: Mexico, Guerrero, Venta de Zopilote.

Genus COMYOPSIS Townsend

COMYOPSIS Townsend, 1919b:176. Type-species, *fumata* Townsend (orig. des.).

FUMATA Townsend, 1919b:176. Type-locality: Nicaragua, Chinandega. Distr. — Nicaragua, Trinidad. Ref. — Thompson, 1963a:476.

Genus EBENIA Macquart

EBENIA Macquart, 1846:299 (1846:171). Type-species, *claripennis* Macquart (orig. des.).

CLARIPENNIS Macquart, 1846:171, pl. 16, fig. 2. Type-locality: "Brazil". Ref. — Thompson, 1963a:478.

TRICHOPODA (Wulp), 1891:261 (*Morinia*). Type-locality: Mexico, Veracruz and Tabasco.

Genus MYIOPHASIOPSIS Townsend

- MYIOPHASIOPSIS Townsend, 1927a:378. Type-species, *flavotegulata* Townsend (orig. des.).
FLAVOTEGULATA Townsend, 1927a:382. Type-locality: Peru, Ushpayacu River.

Genus NEOXANTHOBASIS Blanchard

- NEOXANTHOBASIS Blanchard, 1966b:212. Type-species, *nigra* Blanchard (orig. des.).
NIGRA Blanchard, 1966b:212, fig. 13. Type-locality: Argentina, Rio Negro, Bariloche.

Genus PARAXANTHOBASIS Blanchard

- PARAXANTHOBASIS Blanchard, 1966b:215. Type-species, *tibialis* Blanchard (orig. des.).
TIBIALIS Blanchard, 1966b:215, fig. 14. Type-locality: Argentina, Rio Negro, Bariloche.
UNICOLOR (Aldrich), 1934:112 (*Xanthobasis*). Type-locality: Argentina, Rio Negro, Lake Gutiérrez. Ref. — Blanchard, 1966b:218, fig. 15.

Genus PHAEODEMA Aldrich

- PHAEODEMA Aldrich, 1934:145. Type-species, *mystacina* Aldrich (orig. des.).
MYSTACINA Aldrich, 1934:145. Type-locality: Chile, Llanquihue, Puerto Montt.

Genus PHASIOPHYTO Townsend

- PHASIOPHYTO Townsend, 1919a:548. Type-species, *fumifera* Townsend (orig. des.).
FUMIFERA Townsend, 1919a:548. Type-locality: Peru, Charape River.

Genus PROXANTHOBASIS Blanchard

- PROXANTHOBASIS Blanchard, 1966b:219. Type-species, *rufipes* Blanchard (orig. des.).
ALDRICHI Blanchard, 1966b:222, fig. 17. Type-locality: Argentina, Rio Negro, Bariloche.
RUFESCENS Blanchard, 1966b:222, fig. 18. Type-locality: Argentina, Rio Negro, Bariloche.
RUFIPES Blanchard, 1966b:219, fig. 16. Type-locality: Argentina, Rio Negro, Bariloche.

Genus XANTHOBASIS Aldrich

- XANTHOBASIS Aldrich, 1934:110. Type-species, *angustifrons* Aldrich (orig. des.).
ANGUSTIFRONS Aldrich, 1934:111. Type-locality: Argentina, Rio Negro, Lake Gutiérrez.
NEOPOLLINOSA Blanchard, 1966b:226, fig. 19. Type-locality: Argentina, Moreno, Bariloche.
POLLINOSA Aldrich, 1934:112. Type-locality: Argentina, Rio Negro, Bariloche.

UNPLACED SPECIES OF EBENINI

- FUMATA Wulp, 1895:261 (*Morinia*). Type-locality: Mexico, Tabasco.

Tribe SOPHIINI

Genus CORDYLIGASTER Macquart

- CORDYLIGASTER Macquart, 1843:247 (1843:90). Type-species, *Dexia petiolata* Wiedemann (orig. des.).
- Megistogaster* Macquart, 1851:185 (1851:212). Type-species, *fuscipennis* Macquart (Townsend, 1916:7) = *petiolata* (Wiedemann).
- Cordylidexia* Giglio-Tos, 1893:67 (unnecessary nom. nov. for *Cordyligaster* Macquart). Type-species, *Dexia petiolata* Wiedemann (aut.).
- Eucordylidexia* Townsend, 1915i:41. Type-species, *ategulata* Townsend (orig. des.) = *petiolata* (Wiedemann).
- Ref. — Aldrich, 1927d:85 (key).
- ANALIS (Macquart), 1851:187 (1851:214) (*Megistogaster*). Type-locality: Brazil, Amazonas.
- FUSCIFACIES Bigot, 1888a:101. Type-locality: Unknown (presumably South America, cf. Brauer, 1897:365).
- PETIOLATA (Wiedemann), 1830:374 (*Dexia*). Type-locality: "Brazil". Distr. — Central to South America.
- fuscipennis* Macquart, 1851:186 (1851:213), pl. 19, fig. 7 (*Megistogaster*). Type-locality: "South America (as "Java", error).
- ategulata* Townsend, 1915i:41 (*Eucordylidexia*). Type-locality: Guatemala, Puerto Barrios.
- TIPULIFORMIS Walker, 1857:205. Type-locality: "South America".

Genus EUANTHA Wulp

- EUANTHA Wulp, 1885:198. Type-species, *Dexia dives* Wiedemann (mon.) = *liturata* (Olivier).
- Neoeuanta* Townsend, 1931a:336. Type-species, *Dexia aucta* Wiedemann (orig. des.). N. SYN.
- Ref. — Aldrich, 1927c:28 (key).
- AUCTA (Wiedemann), 1830:377 (*Dexia*). Type-locality: "Brazil". Ref. — Aldrich, 1927c:27. N. COMB.
- INTERRUPTA Aldrich, 1927c:28. Type-locality: Costa Rica, San Mateo.
- nigra* Townsend, 1931a:338 (*Sophia*). Type-locality: Panamá, Chiriqui. N. SYN.
- LITTURATA (Olivier), 1812:423 (*Ocyptera*). Type-locality: U.S.A., Carolina. Distr. — U.S.A., Mexico.
- liturata*, emend. or error.
- PULCHRA Wulp, 1891:249. Type-locality: Mexico, Guerrero, Amulango (formely Amula).

Genus EUANTHOIDES Townsend

- EUANTHOIDES Townsend, 1931a:337. Type-species, *petiolata* Townsend (orig. des.).
- PETIOLATA Townsend, 1931a:337. Type-locality: "Mexico".

Genus EUCORDYLIGASTER Townsend

- EUCORDYLIGASTER Townsend, 1917c:123. Type-species, *Cordyligaster septentrionalis* Townsend (orig. des.).
- Cordyligaster*, authors, not Macquart.
- Ref. — Aldrich, 1927c.

- MINUSCULA (Wulp), 1891:252, pl. 6, fig. 7-7a (*Cordyligaster*). Type-locality: Mexico, Guerrero and Teapa.
- NYOMULA (Townsend), 1914:93 (*Cordyligaster*). Type-locality: Peru, Piura Valley, Nomala.

Genus LEPTIDOSOPHIA Townsend

- LEPTIDOSOPHIA Townsend, 1931a:335. Type-species, *lutescens* Townsend (orig. des.).
- FLAVA (Aldrich), 1929:6 (*Euantha*). Type-locality: Peru, Chanchamayo (as Couchamayo).
- LUTESCENS Townsend, 1931a:336. Type-locality: Peru, Cuzco, Pruiipini Valley, Callanga.

Genus SOPHIA Robineau-Desvoidy

- SOPHIA Robineau-Desvoidy, 1830:317. Type-species, *filipes* Robineau-Desvoidy (Townsend, 1916:9).
- DESVOIDYI Townsend, 1931a:338. Type-locality: Brazil, Espírito Santo.
- FILIPES Robineau-Desvoidy, 1830:318. Type-locality: "Brazil". Unrecognized.

Tribe SHANNONIELLINI

Townsend proposed this tribe for some aberrant forms of Tachinidae with undeveloped post-scutellum. He placed this group in the Melanophoridae (= Rhinophorinae). Shannoniellini seems to be a mixture of some forms close to the Minthoini. At present I prefer to maintain this tribe as distinct.

Genus BEZZIMYA Townsend

- BEZZIMYA Townsend, 1919a:591. Type-species, *busckii* Townsend (orig. des.). Ref. — Sabrosky, 1963:3 (key).
- BUSCKII Townsend, 1919a:592. Type-locality: Panama, Trinidad River. Distr. — U.S.A., Panama.

Genus SHANNONIELLA Townsend

- SHANNONIELLA Townsend, 1939b:249. Type-species, *cuspidata* Townsend (orig. des.).
- CUSPIDATA Townsend, 1939b:251. Type-locality: Brazil, Guanabara, Rio de Janeiro.

Genus TRYPETIDOMIMA Townsend

- TRYPETIDOMIMA Townsend, 1935b:68. Type-species, *lutea* Townsend (orig. des.).
- LUTEA Townsend, 1935b:68. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Tribe MINTHOINI
(ERIOTHRIXINI, IN PART)

The recent studies of Verbeke (1962) and Mesnil (1966) indicate that the tribe Eriothrixini *sensu* Townsend is a mixture. Townsend (1936:59, 60) considered in that tribe the genera *Eriothrix* Meigen and *Mintho* Robineau-Desvoidy, but he too questioned whether these two groups belonged in the same tribe (*loc. cit.*, pg. 61).

Verbeke (1966:99) showed that *Eriothrix* Meigen belongs to a group close to *Thelaira* Meigen on the basis of male genitalia characters. He placed *Mintho* Robineau-Desvoidy, which has a different genitalic pattern (*loc. cit.*, pg. 129) in his Echinomyiinae which includes also forms belonging to the tribes Aphriini, Leskiini and Siphonini of the present catalogue.

Mesnil (1966:882) supports Verbeke's views and considers the two genera *Eriothrix* Meigen and *Mintho* Robineau-Desvoidy in distinct subtribes.

Externally these two forms possess certain peculiar characters in common and run in Townsend's keys to Eriothrixini: however, the New World forms of his tribe are closer to the *Mintho* group rather than the *Eriothrix* one, on the basis of genitalic structure.

I have not had the opportunity to study and rearrange thoroughly all the Neotropical components assigned here. I have assigned to this group forms with ovate, usually laterally compressed abdomen, a characteristic head shape, and a very peculiar, usually elongate front leg with swollen and flattened tarsi as shown in *Actinochaeta* Brauer & Bergenstamm.

There is strong evidence that the genera *Shannoniella* Townsend and *Bezzimyia* Townsend placed as Shannoniellini may represent aberrant forms of this complex.

Genus ACTINOCHAETA Brauer & Bergenstamm

- ACTINOCHAETA Brauer & Bergenstamm, 1889:136 (1889:69). Type-species, *columbiae* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.).
- Actinochaetopsis* Townsend, 1934a:393. Type-species, *amazonica* Townsend (orig. des.). N. SYN.
- Paractinochaeta* Townsend, 1934a:393. Type-species, *Thelairodes carlosalbertoi* Lima (orig. des.). N. SYN.
- Tapajomintho* Townsend, 1934a:394. Type-species, *nigriventris* Townsend (orig. des.). N. SYN.
- AMAZONICA (Townsend), 1934a:393 (*Actinochaetopsis*). Type-locality: Brazil, Pará, Boa Vista, Tapajós River. N. COMB.
- CARLOSALBERTOI (Lima), 1926:552 (*Thelairodes*). Type-locality: Brazil, Rio de Janeiro, Guanabara, Tijuca, Alto da Boa Vista. Ref. — Fischer, 1935 (notes). N. COMB.
- COLUMBIAE Brauer & Bergenstamm, 1889:136 (1889:69). Type-locality: "Colombia". Distr. — Costa Rica, Panama, Colombia, Peru. Ref. — Aldrich, 1927e:3 (notes).
- NIGRIVENTRIS (Townsend), 1934a:394 (*Tapajomintho*). Type-locality: Brazil, Pará, Boa Vista, Tapajós River. N. COMB.

Genus ACTINOMINTHELLA Townsend

- ACTINOMINTHELLA Townsend, 1928c:158. Type-species, *atrophopodella* Townsend (orig. des.).
ATROPHOPODELLA Townsend, 1928c:158. Type-locality: Peru, Chosica Canyon.

Genus ATROPHOPALPUS Townsend

- ATROPHOPALPUS Townsend, 1892a:130. Type-species, *angusticornis* Townsend (orig. des.). Ref. — Reinhard, 1934 (rev., in *Ceratomyiella*).
RUFIPES Curran, 1926a:106. Type-locality: Jamaica, Hill Gardens.
RUFOPALPUS Curran, 1926a:105. Type-locality: Jamaica, Hill Gardens.

Genus CATEMOPHRYS Townsend

- Wulpia* Brauer & Bergenstamm, 1893:128, 188 (1893:40, 100) (preocc. Bigot, 1886). Type-species, *aperta* Brauer & Bergenstamm (orig. des., as gen. n. sp. n.) = *sequens* (Townsend).
CATEMOPHRYS Townsend, 1908:65. Type-species, *Vanderwulpia sequens* Townsend (orig. des.).
Brauerimyia Townsend, 1908:65 (nom. nov. for *Wulpia* Brauer & Bergenstamm). Type-species, *Wulpia aperta* Brauer & Bergenstamm (aut.) = *sequens* (Townsend).
SEQUENS (Townsend), 1892a:172 (*Vanderwulpia*). Type-locality: U.S.A., New Mexico. Distr. — U.S.A., Mexico.
aperta Brauer & Bergenstamm, 1893:128, 188 (1893:40, 100) (*Wulpia*). Type-locality: "Mexico".

Genus CERATOMYIELLA Townsend

- CERATOMYIELLA Townsend, 1891b:379. Type-species, *conica* Townsend (orig. des.).
Oedemapeza Townsend, 1908:66. Type-species, *Atrophopoda townsendi* Williston (orig. des.).
Ref. — Reinhard, 1934a (rev.).
BICINCTA Reinhard, 1934a:15. Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico.
TOWNSENDI (Williston), 1896:356, pl. 11, figs. 93a, 93b, 94 (*Atrophopoda*). Type-locality: St. Vincent, W. I.

Genus DIAPHOROPEZA Townsend

- DIAPHOROPEZA Townsend, 1908:66. Type-species, *Atrophopoda braueri* Williston (orig. des.).
BRAUERI (Williston), 1896:357, pl. 11, figs. 93c, 94, 94a, 94b (*Atrophopoda*). Type-locality: St. Vincent, W. I.
MAYENSIS Townsend, 1929:369. Type-locality: Peru, Shanao, Mayo River.
PERUANA Townsend, 1911b:147 (also, 1912a:308). Type-locality: Peru, Piura, Sullana, Río Chira Valley.
PERUVIANA Townsend, 1911b:147. Type-locality: Peru, Sullana.

Genus HYPERTROPHOCERA Townsend

- HYPERTROPHOCERA Townsend, 1891b:360. Type-species, *parvipes* Townsend (orig. des.).

PARVIPES Townsend, 1891b:361. Type-locality: U.S.A., N. Mexico. Distr. — U.S.A., Mexico (Chihuahua).

Genus NEOMETACHAETA Townsend

NEOMETACHAETA Townsend, 1915m:415. Type-species, *polita* Townsend (orig. des.).

POLITA Townsend, 1915m:416. Type-locality: Peru, Chosica Canyon.

Genus PARADIDYMA Brauer & Bergenstamm

PARADIDYMA Brauer & Bergenstamm, 1891:382 (1891:78). Type-species, *Didyma validinervis* Wulp (mon.).

Atrophopoda Townsend, 1891b:373. Type-species, *singularis* Townsend (orig. des.).

Microchira Brauer & Bergenstamm, 1893:128 (1893:40). Type-species, *mexicana* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.).

Lachnomopsis Townsend, 1915m:421. Type-species, *armata* Townsend (orig. des.).

Ref. — Reinhard, 1934:18 (rev.).

AFFINIS Reinhard, 1934a:35. Type-locality: U.S.A., Texas. Distr. — U.S.A., Guatemala.

ALDRICHI Reinhard, 1934a:30. Type-locality: Guatemala, Ingenio R. R. Station. Distr. — Central America.

APICALIS Reinhard, 1934a:33. Type-locality: U.S.A., Texas. Distr. — U.S.A., Guatemala.

ARMATA (Townsend), 1915m:421 (*Lachnomopsis*). Type-locality: "Peru".

BRASILIANA Reinhard, 1934a:38. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

CRASSISETA Reinhard, 1934a:27. Type-locality: Mexico, Chihuahua, Mound Valley.

DERELICTA Reinhard, 1934a:25. Type-locality: Mexico, Chihuahua, Mound Valley.

MERISTA Reinhard, 1953a:95. Type-locality: Mexico, Amanalco.

MEXICANA (Brauer & Bergenstamm), 1893:128 (1893:40) (*Microchira*). Type-locality: "Mexico".

aperta Brauer & Bergenstamm, 1893:127 (1893:39). Type-locality: "Mexico".

ORBITALIS Coquillett, 1904:92. Type-locality: Nicaragua, Chinandega.

PERUANA (Townsend), 1928c:159 (*Atrophopoda*). Type-locality: Peru, Piura, Cacaturo.

PERUVIANA Townsend, 1928c:159. Type-locality: Peru, Chosica.

PILIVENTRIS Reinhard, 1934a:32. Type-locality: Peru, Huariaca.

RECINCTA Reinhard, 1964:19. Type-locality: Mexico, Morelos, Cuernavaca.

SINGULARIS (Townsend), 1891b:374 (*Atrophopoda*). Type-locality: U.S.A., Illinois. Distr. — Canada, U.S.A., Mexico.

VALIDINERVIS (Wulp), 1890:164 (*Didyma*). Type-locality: Mexico, Guerrero.

Genus VANDERWULPIA Townsend

VANDERWULPIA Townsend, 1891b:381. Type-species, *atrophopodoides* Townsend (orig. des.).

ATROPHOPODOIDES Townsend, 1891b:381. Type-locality: U.S.A., Mexico. Distr. — U.S.A., Mexico (Chihuahua).

Tribe LESKIINI

The hosts of Leskiini are various lepidopterous larvae, especially those of concealed habits such as leaf-miners, case-makers, leaf-rollers, etc.

Genus BESKIOLESKIA Townsend

- BESKIOLESKIA Townsend, 1919a:562. Type-species, *busckii* Townsend (orig. des.).
BUSCKII Townsend, 1919a:562. Type-locality: Panama,, Cabima and Taboga Is.

Genus DEJENIOPALPUS Townsend

- DEJENIOPALPUS Townsend, 1916f:312. Type-species, *texensis* Townsend (orig. des.).
BRASILIENSIS Townsend, 1929:368. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Ref. — James, 1947:106.
LONGIPALPIS (Wulp), 1890:138 (*Myobia*). Type-locality: Mexico, Guerrero Chilpancingo. Ref. — Aldrich, 1924a:214.
TENUIROSTRIS James, 1947:105. Type-locality: Mexico, Tapachula.

Genus EUMYOBIA Townsend

- EUMYOBIA Townsend, 1911b:146 (also, 1912a:312). Type-species, *flava* Townsend (orig. des., as gen. n., sp. n.).
EIBENS (Wiedemann), 1830:249 (*Stomoxys*). Type-locality: "Brazil". Ref. — Townsend, 1931b:90.
FLAVA Townsend, 1911b:146, 147 (also, 1912a:312). Type-locality: Peru, Río Chira Valley, Sullana. Ref. — Townsend, 1931b:90.

Genus GALAPAGOSIA Curran

- GALAPAGOSIA Curran, 1934a:171. Type-species, *minuta* Curran (orig. des.).
MINUTA Curran, 1934a:172. Type-species, Galápagos Is., Indefatigable I.

Genus GENEIA Rondani

- GENEIA Rondani, 1850b:172. Type-species, *maculiventris* Rondani (mon.) = *trifaria* (Wiedemann).
Geneopsis Townsend, 1927a:212. Type-species, *major* Townsend (orig. des.).
Geneoglossa Townsend, 1935d:225. Type-species, *glossata* Townsend (orig. des.).
Ref. — James, 1947:111 (key).
GLOSSATA (Townsend), 1935d:225 (*Geneoglossa*). Type-locality: Brazil, Pernambuco, Tapera.
GRACILIS James, 1947:111. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
MAJOR (Townsend), 1927a:311 (*Geneopsis*). Type-locality: Brazil, São Paulo, Itaquaquecetuba, and Paraguay.
PELLUCENS (Curran), 1925a:261 (*Leskia*). Type-locality: Honduras, Corocito.

TRIFARIA (Wiedemann), 1824:41 (*Stomoxys*). Type-locality: "South America". Distr. — Neotropical. Ref. — James, 1947:114.
maculiventris Rondani, 1850b:173, pl. 4, figs. 4-5. Type-locality: "Venezuela". Ref. — Aldrich, 1924b:210.

Genus GENEODES Townsend

GENEODES Townsend, 1934:394. Type-species, *griseescens* Townsend (orig. des.).
GRISESCENS Townsend, 1934:394. Type-locality: Brazil, Pará, Urucurituba, Tapajós River.

Genus JAYNESLESKIA Townsend

JAYNESLESKIA Townsend, 1934:395. Type-species, *Leskiomima jaynesi* Aldrich (orig. des.).
JAYNESI (Aldrich), 1924:17 (*Leskiomima*). Type-locality: Argentina, Tucumán. Distr. — Argentina, Brazil.

Genus LESKIOMIMA Brauer & Bergenstamm

LESKIOMIMA Brauer & Bergenstamm, 1891:372, 406 (1891:68, 102) (as *Leskiomima*, p. 372 (68), in part). Type-species, *Stomoxys tenera* Wiedemann (mon.).
Ref. — James, 1947:100 (key).
AUSTRALIS Townsend, 1929:368. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus METAMYOBIA Townsend

METAMYOBIA Townsend, 1927a:213. Type-species, *filipalpis* Townsend (orig. des.).
FILIPALPIS Townsend, 1927a:328. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus MICROLESKIA Thompson

MICROLESKIA Thompson, 1968:143. Type-species, *longipes* Thompson (orig. des.).
LONGIPES Thompson, 1968:143. Type-locality: Trinidad, "Simla".

Genus MICROLESKIA Thompson

MINTHOLESKIA Townsend, 1934:395. Type-species, *melanopyga* Townsend (orig. des.).
MELANOPYGA Townsend, 1934:395. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus MYOBIOPSIS Townsend

MYOBIOPSIS Townsend, 1916e:628. Type-species, *similis* Townsend (orig. des.).
Leskiopalpus Townsend, 1916e:629. Type-species, *calidus* Townsend (orig. des.) = *depilis* Coquillett.
Ref. — Curran, 1934c:507 (key, as *Myobia*).

- ANGUSTA (Walker), 1852:314 (*Dexia*). Type-locality: "Brazil". Unrecognized.
- AURIFRONS (Macquart), 1846:297 (1846:169), pl. 15, fig. 8 (*Myobia*). Type-locality: "Venezuela".
- DIADEMA (Wiedemann), 1830:382 (*Dexia*). Type-locality: "Brazil". Ref. — Thompson, 1963a:347 (larva).
- FAMELICA (Wiedemann), 1830:250 (*Stomoxys*). Type-locality: "Brazil". Ref. — Aldrich, 1929a:12.
- FLAVIPENNIS (Wiedemann), 1830:380 (*Dexia*). Type-locality: "Brazil".
- FLAVESCENS (Townsend), 1929:368 (*Leskiopalpus*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- SANCTAECRUCIS (Thompson), 1963a:348, fig. 21 (*Leskiopalpus*). Type-locality: Trinidad, Santa Cruz Valley. N. COME.
- SIPHONINA (Villeneuve), 1937:209 (*Myobia*). Type-locality: Guyana, Essequibo.
- PENALTIS (Curran), 1934c:508 (*Myobia*). Type-locality: Guyana, Bartica, Kartabo.
- PERPECTA (Walker), 1860:307 (*Dexia*). Type-locality: "Mexico".
- PERTINAX (Curran), 1934c:507 (*Myobia*). Type-locality: Brazil, Roraima, Vista Alegre (as "British Honduras", error).
- VERNA (Curran), 1934c:509 (*Myobia*). Type-locality: Guyana, Bartica, Kartabo.

Genus PARTHENOLESKIA Townsend

- PARTHENOLESKIA Townsend, 1941b:339. Type-species, *parkeri* Townsend (orig. des.).
- PARKERI Townsend, 1941b:340. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Ref. — Parker, 1953:53 (immature stages).

Genus PROLESKIOMIMA Townsend

- PROLESKIOMIMA Townsend, 1934:395. Type-species, *frontalis* Townsend (orig. des.).
- FRONTALIS Townsend, 1934:396. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Ref. — James, 1947:95.

Genus SIPHOLESKIA Townsend

- SIPHOLESKIA Townsend, 1916c:628. Type-species, *Drepanoglossa occidentalis* Coquillett (orig. des.).
- LORIOLA Reinhard, 1955c:126. Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico.
- OCCIDENTALIS (Coquillett), 1895a:126 (*Drepanoglossa*). Type-locality: U.S.A., California. Distr. — U.S.A., Mexico (Chihuahua).

Genus SPATHIPALPUS Rondani

- SPATHIPALPUS Rondani, 1863:20. Type-species, *philippii* Rondani (Townsend, 1916c:9). Ref. — Cortés, 1951a:62.
- PHILIPPI Rondani, 1863:21. Type-locality: Chile, Valdivia. Distr. — Chile (Angol), Argentina (Rio Negro, Lake Nahuel Huapi).
- ? *flavifrons* Rondani, 1863:21. Type-locality: Chile, Valdivia.

Genus STOMATODEXIA Brauer & Bergenstamm

- STOMATODEXIA Brauer & Bergenstamm, 1889:125 (1889:57). Type-species, *Stomoxys cothurnata* Wiedemann (mon.).
Eustomatodexia Townsend, 1892c:167. *Nomen nudum*.
COTHURNATA (Wiedemann), 1830:249 (*Stomoxys*). Type-locality: "Brazil". Ref. — Aldrich, 1929:9.
insulensis (Townsend), 1892c:167 (*Eustomatodexia*). *Nomen nudum*.
LONGITARSIS (Macquart), 1843:249 (1843:92), pl. 11, fig. 2 (*Prosenia*). Type-locality: Colombia, Bogotá, Santa Fé. Unrecognized.
MACULIFERA (Bigot), 1889:264 (*Prosenia*). Type-locality: "Mexico". Unrecognized.
OBSCURA (Walker), 1852:307 (*Dexia*). Type-locality: "Brazil". Unrecognized.
QUADRIMACULATA (Walker), 1852:319 (*Dexia*). Type-locality: "Brazil". Unrecognized.
SIMILIGENA Wulp, 1891:239. Type-locality: Mexico, Guerrero, Amulango (formely Amula).
TINCTISQUAMAE Curran, 1926a:104. Type-locality: Jamaica, Cinchona.

Genus TAPAJOLESKIA Townsend

- TAPAJOLESKIA Townsend, 1934:396. Type-species, *taurea* Townsend (orig. des.).
TAUREA Townsend, 1934:397. Type-locality: Brazil, Pará, Urucurituba, Tapajós River.

Genus TIPULOLESKIA Townsend

- TIPULOLESKIA Townsend, 1931a:331. Type-species, *mima* Townsend (orig. des.).
MIMA Townsend, 1931a:332. Type-locality: "Brazil".

Genus TRICHOPYRRHOSIA Townsend

- TRICHOPYRRHOSIA Townsend, 1927a:243. Type-species, *uruhuasi* Townsend (orig. des.).
URUHUASI Townsend, 1927a:362. Type-locality: Peru, Uruhuasi.

Genus TROCHIOGLOSSA Townsend

- TROCHIOGLOSSA Townsend, 1919a:561. Type-species, *tropica* Townsend (orig. des.).
AUREA Thompson, 1963a:350. Type-locality: Trinidad, St. Augustine.
TROPICA Townsend, 1919a:561. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).

Genus TROCHIOLESKIA Townsend

- TROCHIOLESKIA Townsend, 1917b:226. Type-species, *flava* Townsend (orig. des.).
FLAVA Townsend, 1917b:227. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).

Genus URULESKIA Townsend

- URULESKIA Townsend, 1934a:397. Type-species, *aurescens* Townsend (orig. des.).
AURESCENS Townsend, 1934a:397. Type-locality: Brazil, Pará, Urucurituba, Tapajós River.

Genus URUMYOBIA Townsend

- URUMYOBIA Townsend, 1934a:396. Type-species, *aurata* Townsend (orig. des.).
AURATA Townsend, 1934a:396. Type-locality: Brazil, Pará, Urucurituba, Tapajós River.

UNRECOGNIZED SPECIES OF LESKIINI

- ANGULATA Wulp, 1890:136 (*Myobia*). Type-locality: Mexico, Guerrero, Chilpancingo.
LEPIDA Wulp, 1890:135 (*Myobia*). Type-locality: Mexico, Guerrero.
LONGICORNIS Macquart, 1843:222 (1843:65), pl. 7, fig. 3 (*Myobia*). Type-locality: "Brazil or Chile".
OPIMA Wulp, 1890:136 (*Myobia*). Type-locality: Mexico, sev. places.
PLANA Walker, 1852:315 (*Dexia*). Type-locality: "Brazil".
RUFOPYGATA Bigot, 1889:262 (*Viviania*). Type-locality: "Mexico".

Tribe TELOTHYRIINI

Some species of Telothyriini have been reared from lepidopterous larvae.

Genus COMATACTA Coquillett

- COMATACTA Coquillett, 1902a:199. Type-species, *Brachycoma pallidula* Wulp (orig. des.) = *variegata* (Fabricius).
INSULARIS Curran, 1927c:12. Type-locality: Puerto Rico, San Juan.
MICROPALPUS (Curran), 1925e:9 (*Ptilomya*). Type-locality: Bolivia, Paraguay River, Piedra Blanca (4 miles west of Corumbá, Mato Grosso, Brazil). Ref. — Townsend, 1933:527.
NAUTLANA Townsend, 1908:101. Type-locality: Mexico, Veracruz, San Rafael Yucatán, Temax.
TRICINCTA (Fabricius), 1805:301 (*Musca*). Type-locality: "South America".
VARIEGATA (Fabricius), 1805:281 (*Stomoxys*). Type-locality: "South America".
pallidula Wulp, 1890:95 (*Brachycoma*). Type-locality: Mexico, North

Yucatán, Temax.

Genus EUPTILOMYIA Townsend

- EUPTILOMYIA Townsend, 1939c:451. Type-species, *frontalis* Townsend (orig. des.).
FRONTALIS Townsend, 1939c:451. Type-locality: Brazil, São Paulo, Itaquecetuba.

Genus EUTELOTHYRIA Townsend

- EUTELOTHYRIA Townsend, 1931a:332. Type-species, *itaquecetubae* Townsend (orig. des.).

- ITAQUAQUECETUBAE Townsend, 1931a:333. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
TRINITATIS Thompson, 1963a:484, figs. 145, 146, 149. Type-locality: Trinidad, Brazil village.

Genus FLORADALIA Thompson

- FLORADALIA Thompson, 1963a:486. Type-species, *major* Thompson (orig. des.).
MAJOR Thompson, 1963a:486, figs. 142, 147. Type-locality: Trinidad, Maracas Valley.
MINOR Thompson, 1963a:488, figs. 138, 141. Type-locality: Trinidad, St. Augustine.

Genus LESKIOPSIS Townsend

- LESKIOFISIS Townsend, 1916c:627. Type-species, *Myobia thecata* Coquillett (orig. des.).
BRASILIENSIS Townsend, 1929:369. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus PTILOMYIOPSIS Townsend

- Ptilomyia* Curran, 1925e:8 (preocc. Coquillett, 1910). Type-species, *plumata* Curran (orig. des.).
PTILOMYIOPSIS Townsend, 1933:527 (nom. nov. for *Ptilomyia* Curran). Type-species, *Ptilomyia plumata* Curran (aut.).
PLUMATA (Curran), 1925e:8 (*Ptilomyia*). Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).

PTILOMYOIDES Curran

- PTILOMYOIDES Curran, 1928c:112. Type-species, *Ptilomyia bequaerti* Curran (mon.).
Ptilomyioides Curran, 1934:421, error or emendation.
BEQUAERTI (Curran), 1925d:352 (*Ptilomyia*). Type-locality: Brazil, Roraima, São Alberto.

Genus TELOTHYRIA Wulp

- TELOTHYRIA Wulp, 1890:167. Type-species, *cupreiventris* Wulp (Brauer & Bergenstamm, 1893:44) (1893:132).
Thereuops Brauer & Bergenstamm, 1891:378 (1891:74). Type-species, *Miltogramma brevipennis* Schiner (Brauer & Bergenstamm, 1893:132; 1894:44).
BREVIPENNIS (Schiner), 1868:324 (*Miltogramma*). Type-locality: "Brazil".
Ref. — Aldrich, 1929:7 (type redescription).
CUPREIVENTRIS Wulp, 1890:182, pl. 4, fig. 14-14a. Type-locality: Mexico, Veracruz and Tabasco.
ILLUCENS Wulp, 1890:183. Type-locality: Mexico, Tabasco.
PLACIDA Wulp, 1890:182. Type-locality: Mexico, Tabasco.
RELICTA Wulp, 1890:171. Type-locality: Mexico, Veracruz.
RUFOSTRIATA Wulp, 1890:172. Type-locality: Mexico, Veracruz and Tabasco.

UNRECOGNIZED SPECIES OF DEXIINAE

- BASALIS Walker, 1852:311 (*Dexia*). Type-locality: "South America".
 CARBONARIA Giglio-Tos, 1893:4 (*Chaetogena*). Type-locality: "Mexico".
 CHRYSOCEPHALA Bigot, 1889:261 (*Ceromasia*). Type-locality: "Mexico".
 CINCTA Giglio-Tos, 1893:4 (*Chaetona*). Type-locality: "Mexico".
 CONVEXA Walker, 1852:312 (*Dexia*). Type-locality: "Brazil".
 FUSCANIPENNIS Macquart, 1846:316 (1846:188), pl. 20, fig. 11 (*Dexia*). Type-locality: Mexico, Yucatán, Merida.
 GENUINA Wulp, 1891:243 (*Dexia*). Type-locality: Mexico, Guerrero, Omilteme.
 GORTYS Walker, 1849:839 (*Dexia*). Type-locality: "Brazil".
 GRACILIS Giglio-Tos, 1893:4 (*Chaetogena*). Type-locality: "Mexico".
 PARVICORNIS Wulp, 1883:33, pl. 2, fig. 3 (*Dexia*). Type-locality: "Argentina".
 QUADRIVITTATA Bigot, 1889:261 (*Ceromasia*). Type-locality: "Mexico".
 SUAVIS Wulp, 1883:33 (*Dexia*). Type-locality: "Guadeloupe".
 TENUICORNIS Wulp, 1883:32, pl. 2, fig. 5, 6 (*Dexia*). Type-locality: "Argentina".

Subfamily GONIINAE

(Exoristinae)

Tribe OPHIRIONINI

(Anacamptomyiini, in part)

This tribe includes some Neotropical genera placed by Townsend (1936:78) in his tribe Anacamptomyiini. The tribal name is based on the group-unit Ophirioniae (Townsend, 1913a:53). Species of this group have been recorded as parasites of social wasps of the genus *Polybia*. This group merits tribal status both morphologically and biologically and lies near Blondeliini.

Genus OPHIRION Townsend

- OPHIRION Townsend, 1911b:134 (also, 1912a:310). Type-species, *mirabile* Townsend (orig. des., as gen. n., sp. n.).
 MIRABILE Townsend, 1911b:134 (also, 1912a:311). Type-locality: Peru, Piura.

Genus OPHIRIONOPSIS Townsend

- OPHIRIONOPSIS Townsend, 1927a:217. Type-species, *brasiliensis* Townsend (orig. des.).
 BRASILIENSIS Townsend, 1927a:340. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
 POLYBIA (Curran), 1937:1 (*Telothyriosoma*). Type-locality: Trinidad, Marine Club. N. comb.

Genus POLYBIOPHILA Curran

- POLYBIOPHILA Curran, 1937:2. Type-species, *fitzgeraldi* Curran (orig. des.).
 FITZGERALDI Curran, 1937:3. Type-locality: Trinidad, St. Benedict Mt.

Tribe ACEMYINI

The species of the tribe Acemyini parasitize members of the order Orthoptera.

Genus BACULOCAPTUS Cortés

BACULOCAPTUS Cortés, 1968a:106. Type-species, *valparadisi* Cortés (orig. des.).

VALPARADISI Cortés, 1968:108a. Type-locality: Chile, Viña del Mar (Valparaiso).

Genus CERACIA Rondani

CERACIA Rondani, 1865:221 (1865:49). Type-species, *mucronifera* Rondani (mon.).

Myothyria Wulp, 1890:208. Type-species, *majorina* Wulp (Brauer & Bergenstamm, 1891:54 (1891:358)).

Acemyiopsis Townsend, 1915m:433. Type-species, *punensis* Townsend (orig. des.).

Clythopsis Townsend, 1927a:276. Type-species, *confundens* Townsend (orig. des.) = *brachyptera* (Thomson).

Ref. — Aldrich, 1934:136.

BRACHYPTERA (Thomson), 1869:527 (*Myobia*). Type-locality: Brazil, Guanabara, Rio de Janeiro. Ref. — Townsend, 1931b:89 (notes on type).
confundens Townsend, 1927a:298 (*Clythopsis*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.

DENTATA (Coquillett), 1895b:311 (*Acemyia*). Type-locality: U.S.A., California. Distr. — Canada, U.S.A., Mexico, ? Chile.

MAJORINA (Wulp), 1891:209, pl. 4, fig. 19 (*Myothyria*). Type-locality: Mexico, Guerrero and Tabasco. Ref. — Townsend, 1931b:89 (notes on type).

maldonadoi Blanchard, in Lloyd, 1951:217. *Nomen nudum*.

PUNENSIS (Townsend), 1915m:433 (*Acemyiopsis*). Type-locality: Peru, Puno.

SUBANDINA Blanchard, 1943c:19, fig. 20. Type-locality: Argentina, Rio Negro, Comallo, Chile (Angol, Concepción, Santiago).

Tribe BLONDELIINI

(Compsilurini, Trypherini in part)

The food habits of the tribe Blondeliini are varied, including larvae of Lepidoptera, Coleoptera and Hymenoptera (Tenthredinidae).

Genus ACEMEIGENIA Townsend

ACEMEIGENIA Townsend, 1927a:241. Type-species, *inca* Townsend (orig. des.).

INCA Townsend, 1927a:282. Type-locality: Peru, Cuzco.

Genus ACTINODORIA Townsend

ACTINODORIA Townsend, 1927a:272. Type-species, *cuprea* Townsend (orig. des.).

ARGENTEA Thompson, 1964:126, figs. 18-20. Type-locality: Trinidad, Melajo River.

ARGENTIFRONS (Wulp), 1890:183 (*Telothyria*). Type-locality: Mexico, Guerrero. N. COMB.

CUPREA Townsend, 1927a:283. Type-locality: Panama, Taboga I. Distr. — Panama, Trinidad. Ref. — Thompson, 1964:98.

Genus ACTINOTACHINA Townsend

ACTINOTACHINA Townsend, 1927a:278. Type-species, *angusta* Townsend (orig. des.).

Lydellactia Townsend, 1927a:279. Type-species, *clausa* Townsend (orig. des.). N. SYN.

Ptilolydella Townsend, 1927a:278. Type-species, *aristalis* Townsend (orig. des.). N. SYN.

Biohypostena Townsend, 1927a:280. Type-species, *brasiliانا* Townsend (orig. des.) N. SYN.

Actinophryno Townsend, 1928b:148. Type-species, *angusta* Townsend (orig. des.) = *townsendi* Guimarães. N. SYN.

ABERRANS (Townsend), 1929:372 (*Ptilolydella*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. N. COMB.

ANGUSTA Townsend, 1927a:284. Type-locality: Peru, Yahuar Mayo.

APISTALIS (Townsend), 1927a:354 (*Ptilolydella*). Type-locality: Brazil, São Paulo, São Paulo, Cantareira. N. COMB.

BRASILIANA (Townsend), 1927a:290 (*Biohypostena*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. N. COMB.

CLAUSA (Townsend), 1927a:176, (*Lydellactia*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. N. COMB.

TENUIS Blanchard, 1959:176, fig. 13. Type-locality: Argentina, Buenos Aires, José C. Paz.

TOWNSENDI Guimarães, nom. nov. for *angusta* Townsend. Type-locality: Paraguay, Santa Trinidad.

angusta Townsend, 1928b:148 (*Actinophryno*; preocc. Townsend, 1928). Type-locality: Paraguay, Santa Trinidad. N. COMB.

Genus ADORYPHOROPHAGA Townsend

ADORYPHOROPHAGA Townsend, 1931:469. Type-species, *Doryphorophaga aberrans* Townsend (orig. des.).

OCHRIFRONS (Wulp), 1890:180 (*Telothyria*). Type-locality: Mexico, Guerrero. N. COMB.

OVATA (Wulp), 1890:182 (*Telothyria*). Type-locality: Mexico, Guerrero. N. COMB.

Genus ANAMETOPOCHAETA Townsend

ANAMETOPOCHAETA Townsend, 1919a:571. Type-species, *olindoides* Townsend (orig. des.).

OLINDOIDES Townsend, 1919a:571. Type-locality: Peru, Chaquimayo.

Genus ANAPHORINIA Townsend

ANAPHORINIA Townsend, 1927a:266. Type-species, *ourata* Townsend (orig. des.).

AURATA Townsend, 1927a:285. Type-locality: Brazil, São Paulo, Itaquaquecetuba.



Genus ANISIA Wulp

- ANISIA Wulp, 1890:186. Type-species, *inflexa* Wulp (Coquillett, 1910:507).
 ABERRANS Wulp, 1890:198. Type-locality: Mexico, Guerrero (sev. places).
 ACCEDENS Wulp, 1890:196. Type-locality: Mexico, Guerrero.
 AEGROTA Wulp, 1890:192. Type-locality: Mexico, Guerrero, Sierra de las Aguas Escondidas.
 CANDICANS Wulp, 1890:194. Type-locality: Mexico, Morelos, Cuernavaca.
 CILIATA Wulp, 1890:203. Type-locality: Mexico, Morelos, Cuernavaca.
 CINERARIA Wulp, 1890:189. Type-locality: Mexico, Guerrero, Chilpancingo.
 CINEREA Brêthes, 1909:100. Type-locality: Argentina, Mendoza, Punta del Inca. Unrecognized.
 CONGERENS Wulp, 1890:194. Type-locality: Mexico, Guerrero (sev. places).
 CONSPERSA Wulp, 1890:199. Type-locality: Mexico, Guerrero, Omilteme.
 FATUA Wulp, 1890:197. Type-locality: Mexico, Guerrero, Venta de Zopilote.
 FULVIPENNIS Wulp, 1890:190, pl. 4, figs. 15, 15a. Type-locality: Mexico, Guerrero and Tabasco.
 GAGATINA Wulp, 1890:201. Type-locality: Mexico, Guerrero, Omilteme.
 INEPTA Wulp, 1890:195. Type-locality: Mexico, Guerrero (sev. places).
 INFLEXA Wulp, 1890:188. Type-locality: Mexico, Guerrero, Xucumanatlán.
 INTRUSA Wulp, 1890:193. Type-locality: Mexico, Guerrero.
 MACROPTERA Wulp, 1890:198. Type-locality: Mexico, Mexico City.
 MORIONELLA Wulp, 1890:195. Type-locality: Mexico, Guerrero.
 MUCOREA Wulp, 1890:199. Type-locality: Mexico, Guerrero and Morelos.
 NEGLECTA Wulp, 1890:191. Type-locality: Mexico, Guerrero, Sierra de las Aguas Escondidas.
 NIGELLA Wulp, 1890:193. Type-locality: Mexico, Tabasco, Teapa.
 NIGRITHORAX Wulp, 1890:197. Type-locality: Mexico, Veracruz and Tabasco.
 NIGROCINCTA Wulp, 1890:201. Type-locality: Mexico, Guerrero (sev. places).
 OBSCURIFRONS Wulp, 1890:197. Type-locality: Mexico, Tabasco.
 OPACA Wulp, 1890:200. Type-locality: Mexico, Guerrero.
 OPHTHALMICA Wulp, 1890:203. Type-locality: Mexico, Guerrero.
 PALLIDIPALPIS Wulp, 1890:190. Type-locality: Mexico, Tabasco.
 PALPOSA Wulp, 1890:202. Type-locality: Mexico, Veracruz.
 PEREGRINA Wulp, 1890:196. Type-locality: Mexico, Guerrero, Xucumanatlán.
 PULICARIA Wulp, 1890:191. Type-locality: Mexico, Orizaba.
 PULLATA Wulp, 1890:195. Type-locality: Mexico, Guerrero.
 REMISSA Wulp, 1890:201. Type-locality: Mexico, Guerrero.
 RUBRIPES Wulp, 1890:189. Type-locality: Mexico, Guerrero, Veracruz.
 RUFICOXA Wulp, 1890:190. Type-locality: Mexico, Tabasco, Teapa.
 SIGNATA Wulp, 1890:193. Type-locality: Mexico, Veracruz, Atoyac.
 SIMILIS Wulp, 1890:203. Type-locality: Mexico, Guerrero.
 STOLIDA Wulp, 1890:192. Type-locality: Mexico, Guerrero.
 IMBRINA Wulp, 1890:200. Type-locality: Mexico, Tabasco.
 VANDERWULPI Townsend, 1892d:81. Type-locality: Jamaica, Portland.

Genus ANOXYNOPS Townsend

- ANOXYNOPS Townsend, 1927a:274. Type-species, *conicus* Townsend (orig. des.).
 AURIFRONS Thompson, 1968:57, figs. 29-32. Type-locality: Trinidad, Santa Cruz Valley.
 CONICUS Townsend, 1927a:286. Type-locality: Brazil, São Paulo, Itaquaquecetuba.



FLAVOCALYPTRATUS (Wulp), 1890:147, pl. 4, fig. 8 (*Hypostena*). Type-locality: Costa Rica, Volcán de Irazú. N. comb.

Genus ANOXYNOPSELLA Townsend

ANOXYNOPSELLA Townsend, 1935d:226. Type-species, *argentescens* Townsend (orig. des.).

ARGENTESCENS Townsend, 1935d:226. Type-ocality: Brazil, Pernambuco, Tapera.

Genus ANOXYNOPSOIDEA Thompson

ANOXYNOPSOIDEA Thompson, 1968:61. Type-species, *grisea* Thomson (orig. des.).

GRISEA Thompson, 1968:61, figs. 33, 34. Type-locality: Trinidad, I.C.T.A.

Genus APLOMYIOPSIS Villeneuve

APLOMYIOPSIS Villeneuve, 1933:125. Type-species, *galerucellae* Villeneuve (mon.).

Synaplomyia Villeneuve, 1934:181 (unjustified nom. nov. for *Aplomyiopsis* Villeneuve). Type-species, *Aplomyiopsis galerucellae* Villeneuve (aut.).

EPIACHNAE (Aldrich), 1923:95 (*Paradexodes*). Type-locality: Mexico, D.F., Coapa. Ref. — Landis & Howard, 1940.

Genus ARGYRODORIA Townsend

ARGYRODORIA Townsend, 1927a:265. Type-species, *hemiargyroides* Townsend (orig. des.).

Hypophorinia Townsend, 1927a:279. Type-species, *hyphena* Townsend (orig. des.). N. SYN.

HEMIARGYROIDES Townsend, 1927a:288. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

HYPHENA (Townsend), 1927a:318 (*Hypophorinia*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. N. comb.

REFUGA (Wulp), 1890:185 (*Telothyria*). Type-locality: Mexico, Guerrero. N. comb.

Genus BLONDELIA Robineau-Desvoidy

BLONDELIA Robineau-Desvoidy, 1830:122. Type-species, *Tachina nigripes* Fallén (Coquillett, 1910:515; suspension of the rules required, cf. Sabrosky & Arnaud, 1965:1039).

Anetia, authors, not Robineau-Desvoidy.

Lydella, authors, not Robineau-Desvoidy.

ALBOPILOSA (Curran), 1926a:112 (*Masicera*). Type-locality: Jamaica, Cinchona.

FLAVIVENTRIS (Macquart), 1843:218 (1843:61) (*Lydella*). Type-locality: "Guiana".

FRUGALE (Curran), 1934c:511 (*Lydella*). Type-locality: Guyana, Bartica, Kartabo.

INCOMPLETA (Curran), 1928a:43 (*Lydella*). Type-locality: Jamaica, Hope Gardens.

- POLITA (Townsend), 1919a:572 (*Phrynolydella*). Type-locality: U.S.A., Arizona. Distr. — U.S.A., Mexico (Chihuahua).
 PRUDENS (Curran), 1934c:511 (*Lydella*). Type-locality: Guyana, Bartica, Kartabo.
 PULCHELLA (Curran), 1934c:511 (*Lydella*). Type-locality: Guyana, Bartica, Kartabo.
 SODALIS (Wulp), 1890:106 (*Masicera*). Type-locality: Mexico, Guerrero, Tabasco.
 TANTILLA (Wulp), 1890:106 (*Masicera*). Type-locality: Mexico, Tabasco, Teapa. Unrecognized.
 VERTICALE (Curran), 1934c:511 (*Lydella*). Type-locality: Guyana, Bartica, Kartabo.

Genus BOLODORIA Townsend

- BOLODORIA Townsend, 1927a:262. Type-species, *yahuarmayana* Townsend (orig. des.).
 YAHUARMAYANA Townsend, 1927a:290. Type-locality: Peru, Yahuarmayo.

Genus BONNANIOPS Townsend

- BONNANIOPS Townsend, 1935d:231. Type-species, *aberrans* Townsend (orig. des.).
 ABERRANS Townsend, 1935d:231. Type-locality: "Trinidad".

Genus CAENISOMA Townsend

- CAENISOMA Townsend, 1927a:269. Type-species, *charapense* Townsend (orig. des.).
 CHARAPENSE Townsend, 1927a:292. Type-locality: Peru, Charape River.

Genus CAENISOMOPSIS Townsend

- CAENISOMOPSIS Townsend, 1934a:398. Type-species, *paraensis* Townsend (orig. des.).
 PARAENSIS Townsend, 1934a:398. Type-locality: Brazil, Pará, Boa Vista, Rio Tapajós.

Genus CALOLYDELLA Townsend

- CALOLYDELLA Townsend, 1927a:278. Type-species, *geminata* Townsend (orig. des.).
 BLANDITA (Wulp), 1890:142, pl. 4, fig. 4-4a (*Hypostena*). Type-locality: Mexico (sev. places), and Costa Rica. N. comb.
 CONCINNA (Wulp), 1890:142 (*Hypostena*). Type-locality: Mexico, Guerrero. N. comb.
 GEMINATA Townsend, 1927a:293. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus CALPODOMYIA Townsend

- CALPODOMYIA Townsend, 1915m:414. Type-species, *linearis* Townsend (orig. des.).
 LINEARIS Townsend, 1915m:415. Type-locality: Puerto Rico, Mayaguez.



Genus CANALIA Curran

CANALIA Curran, 1934d:431. Type-species, *fasciata* Curran (orig. dcs.).
FASCIATA Curran, 1934d:465, figs. 169, 191. Type-locality: Panama.

Genus CARMODYMYIA Thompson

CARMODYMYIA Thompson, 1968:129. Type-species, *ancylostomiae* Thompson (orig. des.).
ANCYLOSTOMIAE Thompson, 1968:130, figs. 80-84. Type-locality: Trinidad, Paradise Mt.

Genus CATAPHORINIA Townsend

CATAPHORINIA Townsend, 1927a:280. Type-species, *angusta* Townsend (orig. des.).
ANGUSTA Townsend, 1927a:294. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus BRAZILIOMYIA Thompson

BRAZILIOMYIA Thompson, 1968:98. Type-species, *obscura* Thompson (orig. des.).
OBSCURA Thompson, 1968:98, figs. 54, 55. Type-locality: Trinidad, Forest, 1 mile west of Arima.

Genus CATAPHORINIOPSIS Thompson

CATAPHORINIOPSIS Thompson, 1968:133. Type-species, *fumipennis* Thompson (orig. dcs.).
FUMIPENNIS Thompson, 1968:134, figs. 85b, 86, 88. Type-locality: Trinidad, "Simla".
GALBAE Thompson, 1968:137, figs. 85a-87. Type-locality: Trinidad, St. Augustine.

Genus CELATORIA Coquillett

CELATORIA Coquillett, 1890:235. Type-species, *crawii* Coquillett (orig. dcs.) = *diabroticae* (Shimer).
BOSQI Blanchard, 1937:47, figs. 6a-d. Type-locality: Argentina, Buenos Aires, Baradero.
BRASILIENSIS Townsend, 1929:375. Type-locality: Brazil, Minas Gerais, Viçosa.
COMPRESSA (Wulp), 1890:153, pl. 4, fig. 9 (*Degeeria*). Type-locality: Mexico (Guerrero, Veracruz).
DIABROTICAE (Shimer), 1871:219 (*Tachina*). Type-locality: U.S.A., Illinois. Distr. — U.S.A., Mexico.
MARACASI Thompson, 1968:172, fig. 109. Type-locality: Trinidad, Maracas Valley.
NIGRICANS (Wulp), 1890:153 (*Degeeria*). Type-locality: Mexico, Guerrero, Chilpancingo.



Genus CELATORIOPSIS Blanchard

- CELATORIOPSIS Blanchard, 1963:223. Type-species, *eucelatorioides* Blanchard (orig. des.).
EUCELATORIODES Blanchard, 1963:228, fig. 33. Type-locality: Argentina, Misiones, Cerro Azul.

Genus CHAETODORIA Townsend

- CHAETODORIA Townsend, 1927a:259. Type-species, *conica* Townsclad (orig. des.).
CONICA Townsend, 1927a:296. Type-locality: Peru, Charape River. Ref. — Townsend, 1934:405 (generically valid from *Conactia* Townsend).

Genus CHAETOLIXOPHAGA Blanchard

- CHAETOLIXOPHAGA Blanchard, 1940:241. Type-species, *laspeyresiae* Blanchard (orig. des.).
LASPEYRESIAE Blanchard, 1940:241, figs. 5a-f. Type-locality: Argentina, La Plata.

Genus CHAETONODEXODES Townsend

- CHAETONODEXODES Townsend, 1916g:321. Type-species, *rafaeli* Townsend (orig. des.). Ref. — Aldrich, 1931a:205 (notes).
MARSHALLI Aldrich, 1931a:207. Type-locality: "Trinidad".
RAFAELI Townsend, 1916g:322. Type-locality: Mexico, Veracruz, San Rafael.

Genus CHAETOXYNOPS Townsend

- CHAETOXYNOPS Townsend, 1928b:148. Type-species, *chaetosus* Townsend (orig. dcs.).
CHAETOSUS Townsend, 1928b:149. Type-locality: Paraguay, Sta. Trinidad.
paraguayana Townsend, 1928b:150 (*Anaphorinia*). Type-locality: Paraguay, Sta. Trinidad. N. SYN.

Genus CHAQUIMAYOIA Townsend

- CHAQUIMAYOIA Townsend, 1927a:235. Type-species, *plumosula* Townsend (orig. des.).
PLUMOSULA Townsend, 1927a:297. Type-locality: Peru, Chaquimayo.

Genus CHRYSOPROSPHERYSA Townsend

- CHRYSOPROSPHERYSA Townsend, 1928b:147. Type-species, *croesus* Townsend (orig. des.).
CROESUS Townsend, 1928b:148. Type-locality: Peru, Ushpayacu River.

Genus CLYTHOXYNOPS Townsend

- CLYTHOXYNOPS Townsend, 1927a:272 (as *Clithoxynops*, misprint, changed in Errata). Type-species, *orbitalis* Townsend (orig. des.).
ORBITALIS Townsend, 1927a:299. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus CONACTIA Townsend

- CONACTIA Townsend, 1927a:257. Type-species, *reclinata* Townsend (orig. des.).
RECLINATA Townsend, 1927a:299. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus CONACTIODORIA Townsend

- CONACTIODORIA Townsend, 1934:402. Type-species, *aurea* Townsend (orig. des.).
ALEURITES Townsend, 1940b:893. Type-locality: Brazil, São Paulo, Tietê.
AUREA Townsend, 1934a:403. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus COROZALIA Curran

- COROZALIA Curran, 1934:429. Type-species, *longula* Curran (orig. des.).
LONGULA Curran, 1934:465. Type-locality: Panama, Canal Zone, Corozal (cf. Arnaud, 1963:112).

Genus CROESOACTIA Townsend

- CROESOACTIA Townsend, 1927a:269. Type-species, *cincta* Townsend (orig. des.).
CINCTA Townsend, 1927a:300. Type-locality: Peru, Charape River.

Genus CRYPTOMEIGENIA Brauer & Bergenstamm

- CRYPTOMEIGENIA Brauer & Bergenstamm, 1891:311 (1891:7). Type-species, *setifacies* Brauer & Bergenstamm (mon.).
AURIFACIES Walton, 1912:199, pl. 10. Type-locality: Puerto Rico, Anasco.
ELEGANS (Wulp), 1890:143 (*Hypostena*). Type-locality: Mexico, Guerrero, Xucumanatlán. N. comb. (C. W. Sabrosky, *in litt.*).
SETIFACIES Brauer & Bergenstamm, 1891:311 (1891:6). Type-locality: "Brazil".
MERIDIONALIS (Townsend), 1912a:305 (*Eumyothyria*). Type-locality: Peru, Piura.

Genus CRYPTOMEIGENIOIDEA Thompson

- CRYPTOMEIGENIOIDEA Thompson, 1968:40. Type-species, *longipes* Thompson (orig. des.).
LONGIPES Thompson, 1968:40, figs. 12-15, 17a. Type-locality: Trinidad, Simla.

Genus CUPARYMYIA Townsend

- CUPARYMYIA Townsend, 1934:397. Type-species, *anametopochaetoides* Townsend (orig. des.).
ANAMETOPOCHAETOIDES Townsend, 1934:397. Type-locality: Brazil, Boa Vista, Rio Tapajós, Cupari trail.



Genus DEXODIOPSIS Thompson

- DEXODIOPSIS Thompson, 1968:202. Type-species, *aurea* Thompson (orig. des.).
AUREA Thompson, 1968:202, figs. 138-140. Type-locality: Trinidad, Curepe.
CLARIPALPIS Thompson, 1968:205, fig. 37. Type-locality: Trinidad, P. Highway.

Genus DEXODIMYIA Thompson

- DEXODIMYIA Thompson, 1968:181. Type-species, *discalis* Thompson (orig. des.).
DISCALIS Thompson, 1968:181. Type-locality: Trinidad, New Farm.

Genus DEXODIOIDEA Thompson

- DEXODIOIDEA Thompson, 1968:80. Type-species, *aurea* Thompson (orig. des.).
AUREA Thompson, 1968:81, fig. 45. Type-locality: Trinidad, Tunapuna.

Genus DEXODOMINTHO Townsend

- DEXODOMINTHO Townsend, 1935d:226. Type-species, *fumipennis* Townsend (orig. des.).
FUMIPENNIS Townsend, 1935d:226. Type-locality: Brazil, São Paulo, São Paulo, Vila Ema.

Genus DIDYMA Wulp

- DIDYMA Wulp, 1890:156. Type-species, *albomicans* Wulp (Brauer & Bergenstamm, 1893:30) (1893:118).
ALBOMICANS Wulp, 1890:162. Type-locality: Mexico, Orizaba and Guerrero. Ref. — Townsend, 1931b:179 (note on type).
AMBULATRIX Wulp, 1890:158. Type-locality: Mexico, Guerrero, Sierra de las Aguas Escondidas.
APICALIS Brêthes, 1920a:32. Type-locality: "Peru". Unrecognized.
CALYPTRATA Williston, 1896:359, pl. 11, fig. 95. Type-locality: St. Vincent I., West Indics.
EXIGUA Wulp, 1890:160. Type-locality: Mexico, Guerrero, La Venta.
INCONSPICUA Wulp, 1890:163. Type-locality: Mexico, Guerrero, Amulango (formely Amula).
MODESTA Wulp, 1890:160. Type-locality: Mexico, Guerrero, Sierra de las Aguas Escondidas.
NIGRICOLOR Wulp, 1890:158. Type-locality: Mexico, Veracruz and Tabasco; Costa Rica.
OTIOSA Wulp, 1890:159. Type-locality: Mexico, Guerrero, Amulango (formely Amula).
PAVIDA Wulp, 1890:165. Type-locality: Mexico, Guerrero.
PROMPTA Wulp, 1890:161. Type-locality: Mexico, Guerrero.
TIMIDA Wulp, 1890:161. Type-locality: Mexico, Guerrero, Amulango (formely Amula).



Genus DIDYMOPS Townsend

- DIDYMOPS Townsend, 1927a:262. Type-species, *yahuarmayensis* Townsend (orig. des.).
YAHUARMAYENSIS Townsend, 1927a:302. Type-locality: Peru, Yahuarmayo River.

Genus DORYPHOROPHAGA Townsend

- DORYPHOROPHAGA Townsend, 1912d:164. Type-species, *Lydella doryphorae* Riley (orig. des.).
Austrolydella Townsend, 1919a:573. Type-species, *assimilis* Townsend (orig. des.). N. SYN.
Ref. — Reinhard, 1935b (rev.).
ASSIMILIS (Townsend), 1919a:573 (*Austrolydella*). Type-locality: Peru, Santa Eulalia. N. COMB.
AUSTRALIS Reinhard, 1935b:389. Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico.
DEJECTA (Wulp), 1890:113 (*Masicera*). Type-locality: Mexico, Guerrero. Unrecognized.
DORYPHORAE (Riley), 1869:111 (*Lydella*). Type-locality: U.S.A., Missouri. Distr. — U.S.A., Mexico.
HYALINIPENNIS (Wulp), 1890:152 (*Degeeria*). Type-locality: Mexico, Guerrero, Amulango (formely Amula). N. COMB.
magnicornis Wulp, 1890:153 (*Degeeria*). Type-locality: Mexico, Guerrero and Veracruz. N. SYN.
MACELLA Reinhard, 1935b:390. Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico.
NITIDIUSCULA (Wulp), 1890:151 (*Degeeria*). Type-locality: Mexico, Guerrero, Chilpancingo. N. COMB.
SEROTINA (Wulp), 1890:124 (*Prospheerisa*). Type-locality: Mexico, Mexico City. N. COMB.

Genus EGAMEIGENIA Townsend

- EGAMEIGENIA Townsend, 1927a:278. Type-species, *amazonica* Townsend (orig. des.).
AMAZONICA Townsend, 1927a:303. Type-locality: Brazil, Amazonas, Tefé.

Genus EPIDEXIA Townsend

- EPIDEXIA Townsend, 1912b:112. Type-species, *filamentosa* Townsend (orig. des.) = *pulverea* (Coquillett).
MIMELA Reinhard, 1953a:96. Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico (Morelos).

Genus EPIMEIGENIA Townsend

- EPIMEIGENIA Townsend, 1931a:463. Type-species, *argentina* Townsend (orig. des.).
ARGENTINA Townsend, 1931a:463. Type-locality: Argentina, Mendoza.



Genus EPIPHANOCERA Townsend

EPIPHANOCERA Townsend, 1915m:419. Type-species, *costalis* Townsend (orig. des.).

COSTALIS Townsend, 1915m:419. Type-locality: Peru, Canyon of Río Huallaga, Hariaca.

Genus EPIPROSPHERYSA Townsend

EPIPROSPHERYSA Townsend, 1927a:277. Type-species *charapensis* Townsend (orig. des.).

CHARAPENSIS Townsend, 1927a:305. Type-locality: Peru, Charape River.

Genus ERYTHROARGYROPS Townsend

ERYTHROARGYROPS Townsend, 1917c:124. Type-species, *elegans* Townsend (orig. des.).

ELEGANS Townsend, 1917c:124. Type-locality: Peru, Jaen, Huascaray.

Genus EUANISIA Blanchard

EUANISIA Blanchard, 1947a:12. Type-species, *mesacarrioni* Blanchard (orig. des.).

MESACARRIONI Blanchard, 1947a:13, fig. 6. Type-locality: Uruguay, Montevideo. Ref. — Blanchard, 1958:40 (biol. notes).

Genus EUCELATORIA Townsend

EUCELATORIA Townsend, 1908:249. Type-species, *Tachina armigera* Coquillett (orig. des.).

ARMIGERA (Coquillett), 1889:332 (*Tachina*). Type-locality: U.S.A., California. Distr. — U.S.A., Mexico, West Indies.

AURESCENS Townsend, 1917b:226 (as *australis* ssp.). Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães) and Corumbá.

AUSTRALIS Townsend, 1911b:140 (also, 1912a:315). Type-locality: Peru, Piura. Distr. — West Indies (St. Vincent), Puerto Rico, Peru.

oppugnator Walton, 1914b:93, fig. 7 (*Compsilura*). Type-locality: "Puerto Rico". Ref. — Aldrich, 1927a:19 (synonymy).

COMOSA (Wulp), 1890:119 (*Prospberrysa*). Type-locality: Mexico, Tabasco, Teapa.

CORA (Bigot), 1889:259 (*Degeeria*). Type-locality: "Mexico".

MONTANA Townsend, 1929:371. Type-locality: Peru, Ushpayacu River.

NIGRIPALPIS (Bigot), 1889:258 (*Chaetolyga*). Type-locality: "Mexico".

POLLENS (Wulp), 1890:174 (*Telothyria*). Type-locality: Mexico, Guerrero. N. comb.

Genus EUCELATORIOIDEA Thompson

EUCELATORIOIDEA Thompson, 1968:176. Type-species, *nigripalpis* Thompson (orig. des.).

NIGRIPALPIS Thompson, 1968:177. Type-locality: "Trinidad".

PHYSONOTAE Thompson, 1968:179. Type-locality: Trinidad, Caroni.

Genus EUCELATORIOPSIS Townsend

EUCELATORIOPSIS Townsend, 1927a:276. Type-species, *teffeensis* Townsend (orig. des.).

Tachinophytopsis Townsend, 1927a:277. Type-species, *carinata* Townsend (orig. des.).

This genus has been recorded as parasite of several cassidine Chrysomelidae.

CARINATA (Townsend), 1927a:359 (*Tachinophytopsis*). Type-locality: Brazil, São Paulo (Itaquaquecetuba and Cantareira).

DIMMOCKI (Aldrich), 1932a:5 (*Anetia*). Type-locality: U.S.A., Massachusetts. Distr. — U.S.A., Mexico.

DISSEPTA (Wulp), 1890:176 (*Telothyria*). Type-locality: Mexico, Guerrero. N. comb.

MERIDIONALIS (Townsend), 1912a:316 (*Dexodes*). Type-locality: Peru, Piura.

OCCULTA (Wulp), 1890:184 (*Telothyria*). Type-locality: Mexico, Guerrero. N. comb.

PARKERI Sabrosky, 1952:325. Type-locality: Uruguay, Montevideo.

STRIOLATA (Wulp), 1890:179 (*Telothyria*). Type-locality: Mexico, Veracruz and Tabasco. N. comb.

TEFFEENSIS Townsend, 1927a:306. Type-locality: Brazil, Amazonas, Tefé (formerly spelled as Tefé).

Genus EUCLAUSICELLA Townsend

EUCLAUSICELLA Townsend, 1927a:269. Type-species, *uruhuasi* Townsend (orig. des.).

Eclausicella, misprint.

URUHUASI Townsend, 1927a:306. Type-locality: Peru, San Gabán Valley, Uruhuasi.

Genus EUMACHAERAEA Townsend

EUMACHAERAEA Townsend, 1927a:263. Type-species, *auricephala* Townsend (orig. des.).

AURICEPHALA Townsend, 1927a:308. Type-locality: Peru, El Potrero and Cavalera Mts.

Genus EUPROSPHERYSA Townsend

EUPROSPHERYSA Townsend, 1928b:147. Type-species, *fumpennisi* Townsend (orig. des.).

FUMPENNISI Townsend, 1928b:147. Type-locality: Peru, Las Flores, Mayo River. Ref. — Thompson, 1968:94 (immature stages).

Genus EUPTILODEGEERIA Townsend

EUPTILODEGEERIA Townsend, 1931a:465. Type-species, *Hypostena obumbrata* Wulp (orig. des.).

OBUMBRATA (Wulp), 1890:143 (*Hypostena*). Type-locality: Mexico, Guerrero and Veracruz.

Genus EUTHELYCONYCHIA Townsend

- EUTHELYCONYCHIA Townsend, 1927a:279. Type-species, *clausa* Townsend (orig. des.).
CLAUSA Townsend, 1927a:309. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus GYMNOGORIA Townsend

- GYMNOGORIA Townsend, 1927a:260. Type-species, *capitata* Townsend (orig. des.).
CAPITATA Townsend, 1927a:311. Type-locality: Peru, Charape River.

Genus HELIODEXODES Thompson

- HELIODEXODES Thompson, 1968:197. Type-species, *argenteus* Thompson (orig. des.).
ARGENTUS Thompson, 1968:197, figs. 130-133. Type-locality: Trinidad, American Road, Morvan Hill.

Genus HELIOLYDELLA Townsend

- HELIOLYDELLA Townsend, 1927a:277. Type-species, *aurata* Townsend (orig. des.).
AURATA Townsend, 1927a:314. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
OBSCURATA (Wulp), 1890:62 (*Exorista*). Type-locality: "Mexico". N. comb.

Genus HEMIARGYROPHYLAX Townsend

- HEMIARGYROPHYLAX Townsend, 1927a:265. Type-species, *punctilucis* Townsend (orig. des.).
PUNCTILUCIS Townsend, 1927a:314. Type-locality: Peru, Chachapoyas.

Genus HEMIARGYROPSIS Townsend

- HEMIARGYROPSIS Townsend, 1927a:260. Type-species, *frontalis* Townsend (orig. des.).
FRONTALIS Townsend, 1927a:315. Type-locality: Peru, Charape River.

Genus HEMILYDELLA Townsend

- HEMILYDELLA Townsend, 1927a:278. Type-species, *fasciata* Townsend (orig. des.).
FASCIATA Townsend, 1927a:315. Type-locality: Peru, La Tina, Macará River.

Genus HYPODORIA Townsend

- HYPODORIA Townsend, 1927a:274. Type-species, *orbitalis* Townsend (orig. des.).
ORBITALIS Townsend, 1927a:317. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus HYPOHOUGHIA Townsend

- HYPOHOUGHIA Townsend, 1927a:255. Type-species, *reclinata* Townsend (orig. des.).
RECLINATA Townsend, 1927a:318. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus HYPOMYOTHYRIA Townsend

- HYPOMYOTHYRIA Townsend, 1927a:276. Type-species, *hypodermica* Townsend (orig. des.).
HYPODERMICA Townsend, 1927a:318. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus HYPOPHYLAX Townsend

- HYPOPHYLAX Townsend, 1935d:232. Type-species, *prosperyx* Townsend (orig. des.).
PROSPERYX Townsend, 1935d:232. Type-locality: Guyana, Pacarayma Mts., Upper Ireng River.

Genus HYPOPROXYNOPS Townsend

- HYPOPROXYNOPS Townsend, 1927a:279. Type-species, *rufiventris* Townsend (orig. des.).
RUFIVENTRIS Townsend, 1927a:319. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
xanthipes Blanchard in Marchionatto, [1940]:44. *Nomen nudum*.

Genus INCAMYIA Townsend

- INCAMYIA Townsend, 1912a:317. Type-species, *cuzcensis* Townsend (orig. des.). Ref. — Cortés, 1968b (key).
Sphalloglandulus Townsend, 1915m:438. Type-species, *unicus* Townsend (orig. des.).
Prophrynopsis Townsend, 1927a:273. Type-species, *peruviana* Townsend (orig. des.). N. SYN.
Ref. — Aldrich, 1928:15 (key); 1934:65 (key to Patagonian spp.).
CHARLINI Cortés, 1968b:19. Type-locality: Chile, Santiago, Rinconada de Maipo.
CHILENSIS Aldrich, 1928b:16. Type-locality: Chile, Angol. Distr. — Chile. Juan Fernández I., and Argentina.
Ref. — Caltagirone, 1953 (biol. immature stage); Parker, 1953 (immature stages).
CUZCENSIS Townsend, 1912a:317. Type-locality: Peru, Cuzco.
NUDA Aldrich, 1934:66. Type-locality: Argentina, Rio Negro, Bariloche.
PERUVIANA (Townsend), 1927a:353 (*Prophrynopsis*). Type-locality: Peru, Matucana. N. COMB.
SPINICOSTA Aldrich, 1928b:15. Type-locality: Chile, Valparaíso, Perales.
STRIATA Aldrich, 1928b:16. Type-locality: Peru, Oroya.
UNICA (Townsend), 1915m:438 (*Sphalloglandulus*). Type-locality: Peru, Oroya, Río Mantaro Valley.



Genus INCAMYIOPSIS Townsend

- INCAMYIOPSIS Townsend, 1919a:587. Type-species, *imitatrix* Townsend (orig. des.).
IMITATRIX Townsend, 1919a:587. Type-locality: Peru, Oroya.

Genus ITABIOMYIA Townsend

- ITASIOMYIA Townsend, 1927a:280. Type-species, *fulvescens* Townsend (orig. des.).
FULVESCENS Townsend, 1927a:319. Type-locality: Brazil, São Paulo, São Paulo, Cantareira.

Genus ITALISPIDEA Townsend

- ITALISPIDEA Townsend, 1927a:268. Type-species, *antennalis* Townsend (orig. des.).
ANTENNALIS Townsend, 1927a:320. Type-locality: Brazil, São Paulo, Itaguaquecetuba and Cantareira.

Genus ITALYDELLA Townsend

- ITALYDELLA Townsend, 1927a:278. Type-species, *geminata* Townsend (orig. des.).
CHACHAPOYANA Townsend, 1929:372. Type-locality: Peru, Chachapoyas.
CEMINATA Townsend, 1927a:320. Type-locality: Brazil, São Paulo, Itaguaquecetuba.

Genus ITAPLECTOPS Townsend

- TAPLECTOPS Townsend, 1927a:265. Type-species, *antennalis* Townsend (orig. des.).
ANTENNALIS Townsend, 1927a:321. Type-locality: Brazil, São Paulo, São Paulo, Cantareira.

Genus LEPTOSTYLUM Macquart

- LEPTOSTYLUM Macquart. 1851:180 (1851:207). Type-species, *pulchellum* Macquart (orig. des.) = *busckii* (Townsend).
Argyreomyia Townsend, 1915m:426. Type-species, *busckii* Townsend (orig. des.).
BUSCKII (Townsend), 1915m:426 (*Argyreomyia*). Type-locality: Panama, Porto Bello.
ITAQUAQUECETUBAE (Townsend), 1929:375 (*Argyreomyia*). Type-locality: Brazil, São Paulo, Itaguaquecetuba.
LEUCONOTUM (Wulp), 1892:195 (*Exorista*). Type-locality: Mexico, Guerrero.
PULCHELLUM Macquart. 1851:181 (1851:208), pl. 19, fig. 2. Type-locality: Brazil, Bahia. Ref. — Townsend, 1931b:179 (notes on types).

Genus LESKIOLYDELLA Townsend

- LESKIOLYDELLA Townsend, 1927a:269. Type-species, *aurata* Townsend (orig. des.).



AURATA Townsend, 1927a:324. Type-locality: Brazil, São Paulo, São Paulo, Cantarcira.

Genus LIXINIA Curran

LIXINIA Curran, 1926a:108. Type-species, *jamaicensis* Curran (orig. des.).

CARINATA Curran, 1926a:108. Type-locality: Jamaica, Hope Gardens.

JAMAICENSIS Curran, 1926a:108. Type-locality: Jamaica, Hill Gardens.

Genus LIXOPHAGA Townsend

LIXOPHAGA Townsend, 1908:86. Type-species, *parva* Townsend (orig. des.).

Euzenilliopsis Townsend, 1916:76. Type-species, *diatraeae* Townsend (orig. des.).

Ref. — Aldrich, 1925 (rev.); Curran, 1935 (key to males).

DIATRAEAE (Townsend), 1915d:76 (*Euzenilliopsis*). Type-locality: "Cuba" (as Louisiana, error). Distr. — West Indies, Neotropical. Ref. — Thompson, 1960:209; Box, 1928 (biol.).

FAMELICA (Wiedemann), 1830:331 (*Tachina*). Type-locality: "Brazil". Ref. — Aldrich, 1927c:17.

GRISEA (Curran), 1926a:106 (*Hypostena*). Type-locality: Jamaica, Dcnbigh.

REMORA Reinhard, 1953:102. Type-locality: Mexico, Morelos, Acatlipa.

SCINTILLA Reinhard, 1953:101. Type-locality: Mexico, Morelos, Acatlipa.

SOLITARIA (Curran), 1926a:107 (*Hypostena*). Type-locality: Jamaica, Blue Castle.

STENOMAE Curran, 1935:23. Type-locality: Costa Rica, San Pedro de Montes de Oca.

Genus LYDELLOHOUGHIA Townsend

LYDELLOHOUGHIA Townsend, 1927a:270. Type-species, *nana* Townsend (orig. des.).

NANA Townsend, 1927a:324. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus LYDELLOTHELAIRA Townsend

LYDELLOTHELAIRA Townsend, 1919a:558. Type-species, *collaris* Townsend (orig. des.).

COLLARIS Townsend, 1919a:559. Type-locality: Peru, Casahuiri.

Genus LYDINOLYDELLA Townsend

LYDINOLYDELLA Townsend, 1927a:278. Type-species, *metallica* Townsend (orig. des.).

Myiodoria Townsend, 1927a:279. Type-species, *discahis* Townsend (orig. des.). N. SYN.

DISCALIS (Townsend), 1927a:331 (*Myiodoria*). Type-locality: Peru, Chaquimayo River. N. COMB.

HUMERALIS (Wulp), 1890:173 (*Telothyria*). Type-locality: Mexico, Tabasco. N. COMB.

- METALLICA Townsend, 1927a:325. Type-locality: Brazil, São Paulo, Itaqua-
qucetuba. Ref. — Berry & Parker, 1949:97 (biol.).
? *bruchii* Blanchard, 1943a:92, fig. 1. Type-locality: Argentina,
Tucumán.
RASLIS (Wulp), 1890:175 (*Telothyria*). Type-locality: Mexico, Guerrero.
N. COMB.

Genus MACHAIROMASICERA Townsend

- MACHAIROMASICERA Townsend, 1919a:577. Type-species, *carinata* Townsend
(orig. dcs.).
CARINATA Townsend, 1919a:578. Type-locality: Ecuador, Manchi.

Genus MARACASIMYIA Thompson

- MARACASIMYIA Thompson, 1968:140. Type-species, *antennalis* Thompson
(orig. des.).
ANTENNALIS Thompson, 1968:141, fig. 89. Type-locality: Trinidad, Maracas
Valley.

Genus MAYOPHORINIA Townsend

- MAYOPHORINIA Townsend, 1927a:263. Type-species, *angusta* Townsend (orig.
des.).
ANGUSTA Townsend, 1927a:326. Type-locality: Peru, Yahuar Mayo River.

Genus MEIGENIELLOIDES Townsend

- MEIGENIELLOIDES Townsend, 1919a:573. Type-species, *cinerea* Townsend (orig.
des.).
CINEREA Townsend, 1919a:574. Type-locality: U.S.A., N. Mexico. Distr. —
U.S.A., Mexico.

Genus MELANACTIA Townsend

- MELANACTIA Townsend, 1927a:258. Type-species, *macrocera* Townsend (orig.
des.).
MACROCERA Townsend, 1927a:326. Type-locality: Brazil, São Paulo, Itaqua-
qucetuba.

Genus MELANODORIA Townsend

- MELANODORIA Townsend, 1927a:280. Type-species, *nigrisquamis* Townsend
(orig. des.).
NIGRISQUAMIS Townsend, 1927a:327. Type-locality: Peru, Yahuar Mayo.

Genus METADORIA Brauer & Bergenstamm

- METADORIA Brauer & Bergenstamm, 1893:117 (1893:29). Type-species,
mexicana Brauer & Bergenstamm (orig. dcs., as gen. n., sp. n.)
= *barbata* (Bigot).
Ref. — Townsend, 1931:180 (notes on types).

- BARBATA (Bigot), 1889:260 (*Phorocera*). Type-locality: "Mexico".
melanocephala Bigot, 1889:260 (*Phorocera*). Type-locality: "Mexico".
atricaps Wulp, 1890:79 (*Phorocera*). Type-locality: Mexico, Guerrero.
mexicana Brauer & Bergenstamm, 1893:117 (1893:20). Type-locality:
 "Mexico".
 BASILARIS (Wulp), 1890:159 (*Didyma*). Type-locality: Mexico, Guerrero,
 Omilteme.
 NIGRIFACIES Bigot, 1889:268 (*Oplisa*). Type-locality: "Mexico".
 PARVULA (Wulp), 1890:78 (*Phorocera*). Type-locality: Mexico, Orizaba.
 N. COMB.

Genus MELANORLOPTERYX Townsend

- MELANORLOPTERYX Townsend, 1927a:273. Type-species, *costalis* Townsend
 (orig. des.).
 COSTALIS Townsend, 1927a:328. Type-locality: Peru, Chachapoyas.

Genus MELANOROMINTHO Townsend

- MELANOROMINTHO Townsend, 1935d:226. Type-species, *barbiellini* Townsend
 (orig. des.).
 BARBIELLINI Townsend, 1935d:227. Type-locality: Brazil, São Paulo, São
 Paulo, Vila Ema.

Genus MESOCHAETA Brauer & Bergenstamm

- MESOCHAETA Brauer & Bergenstamm, 1891:341 (1891:37). Type-species,
Didyma commixta (Wulp) (as *connexa*, error) (mon.).
 COMMIXTA (Wulp), 1890:163 (*Didyma*). Type-locality: Mexico, Tabasco,
 Teapa.
 MERIDIONALIS Townsend, 1929:376. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
 PAULISTA Townsend, 1929:376. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus METARRHINOMYIA Townsend

- METARRHINOMYIA Townsend, 1927a:279. Type-species, *angusta* Townsend
 (orig. des.).
 ANGUSTA Townsend, 1927a:329. Type-locality: Peru, Río Charape.

Genus METOPOACTIA Townsend

- METOPOACTIA Townsend, 1927a:268. Type-species, *andina* Townsend (orig.
 des.).
 ANDINA Townsend, 1927a:329. Type-locality: Peru, Chachapoyas.

Genus MICROTOWNSENDIA Curran

- MICROTOWNSENDIA Curran, 1934d:463. Type-species, *nitens* Curran (orig.
 des.).
 NITENS Curran, 1934d:471. Type-locality: Panama, Canal Zone, Corozal
 (cf. Arnaud, 1963a:121).



Genus MINTHOMYIA Townsend

- MINTHOMYIA Townsend, 1919a:564. Type-species, *obdominalis* Townsend (orig. des.).
ABDOMINALIS Townsend, 1919a:564. Type-locality: Peru, Jaen, Charape River.

Genus MINTHOPSIS Townsend

- MINTHOPSIS Townsend, 1915m:417. Type-species, *vittata* Townsend (orig. des.).
VITTATA Townsend, 1915m:417. Type-locality: "Peru". Ref. — Townsend, 1934:405 (notes on types).

Genus MINTHOTACHINA Townsend

- MINTHOTACHINA Townsend, 1935d:227. Type-species, *miscella* Townsend (orig. des.).
MISCELLA Townsend, 1935d:227. Type-locality: "Trinidad".

Genus MUSCINOTHELAIRA Townsend

- MUSCINOTHELAIRA Townsend, 1916g:310. Type-species, *lutzi* Townsend (orig. des.).
LUTZI Townsend, 1916g:310. Type-locality: Brazil, São Paulo.

Genus MYIODORIOPS Townsend


- MYIODORIOPS Townsend, 1935d:227. Type-species, *marginalis* Townsend (orig. des.).
MARGINALIS Townsend, 1935d:227. Type-locality: Guyana, Parika, Rupununi

Genus MYIOMINTHO Brauer & Bergenstamm

- MYIOMINTHO Brauer & Bergenstamm, 1889:138 (1889:70). Type-species, *elata* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.). Ref. — Aldrich, 1925c:118 (type redeser.).
ELATA Brauer & Bergenstamm, 1889:138 (1889:70). Type-locality: "Venezuela".

Genus MYIOPHARUS Brauer & Bergenstamm

- MYIOPHARUS Brauer & Bergenstamm, 1889:161 (1889:93). Type-species, *metopia* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.).
Hemiargyra Townsend, 1908:88. Type-species, *nigra* Townsend (orig. des.).
Ref. — Aldrich, 1924a:216 (notes on types); idem, 1934:62 (Chilean spp.).
CARBONARIUS (Wulp), 1890:78 (*Phorocera*). Type-locality: Mexico, Guerrero. N. comb.
CONNEXUS (Wulp), 1890:173 (*Telothyria*). Type-locality: Mexico, Guerrero. N. comb.
HUASCARAYUS (Townsend), 1917c:125 (*Hemiargyra*). Type-locality: Peru, Huascaray Ridge.



- METOPIA Brauer & Bergenstamm, 1889:161 (1889:93). Type-locality: Mexico, Tacubaya and Yucatan (Merida).
 MOESTUS (Wulp), 1890:158 (*Didyma*). Type-locality: Mexico, Guerrero.
 MURINUS (Wulp), 1890:180 (*Telothyria*). Type-locality: Mexico, Guerrero.
 N. COMB.
 NIGER (Townsend), 1908:90 (*Hemiargyra*). Type-locality: "Costa Rica".
 PIRIONI Aldrich, 1934:64. Type-locality: Chile, Valparaiso, Perales. Distr. — Chile (Valparaiso, Concepción).
 SUBAENEUS Aldrich, 1934:63. Type-locality: Chile, Valparaiso, Perales.
 UNICOLOR (Wulp), 1890:63 (*Exorista*). Type-locality: Mexico (sev. places).
 N. COMB.

Genus MYIOXYNOPS Townsend

- MYIOXYNOPS Townsend, 1927a:278. Type-species, *palpalis* Townsend (orig. des.).
 PALPALIS Townsend, 1927a:331. Type-locality: Peru, Yahuarmayo.

Genus NEOARGYROPHYLAX Townsend

- NEOARGYROPHYLAX Townsend, 1927a:265. Type-species, *argentescens* Townsend (orig. des.).
 ARGENTESCENS Townsend, 1927a:332. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus NEOMINTHOPSIS Townsend

- NEOMINTHOPSIS Townsend, 1915m:418. Type-species, *discalis* Townsend (orig. des.).
 DISCALIS Townsend, 1915m:418. Type-locality: Peru, Pachacayo.

Genus NEOXYNOPSIDEA Thompson


- NEOXYNOPSIDEA Thompson, 1968:149. Type-species, *claripalpis* Thompson (orig. des.).
 CLARIPALPIS Thompson, 1968:149, figs. 95, 96. Type-locality: Trinidad, Maracas Valley.

Genus OEDEMAMEDINA Townsend

- OEDEMAMEDINA Townsend, 1927a:269. Type-species, *costalis* Townsend (orig. des.).
 COSTALIS Townsend, 1927a:337. Type-locality: Peru, Uruhuasi.

Genus OLINDOPSIS Townsend

- OLINDOPSIS Townsend, 1927a:274. Type-species, *andinensis* Townsend (orig. des.).
 ANDINENSIS Townsend, 1927a:339. Type-locality: Peru, near Tabalosas.



Genus OLLACHACTIA Townsend

OLLACHACTIA Townsend, 1927a:262. Type-species, *mucronata* Townsend (orig. des.).

MUCRONATA Townsend, 1927a:339. Type-locality: Peru, Ollachea.

Genus OLLACHEA Townsend

OLLACHEA Townsend, 1919a:576. Type-species, *elongata* Townsend (orig. des.).

ELONGATA Townsend, 1919a:577. Type-locality: Peru, Ollachea.

Genus OOMEIGENIA Townsend

OOMEIGENIA Townsend, 1915m:434. Type-species, *chosica* Townsend (orig. des.).

CHOSICA Townsend, 1915m:435. Type-locality: Peru, Chosica.

Genus OROPHOROCERA Townsend

OROPHOROCERA Townsend, 1927a:267. Type-species, *ocellaris* Townsend (orig. des.).

OCELLARIS Townsend, 1927a:342. Type-locality: Peru, Uruhuasi.

Genus OSWALDIA Robineau-Desvoidy

OSWALDIA Robineau-Desvoidy, 1863a:840. Type-species, *muscaria* Robineau-Desvoidy (orig. des.) = *muscaria* (Fallén).

IMMISSA (Reinhard), 1959b:232 (*Lydella*). Type-locality: Mexico, Michoacán, Zamora.

Genus OXYAPORIA Townsend

OXYAPORIA Townsend, 1919b:170. Type-species, *Gymnostylia ornata* Brauer & Bergenstamm (orig. des.).

Gymnostylia Macquart of Brauer & Bergenstamm.

ARGENTINA (Brèthes), 1922:21 (*Gymnostylia*). Type-locality: Argentina, Buenos Aires, Station Condarco.

ORNATA (Brauer & Bergenstamm), 1889:128 (1889:60) (also, 1891:374; 1891:70) (*Gymnostylia*). Type-locality: "Venezuela". Ref. — Aldrich, 1927:16 (notes on types).

Genus OXYNOPS Townsend

OXYNOPS Townsend, 1912b:10. Type-species, *serratus* Townsend (orig. des.) = *anthracinus* (Bigot).

ANTHRACINUS (Bigot), 1889:259 (*Degeeria*). Type-locality: "Mexico".

MONOCHROMA (Wulp), 1890:154 (*Degeeria*). Type-locality: Mexico, Veracruz, Atoyac.

Genus OXYNOPSIS Townsend

- OXYNOPSIS Townsend, 1927a:270. Type-species, *brasiliensis* Townsend (orig. des.).
BRASILIENSIS Townsend, 1927a:343. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus OXYNOPSALIA Curran

- OXYNOPSALIA Curran, 1934c:444. Type-species, *nitida* Curran (orig. des.). *Oxsnopsalia*, misprint.
NITIDA Curran, 1934c:467, figs. 170, 194. Type-locality: Panama, Canal Zone, Barro Colorado I. (cf. Arnaud, 1963:123).

Genus OXYOPHIRION Townsend

- OXYOPHIRION Townsend, 1927a:270. Type-species, *punctigerum* Townsend (orig. des.).
PUNCTIGERUM Townsend, 1927a:343. Type-locality: Brazil, São Paulo, São Paulo, Cantareira.

Genus PANACEMYIA Townsend

- PANACEMYIA Townsend, 1919b:164. Type-species, *panamensis* Townsend (orig. des.).
PANAMENSIS Townsend, 1919b:164. Type-locality: Panama, Taboga I.
STIPATUS (Reinhard), 1958b:225 (*Dolichotarsus*). Type-locality: Mexico, W. Slope of Popocatepetl. N. comb.
TRINITATIS Thompson, 1963a:459, fig. 131. Type-locality: Trinidad, St. Augustine.

Genus PARADORIA Brauer & Bergenstamm

- PARADORIA Brauer & Bergenstamm, 1891:339 (1891:35). Type-species, *nigra* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.).
Euhemiargyra Townsend, 1927a:260. Type-species, *parva* Townsend (orig. des.).
Dactylodidyma Townsend, 1927a:260. Type-species, *dubia* Townsend (orig. des.).
Ref. — Aldrich, 1925d:457.
DUBIA (Townsend), 1927a:301 (*Dactylodidyma*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
NIGRA Brauer & Bergenstamm, 1891:339 (1891:35). Type-locality: "Venezuela". Ref. — Aldrich, 1925:457.
PARVA (Townsend), 1927a:307 (*Euhemiargyra*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus PARALISPE Brauer & Bergenstamm

- PARALISPE Brauer & Bergenstamm, 1891:337 (1891:33). Type-species, *brasiliana* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.).
Stomatolydella Townsend, 1919a:570. Type-species, *infernalis* Townsend (orig. des.).

- ALBIFACIES (Wulp), 1892:194 (*Meigenia*). Type-locality: "Mexico".
 ATRATULA (Walker), 1852:304 (*Tachina*). Type-locality: "Brazil".
 BRASILIANA Brauer & Bergenstamm, 1891:337 (1891:33). Type-locality: "Brazil". Ref. — Aldrich, 1925:113.
 CONTIGUA (Wulp), 1890:120 (*Prospberrya*). Type-locality: Mexico, Cuernavaca.
 INFERNALIS (Townsend), 1919a:570 (*Stomatohydella*). Type-locality: U.S.A., N. Mexico. Distr. — U.S.A., Mexico.
 JAMAICENSIS Curran, 1928a:30. Type-locality: Jamaica, Hope Gardens.
 NIGRITA (Wulp), 1890:77, pl. 3, fig. 11 (*Phorocera*). Type-locality: Costa Rica, Volcán de Irazú.
 PALPALIS Townsend, 1927a:376. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
 TRIFASCIATA (Wulp), 1890:118 (*Prospberrya*). Type-locality: Mexico, Guerrero, Omilteme.

Genus PARANETIA Townsend

- PARANETIA Townsend, 1927a:277. Type-species, *punctata* Townsend (orig. des.).
 PUNCTATA Townsend, 1927a:344. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus PARANOXYNOPS Thompson

- PARANOXYNOPS Thompson, 1968:64. Type-species, *curepeiensis* Thompson (orig. des.).
 CUREPEIENSIS Thompson, 1968:64, fig. 35. Type-locality: Trinidad, Curepe.

Genus PARAPROSPHERYSA Townsend

- PARAPROSPHERYSA Townsend, 1927a:280. Type-species, *fumipennis* Townsend (orig. des.).
 FUMIPENNIS Townsend, 1927a:345. Type-locality: Brazil, São Paulo, São Paulo, Cantareira.

Genus PARATHELAIRA Townsend


- PARATHELAIRA Townsend, 1919a:558. Type-species, *panamensis* Townsend (orig. des.).
 PANAMENSIS Townsend, 1919a:558. Type-locality: Panama, Portobelo.

Genus PHYLLOPHRYNO Townsend

- PHYLLOPHRYNO Townsend, 1927a:263. Type-species, *antennalis* Townsend (orig. des.).
 ANTENNALIS Townsend, 1927a:347. Type-locality: Peru, Huasearay.

Genus PIXIMACTIA Townsend

- PIXIMACTIA Townsend, 1927a:279. Type-species, *uruhuasi* Townsend (orig. des.).
 URUHUASI Townsend, 1927a:348. Type-locality: Peru, Uruhuasi.



Genus PLAXACTIA Townsend

- PLAXACTIA Townsend, 1931a:476. Type-species, *facialis* Townsend (orig. des.).
FACIALIS Townsend, 1931a:477. Type-locality: British Honduras, Manatee.

Genus PRODEXODES Townsend

- PRODEXODES Townsend, 1927a:280. Type-species, *rufiventris* Townsend (orig. des.).
RUFIVENTRIS Townsend, 1927a:350. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus PROPHAENOPSIS Townsend

- PROPHAENOPSIS Townsend, 1927a:273. Type-species, *nitens* Townsend (orig. des.).
NITENS Townsend, 1927a:351. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus PROPHORINIA Townsend

- PROPHORINIA Townsend, 1927a:277. Type-species, *proletaria* Townsend (orig. des.).
PROLETARIA Townsend, 1927a:352. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus PROROGLUTEA Townsend

- PROROGLUTEA Townsend, 1919a:574. Type-species, *piligera* Townsend (orig. des.).
PILIGERA Townsend, 1919a:574. Type-locality: Costa Rica, San Carlos.

Genus PROSPHERYSODORIA Townsend

- PROSPHERYSODORIA Townsend, 1928b:149. Type-species, *paraguayensis* Townsend (orig. des.).
PARAGUAYENSIS Townsend, 1928b:149. Type-locality: Paraguay, Santa Trinidad.

Genus PSEUDOCELATORIA Thompson

- PSEUDOCELATORIA Thompson, 1968:190. Type-species, *robusta* Thompson (orig. des.).
ROBUSTA Thompson, 1968:190, figs. 122, 123. Type-locality: Trinidad, Maracas Valley.

Genus PSEUDOMYOTHYRIA Townsend

- PSEUDOMYOTHYRIA Townsend, 1892d:131. Type-species, *indecisa* Townsend (orig. des.) = *ancilla* (Walker).
ANCILLA (Walker), 1852:299 (*Tachina*). Type-locality: "U.S.A." Distr. — U.S.A., Mexico.



PERPLEXA Townsend, 1911b:135, 148 (also, 1912a:319). Type-locality: Peru, Piura.

Genus PSEUDOPROSPHERYSA Thompson

PSEUDOPROSPHERYSA Thompson, 1968:74. Type-species, *santacruzi* Thompson (orig. des.).

CLARIPALPIS Thompson, 1968:79. Type-locality: Trinidad, Aripo Valley.

SANTACRUZI Thompson, 1968:74, figs. 42, 43. Type-locality: Trinidad, Santa Cruz.

SIMILIS Thompson, 1968:77. Type-locality: Trinidad, St. Augustine.

Genus PSEUDOREDTEMBACHERIA Brauer & Bergenstamm

PSEUDOREDTEMBACHERIA Brauer & Bergenstamm, 1889:138 (1889:70). Type-species, *Redtenbacheria brasiliensis* Schiner (mon.).

Neommasicera Townsend, 1927a:275. Type-species, *fulvipes* Townsend (orig. des.) = *brasiliensis* (Schiner).

Ref. — Townsend, 1931b:179 (gen. synonym).

BRASILIENSIS (Schiner), 1868:323 (*Redtenbacheria*). Type-locality: "Brazil".

Ref. — Aldrich, 1925:119 (type redescri.).

fulvipes Townsend, 1927a:334 (*Neommasicera*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.

fulvipes Townsend, 1927a:338 (*Oligolydella*; preocc. Townsend, 1927a:275). Type-locality: Brazil, São Paulo, Itaquaquecetuba.

SUCCINCTA (Wulp), 1890:133 (*Myobia*). Type-locality: Mexico, Guerrero, Sierra de las Aguas Escondidas. N. comb.

Genus PTILODEGEERIA Brauer & Bergenstamm

PTILODEGEERIA Brauer & Bergenstamm, 1891:375 (1891:71). Type-species, *Hypostena obumbrata* Wulp (mon.; misident) = *lindigi* Townsend.

IMMUNDA (Wulp), 1890:148 (*Hypostena*). Type-locality: Mexico, Veracruz and Tabasco.

LEUCOPHAEA (Wulp), 1890:141 (*Hypostena*). Type-locality: Mexico, Guerrero.

LINDIGI Townsend, 1931a:465. Type-locality: "Venezuela".

obumbrata Wulp of Brauer & Bergenstamm.

TRIANGULIFERA (Bigot), 1889:268 (*Homodexia*). Type-locality: "Mexico".

UMBRIFERA (Walker), 1852:294 (*Tachina*). Type-locality: "Brazil". Unrecognized.

UMBRIPENNIS (Wulp), 1890:144 (*Hypostena*). Type-locality: Mexico, Tabasco.

Genus PYGOPHORINIA Townsend

PYGOPHORINIA Townsend, 1927a:274. Type-species, *peruviana* Townsend (orig. des.).

PERUVIANA Townsend, 1927a:355. Type-locality: Peru, Casahuiri. Ref. — Thompson, 1968:66 (immature stages).



Genus RHOMBOTHYRIOPS Townsend

- RHOMBOTHYRIOPS Townsend, 1915m:419. Type-species, *elegans* Townsend (orig. des.).
ELEGANS Townsend, 1915m:420. Type-locality: Peru, Río Huallaga, Huariaca Canyon.

Genus SANTACRUZIA Thompson

- SANTACRUZIA Thompson, 1968:125. Type-species, *dubiosa* Thompson (orig. des.).
DUBIOSA Thompson, 1968:125, fig. 75. Type-locality: Trinidad, Santa Cruz Valley.

Genus SCHIZOCEROPHAGA Townsend


- SCHIZOCEROPHAGA Townsend, 1916:77. Type-species, *leiby* Townsend (orig. des.).
CARINATA (Wulp), 1890:184 (*Telothyria*). Type-locality: Mexico, Veracruz. N. comb.

Genus SITOPHAGA Gistel

- Fabricia* Meigen, 1838:250 (preocc. Blainville, 1828). Type-species, *Tachina pacta* Meigen (mon.) = *cinerea* (Fallén).
SITOPHAGA Gistel, 1848:ix (nom. nov. for *Fabricia* Meigen). Type-species, *Tachina pacta* Meigen (aut.) = *cinerea* Fallén.
ADSCRIPTA Wulp, 1890:170 (*Telothyria*). Type-locality: Mexico, Veracruz, Atoyac. N. comb.
ANGUSTIFRONS (Reinhard), 1930a:104 (*Viviania*). Type-locality: U.S.A., Texas. Distr. — U.S.A., Cuba.
NUBECULA (Wulp), 1890:170 (*Telothyria*). Type-locality: Mexico, Tabasco, Teapa. N. comb.
ROBUSTA (Wulp), 1890:92 (*Brachycoma*). Type-locality: Mexico, Guerrero and Veracruz.
SORDIDA (Walker), 1852:297 (*Tachina*). Type-locality: "South America".
SUBLUCENS (Wulp), 1890:96 (*Brachycoma*). Type-locality: Mexico, Mexico City.

Genus SPATHIMEIGENIA Townsend

- SPATHIMEIGENIA Townsend, 1915b:19. Type-species, *spinigera* Townsend (orig. des.).
Jicaltepecia Townsend, 1917:49. Type-species, *rafaela* Townsend (orig. des.). N. SYN.
Ref. — Aldrich, 1931d (rev., key).
ALBOPICTA (Bigot), 1889:258 (*Chaetolyga*). Type-locality: "Mexico". Ref. — Aldrich, 1931:10.
CURVA (Wulp), 1890:177 (*Telothyria*). Type-locality: Mexico, Guerrero. N. comb.
DIELOCERI (Townsend), 1942d:438 (*Hylotomomyia*). Type-locality: Brazil, Minas Gerais, Florestal, Pará de Minas. N. comb.



- ITAQUAQUECETUBAE (Townsend), 1929:372 (*Jicaltepecia*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. N. comb.
- MEXICANA Aldrich, 1931d:5. Type-locality: Mexico, Michoacán, Erongaricaro. Ref. — Reinhard, 1958:207 (note).
- OBSCURA Aldrich, 1931d:6. Type-locality: "Mexico".
- RAFAELA (Townsend), 1917a:49 (*Jicaltepecia*). Type-locality: Mexico, Veracruz, Jicaltepec, San Rafael. N. comb.
- REMOTA (Wulp), 1890:181 (*Telothyria*). Type-locality: Mexico, Guerrero. N. comb.
- SUBLINEATA (Wulp), 1890:181 (*Telothyria*). Type-locality: Mexico, Guerrero. N. comb.
- TEXENSIS Aldrich, 1931d:9. Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico.
- VACIVA (Wulp), 1890:176 (*Telothyria*). Type-locality: Mexico, Guerrero. N. comb.
- ZONATA (Bigot), 1889:261 (*Ceromasia*). Type-locality: "Mexico". N. comb.

Genus SPHAERINA Wulp

- SPHAERINA Wulp, 1890:205. Type-species, *nitidula* Wulp (orig. des.).
- Proxynops* Townsend, 1927a:274. Type-species, *proximus* Townsend (orig. des.) = *nitidula* (Wulp).
- NITIDULA Wulp, 1890:205, pl. 4, figs. 16-16a. Type-locality: Mexico, Tabasco, Teapa. Ref. — Townsend, 1931b:179.
- proximus* Townsend, 1927a:353 (*Proxynops*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus SPATHIMYIA Townsend

- SPATHIMYIA Townsend, 1912a:318. Type-species, *ferox* Townsend (orig. des.).
- FEROX Townsend, 1912a:319. Type-locality: Peru, Jaen, Charape River.

Genus STENOCHAETA Thompson

- STENOCHAETA Thompson, 1968:159. Type-species, *claripalpis* Thompson (orig. des.).
- CLARIPALPIS Thompson, 1968:159. Type-locality: Trinidad, Legerville Mt.

Genus STOLATOSOMA Reinhard

- STOLATOSOMA Reinhard, 1953b:96. Type-species, *cidaris* Reinhard (orig. des.).
- CIDARIS Reinhard, 1953b:97. Type-locality: Mexico, Morelos, Acatlipa.

Genus SUCCINGULODES Townsend

- SUCCINGULODES Townsend, 1935d:225. Type-species, *elodioides* Townsend (orig. des.).
- ELODIOIDES Townsend, 1935d:225. Type-locality: Guyana, Pacaraima Mts., head of Mazaruni River.

Genus TACHINOPHYTO Townsend

- TACHINOPHYTO Townsend, 1892d:130. Type-species, *floridensis* Townsend (orig. des.).
Neoxynops Townsend, 1934a:403. Type-species, *nana* Townsend (orig. des.).
FLORIDENSIS Townsend, 1892d:131. Type-locality: U.S.A., Florida. Distr. — U.S.A., Mexico.
NANA (Townsend), 1934a:403 (*Neoxynops*). Type-locality: Brazil, Pará, Urucurituba, Tapajós River.

Genus TALPAROMYIA Thompson

- TALPAROMYIA Thompson, 1968:101. Type-species, *pollinosa* Thompson (orig. des.).
POLLINOSA Thompson, 1968:101, fig. 56. Type-locality: Trinidad, Santa Cruz Valley.

Genus TELOTHYRIOSOMA Townsend

- TELOTHYRIOSOMA Townsend, 1919a:564. Type-species, *tersum* Townsend (orig. des.).
TERSUM Townsend, 1919a:565. Type-locality: Guatemala, Cayuga.

Genus THELAIRODORIA Townsend

- THELAIRODORIA Townsend, 1927a:266. Type-species, *thrix* Townsend (orig. des.).
EXILIS (Wulp), 1890:71 (*Exorista*). Type-locality: Mexico, Guerrero, Amula. N. comb.
THRIX Townsend, 1927a:360. Type-locality: Peru, Chaquimayo.

Genus THELAIRODORIOPSIS Thompson

- THELAIRODORIOPSIS Thompson, 1968:53. Type-species, *maracasi* Thompson (orig. des.).
MARACASI Thompson, 1968:54, figs. 27, 28. Type-locality: Trinidad, "Simla".

Genus THELYOXYNOPS Townsend

- THELYOXYNOPS Townsend, 1927a:273. Type-species, *orbitalis* Townsend (orig. des.).
ORBITALIS Townsend, 1927a:360. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus THELYPHAENOPSIS Townsend

- THELYPHAENOPSIS Townsend, 1927a:262. Type-species, *atra* Townsend (orig. des.).
ATRA Townsend, 1927a:360. Type-locality: Brazil, São Paulo, Itaquaquecetuba.



Genus THEMATHECA Reinhard

- THEMATHECA Reinhard, 1961:207. Type-species, *medeola* Reinhard (orig. des.).
MEDEOLA Reinhard, 1961:208. Type-locality: Mexico, Oaxaca, Río Tehuantepec.

Genus TINALYDELLA Townsend

- TINALYDELLA Townsend, 1927a:265. Type-species, *tinensis* Townsend (orig. des.).
ELONGATA (Wulp), 1890:65, pl. 3, fig. 8-8a (*Exorista*). Type-locality: Costa Rica, Caché. N. comb.
ORDINARIA (Wulp), 1890:64 (*Exorista*). Type-locality: Mexico, Guerrero. N. comb.
TINENSIS Townsend, 1927a:361. Type-locality: Peru, Macará River, La Tina.

Genus TRICHINOAETA Townsend

- TRICHINOAETA Townsend, 1917a:224. Type-species, *orbitalis* Townsend (orig. des.).
ORBITALIS Townsend, 1917a:224. Type-locality: Brazil, Mato Grosso, Chapada (dos Guimarães).

Genus TRYPPHEROMYIA Reinhard

- TRYPPHEROMYIA Reinhard, 1945:32. Type-species, *pallens* Reinhard (orig. des.).
PALLENS Reinhard, 1945:32. Type-locality: U.S.A., Ohio. Distr. — U.S.A., Mexico.

Genus URODEXODES Townsend

- URODEXODES Townsend, 1919a:572. Type-species, *charapensis* Townsend (orig. des.).
CHARAPENSIS Townsend, 1919:572. Type-locality: Peru, Jaen, Charapc River.
CINGULATUS (Schiner), 1868:327 (*Meigenia*). Type-locality: "Brazil". Ref. — Aldrich, 1927c:16.

Genus VELOCIA Robineau-Desvoidy

- VELOCIA Robineau-Desvoidy, 1863:950. Type-species, *cursoria* Robineau-Desvoidy (Townsend, 1916c:9) = *luctuosa* (Meigen).
HAMATA (Wulp), 1890:173 (*Telothyria*). Type-locality: Mexico, Guerrero. N. comb.

Genus XIPHOMYIA Townsend

- XIPHOMYIA Townsend, 1917c:125. Type-species, *gladiatorix* Townsend (orig. des.).
GLADIATRIX Townsend, 1917c:126. Type-locality: Panama, Cabima.
LUCTUOSA (Wulp), 1890:105 (*Masicera*). Type-locality: Mexico (sev. places). N. comb.
RIVALIS Reinhard, 1953a:101. Type-locality: Mexico, Morelos, Acatlipa.

UNPLACED SPECIES OF BLONDELIINI

- ABBREVIATA Bigot, 1889:262 (*Ceromasia*). Type-locality: "Mexico".
 AENEA Bigot, 1889:259 (*Chaetolyga*). Type-locality: "Mexico".
 ALBIDULA Wulp, 1890:59, pl. 3, figs. 6-6a (*Meigenia*). Type-locality: Mexico, Tabasco, Teapa.
 ALBOCINGULATA Wulp, 1890:148 (*Hypostena*). Type-locality: Costa Rica, Caché.
 ALBOMARGINATA Wulp, 1890:155 (*Degeeria*). Type-locality: Mexico (sev. places).
 AMERICANA Bigot, 1889:260 (*Prosopea*). Type-locality: "Mexico".
 ASSIMULATA Wulp, 1891:180 (*Telothyria*). Type-locality: Mexico, Guerrero.
 BASALIS Wulp, 1890:152 (*Degeeria*). Type-locality: Mexico, Guerrero, Omilteme.
 BIGEMINATA Curran, 1927a:10 (*Lydella*). Type-locality: West Indies, St. Croix.
 BILINEATA Wulp, 1890:112 (*Masicera*). Type-locality: Mexico, Yucatán.
 CALYPTRATA Aldrich, 1934:73 (*Phorocera*). Type-locality: Chile, Llanquihue, Lake Correntoso.
 CIRRYSOCEPHALA Bigot, 1889:261 (*Ceromasia*). Type-locality: "Mexico".
 CINCTA Walker, 1852:303 (*Tachina*). Type-locality: "Brazil".
 COMATA Wulp, 1890:177 (*Telothyria*). Type-locality: Mexico, Guerrero.
 COSTALIS Walker, 1852:344 (*Musca*). Type-locality: "South America". (preocc. Fabricius, 1794:360).
 COSTALIS Wulp, 1890:178 (*Telothyria*). Type-locality: Mexico, Guerrero and Veracruz.
 CRURALIS Giglio-Tos, 1893:7 (*Degeeria*). Type-locality: "Mexico".
 CYLINDRIVENTRIS Wulp, 1890:145 (*Hypostena*). Type-locality: Mexico, Guerrero.
 DEGEERIOIDES Wulp, 1890:209 (*Myothyria*). Type-locality: Mexico, Veracruz.
 DEPLANATA Wulp, 1890:147 (*Hypostena*). Type-locality: Costa Rica, Río Sucio.
 DICAX Giglio-Tos, 1893:7 (*Degeeria*). Type-locality: "Mexico".
 ELONGATA Robineau-Desvoidy, 1830:139 (*Medina*). Type-locality: Brazil, Paraná, Guaratuba.
 EXPERGITA Walker, 1860:304 (*Masicera*). Type-locality: "Mexico".
 FASCIATA Wulp, 1890:179 (*Telothyria*). Type-locality: Mexico, Tabasco.
 FIMBRICURA Wulp, 1890:172 (*Telothyria*). Type-locality: Mexico, Guerrero.
 FERTORIA Walker, 1861:300 (*Eurigaster*). Type-locality: "Mexico".
 FLAVESCENS Wulp, 1890:112 (*Masicera*). Type-locality: "Mexico".
 FLAVIVENTRIS Wulp, 1890:59 (*Meigenia*). Type-locality: Mexico, Orizaba.
 FORTICULA Wulp, 1890:174 (*Telothyria*). Type-locality: Mexico, Guerrero.
 GENTICA Walker, 1860:302 (*Masicera*). Type-locality: "Mexico".
 GRATIOSA Wulp, 1890:60 (*Meigenia*). Type-locality: Mexico, Guerrero, Xucumanatlán.
 INSECTA Giglio-Tos, 1893:7 (*Degeeria*). Type-locality: "Mexico".
 LEUCOCYLA Wulp, 1890:154 (*Degeeria*). Type-locality: Mexico (sev. places).
 LONGIPES Wulp, 1890:155 (*Degeeria*). Type-locality: Mexico (sev. places).
 LUGENS Wulp, 1890:177 (*Telothyria*). Type-locality: Mexico, Morelos and Veracruz.
 MELALEUCA Wulp, 1890:145, pl. 4, fig. 6 (*Hypostena*). Type-locality: Mexico, Tabasco, Teapa.
 MEXICANA Giglio-Tos, 1893:7 (*Degeeria*). Type-locality: "Mexico".
 MINIMA Wulp, 1890:148 (*Hypostena*). Type-locality: Mexico, Tabasco, Teapa.



- NEGRENSIS Aldrich, 1934:72 (*Phorocera*). Type-locality: Argentina, Rio Negro, Lake Gutiérrez.
- NERVOSA Wulp, 1890:155 (*Degeeria*). Type-locality: Mexico, Guerrero, Omilteme.
- NIGRA Bigot, 1889:259 (*Cestonia*). Type-locality: "Mexico".
- NIGRIVENTRIS Williston, 1896:259, pl. 11, fig. 96 (*Degeeria*). Type-locality: St. Vincent, W. I.
- NIGROCOSTALIS Wulp, 1890:151, pl. 4, fig. 10 (*Degeeria*). Type-locality: Mexico (sev. places).
- NUBILOSA Wulp, 1890:149 (*Hypostena*). Type-locality: Mexico, Guerrero, Amula.
- PACATA Wulp, 1890:185 (*Telothyria*). Type-locality: Mexico, Guerrero.
- PILOSA Wulp, 1890:144 (*Hypostena*). Type-locality: Mexico (sev. places).
- QUADRISTRATA Wulp, 1890:146, pl. 4, fig. 5-5a (*Hypostena*). Type-locality: Costa Rica, Volcán de Irazú.
- RAVA Wulp, 1890:178 (*Telothyria*). Type-locality: Mexico, Guerrero.
- RECONDITA Wulp, 1890:176 (*Telothyria*). Type-locality: Mexico, Guerrero.
- SCITA Walker, 1852:302 (*Tachina*). Type-locality: "Brazil".
- SPINIPES Bigot, 1889:262 (*Ceromastia*). Type-locality: "Mexico".
- STRIGATA Wulp, 1890:105 (*Masicera*). Type-locality: Mexico (sev. places).
- SUBTILIS Wulp, 1890:149 (*Hypostena*). Type-locality: Mexico, Guerrero.
- TRICHOSOMA Wulp, 1890:208 (*Myothyria*). Type-locality: Mexico, Veracruz, Atoyac.
- TRIFASCIATA Walker, 1837:350 (*Tachina*). Type-locality: no definite locality, but southern South America. Cf. Austen, 1907:329.
- TRIFURCA Wulp, 1890:175 (*Telothyria*). Type-locality: Mexico, Guerrero.
- TURBINATA Wulp, 1890:146 (*Hypostena*). Type-locality: Mexico, near Veracruz.
- VICINA Wulp, 1890:184 (*Telothyria*). Type-locality: Mexico, Guerrero and Tabasco.
- VULGATA Walker, 1852:300 (*Tachina*). Type-locality: "South America".

Tribe CALODEXIINI, New Tribe

The genus *Calodexia* Wulp placed by Townsend among of the Urodexiini represent a highly specialized group of flies which the females are commonly found associated to swarm raids of army ants.

This group is larviparous and has been recorded as parasite of Orthoptera and Blattodea. Judging by the males characters these flies go to Blondeliini. In the same species often occurs that some specimens have bare prosternum and other bristled. The most distinctive character of this tribe is the structure of the female abdomen of which the fifth tergite is convex and rounded in posterior view and the tubular ovopositor arising on the ventral side of abdomen. This group may prove to be morphologically and biologically justified and appears to merit tribal status.

Ref. — Rettenmeyer, 1961 (biol.).

Genus CALODEXIA Wulp

- CALODEXIA Wulp, 1891:257. Type-species, *majuscula* Townsend (Coquillett, 1910:517).
- Oestrogaster* Townsend, 1912a:309. Type-species, *fumosus* Townsend (orig. des.).
- Oestrogastropsis* Townsend, 1915m:424. Type-species, *mexicana* Townsend (orig. des.).

- Oestrogastroles* Townsend, 1915m:425. Type-species, *similis* Townsend (orig. des.).
- Nicetria* Reinhard, 1953a:99. Type-species, *peponis* Reinhard (orig. des.). N. SYN.
- Ref. — Curran, 1934b: (rev.); Thompson, 1968 (Key to Trinidad spp.).
- AGILIS Curran, 1934b:13. Type-locality: Panama, Canal Zone, Barro Colorado I.
- ALDRICHI Curran, 1934b:4. Type-locality: Brazil, Mato Grosso, Chapada.
- APICALIS Curran, 1934b:17. Type-locality: Panama, Canal Zone, Barro Colorado I.
- BELLA Curran, 1934b:8. Type-locality: Panama, Canal Zone, Barro Colorado I. Ref. — Thompson, 1968:119 (larvae).
- BEQUAERTI Curran, 1934b:19. Type-locality: Guatemala, Guatallon, Moca.
- CALLANI Thompson, 1968:109. Type-locality: Trinidad, St. Augustine.
- CAUDATA Curran, 1934b:16. Type-locality: Panama, Cabima.
- CONTINUA Curran, 1934b:10. Type-locality: Panama, Canal Zone, Barro Colorado I. Distr. — Guatemala, Panama.
- DIVES Curran, 1934b:6. Type-locality: Panama, Canal Zone, Barro Colorado I.
- FASCIATA Curran, 1934b:5. Type-locality: Panama, Canal Zone, Barro Colorado I.
- FLAVIPES (Schiner), 1868:326 (*Mcigenia*). Type-locality: Brazil. Ref. — Aldrich, 1929a:21.
- FULVIBASIS Curran, 1934b:7. Type-locality: Peru, Jaen, Huascaray.
- FUMOSA (Townsend), 1912a:310 (*Oestrogaster*). Type-locality: Peru, Jaen, Charape River. Distr. — Panama, Peru, Brazil. Ref. — Thompson, 1968 (larvae).
- GLOBOSA (Reinhard), 1953a:100 (*Nicetria*). Type-locality: Mexico, Veracruz, Las Tuxtlas, Volcano San Martin. N. COMB.
- GRATA (Wulp), 1890:134 (*Myobia*). Type-locality: Mexico, Guerrero. N. COMB.
- INSOLITA Curran, 1934b:506. Type-locality: Guyana, Bartica, Kartabo.
- INTERRUPTA Curran, 1934b:9. Type-locality: Panama, Canal Zone, Barro Colorado I.
- MAJOR Curran, 1934b:12. Type-locality: Panama, Canal Zone, Barro Colorado I. Distr. — Panama, Costa Rica.
- MAJUSCULA Wulp, 1891:257, pl. 6, fig. 12. Type-locality: Mexico, Tabasco, Teapa. Distr. — Mexico.
- MEXICANA (Townsend), 1915m:425 (*Oestrogastropsis*). Type-locality: Mexico, Veracruz, San Rafael.
- NIGRIPES Thompson, 1968:115, figs. 66, 68. Type-locality: Trinidad, Maracas Valley.
- OBSCURIPES Wulp, 1891:258. Type-locality: Mexico, Veracruz.
- PANAMENSIS Curran, 1934b:18. Type-locality: Panama, Canal Zone, Barro Colorado I.
- PEPONIS (Reinhard), 1953a:99 (*Nicetria*). Type-locality: Mexico, Veracruz, Las Tuxtlas, Volcano San Martin. N. COMB.
- RUBRIPES Thompson, 1968:114, figs. 65, 69. Type-locality: Trinidad, St. Joseph.
- SCURRA (Wulp), 1890:134 (*Myobia*). Type-locality: Mexico, Guerrero, Amula. N. COMB.
- SIMILIS (Townsend), 1915m:425 (*Oestrogastroles*). Type-locality: Panama, Arraigán (as "Araján"). Distr. — Mexico (Veracruz), Panama.

- TOWNSENDI Curran, 1934b:7. Type-locality: Peru, Jaen, Huascaray.
 VALERA Curran, 1934b:19. Type-locality: Venezuela, Valera.
 VARIA Curran, 1934b:20. Type-locality: Panama, Canal Zone, Barro Colorado I.
flavipes of Aldrich, not Schiner. Distr. — Panama, Costa Rica.
 VENTERIS Curran, 1934b:15. Type-locality: Panama, Canal Zone, Barro Colorado I. Distr. — Panama, Costa Rica.

Genus EUOESTROGASTER Townsend

- EUOESTROGASTER Townsend, 1935d:224. Type-species, *fuscus* Townsend (orig. des.).
 FUSCUS Townsend, 1935d:224. Type-locality: Guyana, Pacaraima Mts., upper Ireng River.

Genus EUOESTROGASTRODES Townsend

- EUOESTROGASTRODES Townsend, 1935d:224. Type-species, *flavescens* Townsend (orig. des.).
 FLAVESCENS Townsend, 1935d:224. Type-locality: "Trinidad".

Tribe NEOMINTHOINI

Genus ASILIDOTACHINA Townsend

- ASILIDOTACHINA Townsend, 1931a:461. Type-species, *elongata* Townsend (orig. des.).
 ELONGATA Townsend, 1931a:462. Type-locality: Peru, Oxapampa.

Genus CRYPTOCLADOCERA Bezzi

- CRYPTOCLADOCERA Bezzi, 1923:653. Type-species, *prodigiosa* Bezzi (orig. des.).
 Ref. — Arnaud, 1963b (key).
 BEZZII Arnaud, 1963b:9, figs. 1, 2-4, 7-8. Type-locality: Panama, Canal Zone, Barro Colorado I.
 MOJINGENSIS Arnaud, 1963b:13, figs. 1, 5, 11, 12. Type-locality: Panama, Fort Sherman, Mogina Swamp.
 PICHILINGUENSIS Arnaud, 1963b:17, figs. 1, 6, 9, 10. Type-locality: Ecuador, Los Rios, Pichilingue.
 PRODIGIOSA Bezzi, 1923:655, figs. 5, 7. Type-locality: Surinam.

Genus CEROTACHINA Arnaud

- CEROTACHINA Arnaud, 1963b:20. Type-species, *elegantula* Arnaud (orig. des.).
 ALBULA Arnaud, 1963b:30, figs. 1, 24. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
 ELEGANTULA Arnaud, 1963b:22, figs. 1, 13, 23. Type-locality: Brazil, Santa Catarina, Nova Teutônia.

Genus EUPELECOTHECA Townsend

- EUPELECOTHECA Townsend, 1919b:169. Type-species, *celer* Townsend (orig. des.).
CYLINDRATA (Wulp), 1892:82 (*Phorocera*). Type-locality: Mexico, Guerrero, Chilpancingo. N. comb.

Genus EUTHELAIIRA Townsend

- EUTHELAIIRA Townsend, 1912a:305. Type-species, *inambarica* Townsend (orig. des.).
INAMBARICA Townsend, 1912a:306. Type-locality: Peru, Inambari River.
RUFILABRIS (Wulp), 1890:70 (*Phorocera*). Type-locality: Mexico, Guerrero.

Genus ITEUTHELAIIRA Townsend

- ITEUTHELAIIRA Townsend, 1927a:264. Type-species, *intermedia* Townsend (orig. des.) = *esuriens* (Fabricius).
CHAETOSA Townsend, 1929:369. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
ESURIENS (Fabricius), 1805:301 (*Musca*). Type-locality: "South America". Ref. — Townsend, 1931b:170, 468.
intermedia Townsend, 1927a:323. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus NEOMINTHO Brauer & Bergenstamm

- NEOMINTHO Brauer & Bergenstamm, 1891:339 (1891:35). Type-species, *Tachina macilentata* Wiedemann (Townsend, 1916:8).
Euthelairopsis Townsend, 1927a:258. Type-species, *brasilensis* Townsend (orig. des.).
MACILENTA (Wiedemann), 1830:305 (*Tachina*). Type-locality: Brazil. Ref. — Aldrich, 1927c:19 (type redescr., genitalia).
brasilensis Townsend, 1927a:309 (*Euthelairopsis*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus NEOMINTHOIDEA Thompson

- NEOMINTHOIDEA Thompson, 1968:30. Type-species, *trinidadensis* Thompson (orig. des.).
TRINIDADENSIS Thompson, 1968:30, fig. 10. Type-locality: Trinidad, Brazil (village).

Genus PELECOTHECA Townsend

- PELECOTHECA Townsend, 1919b:168. Type-species, *panamensis* Townsend (orig. des.).
Adercomyia Arnaud, 1963b:32. Type-species, *sabroskyi* Arnaud (orig. des.).
N. syn.

- BISETA (Arnaud), 1963b:40, figs. 1, 33, 34 (*Adercomyia*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. N. comb.
- FLAVIPES Thompson, 1968:46, figs. 22-24. Type-locality: Trinidad, Maracas Valley.
- MACILENTA (Wulp), 1890:122 (*Prospberrya*). Type-locality: Mexico, Orizaba. Ref. — Arnaud, 1963b:52.
- MACRA (Wulp), 1890:84 (*Phorocera*). Type-locality: Mexico, Tabasco, Teapa. N. comb. Ref. — Arnaud, 1963b:53.
- PANAMENSIS Townsend, 1919b:169. Type-locality: Panama, Canal Zone, Las Cascadas. Ref. — Arnaud, 1963b:47.
- PAULENSIS Townsend, 1929:369. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- SABROSKYI (Arnaud), 1963b:34, figs. 1, 25, 32 (*Adercomyia*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. N. comb.
- TRINIDADENSIS Thompson, 1968:44. Type-locality: Trinidad, Floradale.

Genus SYNORBITALIA Thompson

- SYNORBITALIA Thompson, 1968:32. Type-species, *flavipes* Thompson (orig. des.).
- FLAVIPES Thompson, 1968:32. Type-locality: Trinidad, "Simla".

Genus THELAIROCHAETONA Townsend

- THELAIROCHAETONA Townsend, 1919a:559. Type-species, *thrix* Townsend (orig. des.).
- THRIX Townsend, 1919a:559. Type-locality: Panama, Porto Bello.

Genus VIVIANOIDEA Thompson

- VIVIANOIDEA Thompson, 1968:37. Type-species, *grisea* Thompson (orig. des.).
- GRISEA Thompson, 1968:37, fig. 11. Type-locality: Trinidad, Simla.

Tribe EXORISTINI

(including Phorocerini, and Phoriniini of Townsend)

The species of the tribe Exoristini parasitizes a great variety of lepidopterous larvae, some attack tenthredinid larvae.

Genus EUPHOROCERA Townsend

- EUPHOROCERA Townsend, 1892a:112. Type-species, *tachinomoides* Townsend (orig. des.).
- Plagiotachina* Townsend, 1927a:261. Type-species, *peruviana* Townsend (orig. des.).
- BISERIALIS (Schiner), 1868:326 (*Phorocera*). Type-locality: "South America". Ref. — Aldrich, 1927c:22 (type redes.).
- APPENDICULATA (Wulp), 1890:85 (*Phorocera*). Type-locality: Mexico, Guerrero. N. comb.



- FALTEATA (Wulp), 1890:124 (*Prospberrya*). Type-locality: Mexico, Guerrero. N. comb.
- CARIDEI (Brèthes in Massini & Brèthes), 1918:214, 3 figs., 1 col. plate (*Paraxorista*). Type-locality: Argentina, Córdoba. Ref. — Blanchard, 1935:11.
- CINEHEA (Wulp), 1890:81 (*Phorocera*). Type-locality: Brazil. N. comb.
- CLARIPENNIS (Macquart), 1840:209, pl. 5, fig. 8 (*Phorocera*). Type-locality: U.S.A. Distr. — Alaska, Canada.
- CUMUTOENSIS Thompson, 1968:16, fig. 4. Type-locality: Trinidad, Mexico (distr. confused with *edwardsii*).
- EUROTAE Blanchard, 1937:53, figs. 8 a-e. Type-locality: Argentina, Corrientes.
- FLORIDENSIS Townsend, 1916d:217. Type-locality: U.S.A., Florida. Distr. — U.S.A., Mexico.
- HAYWARDI (Blanchard), 1947a:16, figs. 7 a-f (*Plagiotachina*). Type-locality: Argentina, Buenos Aires, El Tigre. Distr. — Uruguay, Argentina.
- HELICONUNARUM (Townsend), 1929:373 (*Plagiotachina*). Type-locality: Peru, Santa Eulalia, Piura River Valley.
- MINOR Townsend, 1912a:303. Type-locality: Peru, Piura and Chaipara.
- NOERA (Reinhard), 1957:109 (*Phorocera*). Type-locality: Mexico, Cuernavaca. N. comb.
- ORIENTALIS (Townsend), 1929:373 (*Plagiotachina*). Type-locality: Peru, Ucayali River, Yarinacocha.
- PERUVIANA Townsend, 1912:303. Type-locality: Peru, Piura.
- PLATENSIS (Brèthes), 1922:21 (*Parasetigena*). Type-locality: Argentina, Buenos Aires. Unrecognized. Ref. — Blanchard, 1935:12.
- SCUTELLARIS (Wulp), 1890:85 (*Phorocera*). Type-locality: Mexico, Guerrero.
- TACHINOMOIDES Townsend, 1892:112. Type-locality: U.S.A. Distr. — U.S.A., Mexico.
- TOWNSENDI Guimarães, nom. nov. for *peruviana* Townsend. Type-locality: Peru, Casma.
- peruviana* Townsend, 1927a:348 (*Plagiprospherysa*; preocc. Townsend, 1912). Type-locality: Peru, Casma.
- VIVIDA (Wiedemann), 1830:312 (*Tachina*). Type-locality: Brazil. Ref. — Aldrich, 1925d:468, fig. 1.

Genus EUPHOROCEROPSIS Townsend

- EUPHOROCEROPSIS Townsend, 1917a:49. Type-species, *alba* Townsend (orig. des.).
- ALBA Townsend, 1917a:50. Type-locality: Mexico, Tampico.

Genus EPIPLAGIOPS Blanchard

- EPIPLAGIOPS Blanchard, 1943d:450. Type-species, *littoralis* Blanchard (orig. des.).
- LITTORALIS Blanchard, 1943d:451, figs. a-f. Type-locality: Argentina, Santa Fé. Ref. — Parker, et al., 1950:16 (immature stages, biology).

Genus GOPAULIA Thompson

- GOPAULIA Thompson, 1968:70. Type-species, *nigrifrons* Thompson (orig. des.).
- NIGRIFRONS Thompson, 1968:7, figs. 39-41. Type-locality: Trinidad, "Nursery".

Genus MACROHOUGHIOPSIS Townsend

MACROHOUGHIOPSIS Townsend, 1927a:261. Type-species, *similis* Townsend (orig. des.).

SIMILIS Townsend, 1927a:325. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus METAPHOROCERA Thompson

METAPHOROCERA Thompson, 1968:18. Type-species, *maracasi* Thompson (orig. des.).

MARACASI Thompson, 1968:18, fig. 5. Type-locality: Trinidad, Curepe.

Genus MYERSIMYIA Townsend

MYERSIMYIA Townsend, 1935d:221. Type-species, *mixta* Townsend (orig. des.).

MIXTA Townsend, 1935d:221. Type-locality: Trinidad. Ref. — Thompson, 1968:128.

Genus PALPEXORISTA Townsend

PALPEXORISTA Townsend, 1926:28. Type-species, *phoroceroideis* Townsend (orig. des.) = *imitator* (Aldrich & Webber).

Yahuartachina Townsend, 1927a:261. Type-species, *yahuarphrynoideis* Townsend (orig. des.).

HEROS (Schiner), 1868:325 (*Phorocera*). Type-locality: Brazil, Rio de Janeiro. Ref. — Aldrich, 1927:21.

LONGIUSCULA (Walker), 1857:198 (*Masicera*). Type-locality: "South America".

MERIDIONALIS Thompson, 1968:21, figs. 6, 7. Type-locality: Trinidad, Mt. St. Benedict.

MINOR Thompson, 1968:24. Type-locality: Trinidad, Santa Cruz Valley.

RUSTI (Aldrich), 1929:3 (*Phorocera*). Type-locality: Argentina, Tucumán.

YAHUARPHRYNOIDES (Townsend), 1927a:369 (*Yahuartachina*). Type-locality: Peru, Yahuarmayo.

Genus PARAMESOCHAETA Brauer & Bergenstamm

PARAMESOCHAETA Brauer & Bergenstamm, 1891:341 (1891:37). Type-species, *Mystacella fuscicostalis* Wulp (mon.).

FUSCICOSTALIS (Wulp), 1890:57, pl. 3, fig. 5a (*Mystacella*). Type-locality: Mexico and Costa Rica.

Genus PARASÉTIGENA Brauer & Bergenstamm

PARASÉTIGENA Brauer & Bergenstamm, 1891:339, 401 (1891:35, 97). Type-species, *Chaetona segregata* Rondani (mon.; misident) = *agilis* (Robineau-Desvoidy).

HICHINSI Cortés, 1967b:13. Type-locality: Chile, Santiago, Maipú, La Rinconada.

Genus PROSPALAEA Aldrich

PROSPALAEA Aldrich, 1925c:111. Type-species, *Prosopaea insularis*, Brauer & Bergenstamm (mon.).

INSULARIS (Brauer & Bergenstamm), 1891:334 (1891:30) (*Prosopaea*)
Type-locality: West Indies, St. Thomas.

Genus STOMATOMYIA Brauer & Bergenstamm

STOMATOMYIA Brauer & Bergenstamm, 1889:98 (1889:30). Type-species, *Chaetona filipalpis* Rondani (mon.).

Plagiprospherysa Townsend, 1892:113. Type-species, *valida* Townsend (orig. des.) = *parvipalpis* (Wulp).

filipalpis (Blanchard) in Denier, 1939:560 (*Plagiprospherysa*). *Nomen nudum*.

MERIDIONALIS (Townsend), 1912a:304 (*Plagiops*). Type-locality: Peru, Piura.
PARVIPALPIS (Wulp), 1890:124, pl. 3, figs. 15-15a (*Prospherysa*). Type-locality: Mexico, Guerrero (sev. places).

PUER (Williston), 1896:354, pl. 11, fig. 91 (*Phorocera*). Type-locality: St. Vincent, W. I.

TRINITATIS Thompson, 1968:91. Type-locality: "Trinidad".

Genus STOMATOTACHINA Townsend

STOMATOTACHINA Townsend, 1931a:464. Type-species, *splendida* Townsend (orig. des.) = *porteri* (Brèthes).

PORTERI (Brèthes), 1920:12, fig. 1 (*Parasetigena*). Type-locality: Chile.

splendida Townsend, 1931a:464. Type-locality: Chile, Concepción.

Genus TACHINOMYIA Townsend

TACHINOMYIA Townsend, 1892:96. Type-species, *robusta* Townsend (orig. des.) = *panaetius* (Walker).

VARIATA Curran, 1926b:169. Type-locality: Canada, Ontario. Distr. — Canada, U.S.A., Mexico.

Genus TETRAGRAPHIA Brauer & Bergenstamm

TETRAGRAPHIA Brauer & Bergenstamm, 1891:351 (1891:47). Type-species, *tessellata* Brauer & Bergenstamm (mon.).

TESSELLATA Brauer & Bergenstamm, 1891:351 (1891:47). Type-locality: Cuba.

UNRECOGNIZED SPECIES OF EXORISTINI

ELEPHARIPUS Brauer & Bergenstamm, 1891:342 (1891:38) (*Ctenophorocera*). Type-locality: unknown.

BOTYVORA Robineau-Desvoidy, 1830:138 (*Phorocera*). Type-locality: "Cuba".

BRUNISQUAMIS Robineau-Desvoidy: 1830:157 (*Elaphoria*). Type-locality: "Brazil".



- CASANUEVAI Cortés, 1945 (*Phorocera*). Type-locality: Chile, Valparaíso.
- CHRYSOSELUS Walker, 1852:296 (*Tachina*). Type-locality: "Brazil".
- CHRYSOSELUS Walker, 1852:329 (*Sarcophaga*). Type-locality: "South America". Ref. — Aldrich, 1930:19.
- CIRRATA Robineau-Desvoidy, 1830:138 (*Phorocera*). Type-locality: Brazil.
- ELISAE Cortés, 1945:162 (*Phorocera*). Type-locality: Chile, Angol, Malleco.
- GUIANICA Curran, 1934c:513 (*Phorocera*). Type-locality: Guyana, Bartica, Kartabo.
- IMMACULATA Wulp, 1890:82 (*Phorocera*). Type-locality: Mexico, Guerrero.
- NIGRIFRONS Wulp, 1890:81 (*Phorocera*). Type-locality: Mexico, Guerrero.
- PERUVIANA Townsend, 1912:303 (*Euphorocera*). Type-locality: Peru, Piura.
- SETIGERA Wulp, 1890:84 (*Phorocera*). Type-locality: Mexico, Veracruz and Tabasco.
- SOBRINA Wulp, 1890:84 (*Phorocera*). Type-locality: Mexico, Tabasco.
- TANUMEANA Townsend, 1931:472 (*Macropatelloa*). Type-locality: Chile, Tanumé and Talagante.
- TENUISETA Macquart, 1846:294 (1846:166), pl. 14, fig. 6 (*Phorocera*). Type-locality: French Guiane, Cayenne.
- VENUSA Curran, 1928:29 (*Phorocera*). Type-locality: Jamaica, Hope Gardens.

Tribe EUMASICERINI

(Trypherini, in part)

Genus AGRARIALIA Curran

- AGRARIALIA Curran, 1934d:459. Type-species, *sexualis* Curran (orig. des.).
- SEXUALIS Curran, 1934d:469, figs. 160, 179. Type-locality: Panama, Corozal, Canal Zone (cf. Arnaud, 1963:106).

Genus APLOMYODORIA Townsend

- APLOMYODORIA Townsend, 1928c:160. Type-species, *arida* Townsend (orig. des.).
- ARIDA Townsend, 1928c:161. Type-locality: Peru, Chosica Canyon.

Genus ARGYROCHAETONA Townsend

- ARGYROCHAETONA Townsend, 1919b:167. Type-species, *cubana* Townsend (orig. des.).
- CUBANA Townsend, 1919b:167. Type-locality: Cuba, Santiago de las Vegas. Ref. — Thompson, 1963b:342.
- PERUANA Townsend, 1928c:161. Type-locality: Peru. Trujillo, Cartavio.

Genus ARGYROPHYLAX Brauer & Bergenstamm

- ARGYROPHYLAX Brauer & Bergenstamm, 1889:163 (1889:95). Type-species, *Tachina albincisa* wiedemann (mon.).
- Ref. — Curran, 1928:43 (key to Jamaican sp.).

- ALBINCISA (Wiedemann), 1830:334 (*Tachina*). Type-locality: West Indies, St. Thomas. Ref. — Walton, 1916:189 (redesc.).
- EISETOSA Thompson, 1963b:351, figs. 48-49. Type-locality: "Trinidad" (no type designated).
- GOWDEYI Curran, 1928a:44. Type-locality: Jamaica, Hope Gardens.
- PURPURESCENS Townsend, 1929:375. Type-locality: Peru, Pucalpa, Rio Ucayali. Ref. — Thompson, 1963b:346.
- TRIANGULIFERA Thompson, 1963b:352, figs. 50-51. Type-locality: "Trinidad" (no type designated).
- TRISSETOSA Curran, 1928a:43. Type-locality: Jamaica, Hope Gardens.

Genus CYLINDROMASICERA Townsend

- CYLINDROMASICERA Townsend, 1915d:61. Type-species, *prima* Townsend (orig. des.).
- PRIMA Townsend, 1915d:62. Type-locality: Peru, Sullana.

Genus EULOEWIODORIA Townsend

- EULOEWIODORIA Townsend, 1927a:260. Type-species, *eulalia* Townsend (orig. des.).
- EULALIA Townsend, 1927a:308. Type-locality: Peru, Santa Eulalia.

Genus EURYTHEMYIA Reinhard

- Eurylochus* Reinhard, 1967a:108. Type-species, *dissitus* Reinhard (orig. des.) (preocc. Torre-Bueno, 1914).
- EURYTHEMYIA Reinhard, 1967b:600 (nom. nov. for *Eurylochus* Reinhard). Type-species, *Eurylochus dissitus* (aut.).
- DISSITUS (Reinhard), 1967a:109 (*Eurylochus*). Type-locality: Mexico, Veracruz, Jalapa.

Genus EUMASICERA Townsend

- EUMASICERA Townsend, 1909:249. Type-species, *coccidela* Townsend (orig. des.) = *sternalis* (Coquillett).
- BISTRIGATA (Wulp), 1890:109 (*Masicera*). Type-locality: Mexico, Tabasco, Teapa.
- CALCARATA (Wulp), 1890:114 (*Masicera*). Type-locality: Mexico, Morelos, Cuernavaca.
- IMPEDITA (Wulp), 1890:107 (*Masicera*). Type-locality: Mexico, Cuernavaca.
- OPTICA (Schiner), 1868:327 (*Exorista*). Type-locality: Brazil. Ref. — Aldrich, 1925:465.
- SORDIDA (Wulp), 1890:113 (*Masicera*). Type-locality: Mexico, sev. places.
- TENUISETA (Macquart), 1846:292, pl. 15, fig. 4 (*Masicera*). Type-locality: Venezuela.

Genus HYPERTROPHOMMA Townsend

HYPERTROPHOMMA Townsend, 1915k:99. Type-species, *opacum* Townsend (orig. des.).

SUBITA Reinhard, 1958:240 (as *Hypertrophomma*, error). Type-locality: Mexico, Morelos, Huajitlan.

Genus METOPIOPS Townsend

METOPIOPS Townsend, 1912a:338. Type-species, *mirabilis* Townsend (orig. des.)? = *Pseudochaeta*.

MIRABILIS Townsend, 1912a:339. Type-locality: Peru, Valley of the Rio.

Genus OROHOUGHIA Townsend

OROHOUGHIA Townsend, 1934a:403. Type-species, *aurata* Townsend (orig. des.).

AURATA Townsend, 1934a:403. Type-locality: Brazil, Boa Vista, Tapajós River.

Genus PARARRHINACTIA Townsend

PARARRHINACTIA Townsend, 1935d:232. Type-species, *parva* Townsend (orig. des.).

PARVA Townsend, 1935d:232, 233. Type-locality: Trinidad.

Genus PHAENOPSIS Townsend

PHAENOPSIS Townsend, 1912a:362. Type-species, *arabella* Townsend (orig. des.).

Dimasicera Townsend, 1915d:62. Type-species, *nitida* Townsend (orig. des.).

ARABELLA Townsend, 1912a:363. Type-locality: Peru, Sullana.

NITIDA (Townsend), 1915d:64 (*Dimasicera*). Type-locality: Peru, Sullana.

Genus PSEUDOCOAETA Coquillett

PSEUDOCOAETA Coquillett, 1895:309. Type-species, *argentifrons* Coquillett (orig. des.). Ref. — Reinhard, 1946 (rev.), Thompson, 1964:100 (Key to, Trinidad. sp.).

CUREPEI Thompson, 1964:114, fig. 9. Type-locality: Trinidad.

FLAVIPALPIS Thompson, 1964:101, figs. 4, 7. Type-locality: Trinidad, St. Augustine.

LATITARSUS Thompson, 1964:111, fig. 1. Type-locality: Trinidad, St. Benedict. Mt.

NITENS Thompson, 1964:108, figs. 3, 5. Type-locality: Trinidad, Orange Grove.

SYNGAMIAE Thompson, 1964:105, figs. 2, 8. Type-locality: Trinidad, Curepe.

TRINITATIS Thompson, 1964:117, fig. 10. Type-locality: Trinidad, Caroni.



Genus SANTACRUZIOPSIS Thompson

- SANTACRUZIOPSIS Thompson, 1968:163. Type-species, *fumipennis* Thompson (orig. des.).
CLARIPENNIS Thompson, 1968:165, figs. 103, 105. Type-locality: Trinidad, Santa Augustine.
FUMIPENNIS Thompson, 1968:163, figs. 102, 104. Type-locality: Trinidad, Santa Cruz Valley.

Genus THELAIRALIA Curran

- THELAIRALIA Curran, 1934d:462. Type-species, *fasciata* Curran (orig. des.).
FASCIATA Curran, 1934d:470, figs. 151, 157. Type-locality: Panama, Canal Zone, Corozal (cf. Arnaud, 1963:130).

UNPLACED SPECIES OF EUMASICERINI

- APPROXIMATA Wulp, 1890:202 (*Anisia*). Type-locality: Mexico, Guerrero.
ARGENTICEPS Wulp, 1890:135 (*Myobia*). Type-locality: Mexico, Guerrero.
MINUTA Wulp, 1890:123 (*Prosphegysa*). Type-locality: Mexico.

Tribe SIPHONINI

(Crocutini, Actini)

The hosts of Siphonini with few exceptions are various Microlepidoptera.

Genus ACRONARISTA Townsend

- ACRONARISTA Townsend, 1908:85. Type-species, *mirabilis* Townsend (orig. des.).
ADUNATA Reinhard, 1961:206. Type-locality: Mexico, Tehuacana, Puebla.

Genus ACRONARISTOPSIS Townsend

- ACRONARISTOPSIS Townsend, 1919b:178. Type-species, *bahamensis* Townsend (orig. des.).
BAHAMENSIS Townsend, 1919b:178. Type-locality: Bahamas I., Nassau.

Genus ACTIA Robineau-Desvoidy

- ACTIA Robineau-Desvoidy, 1830:85. Type-species, *pilipennis* Robineau-Desvoidy (van Endem, 1954:63; suspension of the rules required (cf. Sabrosky & Arnaud, 1965:1061) = *lamia* (Meigen).
Uruactia Townsend, 1927a:256. Type-species, *uruhuasi* Townsend (orig. des.).
Ref. — Aldrich, 1934:131 (key to Patagonian sp.).
AMBLYCERA Aldrich, 1934:132. Type-locality: Argentina, Rio Negro, Bariloche.
AMERICANA (Townsend), 1892c:69 (*Tryptocera*). Type-locality: U.S.A. District of Columbia. Distr. — U.S.A., Mexico.



- CORNUTA Aldrich, 1934:131. Type-locality: Chile, Angol. Distr. — South Chile and South Argentina.
- INTERRUPTA Curran, 1933:6. Type-locality: U.S.A., New York. Distr. — U.S.A., Mexico.
- NITIDIVENTRIS Curran, 1933:4. Type-locality: Panama, Canal Zone, Barro Colorado I.
- PANAMENSIS Curran, 1933:3. Type-locality: Panama, Canal Zone, Barro Colorado I.
- URUHUASI (Townsend), 1927a:364 (*Uruactia*). Type-locality: Peru, Uruhuasi. Ref. — Aldrich, 1934:130.

Genus ACTINACTIA Townsend

- ACTINACTIA Townsend, 1927a:248. Type-species, *lutea* Townsend (orig. dés.).
- LUTEA Townsend, 1927a:283. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus ACTINOCROCUTA Townsend

- ACTINOCROCUTA Townsend, 1935d:228. Type-species, *chaetosa* Townsend (orig. dés.).
- CHAETOSA Townsend, 1935d:228. Type-locality: "Trinidad".

Genus ALDRICHIOPA Guimarães, nom. nov.

- Aphelogaster* Aldrich, 1934:22. Type-species, *coracella* Aldrich (orig. des.) (preocc. Kolbe, 1897).
- ALDRICHIOPA Guimarães (n. name for *Aphelogaster* Aldrich). Type-species, *Aphelogaster coracella* Aldrich (aut.).
- CORACELLA (Aldrich), 1934:23, fig. 4 (*Aphelogaster*). Type-locality: Argentina, Lake Gutiérrez. Distr. — South Argentina.

Genus AUSTROPHYTOMYPTERA Blanchard

- AUSTROPHYTOMYPTERA Blanchard, 1962:252. Type-species, *malloi* Blanchard (orig. des.).
- MALLOI Blanchard, 1962:253, fig. 2. Type-locality: Argentina, Córdoba, Cruz del Eje.

Genus BOLBOCHAETA Bigot

- BOLBOCHAETA Bigot, 1885c:xliv. Type-species, *haustellata* Bigot (mon.).
- HAUSTELLATA Bigot, 1885c:xliv. Type-locality: Argentina, Buenos Aires. Ref. — Brauer, 1897:368 (notes on types).

Genus BORGMEIERMYIA Townsend

- BORGMEIERMYIA Townsend, 1935a:293. Type-species *brasiliانا* Townsend (orig. des.). Ref. — Arnaud, 1963d:5 (rev.).



- BRASILIANA Townsend, 1935a:293, figs. 1, 2. Type-locality: Brazil, Rio de Janeiro, Guanabara. Ref. — Arnaud, 1963d:5. Distr. — Brazil (Rio de Janeiro, Santa Catarina).
- PERUANA Arnaud, 1963d:10, fig. 11. Type-locality: Peru, Tingo Maria, Monson Valley.
- ROZENI Arnaud, 1963d:12, figs. 1-4, 6, 7, 9, 10, 15-17. Type-locality: Brazil, Santa Catarina, Nova Teutônia.

Genus CAMPOSODES Cortés

- CAMPOSODES Cortés, 1967a:4. Type-species, *evanescens* Cortés (orig. des.).
- EVANESCENS Cortés, 1967a:4, figs. 1-2. Type-locality: Chile, Santiago, Maipú, Quebrada de la Plata. Distr. — Chile (Santiago, Coquimbo, Valparaíso).

Genus CEROMYA Robineau-Desvoidy

- CEROMYA Robineau-Desvoidy, 1830:86. Type-species, *testacea* Robineau-Desvoidy (Coquillett, 1910:520) = *bicolor* (Meigen).
- SUBOPACA (Aldrich), 1934:133 (*Actia*). Type-locality: Argentina, Rio Negro, Bariloche. N. comb.
- UNICOLOR (Aldrich), 1934:133 (*Actia*). Type-locality: Argentina, Rio Negro, Bariloche. N. comb.

Genus CLAUSICELLA Rondani

- CLAUSICELLA Rondani, 1856:61. Type-species, *suturata* Rondani (orig. des., as gen., sp. n.).
- Epigrimyia*, authors in part, not Townsend.
- Ref. — Reinhard, 1946 (rev.).
- TOWNSENDI (Curran), 1931:22 (*Epigrimyia*). Type-locality: Puerto Rico, Isabela Substation.

Genus CLAUSICELLANA Curran

- CLAUSICELLANA Curran, 1927c:12. Type-species, *mitis* Curran (orig. des.).
- MITIS Curran, 1927a:12. Type-locality: Puerto Rico, Albonito.

Genus CORACOMYIA Aldrich

- CORACOMYIA Aldrich, 1934:21. Type-species, *crassicornis* Aldrich (orig. des.).
- CRASSICORNIS Aldrich, 1934:22. Type-locality: Argentina, Lake Nahuel Huapi.

Genus CROCINOSOMA Reinhard

- CROCINOSOMA Reinhard, 1947:20. Type-species, *cornuale* Reinhard (orig. des.).
- CORNUALE Reinhard, 1947:21. Type-locality: U.S.A., Distr. — U.S.A., Mexico.

Genus DELTOCEROMYIA Townsend

DELTOCEROMYIA Townsend, 1931a:477. Type-species, *delta* Townsend (orig. des.).

DELTA Townsend, 1931a:478. Type-locality: Peru, Oxapampa trail at Rio Paucartambo crossing.

Genus DOLICHOPALPELLUS Townsend

DOLICHOPALPELLUS Townsend, 1927a:258. Type-species, *mirabilis* Townsend (orig. des.).

MIRABILIS Townsend, 1927a:302. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus DREPANOGLOSSA Townsend

DREPANOGLOSSA Townsend, 1891:377. Type-species, *lucens* Townsend (orig. des.).

AMYDRIAE Townsend, 1908:76. Type-locality: Mexico, Cuernavaca.

Genus ELFIA Robineau-Desvoidy

ELFIA Robineau-Desvoidy, 1850:190. Type-species, *Actia cingulata* Robineau-Desvoidy (Robineau-Desvoidy, 1863:672).

AENEA (Coquillett), 1895d:57 (*Hypostena*). Type-locality: U.S.A. Distr. — U.S.A., Mexico.

ATRA (Aldrich), 1934:78 (*Lispidea*). Type-locality: Chile, Casa Panguc.

FRONTALIS (Aldrich), 1934:80 (*Lispidea*). Type-locality: Argentina, Rio Grande, Estancia Viamonte, Tierra del Fuego.

INTERRUPTA (Aldrich), 1934:79 (*Lispidea*). Type-locality: Chile, Ancud.

ROTUNDATA (Aldrich), 1934:79 (*Lispidea*). Type-locality: Argentina, Rio Negro, Lake Nahuel Huapi.

TRIANGULARIS (Aldrich), 1934:76, figs. 11-12 (*Lispidea*). Type-locality: Argentina, Lake Correntoso.

VIBRISATA (Aldrich), 1934:78 (*Lispidea*). Type-locality: Argentina, Lake Correntoso.

Genus EPICORONIMYIA Blanchard

EPICORONIMYIA Blanchard, 1940:245. Type-species, *Epigrymyia mundelli* Blanchard (orig. des.).

MUNDELLI (Blanchard), 1935:8, figs. 2-6 (*Epigrymyia*). Type-locality: Argentina, Santiago del Estero.


Genus GINGLYMIA Townsend

GINGLYMIA Townsend, 1892a:118. Type-species, *acirostris* Townsend (orig. des.).

Ginglymyia, emend.

DEVIA Reinhard, 1962a:221. Type-locality: Mexico, Chiapas, Ocosingo.

FRACIDA Reinhard, 1962a:220. Type-locality: Mexico, Colima.



Genus HAYWARDIAMYIA Blanchard

HAYWARDIAMYIA Blanchard, 1955:23. Type-species, *brevicornis* Blanchard (orig. des.).

BREVICORNIS Blanchard, 1955:23, figs. a-j. Type-locality: Argentina, Tucumán.

Genus IRWINIA Cortés

IRWINIA Cortés, 1967a:7. Type-species, *pollinosa* Cortés (orig. des.).

POLLINOSA Cortés, 1967a:7, figs. 3-4. Type-locality: Chile, Coquimbo, Bosque de Fray Jorge, 15 km SW of Pachingo.

Genus LASIONEURA Coquillett

LASIONEURA Coquillett, 1895d:50. Type-species, *johnsoni* Coquillett (orig. des.).

DEXTELLA Reinhard, 1953b:95. Type-locality: Mexico, Morelos, El Zarco.

JOHNSONI Coquillett, 1895d:50. Type-locality. U.S.A., Washington. Distr. — U.S.A., Mexico.

Genus LISPIDEOSOMA Reinhard

LISPIDEOSOMA Reinhard, 1943c:164. Type-species, *flavipes* Reinhard (orig. des.).

TRISTE Reinhard, 1961:208. Type-locality: Mexico, west slope of Paso del Cortez.

Genus LYGAEOMYIA Aldrich

LYGAEOMYIA Aldrich, 1934:143. Type-species, *tristis* Aldrich (orig. des.).

TRISTIS Aldrich, 1934:144, fig. 18. Type-locality: Argentina, Lake Correntoso, Distr. — South Chile, South Argentina.

Genus MAYOSCHIZOCERA Townsend

MAYOSCHIZOCERA Townsend, 1927a:380. Type-species, *ramata* Townsend (orig. des.).

RAMATA Townsend, 1927a:381. Type-locality: Peru, Mayo River near Huanguera.


Genus NEOCRASPEDOTHRIX Townsend

NEOCRASPEDOTHRIX Townsend, 1927a:257. Type-species, *nova* Townsend (orig. des.) (as *nana* Townsend, 1927:333, misprint, corrected in Errata).

NOVA Townsend, 1927a:333. Type-locality: Peru, Huascaray.

Genus PARAPOLIOPS Blanchard

PARAPOLIOPS Blanchard, 1957:45. Type-species, *grioti* Blanchard (orig. des.).



GRIOTI Blanchard, 1957:45. Type-locality: Argentina, Buenos Aires, José C. Paz.

Genus PHASIOSTOMA Townsend

PHASIOSTOMA Townsend, 1915g:225. Type-species, *aristale* Townsend (orig. des.).

SAXATHIS Reinhard, 1952a:1. Type-locality: Mexico, Rio Frio.

Genus PHANTASIOSIPHONA Townsend

PHANTASIOSIPHONA Townsend, 1915h:92. Type-species, *tropica* Townsend (orig. des.).

KUSCHELI Cortés, 1952:110. Type-locality: Juan Fernandes I., Masaticrra, Cerro Aito.

TROPICA Townsend, 1915h:93. Type-locality: Mexico, Veracruz, Jicaltepec, San Rafael.

Genus PLANOMYIA Aldrich

PLANOMYIA Aldrich, 1934:129. Type-species, *browni* Aldrich (orig. des.).

BROWNI Aldrich, 1934:129. Type-locality: Chile, Concepción.

Genus PLECTOPSIS Townsend

PLECTOPSIS Townsend, 1927a:257. Type-species, *palpalis* Townsend (orig. des.).

PALPALIS Townsend, 1927a:349. Type-locality: Brazil, São Paulo, Cantareira.

Genus POLIOPS Aldrich

POLIOPS Aldrich, 1934:94. Type-species, *striatus* Aldrich (orig. des.).

AURATUS Campos, 1953:27. Type-locality: Chile, Concepción, Tomé.

STRIATUS Aldrich, 1934:94. Type-locality: Argentina, Rio Negro, Lake Nahuel Huapi. Distr. — South Argentina, South Chile.

Genus PROSENACTIA Blanchard

PROSENACTIA Blanchard, 1940:237. Type-species, *liebermanni* Blanchard (orig. des.).

LIEBERMANNI Blanchard, 1940:238, fig. 4. Type-locality: Argentina, San Juan, Jachal.

Genus PSEUDOVIVIANA Brauer & Bergenstamm

PSEUDOVIVIANA Brauer & Bergenstamm, 1891:311 (1891:7). Type-species, *platypoda* Brauer & Bergenstamm (mon.).

PLATYPODA Brauer & Bergenstamm, 1891:312 (1891:8). Type-locality: Venezuela.

Genus SCHIZOTACHINA Walker

- Tachina*, subgenus SCHIZOTACHINA Walker, 1852:264. Type-species, *convecta* Walker (Coquillett, 1910:261).
PERUVIANA Townsend, 1929:374. Type-locality: Peru, Huallaga River, Chipeza.

Genus SIPHOACTIA Townsend

- SIPHOACTIA Townsend, 1927a:212. Type-species, *charapensis* Townsend (orig. des.).
CHARAPENSIS Townsend, 1927a:357. Type-locality: Peru, Charape River.

Genus SIPHOCROCUTA Townsend

- SIPHOCROCUTA Townsend, 1935d:229. Type-species, *trinidadensis* Townsend (orig. des.).
TRINIDADENSIS Townsend, 1935d:229. Type-locality: "Trinidad".

Genus SIPHONA Meigen

- Crocuta* Meigen, 1800:39. Type-species, *Musca geniculata* De Geer (Coquillett, 1910:528). Suppressed by ICZN, 1963:339.
SIPHONA Meigen, 1803:281. Type-species, *Conops irritans* Linnaeus (as *Stomoxys irritans* Fabricius; mon., misident.) = *geniculata* (De Geer).
anthomyiformis Lynch Arribálzaga in Brauer, 1898:505 (1898:13). *Nomen nudum*.
BRASILIENSIS (Townsend), 1929:374 (*Siphonopsis*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
CERES (Curran), 1932:14 (*Bucentes*). Type-locality: Guatemala, Antigua, Finca Pintado (cf. Arnaud, 1963a:109).
DILUTA Wulp, 1890:126. Type-locality: Mexico, Veracruz, Atoyac.
FUTILIS Wulp, 1890:125. Type-locality: Mexico, several places.
SINGULARIS (Wiedemann), 1830:335 (*Tachina*). Type-locality: Brazil. Unrecognized. Ref. — Jacnnicke, 1868:395.
PLUSIAE Coquillett, 1895c:125. Type-locality: U.S.A., California. Distr. — U.S.A., Mexico.
PSEUDOMACULATA Blanchard, 1963:233, fig. 35. Type-locality: Argentina, Santa Fé.

Genus SISYPHOMYIA Townsend

- SISYPHOMYIA Townsend, 1927a:257. Type-species, *pygmaea* Townsend (orig. des.).
PYGMAEA Townsend, 1927a:357. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus TRICHOTOPTERYX Townsend

- TRICHOTOPTERYX Townsend, 1919a:586. Type-species, *tropica* Townsend (orig. des.).
TROPICA Townsend, 1919a:587. Type-locality: Peru, Uruhuasi.



Genus TRICHSCHIZOTACHINA Townsend

TRICHSCHIZOTACHINA Townsend, 1935d:228. Type-species, *trinitas* Townsend (orig. dcs.).

TRINITAS Townsend, 1935d:228. Type-locality: Trinidad.

Genus XEOPROSOPA Townsend

XEOPROSOPA Townsend, 1919a:584. Type-species, *uruhuasi* Townsend (orig. des.).

URUHUASI Townsend, 1919a:584. Type-locality: Peru, Uruhuasi.

Tribe GRAPHOGASTRINI

The flies of the tribe Graphogastrini are parasites of Microlepidoptera larvae.

Genus CLASTONEURA Aldrich

CLASTONEURA Aldrich, 1934:26. Type-species, *brevicornis* Aldrich (orig. des.).

BREVICORNIS Aldrich, 1934:27, fig. 5. Type-locality: Argentina, Lake Nahuel Huapi.

Tribe ADMONTIINI

(Trichoparelini)

The species of Admontiini parasitize tipulid larvae.

Genus ADMONTIA Brauer & Bergenstamm

ADMONTIA Brauer & Bergenstamm, 1889:104 (1889:36). Type-species, *podomyia* Brauer & Bergenstamm (orig. des. as gen., sn., sp. n.).

Ref. — Aldrich, 1934:97 (key to Patagonian sp.).

COMMUNIS Aldrich, 1934:99. Type-locality: Argentina, Rio Negro, Lake Nahuel Huapi. Distr. — South Chile, South Argentina.

var. *albescens* Aldrich, 1934:100. Type-locality: Argentina, Rio Negro, Lake Nahuel Huapi and Bariloche.

DEBILIS Aldrich, 1934:102. Type-locality: Chile, Llanquihue, Casa Pangue.

DISCALIS Aldrich, 1934:101. Type-locality: Argentina, Rio Negro, Bariloche.


DUCALIS Reinhard, 1958:278. Type-locality: Mexico, north slope of Popocatepetl.

FLAVIBASIS Aldrich, 1934:103. Type-locality: Argentina, Rio Negro, Lake Gutierrez.

NIGRITA Thompson, 1968:123. Type-locality: Trinidad, "Simla".

OCCIDENTALIS Wulp, 1892:195. Type-locality: Mexico, Guerrero.

PICTIVENTRIS Aldrich, 1934:100. Type-locality: Chile, Llanquihue, Peulla.



Genus AUSTROSTAUROCHAETA Townsend

AUSTROSTAUROCHAETA Townsend, 1931a:474. Type-species, *Degeeria antarctica* Thomson (orig. gen.).

ANTARCTICA (Thomson), 1869:527 (*Degeeria*). Type-locality: Patagonia.

Tribe OEDEMATOCERINI

(Hyperecteinini, in part)

This tribe includes most of the species referred by Townsend (1936: 158) to his tribe Hyperecteinini. The chief tribal characters common to both tribes is the elongated third antennal segment, the deeply excavated face sometimes bounded by sharp parafacials. The tribe Hyperecteinini *sensu* Townsend seems to be a mixture, biologically speaking; it contains species like *Hyperecteina cinerea* (Aldrich) which lays macrotype eggs, *Oedematocera flaveola* (Coquillett) which lays microtype eggs, and the larviparous *O. dampfi* Ald.

Most of the Oedematocerini have been regularly recorded as parasites of Orthoptera and Phasmida. The puparia possess a very peculiar posterior spiracle.

Genus MIAMIMIYIOPS Townsend

MIAMIMIYIOPS Townsend, 1939c:452. Type-species, *mattoensis* Townsend (orig. des.).

MATTOENSIS Townsend, 1939c:452. Type-locality: Brazil, Mato Grosso, Maracaju.

Genus NEPOPHASMOPHAGA Townsend

NEPOPHASMOPHAGA Townsend, 1927a:268. Type-species, *facialis* Townsend (orig. des.).

FACIALIS Townsend, 1927a:337. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Ref. — Parker, 1953:57 (immature stages, host).

Genus OEDEMATOCERA Townsend

OEDEMATOCERA Townsend, 1916f:621. Type-species, *Hypostena flaveola* Coquillett (orig. des.).

SCHISTOCERCOPHAGA Townsend, 1928a:152. Type-species, *Oedematocera dampfi* Aldrich (orig. des.).

Ref. — Reinhard, 1942b:106 (key as *Oedematocera*).

DAMPFI Aldrich, 1927b:17. Type-locality: Mexico, Veracruz, Córdoba. Distr. — Mexico, Guatemala. Ref. — Curran, 1927:18 (immature stages).

STRIATA Aldrich, 1927a:301. Type-locality: Panama, Cabima.

Genus PHASMOVORA Cortés

PHASMOVORA Cortés, 1968a:102. Type-species, *phasmophagae* Cortés (orig. des.).



PHASMOPHAGAE Cortés, 1968a:105. Type-locality: Chilc, 15 kms. E. of Los Queñes, Cajón del Río Claro (Curicó).

Genus TAMANAMYIA Thompson

TAMANAMYIA Thompson, 1963a:466. Type-species, *trinitatis* Thompson (orig. des.).

TRINITATIS Thompson, 1963a:466. Type-locality: Trinidad, Pluna Mitán Road.

Tribe ERYTHROCERINI

The members of the tribe Erythrocerini are parasites of adult scarabeid beetles, larvae of microlepidoptera and earwigs (*Forficula*).

Genus ERYNNIA Robineau-Desvoidy

ERYNNIA Robineau-Desvoidy, 1830:125. Type-species, *nitida* Robineau-Desvoidy (mon.) = *ocypterata* (Fallén).

Ref. — Reinhard, 1968 (key to species).

Anachaeopsis, authors, not Brauer & Bergenstamm.

TORTRICIS (Coquillett), 1895:55 (*Pseudomyiothyria*). Type-locality: U.S.A., California. Distr. — U.S.A., Mexico.

TRICINCTA Reinhard, 1968:506. Type-locality: Mexico, Durango, Tlahualilc

Tibe GONIINI

The members of the tribe Goniini parasitize a large variety of lepidopterous larvae, chiefly armyworms and cutworms.

Subtribe GONIINA

Genus ACROGLOSSA Williston

ACROGLOSSA Williston, 1889:1916. Type-species, *hesperidarum* Williston (mon.).

Neacroglossa Townsend, 1927a:238. Type-species, *brasiliensis* Townsend (orig. des.).

BRASILIENSIS (Townsend), 1927a:332 (*Neocroglossa*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. Ref. — Thompson, 1963b:274.

HESPERIDARUM Williston, 1889:1917. Type-locality: U.S.A., N. Hampshire. Type-locality: U.S.A., Mexico.

VETULA Reinhard, 1964:20. Type-locality: Mexico, Veracruz, Minatitlán.

Genus ARAUCOSIMUS Aldrich


ARAUCOSIMUS Aldrich, 1934:88. Type-species, *bullocki* Aldrich (orig. des.).

BULLOCKI Aldrich, 1934:88. Type-locality: Chile, Angol.

SUPERBUS Cortés, 1945a:122. Type-locality: Chile, Valparaíso, Perales. Ref. — Campos, 1953:26.

Genus BRACHYCNEPHALIA Townsend

BRACHYCNEPHALIA Townsend, 1927a:236. Type-species, *brasiliensis* Townsend (orig. des.).



BRASILIENSIS Townsend, 1927a:291. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus CHAETOCNEPHALIA Townsend

CHAETOCNEPHALIA Townsend, 1915e:63. Type-species, *alpina* Townsend (orig. des.). Ref. — Aldrich, 1934:89.

ALPINA Townsend, 1915e:63. Type-locality: Peru, Oroya.

AMERICANA (Schiner), 1868:327 (*Cnephalia*). Type-locality: Chile. Ref. — Aldrich, 1934:89.

INNUPTA Cortés, 1945a:120. Type-locality: Chile, Santiago, Las Condes. Ref. — Campos, 1953:26 (allotype descr.).

Genus CHAETOCRANIOPSIS Townsend

CHAETOCRANIOPSIS Townsend, 1915e:68. Type-species, *chilensis* Townsend (orig. des.).

Valpogonia Townsend, 1928c:163. Type-species, *chilensis* Townsend (orig. des.) = *argenteiceps* Aldrich.

Ref. — Cortés, 1945a:116 (key to species).

ARGENTEICEPS Aldrich, 1928b:20. Type-locality: Chile, Valparaíso, Perales. *chilensis* Townsend, 1928c:163 (*Valpogonia*). Type-locality: Chile, Valparaíso (preocc. Townsend, 1915e).

CHILENSIS Townsend, 1915e:69. Type-locality: Chile.

OBLITERATUS Cortés, 1945a:117. Type-locality: Chile, Valparaíso, Perales.

SIMILIS (Townsend), 1928c:163 (*Valpogonia*). Type-locality: Chile, Valparaíso. Distr. — Chile (Valparaíso, Santiago).

Genus CNEPHALODES Townsend

CNEPHALODES Townsend, 1911b:145. Type-species, *pollinosus* Townsend (orig. des. as gen. n., sp. n.).

Cnephalodopsis Townsend, 1912a:345 (change of name without cause for *Cnephalodes* Townsend).

POLLINOSUS Townsend, 1911b:145 (also, 1912a:345). Type-locality: Piura, Peru.

Genus DISTICHONA Wulp

DISTICHONA Wulp, 1890:42, 44. Type-species, *varia* Wulp (mon.).

AURICEPS Coquillett, 1904:186. Type-locality: Mexico, Tabasco, Frontera.

DISCREPANS (Wulp), 1890:115, pl. 3, fig. 17 (*Baumhaueria*). Type-locality: Mexico, Morelos, Cuernavaca.

VARIA Wulp, 1890:44, pl. 3, figs. 2, 2a-2b. Type-locality: Mexico, Morelos, Cuernavaca.

Genus DOLICHOCNEPHALIA Townsend

DOLICHOCNEPHALIA Townsend, 1915d:64. Type-species, *puna* Townsend (orig. des.).

PUNA Townsend, 1915d:66. Type-locality: Peru, Oroya.

Genus DOLICHOGONIA Townsend

DOLICHOGONIA Townsend, 1915d:67. Type-species, *aurea* Townsend (orig. des.).

AUREA Townsend, 1915d:68. Type-locality: Peru, Oroya.

Genus ENCHOMYIA Aldrich

ENCHOMYIA Aldrich, 1934:42. Type-species, *Gonia erythrocer*a Bigot (orig. des.).

ERYTHROCERA (Bigot), 1888a:86 (*Gonia*). Type-locality: Chile. Distr. — Chile, Santiago. Ref. — Cortés, 1951:60.

Genus FRONTOCNEPHALIA Townsend

FRONTOCNEPHALIA Townsend, 1916h:16. Type-species, *angusta* Townsend (orig. des.).

ANGUSTA Townsend, 1916h:16. Type-locality: Brazil, Pará, Santarém.

Genus GERMARIOPSIS Townsend

GERMARIOPSIS Townsend, 1915e:66. Type-species, *andina* Townsend (orig. des.).

ANDINA Townsend, 1915e:67. Type-locality: Peru, Lima, San Cristóbal.

Genus GONIA Meigen

Salmacia Meigen, 1800:38. Type-species, *Musca capitata* De Geer (Coquillett, 1910:602). (Suppressed by I.C.Z.N., 1963:339).

GONIA Meigen, 1803:280. Type-species, *bimaculata* Wiedemann (Sabrosky & Arnaud, 1965:1075).

Phosococephalops Townsend, 1927a:237. Type-species, *fulvus* Townsend (orig. des.) = *pallens* (Wiedemann).

ALPINA (Townsend), 1912a:347 (*Salmacia*). Type-locality: Peru, Tirapata.

CRASSICORNIS (Fabricius), 1794:328 (*Musca*). Type-locality: West Indies. Distr. — U.S.A., Florida, Neotropical. Ref. — Thompson, 1963b:268.

angusta Macquart, 1843:208 (1843:51). Type-locality: Cuba (cf. Robineau-Desvoidy, 1863:750).

chaetosa Townsend, 1912a:347 (*Salmacia*). Type-locality: Peru, Nana. *angustata*, error.

pallens, authors, not Wiedemann.

GENEI Rondani, 1863:14. Type-locality: "Venezuela". Unrecognized.

LINEATA Macquart, 1851:151 (1851:178). Type-locality: "Patagonia". Ref. — Cortés, 1963:249.

LONGIPULVILLI Tothill, 1924:208. Type-locality: U.S.A., British Columbia. Distr. — U.S.A., Mexico.

MEXICANA Wulp, 1890:40, pl. 2, figs. 19, 19a-b, 20a. Type-locality: Mexico, Durango, Ciudad.



- PACIFICA (Townsend), 1912a:346 (*Salmacia*). Type-locality: Peru, Nana.
 PALLENS Wiedemann, 1830:346. Type-locality: Brazil.
chilensis Macquart, 1843:207 (1843:51), pl. 5, fig. 4. Type-locality: "Chile".
fulvus Townsend, 1927a:347 (*Phosococephalops*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
 PERUVIANA (Townsend), 1912a:346 (*Salmacia*). Type-locality: Peru, Piura.
 SEQUAX Williston, 1887:12. Type-locality: U.S.A., California. Distr. — U.S.A., Mexico.
 SIMILLIMA Vimmer & Soukup, 1940:360. Type-locality: Peru, Puno. Unrecognized.
subcompressus Gundlach in Pazos, 1914:1002 (*Gonia*, as *Conia*, error). *Nomen nudum*.
 TEXENSIS Reinhard, 1924:357. Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico.
 VIRESCENS Macquart, 1843:207 (1843:51). Type-locality: Brazil or Chile. Unrecognized.

Genus GONISTYLUM Macquart

- GONISTYLUM Macquart, 1851:153 (1851:180). Type-species, *ruficorne* Macquart (orig. des.).
Gonystylum, error.
 RUFICORNE Macquart, 1851:153 (1851:180), fig. 3. Type-locality: Argentina, Corrientes.

Genus IMAGUNCULA Reinhard

- IMAGUNCULA Reinhard, 1958b:230. Type-species, *tabida* Reinhard (orig. des.).
 TABIDA Reinhard, 1958b:230. Type-locality: U.S.A., N. Mexico. Distr. — U.S.A., Mexico.

Genus ITACNEPHALIA Townsend

- ITACNEPHALIA Townsend, 1927a:236. Type-species, *analís* Townsend (orig. des.).
 ANALIS Townsend, 1927a:319. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus MAYODISTICHONA Townsend

- MAYODISTICHONA Townsend, 1928a:152. Type-species, *facialis* Townsend (orig. des.).
 FACIALIS Townsend, 1928a:152. Type-locality: Peru, Mayo River, Las Flores.

Genus OLENOCHAETA Townsend

- OLENOCHAETA Townsend, 1892a:115. Type-species, *kansensis* Townsend (orig. des.).

CUBENSIS (Curran), 1927:3 (*Distichona*). Type-locality: Cuba, Pinar del Rio, Cabanas.
PERUVIANA Townsend, 1928c:162. Type-locality: Peru, Chosica, Canyon.

Genus PERACROGLOSSA Townsend

PERACROGLOSSA Townsend, 1931a:468. Type-species, *peruviana* Townsend (orig. dcs.).
PERUVIANA Townsend, 1931a:469. Type-locality: Peru, Chosica, Canyon.

Genus PERUMYIA Arnaud

PERUMYIA Arnaud, 1963c:2. Type-species, *embiaphaga* Arnaud (orig. dcs.).
EMBIAPHAGA Arnaud, 1963c:4, figs. 1, 14. Type-locality: Peru. NW of Pucallpa

Genus PHASIATACTA Townsend

PHASIATACTA Townsend, 1911b:329 (also, 1912a:343). Type-species, *elongata* Townsend (orig. des., as gen. n., sp. n.).
ELONGATA Townsend, 1911b:144, 329 (also, 1912a:344). Type-locality: Peru, Río Piura, Chapairá.

Genus PROTOGONIOPS Townsend

Protogonia Townsend, 1912a:347 (preocc. Cope, 1891). Type-species, *ocellaris* Townsend (orig. des.).
PROTOGONIOPS Townsend, 1913f:133 (n. name for *Protogonia* Townsend). Type-species, *Protogonia ocellaris* Townsend (aut.).
OCELLARIS (Townsend), 1912a:348 (*Protogonia*). Type-locality: Peru, Río Sujo.

Genus PTILOGONIA Bischof

PTILOGONIA Bischof, 1904:94. Type-sepcies, *neotropica* Bischof (orig. des.).
NEOTROPICA Bischof, 1904:95. Type-locality: Brazil, Rio Grande do Sul.

Genus RICOSIA Curran

RICOSIA Curran, 1927c:5. Type-species, *setigena* Curran (orig. des.).
SETIGENA Curran, 1927c:5, fig. 3. Type-locality: Puerto Rico, Aibonito.

Genus SPALLANZANIA Robineau-Desvoidy

SPALLANZANIA Robineau-Desvoidy, 1830:78. Type-species, *galtica* Robineau-Desvoidy (Coquillett, 1910:606) = *hebes* (Fallén).
HEBES (Fallén), 1820:11 (*Tachina*). Type-locality: "Sveden". Distr. — Palearctic, U.S.A., Mexico.

104.177

ONUSTA (Wulp), 1890:46, pl. 3, figs. 4-4a (*Cnephalia*). Type-locality: Mexico, Veracruz, Atoyac. Unrecognized.

Genus VIBRISOVORIA Townsend

VIBRISOVORIA Townsend, 1919a:567. Type-species, *petiolata* Townsend (orig. des.).

AUREA Townsend, 1929:379. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

PETIOLATA Townsend, 1919a:567. Type-locality: Panama, Canal Zone, Balboa Prado.

Subtribe CHAETOGAEDIINA

Genus ANADISTICHONA Townsend

ANADISTICHONA Townsend, 1934a:398. Type-species, *aurata* Townsend (orig. des.).

AURATA Townsend, 1934:398. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus ANDESIMYIA Brèthes

ANDESIMYIA Brèthes, 1909:97. Type-species, *scutellata* Brèthes (orig. des.).

SCUTELLATA Brèthes, 1909:97. Type-locality: Argentina, Mendoza, Valle of Tupungato River.

Genus CHAETOGAEDIA Brauer & Bergenstamm

CHAETOGAEDIA Brauer & Bergenstamm, 1891:336 (1891:32). Type-species, *Prospheysa vilis* Wulp (Townsend, 1908:94).

PHRISSOPIA Townsend, 1908:93. Type-species, *Prospheysa crebra* Wulp (mon.).

GAEDIOSTURMIA Blanchard, 1963:235. Type-species, *auricephala* Blanchard (orig. des.). N. SYN.

AURATA Blanchard, 1963:238. Type-locality: Argentina, Santa Fé.

AURICEPHALA (Blanchard), 1963:235, fig. 36 (*Gaediosturmia*). Type-locality: Argentina, Salta, Orán. N. COMB.

CREBRA (Wulp), 1890:120 (*Prospheysa*). Type-locality: Mexico, Guerrero. Distr. — U.S.A., Mexico.

OCHRICAUDA (Wulp), 1890:118 (*Prospheysa*). Type-locality: Mexico, Guerrero, Omilteme.

OCHRICEPS (Wulp), 1892:194 (*Cnephalia*). Type-locality: Mexico, Guerrero. Distr. — U.S.A., Mexico, Trinidad. Ref. — Thompson, 1963b:279.

RUFIFRONS (Wulp), 1890:121 (*Prospheysa*). Type-locality: Mexico, Guerrero, Chilpancingo. Distr. — U.S.A., N. Mexico, Mexico.

RUFOSTYLATA (Bigot), 1887:cxli (*Frontina*). Type-locality: "Mexico". Unrecognized.

TESSELLATA (Giglio-Tos), 1893:5 (*Acroglossa*). Type-locality: Mexico.

VILIS (Wulp), 1890:121 (*Prospheysa*). Type-locality: Mexico, sev. places. Ref. — Townsend, 1931b:177.

Genus CHETOGLOSSA Townsend

- CHAETOGLOSSA Townsend, 1892a:125. Type-species, *picticornis* Townsend (orig. des.).
PICTICORNIS Townsend, 1892a:126. Type-locality: U.S.A., Florida. Distr. — U.S.A., Mexico.

Genus EUGAEDIOPSIS Townsend

- EUGAEDIOPSIS Townsend, 1916c:620. Type-species, *Gaediopsis ocellaris* Coquillett (orig. des.).
Eugaedia Townsend, 1916f:621. Type-species, *Gaediopsis setosa* Coquillett (orig. des.).
RUBENTIS Reinhard, 1961:209. Type-locality: U.S.A., Arizona. Distr. — U.S.A., California.
SETOSA (Coquillett), 1897:136 (*Gaediopsis*). Type-locality: U.S.A., California Distr. — U.S.A., Mexico.
SIMMONDSI Thompson, 1963b:283, figs. 12, 13. Type-locality: Trinidad. St. Augustine.

Genus GAEDIOPHANA Brauer & Bergenstamm

- GAEDIOPHANA Brauer & Bergenstamm, 1893:123 (1893:35, 113). Type-species, *atra* Brauer & Bergenstamm (mon.) = *lugubris* (Wulp).
LUGUBRIS (Wulp), 1890:53 (*Mystacella*). Type-locality: Mexico, Guerrero. Omilteme. Distr. — U.S.A., Mexico.
tenebricosa Wulp, 1890:77 (*Phorocera*). Type-locality: Mexico, Guerrero, Tuxpango.
atra Brauer & Bergenstamm, 1893:123, 201 (1893:35, 113). Type-locality: Mexico.
MONNULA Reinhard, 1951:5. Type-locality: Mexico, Rio Frio. Distr. — Mexico (Mexico and Oaxaca).

Genus GAEDIOPHANOPSIS Blanchard

- GAEDIOPHANOPSIS Blanchard, 1954:31. Type-species, *koehleri* Blanchard (orig. des.).
KOEHLERI Blanchard, 1954:31, fig. 12. Type-locality: Argentina, Salta, Orán.

Genus GAEDIOPSIS Brauer & Bergenstamm

- GAEDIOPSIS Brauer & Bergenstamm, 1891:336, 401 (1890:32, 97) (also, 1893:190; 1893:102). Type-species, *mexicana* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.).
Poliophrys Townsend, 1908:90. Type-species, *sierricola* Townsend (orig. des.).
FLAVICAUDA (Wulp), 1890:83 (*Phorocera*). Type-locality: Mexico, Guerrero, Omilteme.
MEXICANA Brauer & Bergenstamm, 1891:336 (1891:32). Type-locality: Mexico. Ref. — Aldrich, 1925c:112.
postera Wulp, 1890:56 (*Mystacella*). Type-locality: Mexico, Guerrero, Xucumanatlán.
PUNOENSIS Vimmer & Soukup, 1940:221, 372. Type-locality: Peru, Ollachea, Puno.
RUFESCENS Aldrich, 1929b:4. Type-locality: Costa Rica, La Suiza.



SIERRICOLA (Townsend), 1908:93 (*Poliophrys*). Type-locality: Mexico. Chihuahua, Sierra Madre.

UNPLACED SPECIES OF GONIINI

ALBIFACIES Wulp, 1890:121 (*Prospherysa*). Type-locality: Mexico, Talisco.
FYGMAEA Wulp, 1890:95 (*Brachycoma*). Type-locality: Mexico.

Tribe BELVOSIINI

The hosts of Belvosiini are various heterocerous caterpillars.

Genus ATACTA Schiner

ATACTA Schiner, 1868:328. Type-species, *brasiliensis* Schiner (orig. des.).
Atactomima Townsend, 1916h:15. Type-species, *crenatis* Townsend (orig. des.).

Ref. — Aldrich, 1925a:28 (rev.).

ARGENTIFRONS Aldrich, 1925a:31. Type-locality: Brazil, Mato Grosso. Corumbá.

BRASILIENSIS Schiner, 1868:328, pl. 4, figs. 3a-b. Type-locality: "Brazil".
Distr. — U.S.A., Neotropical.

Ref. — Thompson, 1963b:358.

laticeps Wulp, 1890:92 (*Brachychoma*). Type-locality: Mexico, Guerrero and Tabasco.

CRENATIS (Townsend), 1916h:16 (*Atactomima*). Type-locality: Brazil, Mato Grosso, Chapada. Ref. — Thompson, 1965b:363.

Genus ATACTOPSIS Townsend

ATACTOPSIS Townsend, 1917b:229. Type-species, *facialis* Townsend (orig. des.).

FACIALIS Townsend, 1917b:229. Type-locality: Brazil, Mato Grosso, Chapada.

Genus ATACTOSTURMIA Townsend

ATACTOSTURMIA Townsend, 1915k:92. Type-species, *Blepharipa politana* Townsend (orig. des.).

POLITANA (Townsend), 1911b:132 (also 1912a:340) (*Blepharipa*). Type-locality: Peru, Piura.

VITTATA Thompson, 1963b:381, fig. 67. Type-locality: Trinidad, St. Augustine.

Genus BELVOSIA Robineau-Desvoidy

BELVOSIA Robineau-Desvoidy, 1830:103. Type-species, *bicincta* Robineau-Desvoidy (mon.).

- Belvoisia*, error or emend.
- Latreillia* Robineau-Desvoidy, 1830:104 (preocc. Roux, 1830). Type-species, *Musca bifasciata* Fabricius (Coquillett, 1910:558).
- Willistonina* Brauer & Bergcnstamm, 1889:97 (1889:29). Type-species, *Musca esuriens* Fabricius (mon.; misident.) = *aldrichi* (Townsend).
- Latreillimyia* Townsend, 1908:104 (nom. nov. for *Latreillia* Robineau-Desvoidy). Type-species, *Musca trifasciata* Fabricius (aut.).
- Belvosiomima* Townsend, 1915:413. Type-species, *fosteri* Townsend (orig. des.).
- Belvosioipsis* Townsend, 1927a:248. Type-species, *brasiliensis* Townsend (orig. des.) = *weyenberghiana* Wulp.
- Pseudobelvosia* Blanchard, 1954:8. Type-species, *lugubris* Blanchard (orig. des.). N. SYN.
- Parabelvosia* Blanchard, 1954:12. Type-species, *tibialis* Blanchard (orig. des.). N. SYN.
- Neobelvosioipsis* Blanchard, 1954:20. Type-species, *bosqi* Blanchard (orig. des.). N. SYN.
- Eubelvosioipsis* Blanchard, 1954:15. Type-species, *formosana* Blanchard (orig. des.). N. SYN.
- Ref. — Aldrich, 1928a (rev.); Blanchard, 1954 (Argentinean spp.); Thompson, 1963b:302 (key to Trinidad spp.).
- ALDRICHI (Townsend), 1931a:468 (*Willistonina*). Type-locality: Brazil.
- esuriens*, authors, not Fabricius.
- ANALIS Macquart, 1846:288 (1845:160), pl. 14, fig. 4. Type-locality: ? Brazil. Unrecognized.
- ANSATA Reinhard, 1951:2. Type-locality: Mexico, Michoacan, Guadalajara.
- ATRATA (Walker), 1852:284 (*Tachina*). Type-locality: Brazil. Ref. — Austen, 1907:333.
- AURATILIS Reinhard, 1951:1. Type-locality: Mexico, Michoacan, Guadalajara.
- AURIPILOSA (Blanchard), 1954:39, fig. 14 (*Willistonina*). Type-locality: "Argentina".
- AURULENTA (Bigot), 1888a:84 (*Frontina*). Type-locality: "Brazil". Ref. — Townsend, 1931b:177.
- AUSTRALIS Aldrich, 1928:8. Type-locality: Brazil, Rio Grande do Sul.
- BASALIS (Walker), 1852:285 (*Tachina*), (preocc. Walker, 1837:351). Type-locality: "South America". Unrecognized.
- BELLA Giglio-Tos, 1893:3. Type-locality: Mexico.
- Ref. — Aldrich, 1930b:26.
- BICINCTA Robineau-Desvoidy, 1830:103. Type-locality: U.S.A., Carolina. Distr. — U.S.A. to Central America, West Indies, Tropical South America. Ref. — Thompson, 1963b:302.
- BIEZANKOI (Blanchard, in Biezanko), 1961:5 (*Willistonina*). Type-locality: Argentina, Buenos Aires.
- BIFASCIATA (Fabricius), 1775:777 (*Musca*). Type-locality: West Indies (as "America"). Distr. — U.S.A., Mexico, West Indies, Guatemala.
- bifuscata*, error.
- BOSQI (Blanchard), 1954:20, fig. 6 (*Neobelvosioipsis*). Type-locality: Argentina, Misiones, Loreto. N. COMB.
- BRUCHI (Blanchard), 1954:34 (*Belvosiomima*). Type-locality: Argentina, Córdoba, Alta Gracia.
- CANALIS Aldrich, 1928a:44. Type-locality: Panama, Canal Zone, Barro Colorado I.

- CATAMARCENSIS (Blanchard), 1954:37 (*Belvosiomima*). Type-locality: Argentina, Catamarca, Pomansillo.
- CHAETOSA (Blanchard), 1954:28, fig. 11 (*Latreillemyia*). Type-locality: Argentina, Tucumán.
- CHIESAI (Blanchard), 1954:42, fig. 15 (*Willistonina*). Type-locality: Argentina, Córdoba. N. comb.
- CILIATA Aldrich, 1928:22. Type-locality: "Mexico". Distr. — Mexico, Costa Rica, Panama, Brazil.
- CONTERMINA (Walker), 1852:285 (*Tachina*). Type-locality: "South America". Unrecognized. Ref. — Austen, 1907:333.
- ELUSA Aldrich, 1926b:25. Type-locality: Brazil, Mato Grosso, Chapada. Distr. — Brazil (Mato Grosso), Paraguay (San Bernadino).
- FERRUGINOSA Townsends, 1895:71. Type-locality: "Jamaica".
- FORMOSANA (Blanchard), 1954:15, fig. 4 (*Eubelvosioptis*). Type-locality: Argentina, Formosa. N. comb.
- FORMOSA Aldrich, 1928a:23 (as *ciliata* var.). Type-locality: West Indies, St. Clair, W. I. Ref. — Townsend, 1931b:176.
- FOSTERI (Townsends), 1915m:474 (*Belvosiomima*). Type-locality: Paraguay, Sapucay. Ref. — Thompson, 1963b:310. Distr. — Brazil, Paraguay.
- FRONTALIS Aldrich, 1928a:24. Type-locality: Brazil, Mato Grosso, Chapada.
- FUSCISQUAMULA (Blanchard), 1954:44, fig. 16 (*Willistonina*). Type-locality: Argentina, Catamarca, Belém, Hualfin. N. comb.
- LATA Aldrich, 1928a:39. Type-locality: Guatemala, Puerto Barrios. Distr. — Guatemala, Brazil.
- LEUCOPYGA Wulp, 1882:84. Type-locality: Brazil. Distr. — Mexico, Venezuela, Brazil.
- LILLOI (Blanchard), 1954:47, fig. 17 (*Willistonina*). Type-locality: Argentina, Tucumán. N. comb.
- LUGUBRIS (Blanchard), 1954:10, fig. 2 (*Pseudobelvosia*). Type-locality: Argentina, Misiones. N. comb.
- MANNI Aldrich, 1928a:7. Type-locality: Bolivia, Ixiamas.
- MATAMOROSA Reinhard, 1951:3. Type-locality: Mexico, Puebla, Matamoros.
- MIRA Reinhard, 1958b:232. Type-locality: Mexico, Oaxaca, Tehuantepec.
- NIGRIFRONS Aldrich, 1928a:38. Type-locality: El Salvador, Mirasol. Distr. — El Salvador, Honduras.
- OBESULA (Wulp), 1890:46, pl. 3, fig. 3-3a (*Cnephalia*). Type-locality: Mexico, Tabasco, Teapa.
- PIURANA Townsends, 1911b:143 (also, 1912a:349). Type-locality: Peru, Piura. Distr. — Peru, Argentina. Ref. — Townsend, 1931b:176.
- POTENS (Wiedemann), 1830:312 (*Tachina*). Type-locality: Brazil, Rio de Janeiro. Ref. — Aldrich, 1927c:33.
- PROXIMA (Walker), 1852:287 (*Tachina*). Type-locality: Brazil, Pará.
- pfeifferi* Schiner in Brauer, 1897:355. *Nomen nudum*.
- RECTICORNIS (Macquart), 1855:118 (1855:98) (*Gonia*). Type-locality: Unknown. Distr. — Mexico, Panama, Ecuador, Brazil.
- mexicana* Aldrich, 1928a:11. Type-locality: Mexico, D. F.
- RUFICORNIS Aldrich, 1928a:16 (as *recticornis* var.). Type-locality: Brazil, Mato Grosso, Chapada.
- RUFIFRONS Blanchard, 1954:23, fig. 8. Type-locality: Argentina, Córdoba.
- SEMIFLAVA Aldrich, 1928a:11. Type-locality: U.S.A., N. Mexico. Distr. — U.S.A., Mexico.
- SMITHI Aldrich, 1928a:40. Type-locality: Brazil, Mato Grosso, Chapada.
- SOCIA (Walker), 1852:286 (*Tachina*). Type-locality: Brazil. N. comb.
- SPINICOXA Aldrich, 1928a:41. Type-locality: Paraguay, Sapucay. Distr. — Mexico, Bolivia, Brazil (Amazonas, Mato Grosso), Cuba (Guantanamo).



- TIBIALIS (Blanchard), 1954:13, fig. 3 (*Parabelvosia*). Type-locality: Argentina, Misiones. N. comb.
- VANDERWULPI Williston, 1886:303 (spelled as *Belvosia v.d. Wulpi*). Type-locality: Dominican Republic, Santo Domingo.
- VILLARICANA Reinhard, 1951:4. Type-locality: Paraguay, Villarica.
- VITTATA Aldrich, 1928a:24. Type-locality: Paraguay, Sapucay.
- WEYNBERGHIANA Wulp, 1883:26, pl. 1, fig. 16-18. Type-locality: Argentina. Distr. — Mexico to Argentina. Ref. — Blanchard, 1954:8.
- brasilensis* Townsend, 1927a:289 (*Belvosiosis*). Type-locality: "Brazil".
- WIEDEMANNI Aldrich, 1928a:19. Type-locality: Brazil, Santa Catarina, Blumenau.
- WILLIAMSII Aldrich, 1928a:43. Type-locality: Brazil, São Paulo, Campinas. Distr. — Guyana, Brazil.
- WILLINKI (Blanchard), 1954:18, fig. 5 (*Eubelvosiosis*). Type-locality: Argentina, Misiones, Iguazú.

Genus BELVOSIELLA Curran

- BELVOSIELLA Curran, 1934:493 (key), 518. Type-species, *funditor* Curran (orig. des.).
- FUNDITOR Curran, 1934:518. Type-locality: Guyana, Potaro Landing.

Genus BELVOSIOMIMOPS Townsend

- BELVOSIOMIMOPS Townsend, 1935d:229. Type-species, *barbiellinii* Townsend (orig. des.).
- BARBIELLINI Townsend, 1935d:229. Type-locality: Brazil, São Paulo, São Vicente.

Genus BLEPHARIATACTA Townsend

- BLEPHARIATACTA Townsend, 1931a:467. Type-species, *brasiliana* Townsend (orig. des.).
- BRASILIANA Townsend, 1931a:468. Type-locality: Brazil, São Paulo, Varhagen (formerly Ypanema).

Genus BRACHYBELVOSIA Townsend

- BRACHYBELVOSIA Townsend, 1927a:248. Type-species, *brasiliensis* Townsend (orig. des.).
- BRASILIENSIS Townsend, 1927a:291. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus CHOETEPROSOPA Macquart

- CHOETEPROSOPA Macquart, 1851:169 (1851:186). Type-species, *cyanea* Macquart (orig. des.).
- Chaetoprosopa*, emend.
- Paragaedia* Brauer & Bergenstamm, 1891:350 (1891:46). Type-species, *hedemanni* Brauer & Bergenstamm (mon.).

Paratactopsis Townsend, 1917b:230. Type-species, *setosa* Townsend (orig. des.) = *cyanea* (Macquart).

Ref. — Aldrich, 1925:464 (key).

ALBIFACIES (Bigot), 1888a:92 (*Blepharipeza*). Type-locality: Brazil.

AURICEPS Aldrich, 1925d:464. Type-locality: Mexico, Tacubaya.

CYANEA Macquart, 1851:169 (1851:196), pl. 17, fig. 17. Type-locality: Brazil, Minas Gerais. Ref. — Townsend, 1931b:176.

setosa Townsend, 1917b:230 (*Paratactopsis*). Type-locality: Brazil, Mato Grosso, Chapada.

CYANEIVENTRIS (Macquart), 1846:295 (1846:157), pl. 13, fig. 11 (*Blepharipeza*). Type-locality: Brazil, Minas Gerais.

HEDEMANNI (Brauer & Bergenstamm), 1891:350 (1891:45) (*Paragaedia*). Type-locality: Mexico.

Genus CYOSOPROCTA Reinhard

CYOSOPROCTA Reinhard, 1952a:5. Type-species, *funebria* Reinhard (orig. des.).

AURICEPS Reinhard, 1952a:6. Type-locality: Mexico, Rio Frio.

FUNEBSIA Reinhard, 1952a:6. Type-locality: Mexico, Rio Frio.

Genus PHYLLARISTOMYIA Townsend

PHYLLARISTOMYIA Townsend, 1931a:466. Type-species, *fiebrigi* Townsend (orig. des.).

FIEBRIGI Townsend, 1931a:467. Type-locality: Paraguay, San Bernardino. Distr. — Brazil (Mato Grosso), Paraguay.

Genus PROTOGONIOPSIS Townsend

PROTOGONIOPSIS Townsend, 1915m:412. Type-species, *arida* Townsend (orig. des.).

ARIDA Townsend, 1915m:412. Type-locality: Peru, Chosica Canyon, Goatherd Camp.

Genus TRIACHORA Townsend

TRIACHORA Townsend, 1908:105. Type-species, *Latreillia unifasciata* Robineau-Desvoidy (orig. des.).

Goniomima Townsend, 1908:105. Type-species, *Belvosia luteola* Coquillett (mon.).

Ref. — Aldrich, 1928a (rev. in *Belvosia*).

CHRYSOPYGA (Bigot), 1887:cxli (also 1888a:84, as *chrysopygata*, error) (*Frontina*). Type-locality: Mexico. N. comb.

EQUINOCTIALIS Townsend, 1912:348. Type-locality: Peru, Piura. Ref. — Thompson, 1963b:314.

? *insularis* Curran, 1927:151 (*Belvosia*). Type-locality: Puerto Rico, Barrios.

? *antilliana* Curran, 1927:151 (*Belvosia*). Type-locality: not given.

- LUTEOLA (Coquillett), 1900:253 (*Belvosia*). Type-locality: Puerto Rico, Vieques I. Distr. — Puerto Rico, Peru. Ref. — Thompson, 1963b:319.
- OCHRIVENTRIS (Wulp), 1890:47 (*Cnecphalia*). Type-locality: Mexico, Guerrero. Distr. — Mexico, Costa Rica (San Mateo).
- UNIFASCIATA (Robineau-Desvoidy), 1830:105 (*Latreillia*). Type-locality: U.S.A., Pennsylvania. Distr. — U.S.A., Cuba.

Tribe HARRISIINI

The members of the tribe Harrisini parasitize lepidopterous larvae.

Genus COLURUS Reinhard

- COLURUS Reinhard, 1953a:98. Type-species, *downsi* Reinhard (orig. des.).
- DOWNSI Reinhard, 1953a:98. Type-locality: Mexico, Amalcalco.

Genus ECHINOMASICERA Townsend

- ECHINOMASICERA Townsend, 1915m:413. Type-species, *hystrix* Townsend (orig. des.).
- HYSTRIX Townsend, 1915m:413. Type-locality: Peru, Matucana.

Genus HARRISIA Robineau-Desvoidy

- HARRISIA Robineau-Desvoidy, 1830:323. Type-species, *scutellaris* Robineau-Desvoidy (Townsend, 1917:7).
- BRASILIENSIS Robineau-Desvoidy, 1830:324. Type-locality: Brazil, Paraná, Guaratuba.
- SCUTELLARIS Robineau-Desvoidy, 1830:324. Type-locality: "Brazil".

Genus JAENIMYIA Townsend

- JAENIMYIA Townsend, 1912a:350. Type-species, *albicincta* Townsend (orig. des.).
- ALBICINCTA Townsend, 1912a:350. Type-locality: Peru, Charape River.
- PUNCTATA Townsend, 1912a:351. Type-locality: Peru, Jaen, Charape River.

Genus LESCHENAULTIA Robineau-Desvoidy

- LESCHENAULTIA Robineau-Desvoidy, 1830:324. Type-species, *cilipes* Robineau-Desvoidy (Townsend, 1916:7) = *leucophrys* (Wiedemann).
- Blepharipeza* Macquart, 1843:211 (1843:54). Type-species, *rufipalpis* Macquart (mon.) = *leucophrys* (Wiedemann).
- Harrisioptis* Townsend, 1927a:247. Type-species, *spinosa* Townsend (orig. des.).
- CILIATA (Macquart), 1848:209 (1848:49), pl. 5, fig. 9 (*Phorocera*). Type-locality: "Colombia".



- HOSPITA Reinhard, 1952a:7. Type-locality: Mexico, Michoacan, Zamora.
- JURINIOIDES (Townsend), 1895:71 (*Blepharipeza*). Type-locality: Jamaica, Cinchona.
- LEUCOPHRYS (Wiedemann), 1830:308 (*Tachina*). Type-locality: Brazil. Distr. — Canada, U.S.A., Neotropical.
Ref. — Aldrich, 1927c:30.
- cilipes* Robineau-Desvoidy, 1830:325. Type-locality: "Surinam".
- rufipalpis* Macquart, 1843:212 (1843:55), pl. 6, fig. 1 (*Blepharipeza*). Type-locality: "Mexico".
- spinosa* Townsend, 1927a:313 (*Harrisiopsis*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- LATIFRONS (Walker), 1852:284 (*Tachina*). Type-locality: "South America".
- MONTAGNA (Townsend), 1912a:351 (*Blepharipeza*). Type-locality: Peru, Uruhuasi bridge, Rio San Gaban.
- NUDA Thompson, 1963b:288, figs. 14, 17. Type-locality: Trinidad, "Brazil" (Village).
- NIGRISQUAMIS (Townsend), 1892:80 (*Blepharipeza*). Type-locality: "Jamaica".
- TRICHOPSIS Bigot, 1887:cxl. Type-locality: "Mexico". Unrecognized.

Genus MOREIRIA Townsend

- MOREIRIA Townsend, 1932b:107. Type-species, *maura* Townsend (orig. des.).
- MAURA Townsend, 1932b:107. Type-locality: Brazil, Rio de Janeiro.

Genus PARACHAETA Coquillett

- PARACHAETA Coquillett, 1897:123. Type-species, *Blepharipeza bicolor* Macquart (orig. des.; misident.) = *fusca* Townsend.
- BICOLOR (Macquart), 1846:286 (1846:158), pl. 20, fig. 7 (*Blepharipeza*). Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico.
- NIGRICALYPTRATA (Macquart), 1855:120 (1854:100) (*Masicera*). Type-locality: Brazil, Amazonas.

Genus PROPARACHAETA Townsend

- PROPARACHAETA Townsend, 1928b:152. Type-species, *paraguayensis* Townsend (orig. des.).
- Proparachactopsis* Blanchard, 1942a:367. Type-species, *quinquevittata* Blanchard (orig. des.). N. SYN.
- PARAGUAYENSIS Townsend, 1928b:152. Type-locality: Paraguay, Santa Trinidad.
- QUINQUEVITTATA Blanchard, 1942a:367, fig. 12. Type-locality: Argentina, Tucumán. N. COMB.

Genus PTEROPEZA Townsend

- Chaetoprocta* Brauer & Bergenstamm, 1891:341 (1891:37). Type-species, *Blepharipeza tarsalis* Schiner (mon.), (preocc. Nicéville, 1890).
- PTEROPEZA Townsend, 1908:113 (n. name for *Chaetoprocta*). Type-species, *Blepharipeza tarsalis* Schiner (aut.).
- TARSALIS (Schiner), 1868:336 (*Blepharipeza*). Type-locality: "Venezuela".
Ref. — Aldrich, 1925d:461.

Genus THYSANOPSIS Townsend

THYSANOPSIS Townsend, 1917b:231. Type-species, *albicauda* Townsend (orig. des.).

ALBICAUDA Townsend, 1917b:231. Type-locality: Brazil, Mato Grosso, Chapada.

Tribe STURMIINI

The tribe Sturmiini contains numerous parasites of Lepidopterous larvae and some tenthredinid larvae.

Genus ALLOSTURMIA Blanchard

ALLOSTURMIA Blanchard, 1958:36. Type-species, *turicai* Blanchard (orig. des.).

TURICAI Blanchard, 1958:36. Type-locality: Argentina, Buenos Aires, José C. Paz.

Genus ANEMORILLA Townsend

ANEMORILLA Townsend, 1915m:432. Type-species, *rufescens* Townsend (orig. des.).

RUFESCENS Townsend, 1915m:432. Type-locality: Peru, Lima.

Genus AZYGOBOTHRIA Townsend

AZYGOBOTHRIA Townsend, 1911b:142 (also, 1912a:322). Type-species, *aurea* Townsend (orig. des., as gen. n., sp. n.).

AUREA Townsend, 1911b:142 (also, 1912a:322). Type-locality: Peru, Rio Chira Valley, Sullana.

Genus BICRUCIOSTURMIA Townsend

BICRUCIOSTURMIA Townsend, 1932b:106. Type-species, *bicrucis* Townsend (orig. des.).

BICRUCIS Townsend, 1932b:107. Type-locality: Rio de Janeiro.

Genus BOLOMYIA Brauer & Bergenstamm

BOLOMYIA Brauer & Bergenstamm, 1891:341 (1891:37) (also 1891:347; 1891:43). Type-species, *Mystacella violacea* Wulp (mon.) = *rufata* (Bigot).

COMMETANS (Walker), 1860:299 (*Eurigaster*). Type-locality: "Mexico".

RUFATA (Bigot), 1889:257 (*Exorista*). Type-locality: "Mexico". Distr. — U.S.A., Mexico, Central America.

Ref. — Townsend, 1931b:171 (notes on types).

cuprescens (Walker), 1857:196 (*Eurigaster*). Type-locality: unknown (as "Hindustan", error).

violacea Wulp, 1890:53 (*Mystacella*). Type-locality: "Mexico".

Genus CHRYSOSTURMIA Townsend

- CHRYSOSTURMIA Townsend, 1916h:20. Type-species, *orbitalis* Townsend (orig. des.).
ORBITALIS Townsend, 1916h:20. Type-locality: Brazil, Mato Grosso, Chapada.

Genus CUBAEMYIOPSIS Thompson

- CUBAEMYIOPSIS Thompson, 1963b:385. Type-species, *trinitatis* Thompson, 1963 (orig. des.).
TRINITATIS Thompson, 1963b:386, fig. 68. Type-locality: Trinidad, Santa Cruz Valley.

Genus DRINO Robineau-Desvoidy

- DRINO Robineau-Desvoidy, 1863:259. Type-species, *volucris* Robineau-Desvoidy (orig. des.) = *lota* (Meigen).
Zygosturmia Townsend, 1911:142 (also, 1912a:323). Type-species, *unica* Townsend (orig. des., as gen. n., sp. n.).
Anazygosturmia Townsend, 1927a:271. Type-species, *analis* Townsend (orig. des.). N. COMB.
Cubaemyia Townsend, 1931b:179. Type-species, *Tachina cubaecola* Jaenicke (orig. des.).
Sturmia authors, not Robineau-Desvoidy.
Ref. — Thompson, 1966:393 (key to Trinidad species). Most of species, referred by the authors as *Sturmia* are left here until their proper generic position can be established.
ANALIS (Townsend), 1927a:286 (*Anazygosturmia*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. N. COMB.
AUROCAUDA (Bigot), 1888a:90 (*Blepharipeza*). Type-locality: Uruguay, Montevideo. N. COMB.
AUREOCAUDA Thompson, 1966:398. Type-locality: Trinidad, San Juan.
AUSTRINA (Coquillett), 1902b:13 (*Sturmia*). Type-locality: Bahamas, Nassau. N. COMB.
BALLOU (Curran), 1935:22 (*Sturmia*). Type-locality: Costa Rica, San Pedro de Montes de Oca. N. COMB.
CINERACEA (Wulp), 1890:100 (*Brachycoma*). Type-locality: Mexico, Veracruz. N. COMB.
COMPACTA (Walker), 1852:294 (*Tachina*). Type-locality: "Brazil". Unrecognized.
CUREPEI Thompson, 1966:408, figs. 30, 32. Type-locality: Trinidad, Paradise Mt.
CURTA (Wulp), 1890:112 (*Masicera*). Type-locality: Mexico, Tabasco, Teapa. N. COMB.
CRESCENTIS (Reinhard), 1944:69 (*Sturmia*). Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico.
CUBAECOLA (Jaenicke), 1867:382 (1868:74) (*Tachina*). Type-locality: "Cuba". Distr. — U.S.A., Cuba.
DISTINCTA (Townsend), 1929:377 (*Anazygosturmia*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. N. COMB.
DUBIA (Bigot), 1889a:257 (*Chaetolyga*). Type-locality: Mexico. N. COMB.
FRAUDULENTA (Wulp), 1890:110 (*Masicera*). Type-locality: Mexico, Guerrero. N. COMB.
HEINRICHI (Lima), 1947:280, pl. 1 (*Zygosturmia*). Type-locality: Brazil, São Paulo, Tatui. N. COMB.



- INCA (Townsend), 1911b:142 (also, 1912b:323) (*Zygosturmia*). Type-locality: Peru, Rio Chira Valley, Sullana. Distr. — U.S.A., Peru. Ref. — Aldrich, 1927a:19 (notes). N. comb.
- INCOMPTA (Wulp), 1890:99 (*Brachycoma*). Type-locality: Mexico, Guerrero. Distr. — U.S.A., Mexico. N. comb.
- INQUINATA (Wulp), 1890:107 (*Masicera*). Type-locality: Mexico, Morelos, Cuernavaca. N. comb.
- IRREGULARIS (Wulp), 1890:97 (*Brachycoma*). Type-locality: Mexico, Tabasco, Teapa.
- MACARENSIS (Townsend), 1928c:162 (*Anazygosturmia*). Type-locality: Peru, Rio Maracá, La Tina. N. comb.
- MERIDIONALIS (Townsend), 1917b:231 (as *protoparcis* ssp.) (*Zygosturmia*). Type-locality: Brazil, Mato Grosso, Chapada. N. comb.
- MEXICANA (Giglio-Tos), 1893:6 (*Elepharipoda*). Type-locality: Mexico, Tehuacan. N. comb.
- NIGRICAUDA Thompson, 1966:394, figs. 10-21. Type-locality: Trinidad, Bamboo Grove.
- NIGRIPALPIS Thompson, 1966:404, figs. 27-29. Type-locality: Trinidad, St. Augustine.
- ORBITALIS (Curran), 1934c:515 (*Sturmia*). Type-locality: Guyana, Bartica. Kartabo. N. comb.
- PICEIVENTRIS (Walker), 1837:350 (*Tachina*). Type-locality: "South America". Ref. — Cortés, 1944:161, Campos, 1953. N. comb.
- RHOEO (Walker), 1849:778 (*Tachina*). Type-locality: Unknown. Distr. — U.S.A., Mexico, Jamaica.
protoparcis Townsend, 1892c:70 (*Masicera*). Type-locality: "Jamaica".
geminata Curran, 1926a:113 (*Atacta*). Type-locality: "Jamaica", West Indies.
distincta, authors not Wiedemann.
- RUFICAUDA Thompson, 1966:401, figs. 22-26. Type-locality: Trinidad, Bamboo Grove.
- SOCIABILIS (Greene), 1912a:125, fig. 1 (*Sturmia*). Type-locality: Trinidad, Curepe. N. comb.
- TRIFIDA (Wulp), 1890:96 (*Brachycoma*). Type-locality: Mexico, Guerrero. N. comb.
- TRINIDADENSIS Thompson, 1966:411, figs. 33-36. Type-locality: Trinidad, Curepe.
- VARIPENNIS (Curran), 1934:515 (*Sturmia*). Type-locality: Guyana, Bartica. Kartabo.

Genus EUNEMORILLA Townsend

- EUNEMORILLA Townsend, 1919b:177. Type-species, *peruviana* Townsend (orig. des.).
- PERUVIANA Townsend, 1919b:177. Type-locality: Peru, Santa Eulalia.

Genus HELICONIOPHAGA Thompson

- HELICONIOPHAGA Thompson, 1966:388. Type-species, *cranei* Thompson (mon.).
- CRANEI Thompson, 1966:388, figs. 17, 18. Type-locality: Trinidad, Arima Valley (no type designated).

Genus ITASTURMIA Townsend

- ITASTURMIA Townsend, 1927a:276. Type-species, *intermedia* Townsend (orig. des.).

INTERMEDIA Townsend, 1927a:322. Type-locality: Brazil, São Paulo. Itaquaquecetuba.

Genus MILONIUS Reinhard

MILONIUS Reinhard, 1955c:127. Type-species, *scordadus* Reinhard (orig. des.).

SCORDALUS Reinhard, 1955c:127. Type-locality: U.S.A., Texas. Distr. — U.S.A., Mexico.

Genus MIMOLOGUS Reinhard

MIMOLOGUS Reinhard, 1955c:129. Type-species, *effetus* Reinhard (orig. des.).

EMULATUS Reinhard, 1962b:177. Type-locality: Mexico, 10 miles E. Navaloa, Sonora.

Genus MYIOSTURMIA Townsend

MYIOSTURMIA Townsend, 1927a:272. Type-species, *mixta* Townsend (orig. des.).

MIXTA Townsend, 1927a:331. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

NIGRIPALPIS (Aldrich), 1932b:27 (*Achaetoneura*). Type-locality: "Trinidad". N. COMB.

Genus MYIOSTURMIOPSIS Thompson

MYIOSTURMIOPSIS Thompson, 1963b:267, 393. Type-species, *Masicera abdominalis* Wulp (orig. des.).

ABDOMINALIS (Wulp), 1890:106 (*Masicera*). Type-locality: Mexico, North Yuacatan, Temax. Distr. — Mexico, Trinidad.

Genus MYOTHYRIOPSIS Townsend

MYOTHYRIOPSIS Townsend, 1919a:575. Type-species, *bivittata* Townsend (orig. des.) = *picta* (Wulp).

latifasciata Blanchard, in [Marchionatto], 1941:28. *Nomen nudum*.

PICTA (Wulp), 1890:108 (*Masicera*), pl. 3, figs. 13-13a. Type-locality: Mexico, sev. places. Distr. — U.S.A., Mexico. Ref. — Thompson, 1966:384.

bivittata Townsend, 1919a:575. Type-locality: Brazil, Mato Grosso, Chapada.

Genus MYSTACELLA Wulp

MYSTACELLA Wulp, 1890:51. Type-species, *solita* Wulp (Coquillett, 1910:573).

Macromeigenia Brauer & Bergenstamm, 1891:311 (1891:7). Type-species, *Tachina chrysoprocta* Wiedemann (mon.).

AUREA (Townsend), 1916h:21 (*Macromeigenia*). Type-locality: Brazil, Mato Grosso, Chapada.

CHRYSOPECTA (Wiedemann), 1830:309 (*Tachina*). Type-locality: Unknown.
Distr. — U.S.A., Mexico, Costa Rica.
FLAVIFRONS Wulp, 1890:57. Type-locality: Mexico, Tabasco, Teapa.
SOLITA Wulp, 1890:55. Type-locality: Mexico, Guerrero. Ref. — Townsend,
1931b:171 (notes on type).

Genus NEOPODOMYIA Townsend

NEOPODOMYIA Townsend, 1927a:241. Type-species, *oralis* Townsend (orig.
des.).
ORALIS Townsend, 1927a:334. Type-locality: Peru, Charape River.

Genus OPSOSTURMIA Townsend

OPSOSTURMIA Townsend, 1927a:277. Type-species, *tarsalis* Townsend (orig.
des.).
TARSALIS Townsend, 1927a:340. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus PARADRINO Mesnil

PARADRINO Mesnil, 1949b:103 (as subgenus). Type-species, *Sturmia halli*
Curran (Mesnil, 1949a:35).
SOLITARIS Thompson, 1966:410. Type-locality: Trinidad, Maracas Valley.

Genus PODOSTURMIA Townsend

PODOSTURMIA Townsend, 1928b:151. Type-species, *dirphiae* Townsend (orig.
des.).
DIRPHIAE Townsend, 1928:151. Type-locality: Brazil, Pará, Belém.
molinari Blanchard in [Marchionatto], 1941:29. *Nomen nudum*.

Genus PSEUDOSIPHOSTURMIA Thompson

PSEUDOSIPHOSTURMIA Thompson, 1966:355 (key), 415. Type-species, *aberrans*
Thompson (mon.).
ABERRANS Thompson, 1966:415, figs. 37-39. Type-locality: Trinidad, Champ
Fleurs.

Genus PSEUDOSTURMIA Thompson

PSEUDOSTURMIA Thompson, 1966:355 (key), 423. Type-species, *clavipalpis*
Thompson (orig. des.).
CLAVIPALPIS Thompson, 1966:424, figs. 40-43. Type-locality: Trinidad, Sta.
Cruz Valley.

Genus STENOSTURMIA Townsend

STENOSTURMIA Townsend, 1927a:274. Type-species, *stricta* Townsend (orig.
des.).

PERUANA Townsend, 1929:378. Type-locality: Peru, Tabañosas.
STRICTA Townsend, 1927a:358. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus STURMIA Robineau-Desvoidy

STURMIA Robineau-Desvoidy, 1830:171. Type-species, *vanessae* Robineau-Desvoidy (Robineau-Desvoidy, 1863:888) = *bella* (Meigen).
CONSISTENS (Curran), 1927c:10 (*Erycia*). Type-locality: Puerto Rico, Coamo Springs.
FESTIVA Cortés, 1944e:163. Type-locality: Chile, Valparaíso. Perales. Ref. — Campos, 1953.
HARRISSINAE Coquillett, 1897:111. Type-locality: U.S.A. Distr. — U.S.A.. Mexico.
INSICNIS (Wulp), 1882:85 (*Masicera*). Type-locality: "Chile". Ref. — Cortés. 1944e:166.

Genus STURMIELLINA Thompson

STURMIELLINA Thompson, 1963b:261, 376. Type-species, *trinitatis* Thompson (orig. des.).
TRINITATIS Thompson, 1963b:37, figs. 64-66. Type-locality: Trinidad, Mellajo River.

Genus STURMIOACTIA Townsend

STURMIOACTIA Townsend, 1926a:261. Type-species, *auronigra* Townsend (orig. des.).
AURONIGRA Townsend, 1927a:358. Type-locality: Peru, Charape River.

Genus STURMIOMIMA Townsend

STURMIOMIMA Townsend, 1934:404. Type-species, *sturmioides* Townsend (orig. des.).
STURMIOIDES Townsend, 1934:404. Type-locality: Brazil, Boa Vista, Tapajós River.

Genus STURMIOPSOIDEA Thompson

STURMIOPSOIDEA Thompson, 1966:359. Type-species, *obscura* Thompson (mon.).
OBSCURA Thompson, 1966:359, fig. 1. Type-locality: Trinidad, American Road, Mauvan Hill.

Genus THYSANOMYIA Brauer & Bergenstamm

THYSANOMYIA Brauer & Bergenstamm, 1891:340 (1891:36). Type-species, *Brachycoma fimbriata* Wulp (mon.).
FIMBRIATA (Wulp), 1890:97 (*Brachycoma*). Type-locality: Mexico, Guerrero and Veracruz.
PAULISTA Townsend, 1929:379. Type-locality: Brazil, São Paulo.
PERUANA Townsend, 1929:379. Type-locality: Peru, Charape River.

Genus THYSANOSTURMIA Townsend

- THYSANOSTURMIA Townsend, 1927a:271. Type-species, *scutellaris* Townsend (orig. des.).
SCUTELLARIS Townsend, 1927a:360. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus TOPOMEIGENIA Townsend

- TOPOMEIGENIA Townsend, 1919a:575. Type-species, *matutina* Townsend (orig. des.).
Gymnosturmia Townsend, 1927a:248. Type-species, *grisea* Townsend (orig. des.).
ANDINA (Townsend), 1929:377 (*Gymnosturmia*). Type-locality: Peru, Moyabamba trail, Cavallera Tambo.
GRISEA (Townsend), 1927a:312 (*Gymnosturmia*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
MATUTINA Townsend, 1919a:576. Type-locality: Mexico, Sinaloa. Ref. — Townsend, 1931b:175.

Genus TUNAPUNIA Thompson

- TUNAPUNIA Thompson, 1963b:267, 389. Type-species, *ruficauda* Thompson (orig. des.).
RUFICAUDA Thompson, 1963b:390. Type-locality: "Trinidad".

Genus ZYGOZENILLIA Townsend

- ZYGOZENILLIA Townsend, 1927a:268. Type-species, *plumbea* Townsend (orig. des.).
PLUMBEA Townsend, 1927a:369. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus ZIZIPHOMYIA Townsend

- ZIZIPHOMYIA Townsend, 1916f:317. Type-species, *celcr* Townsend (orig. des.) = *limata* (Coquillett).
ANGUSTA Reinhard, 1967:109. Type-locality: Brazil, Santa Catarina, Nova Teutônia.

UNPLACED SPECIES OF STURMIINI.

- ALACRIS Walker, 1860:304 (*Masicera*). Type-locality: Brazil.
ALBIFRONS Walker, 1837:351 (*Tachina*). Type-locality: "unknown", but Southern S. A. — Ref. — Austen, 1907:329.
ANDINA Bigot, 1888a:91 (*Blepharipeza*). Type-locality: Chile.
CHIHUAHUENSIS Townsend, 1892:165 (*Brachycoma*). Type-locality: Mexico, Chihuahua.
FLAVIFACIES Bigot, 1889:263 (*Masicera*). Type-locality: Mexico.
FOVEATA Wulp, 1890:93 (*Brachycoma*). Type-locality: Mexico.
IMPERIALIS Curran, 1926:110 (*Sturmia*). Type-locality: Jamaica, Blue Castle.
MAURA Wulp, 1890:72 (*Exorista*). Type-locality: Mexico, Guerrero.
MONENSIS Curran, 1926a:111 (*Phorocera*). Type-locality: Jamaica, Mona and Hope Gardens.

- NIGRIPALPIS Wulp, 1890:98 (*Brachycoma*). Type-locality: "Mexico" (sev. places).
 POSTICA Walker, 1861:301 (*Eurigaster*). Type-locality: "Mexico".
 PILITARSIS Rondani, 1850:363 (*Blepharipoda*). Type-locality: "Equatorial America".
 ROBUSTA Aldrich, 1934:91 (*Achaetoneura*). Type-locality: Chile, San Cristóbal and Angol.
 SPURIA Wulp, 1890:101 (*Brachycoma*). Type-locality: "Mexico".
 STRIAELLA Wulp, 1890:100 (*Brachycoma*). Type-locality: Mexico, Guerrero and Tabasco.
 SUBVARIA Walker, 1852:299 (*Tachina*). Type-locality: "West Indies".
 XANTHURA Wulp, 1890:80 (*Phorocera*). Type-locality: Mexico, Guerrero, Venta de Zopilote.

Tribe WINTHEMIINI

Ref. — Thompson, 1963c (Genitalia, larvae, key to Trinidad sp.).

Genus AVIBRISSOSTURMIA Townsend

- AVIBRISSOSTURMIA Townsend, 1927a:266. Type-species, *avida* Townsend (orig. des.).
Pelixia Curran, 1934c:492 (key), 512. Type-species, *vexans* Curran (orig. des.).
 AVIDA Townsend, 1927a:283. Type-locality: Peru, near Tabalosas.
 VEXANS (Curran), 1934c:512 (*Pelixia*). Type-locality: Guyana, Bartica, Kartabo. Ref. — Townsend, 1963b:279.

Genus EUWINTHEMIA Blanchard

- EUWINTHEMIA Blanchard, 1963:218. Type-species, *Pronemorilla kreibohmi* Blanchard (orig. des.).
 KREIBOHMI (Blanchard), 1942a:364, fig. 11 (*Pronemorilla*). Type-locality: Argentina, Tucumán.

Genus HEMISTURMIA Townsend

- HEMISTURMIA Townsend, 1927a:262, 316 (as *Humisturmia*, pg. 262). Type-species, *carcelioides* Townsend (orig. des.).
 AMERICANA (Curran), 1927c:8, fig. 4 (*Anacamptomyia*). Type-locality: Puerto Rico, Mayaguez.
 CARCELOIDES Townsend, 1927a:316. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — West Indies (Trinidad), Brazil.
 TORTRICIS (Coquillett), 1897:103 (*Phorocera*). Type-locality: U.S.A. Distr. — U.S.A., Mexico.

Genus NEMORILLA Rondani

- NEMORILLA Rondani, 1856:66. Type-species, *Tachina maculosu* Meigen (orig. des.).

Tinanemorilla Townsend, 1927a:266. Type-species, *angustipennis* Townsend (orig. des.). N. SYN.
 ANGUSTIPENNIS (Townsend), 1927a:361 (*Tinanemorilla*). Type-locality: Peru, Macará River, La Tina. N. COMB.
chacoensis (Blanchard in Denier), 1939:569 (*Tinanemorilla*). *Nomen nudum*.
 PYSTE (Walker), 1849:754 (*Tachina*). Type-locality: U.S.A., Illinois. Distr. — U.S.A., Mexico.
 RUFICORNIS (Thomson), 1869:520 (*Exorista*). Type-locality: Brazil, Rio de Janeiro. Ref. — Townsend, 1931b:174.
 TRIVITTATA (Wiedemann), 1830:300 (*Tachina*). Type-locality: "St. Thomas". Ref. — Townsend, 1931b:174.
 TRIVITTATA (Wulp), 1890:70 (*Exorista*). Type-locality: Mexico, Veracruz (this species is a secondary homonym of the preceding one, but I prefer not rename at present because their identities requires clarification).

Genus PARACHETOLYGA Bischof

PARACHETOLYGA Bischof, 1904:83. Type-species, *metopia* Bischof (orig. des.).
 METOPIA Bischof, 1904:84. Type-locality: Brazil, Rio Grande do Sul. Ref. — Townsend, 1931b:175 (notes on type); Parker, 1953:58 (immature stages).
elegans Townsend, 1929:377. Type-locality: Brazil, São Paulo, Itaquaquecetuba. N. SYN.

Genus PROCARCELIA Townsend

PROCARCELIA Townsend, 1927a:267. Type-species, *brasiliensis* Townsend (orig. des.).
Procarcella, misprint.
 BRASILIENSIS Townsend, 1927a:349. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus PRONEMORILLA Townsend

PRONEMORILLA Townsend, 1935d:229. Type-species, *mima* Townsend (orig. des.).
 MIMA Townsend, 1935d:230. Type-locality: Brazil, São Paulo, São Vicente.

Genus PROTYPOPHAEMYIA Blanchard

PROTYPOPHAEMYIA Blanchard, 1963:241. Type-species, *Ypophaemyia haywardi* Blanchard (orig. des.).
 HAYWARDI (Blanchard), 1942a:370, fig. 13 (*Ypophaemyia*). Type-locality: Argentina, Tucumán.
 TOWNSENDI Blanchard, 1963:245. Type-locality: Argentina, Santa Fé. Distr. — Argentina (Santa Fé, Corrientes).

Genus WINTHEMIA Robineau-Desvoidy

WINTHEMIA Robineau-Desvoidy, 1830:173. Type-species, *Tachina variegata* Meigen (as Fabricius; Robineau-Desvoidy, 1863:207).



- Microtrichodes* Macquart, 1846:288 (1846:160). Type-species, *analis* Macquart (mon.).
- Masipoda* Brauer & Bergenstamm, 1889:162 (1889:94). Type-species, *geminata* Brauer & Bergenstamm (orig. des., as gen. n., sp. n.).
- Okeopsis* Townsend, 1927a:267. Type-species, *palpalis* Townsend (orig. des.). N. SYN.
- Hemimasipoda* Townsend, 1927a:267. Type-species, *brasiliensis* Townsend (orig. des.).
- Prowinthemia* Townsend, 1928b:151. Type-species, *paraguayensis* Townsend (orig. des.).
- Promasipoda* Townsend, 1934a:399. Type-species, *pinguoides* Townsend (orig. des.) N. SYN.
- Winthemiopsis* Blanchard, 1963:212. Type-species, *grioti* Blanchard (orig. des.). N. SYN. Refs. — Reinhard, 1931a (rev.); Thompson, 1963c (genitalia, larva, key to Trinidad spp.).
- ALABAMAE (Townsend), 1940b:892 (*Hemimasipoda*). Type-locality: Brazil, São Paulo, Ribeirão Preto.
- ANALIS (Macquart), 1846:289 (1846:161), pl. 14, fig. 5 (*Microtrichodes*). Type-locality: Brazil, Minas Gerais. Ref. — Reinhard, 1931a:47, pl. 1, fig. 7.
- ANALISELLA Thompson, 1963c:978, fig. 20. Type-locality: Trinidad, Orange Grove.
- AUREONIGRA Thompson, 1963c:978, figs. 37, 38. Type-locality: Trinidad, Maracas Valley.
- BRASILIENSIS (Townsend), 1927a:315 (*Hemimasipoda*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- CILIGERA Robineau-Desvoidy, 1830:173. Type-locality: Brazil N. COMB.
- COMMUNIS Thompson, 1963c:982, figs. 17, 33. Type-locality: Trinidad, Floradale.
- CONSOBRINA (Wulp), 1890:68 (*Exorista*). Type-locality: Mexico, Tabasco, Teapa.
- CUYANA (Blanchard), 1963:223 (*Neowinthemia*). Type-locality: Argentina, La Rioja. N. COMB.
- DATANAE (Townsend), 1892e:288 (*Exorista*). Type-locality: U.S.A., New York. Distr. — Canada, U.S.A., Mexico.
- DEILEPHILAE (Osten-Sacken), 1887:164 (*Tachina*). Type-locality: Unknown. Distr. — Canada, U.S.A., Mexico.
- DUBIOSA Thompson, 1963c:968, fig. 16. Type-locality: Trinidad, Maracas Valley.
- ELEGANS (Bigot), 1857b:810 (pg. 340 in Spanish edition), pl. 20, fig. 7-7a (*Tachina*). Type-locality: "Cuba". Ref. — Reinhard, 1931a:54.
- ERYTHROPYGA (Bigot), 1889:257 (*Chaetolyga*). Type-locality: "Mexico". *rubidapex* Brauer, 1897:348 (*Chaetolyga*). *Nomen nudum*.
- FULVIDAPEX (Bigot), 1889:263 (*Tricoliga*). Type-locality: Unknown (as "Mexico" or "Celebes"). Unrecognized N. COMB.
- GEMINATA (Brauer & Bergenstamm), 1889:102 (1889:94) (*Masipoda*). Type-locality: "Mexico".
- GRIOTI (Blanchard), 1963:212, fig. 28 (*Winthemiopsis*). Type-locality: Argentina, Santa Fé. N. COMB.
- IGNOBILIS (Wulp), 1890:71 (*Exorista*). Type-locality: Mexico, Guerrero, Amulango (formely Amula). Ref. — Reinhard, 1931a:16.
- INTERMEDIA Reinhard, 1931a:41, pl. 1, fig. 4. Type-locality: U.S.A., Texas. Distr. — Central America, Ecuador, Brazil.



- LATEVITTATA (Wulp), 1890:66 (*Exorista*). Type-locality: Mexico, several places. Ref. — Reinhard, 1931a:36.
- LATIMANA (Wulp), 1890:67, pl. 3, fig. 10 (*Exorista*). Type-locality: "Mexico" (sev. places).
- MIMA Reinhard, 1931a:37, pl. 1, fig. 2. Type-locality: Peru, Rio Charape. Ref. — Thompson, 1963c:974.
- MONTANA Reinhard, 1931a:361, pl. 1, fig. 1. Type-locality: U.S.A., Arizona. Distr. — U.S.A., Mexico.
- OBSKURELLA (Wulp), 1890:48 (*Nemoraea*). Type-locality: Mexico, Guerrero. N. COMB.
- OCCIDENTIS Reinhard, 1931a:22. Type-locality: Canada, British Columbia. Distr. — U.S.A., Mexico.
- OKEFENOKEENSIS Smith, 1916:95. Type-locality: U.S.A., Georgia. Distr. — U.S.A., Cuba.
- PARAGUAYENSIS (Townsend), 1928c:151 (*Prowinthemia*). Type-locality: Paraguay, Santa Trinidad.
- PALPALIS Townsend, 1927a:338 (*Okeopsis*). Type-locality: Brazil, São Paulo, Itaquaquecetuba N. COMB.
- bicolor* Reinhard, 1931a:46. Type-locality: Brazil, São Paulo, Itaquaquecetuba N. SYN.
- PATAGONICA (Blanchard), 1963c:221 (*Neowinthemia*). Type-locality: Argentina, Neuquén, Plottier. N. COMB.
- PERUVIANA (Townsend), 1928c:162 (*Okea*). Type-locality: Peru, Lima.
- signata* Reinhard, 1931a:51, pl. 1, fig. 10. Type-locality: Peru, Lima.
- PICEA (Walker), 1852:293 (*Tachina*). Type-locality: "Colombia". Unrecognized. Ref. — Reinhard, 1931a:54.
- PINGUIS (Fabricius), 1805:302 (*Musca*). Type-locality: "South America". Ref. — Townsend, 1931b:174 (notes on types).
- PINGUIOIDES (Townsend), 1934a:399 (*Promasipoda*). Type-locality: Brazil, Pará, Boa Vista, Tapajós River. N. COMB.
- Ref. — Silva, 1944 (biol.).
- POLLINOSA Thompson, 1963:984. Type-locality: Trinidad, Plum Mitán Road.
- PYRRHOPYGA (Wiedemann), 1819:53 (*Tachina*). Type-locality: Brazil, Bahia.
- RUBRA Vimmer & Soukup, 1940:379 (as *Winthemia*, error). Type-locality: Peru (sev. places). Unrecognized.
- RUBRICORNIS (Wulp), 1890:66 (*Exorista*). Type-locality: Mexico, Veracruz, North Yucatan N. COMB.
- RUFICORNIS (Blanchard), 1942b:101 (*Okeopsis*). Type-locality: Argentina, Tucumán N. COMB.
- RUFILATERA (Rondani), 1850b:171 (*Exorista*). Type-locality: "Venezuela". Unrecognized.
- SEXUALIS Curran, 1927c:7. Type-locality: Puerto Rico, Arecibo and Adjuntas.
- SINGULARIS Reinhard, 1931a:38, pl. 1, fig. 3. Type-locality: Argentina, Tucumán. Distr. — Argentina, Trinidad. Ref. — Thompson, 1963c:976.
- SORORCULA (Wulp), 1890:68 (*Exorista*). Type-locality: Mexico, Morelos, Cuernavaca N. COMB.
- SUBPICEA (Walker), 1852:297 (*Tachina*). Type-locality: "Brazil". Unrecognized. Ref. — Reinhard, 1931a:54.
- TESSELLATA (Wulp), 1890:56 (*Mystacella*). Type-locality: Mexico, Guerrero and Morelos N. COMB.
- TRICOLOR (Wulp), 1890:67, pl. 3, fig. 9 (*Exorista*). Type-locality: Mexico (Morelos, Tabasco), Costa Rica (Cache). Distr. — Central America, Brazil (Mato Grosso, São Paulo). Ref. — Reinhard, 1931a:42.

TRINITATIS Thompson, 1963a:971, fig. 29. Type-locality: Trinidad, Chaguanas.
 VARIEGATA (Wulp), 1890:48 (*Nemoraca*). Type-locality: Mexico, Guerrero.
 N. comb. This species is a secondary homonym of *W. variegata* (Meigen), 1824, but I prefer do not rename at this time).
 XANTHOCERA (Wiedemann), 1830:329 (*Tachina*). Type-locality: "Brazil".
 Distr. — Mexico, Central and South America. Ref. — Aldrich, 1927c:14.

UNPLACED SPECIES OF WINTHEMIINI

GLABRICULA Wulp, 1890:73 (*Phorocera*). Type-locality: Mexico, Guerrero.
 INTERSTINCTA Wulp, 1890:69 (*Exorista*). Type-locality: Mexico, Guerrero.
 NITIDIVENTRIS Bigot, 1889:258 (*Chaetolyga*). Type-locality: Mexico.
 OCCIDENTALIS Bigot, 1889:258 (*Chaetolyga*). Type-locality: Mexico.
 TENUIPALPIS Wulp, 1890:73 (*Exorista*). Type-locality: Mexico, Guerrero.

Tribe MASIPHYINI

The members of the tribe Masiphyini are parasites of praying-mantis.

Ref. — Thompson, 1963d (Trinidad sp.); Guimarães, 1966a (Brazilian sp., hosts); Heitzmann-Fontenelle & Guimarães, 1966 (biol.).

Genus ALSOPSYCHE Brauer & Bergenstamm

ALSOPSYCHE Brauer & Bergenstamm, 1891:313 (1891:91). Type-species, *nemoralis* Brauer & Bergenstamm (mon.).
 NEMORALIS Brauer & Bergenstamm, 1891:313 (1891:91). Type-locality: "Venezuela". Ref. — Aldrich, 1925c:110 (type redescri.).

Genus IGNOTOMYIA Reinhard


IGNOTOMYIA Reinhard, 1961:210. Type-species, *cunina* Reinhard (orig. des.).
 AUREA Thompson, 1963d:1308, figs. 24, 26, 28. Type-locality: Trinidad, St. Augustine.

Genus MANTEOMASIPHYA Guimarães

MANTEOMASIPHYA Guimarães, 1966a:221. Type-species, *brasiliensis* Guimarães (orig. des.).
 BRASILIENSIS Guimarães, 1966a:221, figs. 32, 37. Type-locality: Brazil, São Paulo, Barueri.

Genus MASIPHYA Brauer & Bergenstamm

MASIPHYA Brauer & Bergenstamm, 1891:313 (1891:9). Type-species, *brasiliiana* Brauer & Bergenstamm (mon.).
 MISERIATA (Wulp), 1890:95 (*Brachycoma*). Type-locality: "Mexico" (sev. places).



- BRASILIANA Brauer & Bergenstamm, 1891:313 (1891:9). Type-locality: Brazil, São Paulo, Varnhagen (formely Ypanema). Distr. — Brazil (São Paulo, Goiás). Ref. — Guimarães, 1966a:218.
- RUFICAUDA Wulp, 1890:94 (*Brachycoma*). Type-locality: Mexico. N. COMB.

Genus MASIPHYOIDEA Thompson

- MASIPHYOIDEA Thompson, 1963d:1313. Type-species, *chaetosa* Thompson (orig. des.).
- CHAETOSA Thompson, 1963d:1313, figs. 20, 23, 27 (as *Masphyoidea*, misprint). Type-locality: Trinidad, St. Augustine. Ref. — Guimarães, 1966a:215.

Genus MICROMASIPHYA Townsend

- MICROMASIPHYA Townsend, 1934a:399. Type-species, *curta* Townsend (orig. des.).
- CURTA Townsend, 1934a:399. Type-locality: Brazil, Pará, Urucurituba, Tapajós River. Ref. — Guimarães, 1966a:216.

Genus MYSTACOMYOIDEA Thompson

- MYSTACOMYOIDEA Thompson, 1963d:1318. Type-species, *mirabilis* Thompson (orig. des.).
- MIRABILIS Thompson, 1963d:1318, fig. 25. Type-locality: "Trinidad".
- SPINOSA Guimarães, 1966a:226, figs. 45, 47. Type-locality: Brazil, São Paulo, Barueri.

Genus NEOMASIPHYA Guimarães

- NEOMASIPHYA Guimarães, 1966a:207. Type-species, *thompsoni* Guimarães (orig. des.).
- LENKOI Guimarães, 1966a:210, figs. 4, 11. Type-locality: Brazil, São Paulo, Paraná River, Pôrto Cabral.
- THOMPSONI Guimarães, 1966a:208, figs. 1, 3. Type-locality: Brazil, São Paulo, Paraná River, Pôrto Cabral.

Genus OROMASPHYA Townsend

- OROMASIPHYA Townsend, 1927a:249. Type-species, *ornata* Townsend (orig. des.).
- ORNATA Townsend, 1927a:342. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- URBANAE Guimarães, 1966a:212, figs. 12-16. Type-locality: Brazil, São Paulo, Santo Amaro, Cocaia.

Genus PARAPHASIOPSIS Townsend

- PARAPHASIOPSIS Townsend, 1917b:232. Type-species, *mellicornis* Townsend (orig. des.).

MELLICORNIS Townsend, 1917b:232. Type-locality: Brazil, Mato Grosso, Chapada.

TRINITATIS Thompson, 1963d:1311, figs. 18, 19. Type-locality: "Trinidad".

Genus PHASIOPSIS Townsend

PHASIOPSIS Townsend, 1912b:108. Type-species, *floridana* Townsend (orig. des.).

MANTEOPHAGA Guimarães, 1966a:225, figs. 38, 40. Type-locality: Brazil, São Paulo, Paraná River, Pôrto Cabral.

TRIANGULARIS Thompson, 1963d:1306, fig. 17. Type-locality: Trinidad, Floradale.

Genus PROMASIPHYA Townsend

PROMASIPHYA Townsend, 1927a:379. Type-species, *Masiphya confusa* Aldrich (orig. des.).

CONFUSA (Aldrich), 1925c:109 (*Masiphya*). Type-locality: U.S.A., Florida. Distr. — U.S.A., Mexico.

brasiliانا Brauer & Bergenstamm of Coquillett, 1897:81.

SUBTILIPALPIS Wulp, 1890:96 (*Brachycoma*). Type-locality: Mexico, Guerrero and Veracruz. N. comb.

TOWNSENDI (Aldrich), 1925c:110 (*Masiphya*, as *confusa* var.). Type-locality: U.S.A., Arizona; Mexico.

Genus PSEUDOMASIPHYA Thompson

PSEUDOMASIPHYA Thompson, 1963d:1316. Type-species, *petiolata* Thompson (orig. des.).

PETIOLATA Thompson, 1963d:1316. Type-locality: Trinidad, Maracas Valley.

Genus PROPHASIOPSIS Townsend

PROPHASIOPSIS Townsend, 1927a:250. Type-species, *polita* Townsend (orig. des.).

LOPESI Guimarães, 1966a:22, fig. 37. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

POLITA Townsend, 1927a:351. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus STURMIMASIPHYA Townsend

STURMIMASIPHYA Townsend, 1935d:230. Type-species, *ciliata* Townsend (orig. des.).

CILIATA Townsend, 1935d:230. Type-locality: Brazil, São Paulo, São Paulo, Vila Ema.

Genus THELAIROPHASIA Townsend

THELAIROPHASIA Townsend, 1919b:173. Type-species, *transita* Townsend (orig. des.).

TRANSITA Townsend, 1919b:173. Type-locality: Peru, Jaen, Charape River.



Tribe SIPHOSTURMIINI

Ref. — Thompson, 1963d.

Genus MICROSILLUS Aldrich

MICROSILLUS Aldrich, 1926b:20. Type-species, *Houghia baccharis* Townsend (orig. des.).

Promicrosillus Townsend, 1935d:230. Type-species, *Siphosturmia pollinosa* Townsend (orig. des.).

POLLINOSUS (Townsend), 1912a:321 (*Siphosturmia*). Type-locality: Peru, Rio Chira Valley, Somate.

RUFICAUDUS Thompson, 1963d:1294, figs. 6, 7. Type-locality: "Trinidad".

Genus MYSTACOMYIA Giglio-Tos

MYSTACOMYIA Giglio-Tos, 1893:4. Type-species, *Mystacella rubriventris* Wulp (orig. des.).

RUBRIVENTRIS (Wulp), 1890:52 (*Mystacella*). Type-locality: Mexico, Veracruz, Atoyac and Tuxpango. Ref. — Guimarães, 1966a:228.

Genus SIPHOSTURMIA Coquillett

SIPHOSTURMIA Coquillett, 1897:83. Type-species, *Argyrophylax rostrata* Coquillett (orig. des.).

Siphosturmiopsis Townsend, 1915h:91. Type-species, *rafaeli* Townsend (orig. des.). Ref. — Thompson, 1963d.

RAFAELI (Townsend), 1915h:91 (*Siphosturmiopsis*). Type-locality: Mexico, Veracruz, San Rafael.

Tribe CARCELIINI

The species of Carceliini parasitize lepidopterous larvae and a few attack sawflies.

Genus AMAZOHOUGHIA Townsend

AMAZOHOUGHIA Townsend, 1934:400. Type-species, *argentifrons* Townsend (orig. des.).

ARGENTIFRONS Townsend, 1934:400. Type-locality: Brazil, Pará, Boa Vista, Tapajós River. Distr. — Brazil (Pará), Trinidad. Ref. — Thompson, 1954:98.

FLAVIPES Thompson, 1964:120, figs. 14, 17, 27. Type-locality: "Trinidad".

Genus ANADISCALIA Curran

ANADISCALIA Curran, 1934d:446. Type-species, *basalis* Townsend (orig. des.).

BASALIS Curran, 1934d:468. Type-locality: Panama, Corozal, Canal Zone.

Genus ARIDALIA Curran

- ARIDALIA Curran, 1934d:459. Type-species, *lateralis* Curran (orig. des.).
LATERALIS Curran, 1934d:469, figs. 143, 168. Type-locality: Panama, Canal Zone, Corozal.

Genus BRACHYMASICERA Townsend

- BRACHYMASICERA Townsend, 1911b:133, 143 (also, 1912a:340). Type-species, *polita* Townsend (orig. des., as gen. n., sp. n.).
NIGRIPES Townsend, 1928c:159. Type-locality: Peru, Chiclayo, Pomalca.
POLITA Townsend, 1911b:133, 143 (also, 1912a:341). Type-locality: Peru, Piura.
SUBPOLITA Townsend, 1912a:341 (as *polita* ssp.). Type-locality: Peru, Piura.

Genus CALOCARCELIA Townsend

- CALOCARCELIA Townsend, 1927a:266. Type-species, *fasciata* Townsend (orig. des.) = *cingulata* (Fabricius).
AZULIHOUGHIA Townsend, 1934a:401. Type-species, *amazonica* Townsend (orig. des.). N. SYN.
Ref. — Thompson, 1964:130 (key to Trinidad, species).
AMAZONICA (Townsend), 1934a:401 (*Azulioughia*). Type-locality: Brazil, Pará, Boa Vista, Tapajós River. N. COMB.
AUREOCEPHALA Thompson, 1964:136, figs. 28, 32, 33, 39, 41. Type-locality: Trinidad, Santa Cruz Valley.
CINGULATA (Fabricius), 1805:301 (*Musca*). Type-locality: "West Indies".
fasciata Townsend, 1927a:293. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
MINIMA Thompson, 1964:146, figs. 29, 35, 43, 44. Type-locality: Trinidad, Forest, W. of Arima.
ORELLANA Townsend, 1929:376. Type-locality: Peru, Orellana. Ref. — Thompson, 1964:130. Distr. — Trinidad, Peru.
TRINITATIS Thompson, 1964:141, figs. 26, 34, 42, 45, 46. Type-locality: Trinidad, "Brazil" (village).

Genus CARCELIA Robineau-Desvoidy

- CARCELIA Robineau-Desvoidy, 1830:176. Type-species, *bombylans* Robineau-Desvoidy (Townsend, 1916:6).
FLAVIROSTRIS (Wulp), 1890:69, pl. 3, fig. 7-7a (*Exorista*). Type-locality: Mexico (sev. places).
GRISEOMICANS (Wulp), 1890:74 (*Exorista*). Type-locality: Mexico (sev. places), and Costa Rica.
HALLIANA Cortés, 1945c:27. Type-locality: Chile, Malleco, Angol.
INCULTA (Wiedemann), 1830:33 (*Tachina*). Type-locality: "Brazil". Rcf. — Aldrich, 1925:467.
LAGOAE (Townsend), 1891c:159 (*Exorista*). Type-locality: Mexico, Guana-juato. Distr. — U.S.A., Mexico, Jamaica. Ref. — Sellers, 1943.
flavirostris authors, in part, not Wulp.

- RECLINATA (Aldrich & Webber), 1924:32 (*Zenillia*). Type-locality: U.S.A., N. Mexico. Distr. — U.S.A., Mexico.
RUBRELLA Robineau-Desvoidy, 1830:179 (also, 1863:241). Type-locality: Dominican Republic, Santo Domingo. Unrecognized.

Genus CARCELIOCEPHALA Townsend

- CARCELIOCEPHALA Townsend, 1934a:402. Type-species, *crypta* Townsend (orig. des.).
CRYPTA Townsend, 1934a:402. Type-locality: Brazil, Boa Vista, Tapajós River. Ref. — Thompson, 1963b:323.
SIMILLIMA Thompson, 1963b:327, figs. 34, 36. Type-locality: Trinidad. Maracas Valley.

Genus CARCELIODORIA Townsend

- CARCELIODORIA Townsend, 1928b:150. Type-species, *palpalis* Townsend (orig. des.).
PALPALIS Townsend, 1928b:150. Type-locality: Peru, Huallaga.

Genus CHAETOSISYROPS Townsend

- CHAETOSISYROPS Townsend, 1912a:320. Type-species, *montanus* Townsend (orig. des.).
MONTANUS Townsend, 1912a:321. Type-locality: Peru, Matucana.

Genus EUSISYROPA Townsend

- EUSISYROPA Townsend, 1908:97. Type-species, *Tachina blanda* Osten-Sacken (mon.).
Ref. — Sellers, 1943 (key in *Zenillia*); Thompson, 1963b:297.
AURATOFONTALIS (Brèthes), 1908:475 (*Exorista*). Type-locality: Argentina, Buenos Aires. Ref. — Blanchard, 1959.
AUTOGRAPHAE (Sellers), 1943:5, 7, 23 (*Zenillia*). Type-locality: Cuba, Baraguá.
BLANDOIDES Thompson, 1963b:297, figs. 18, 20. Type-locality: Trinidad, Sta. Cruz Valley.
TUCUMANENSIS (Sellers), 1943:21 (*Zenillia*). Type-locality: Argentina, Tucumán.
VIRILIS (Aldrich & Webber), 1924:40 (*Zenillia*, as *blanda* ssp.). Type-locality: U.S.A., Illinois (as N. York. error). Distr. — U.S.A., Mexico.

Genus GYMNOCARCELIA Townsend

- GYMNOCARCELIA Townsend, 1919a:582. Type-species, *ricinorum* Townsend (orig. des.).
RICINORUM Townsend, 1919a:582. Type-locality: U.S.A., Florida. Distr. — U.S.A., Mexico.

Genus MONTSERRATIA Thompson

- MONTSERRATIA Thompson, 1964:98 (key), 149. Type-species, *ovipara* Thompson (mon.).
OVIPIRA Thompson, 1964:149. Type-locality: Trinidad, Montserrat Hill.



Genus NEPOCARCELIA Townsend

NEPOCARCELIA Townsend, 1927a:267. Type-species, *fulva* Townsend (orig. des.).

antarctiae Blanchard, in [Marchionatto], 1940:28. *Nomen nudum*.

FULVA Townsend, 1927a:336. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

PALUSTRAE (Brèthes), 1908:473 (*Exorista*). Type-locality: Argentina, Buenos Aires. Ref. — Blanchard, 1958:39.

Genus OOMASICERA Townsend

OOMASICERA Townsend, 1911b:145 (also, 1912a:337). Type-species, *chaetosa* Townsend (orig. des., as gen. n., sp. n.).

CHAETOSA Townsend, 1911b:145 (also, 1912a:338). Type-locality: Peru, Valle

Genus SISYROPA Brauer & Bergenstamm

SISYROPA Brauer & Bergenstamm, 1889:163 (1889:95). Type-species, *Tachina thermophila* Wiedemann (mon.).

ALYPIAE Sellers, 1943:101. Type-locality: U.S.A. Distr. — U.S.A., Mexico.

MELALEUCA (Wiedemann), 1830:320 (*Tachina*). Type-locality: "Brazil".

PROSOPINA Brauer & Bergenstamm, 1891:347 (1891:43). Type-locality: "Brazil".

RUFIVENTRIS Brauer & Bergenstamm, 1891:346 (1891:42). Type-locality: "Brazil".

UNPLACED SPECIES OF CARCELIINI

ADJUNCTA Wulp, 1890:55 (*Mystacella*). Type-locality: Mexico, Guerrero.

BREVIS Wulp, 1890:64 (*Exorista*). Type-locality: Mexico, Guerrero.

LONGICORNIS Wulp, 1890:99 (*Brachycoma*). Type-locality: Mexico, Veracruz.

NORMULA Wulp, 1890:109 (*Masicera*). Type-locality: "Mexico".

Tribe ERYCIINI

(Masicerini, Lydellini, Phrynoiini)

The species of Eryciini parasitize various lepidopterous larvae, and few attack sawfly larvae.

Genus ABOLODORIA Townsend

ABOLODORIA Townsend, 1934:400. Type-species, *abdominalis* Townsend (orig. des.).

ABDOMINALIS Townsend, 1934:400. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus ACTINOPROSOPA Townsend

- ACTINOPROSOPA Townsend, 1927a:258. Type-species, *facialis* Townsend (orig. des.).
- FACIALIS Townsend, 1927a:283. Type-locality: Brazil, São Paulo. Itaquaquecetuba.
- MARIS Townsend, 1929:374. Type-locality: Brazil, São Paulo. Itaquaquecetuba.

Genus ADIDYMA Townsend

- ADIDYMA Townsend, 1935d:230. Type-species, *adversa* Townsend (orig. des.).
- ADVERSA Townsend, 1935d:231. Type-locality: Brazil, São Paulo, São Vicente.

Genus AMETADORIA Townsend

- AMETADORIA Townsend, 1927a:276. Type-species, *unispinosa* Townsend (orig. des.).
- UNISPINOSA Townsend, 1927a:285. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus APLOMYA Robineau-Desvoidy

- APLOMYA Robineau-Desvoidy, 1830:184. Type-species, *zonata* Robineau-Desvoidy (Robineau-Desvoidy, 1863:459, 460).
- SELLERSI (Thompson), 1966:356 (*Myxexoristops*). Type-locality: Trinidad, Centeno. N. comb.

Genus ANHANGABAHUIA Townsend

- ANHANGABAHUIA Townsend, 1931a:470. Type-species, *analís* Townsend (orig. des.).
- ANALIS Townsend, 1931a:471. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus ANTISTASEOPSIS Townsend

- ANTISTASEOPSIS Townsend, 1934a:401. Type-species, *brasiliensis* Townsend (orig. des.).
- BRASILIENSIS Townsend, 1934a:401. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus APLOMYOPSIS Townsend

- APLOMYOPSIS Townsend, 1927a:268. Type-species, *brasiliensis* Townsend (orig. des.).
- BRASILIENSIS Townsend, 1927a:287. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus BOLOHOUGHIA Townsend

- BOLOHOUGHIA Townsend, 1927a:275. Type-species, *aurometallica* Townsend (orig. des.).
AUROMETALLICA Townsend, 1927a:291. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
mendocina Blanchard, in [Marchionatto] (1941:21). *Nomen nudum*.

Genus BOURQUINIA Blanchard

- BOURQUINIA Blanchard, 1935:5. Type-species, *deaurata* Blanchard (orig. des.).
DEAURATA Blanchard, 1935:6, figs. 1, 4. Type-locality: Argentina.

Genus CAENIOPSIS Townsend

- CAENIOPSIS Townsend, 1927a:275. Type-species, *brevifrons* Townsend (orig. des.).
BREVIFRONS Townsend, 1927a:292. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus CAMPTOPHRYNO Townsend

- CAMPTOPHRYNO Townsend, 1927a:277. Type-species, *orbitalis* Townsend (orig. des.).
ORBITALIS Townsend, 1927a:294. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus CASAHUIRIA Townsend

- CASAHUIRIA Townsend, 1919a:581. Type-species, *cornuta* Townsend (orig. des.).
CORNUTA Townsend, 1919a:582. Type-locality: Peru, Casahuiri.

Genus CEROMASIOPSIS Townsend


- CEROMASIOPSIS Townsend, 1927a:273. Type-species, *brasiliensis* Townsend (orig. des.).
BRASILIENSIS Townsend, 1927a:296. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus CHAETOPHOROCERA Townsend

- CHAETOPHOROCERA Townsend, 1912a:342. Type-species, *andina* Townsend (orig. des.).
ANDINA Townsend, 1912a:342. Type-locality: Peru, Matucana.
FUSCOSA Townsend, 1912a:343. Type-locality: Peru, San Gaban at Uruhuasi bridge.

Genus CHRYSOEXORISTA Townsend

- CHRYSOEXORISTA Townsend, 1915m:435. Type-species, *viridis* Townsend (orig. des.).



- Chrysodoria* Townsend, 1934:402. Type-species, *chalcos* Townsend (orig. des.). N. SYN.
- ANGUSTIFRONS Townsend, 1916h:21 (as *viridis* ssp.). Type-locality: Brazil, Mato Grosso, Chapada.
- CAERULEIVENTRIS (Wulp), 1890:64 (*Exorista*). Type-locality: Mexico, Tabasco. N. COMB.
- CHALCOS (Townsend), 1934:402 (*Chrysodoria*). Type-locality: Brazil, Pará, Boa Vista, Rio Tapajós. N. COMB.
- FACIALIS (Sellers), 1943:28 (*Zenillia*). Type-locality: Mexico, Veracruz.
- LINEATA (Wulp), 1890:54 (*Mystacella*). Type-locality: Mexico, Tabasco.
- NIGRICAUDA (Wulp), 1890:70 (*Exorista*). Type-locality: Mexico, Veracruz, Atoyac. N. COMB.
- OCHRACEA (Wulp), 1890:63 (*Exorista*). Type-locality: "Mexico". Distr. — U.S.A., Mexico, Costa Rica.
- TAGLINOI (Sellers), 1945:25 (*Zenillia*). Type-locality: Argentina, Tucumán, Tafi Viejo.
- VIRIDIS Townsend, 1915m:435. Type-locality: Peru, San Gaban Canyon, Casahuiri. Ref. — Townsend, 1931b:182.

Genus CHRYSOHOUGHIA Townsend

- CHRYSOHOUGHIA Townsend, 1935d:231. Type-species, *chlorescens* Townsend (orig. des.).
- CHLORESCENS Townsend, 1935d:231. Type-locality: "Trinidad".

Genus CHRYSOMETOPIOPS Townsend

- CHRYSOMETOPIOPS Townsend, 1916h:19. Type-species, *smithii* Townsend (orig. des.).
- SMITHII Townsend, 1916h:19. Type-locality: Brazil, Mato Grosso, Chapada.

Genus CHRYSOPHRYNO Townsend

- CHRYSOPHRYNO Townsend, 1927a:267. Type-species, *egensis* Townsend (orig. des.).
- Chrysophryna*, error.
- ANDINENSIS Townsend, 1929:373. Type-locality: Peru, Tabalosos.
- EGENSIS Townsend, 1927a:298. Type-locality: Amazonas, Tefé.

Genus CHRYSOPHRYXE Sellers

- CHRYSOPHRYXE Sellers, 1943:106. Type-species, *tibialis* Sellers (orig. des.).
- TIBIALIS Sellers, 1943:106. Type-locality: "Brazil".

Genus CHRYSOTRYPHERA Townsend

- CHRYSOTRYPHERA Townsend, 1935d:232. Type-species, *conica* Townsend (orig. des.).
- CONICA Townsend, 1935d:232. Type-locality: Brazil, São Paulo, São Vicente.



Genus DORIELLA Townsend

- DORIELLA Townsend, 1931a:471. Type-species, *Tachina distincta* Wiedemann (orig. des.).
DISTINCTA (Wiedemann), 1824:43 (*Tachina*). Type-locality: "West Indies".

Genus EUMACROHOUGHIA Townsend

- EUMACROHOUGHIA Townsend, 1927a:275. Type-species, *nuda* Townsend (orig. des.). Ref. — Thompson, 1963b:329 (key).
MINOR Thompson, 1963b:333, figs. 41. 42. Type-species, Trinidad, "Brazil" (village).
NUDA Townsend, 1927a:309. Type-locality: Brazil. São Paulo, Itaquaquecetuba. Ref. — Thompson, 1963b:329.

Genus FRONTINIELLA Townsend

- FRONTINIELLA Townsend, 1918:21. Type-species, *parancilla* Townsend (orig. des., as gen. n., sp. n.).
PARANCILLA Townsend, 1918:21 (as *pararcilla*, error). Type-locality U.S.A., Michigan. Distr. — U.S.A., Mexico.
LOXOSTEGEI Blanchard, 1962:249, fig. 1 (as *loxotesgei*, error). Type-locality: Argentina, Tucumán.

Genus ICONOFRONTINA Townsend

- ICONOFRONTINA Townsend, 1931a:330. Type-species, *minthoidea* Townsend (orig. des.).
MINTHOIDEA Townsend, 1931a:331. Type-locality: Argentina, Mendoza.

Genus LESFESIA Robineau-Desvoidy

- LESFESIA Robineau-Desvoidy, 1863:567. Type-species, *Erycia ciliata* Macquart (mon.).
Achaetoneura Brauer & Bergenstamm, 1891:334 (1891:30). Type-species, *hesperus* Brauer & Bergenstamm (Coquillett, 1910:502) = *frenchii* (Williston).
Zygofrontina Townsend, 1915m:427. Type-species, *capitis* Townsend (orig. des.).
Ypophaeomyia Townsend, 1916d:75. Type-species, *malacosomae* Townsend (orig. des.) = *archippivora* (Riley).
? *Euparafrontina* Brèthes, 1917:17. Type-species, *martinezi* Brèthes (orig. des.).
Prophryno Townsend, 1927a:262. Type-species, *aurulans* Townsend (orig. des.). N. SYN.
Achaetoneuropsis Townsend, 1927a:272. Type-species, *affinis* Townsend (orig. des.).
Frontina, authors, nota Meigen.
Refs. — Beneway, 1963; Thompson, 1966:372 (key to Trinidad species).
AFFINIS (Townsend), 1927a:282 (*Achaetoneuropsis*). Type-locality: Brazil, São Paulo, Cantareira. Ref. — Lima, 1950 (host).



- AFRA (Wulp), 1890:22 (*Brachycoma*). Type-locality: Mexico. Tabasco. Unrecognized.
- APICALIS (Wulp), 1890:122 (*Prospberrya*). Type-locality: Mexico. Guerrero, Omilteme.
- ARCHIPPIVORA (Riley), 1871:150 (*Tachina*). Type-locality: U.S.A. Distr. — Canada, U.S.A., Mexico.
- AURICEPS (Macquart), 1843:216 (1843:59) (*Masicera*). Type-locality: "Brazil" or "Chile".
- AURULANS (Townsend), 1927a:353 (*Prophryno*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. N. COMB.
- BARBATULA (Wulp), 1890:98 (*Brachycoma*). Type-locality: Mexico. Guerrero.
- BIGEMINATA (Curran), 1927a:9 (*Frontina*). Type-locality: Puerto Rico. Adjuntas.
- bosqi* Blanchard in [Marchionatto], 1940:29 (*Achaetoneura*). *Nomen nudum*.
- BRASILIENSIS (Townsend), 1917:231 (*Ypophaemyia*). Type-locality: Brazil. Mato Grosso, Corumbá (as *malacosomae*, ssp.).
- CAPITIS (Townsend), 1915:427 (*Zygofrontina*). Type-locality: Peru, Casma, San Rafael. Ref. — Thompson, 1966:373.
- CLAVIPALPIS Thompson, 1966:377, figs. 11, 12. Type-locality: Trinidad. "Brazil" (village).
- DANAI (Townsend), 1940b:893 (*Zygofrontina*). Type-locality: "Brazil".
- ERYTHROCAUDA (Curran), 1934c:516 (*Frontina*). Type-locality: Venezuela, Guanoco.
- FLAVICANS (Wulp), 1890:74 (*Exorista*). Type-locality: Mexico. Orizaba. N. COMB.
- GRIOTI (Blanchard), 1963:239, fig. 37 (*Ypophaemyia*). Type-locality: Argentina. Santa Fé.
- LANIIFERA (Webber), 1930:33 (*Achaetoneura*). Type-locality: Mexico. Distr. — U.S.A., Mexico.
- LATA (Wiedemann), 1830:322 (*Tachina*). Type-locality: Uruguay, Montevideo. Ref. — Aldrich, 1927:8.
- MARTINEZI (Brèthes), 1917:18 (*Euparafrontina*). Type-locality: Peru, Lima. Unrecognized.
- MENDESI (Townsend), 1940b:893 (*Zygofrontina*). Type-locality: Brazil, São Paulo. Campinas.
- POLLINOSA Thompson, 1966:380, figs. 13, 14. Type-locality: Trinidad. St. Augustine.
- PUMILA (Wulp), 1890:108 (*Masicera*). Type-locality: Mexico, Guerrero. N. COMB.
- PROTOGINOI (Blanchard), 1966a:28, fig. 1 (*Achaetoneuropsis*). Type-locality: Argentina, Buenos Aires, Quilmes.
- RECTINERVIS (Wulp), 1890:123 (*Prospberrya*). Type-locality: Mexico. Guerrero.
- RUFIFRONS (Roeder), 1885:346 (*Frontina*). Type-locality: "Puerto Rico". N. COMB.
- RUFOMACULATA (Blanchard), 1963:197 (*Rilleyella*). Type-locality: Argentina. Entre Rios, Concordia.
- TEXANA (Webber), 1930:24 (*Achaetoneura*). Type-locality: U.S.A. Distr. — U.S.A., Mexico.
- townsendi* (Blanchard in Griot) 1944:57 (*Ypophaemyia*; as *Ypohaemyia* error). *Nomen nudum*.
- XYCHUS (Walker), 1849:770 (*Tachina*). Type-locality: "Jamaica".

Genus *LYDELLA* Robineau-Desvoidy

- LYDELLA* Robineau-Desvoidy, 1830:112. Type-species, *griscscens* Robineau-Desvoidy (Robineau-Desvoidy, 1863:855).
RADICIS (Townsend), 1916g:19 (*Andrina*). Type-locality: U.S.A., District of Columbia. Distr. — Canada, U.S.A., Mexico.

Genus *MACROHOUGHIA* Townsend

- MACROHOUGHIA* Townsend, 1927a:261. Type-species, *marmorata* Townsend (orig. des.).
MARMORATA Townsend, 1927a:325. Type-locality: Brazil, São Paulo, Itaquaquecetuba and Cantareira.

Genus *MADREMYIA* Townsend

- MADREMYIA* Townsend, 1916g:305. Type-species, *parva* Townsend (orig. des.) = *saundersii* (Williston).
SAUNDERSII (Williston), 1889:1922 (*Phorocera*). Type-locality: Canada, New England. Distr. — Canada, U.S.A., Mexico. Ref. — Townsend, 1931b:182.
parva Townsend, 1916g:306. Type-locality Mexico, Chihuahua.

Genus *METAGONISTYLUM* Townsend

- METAGONISTYLUM* Townsend, 1927a:379. Type-species, *minense* Townsend (orig. des.).
Amazonella Myers, 1934:191. *Nomen nudum*.
MINENSE Townsend, 1927a:381. Type-locality: Brazil, Minas Gerais, Viçosa. Distr. — U.S.A. (introd. not established), West Indies, Panama, Guyana, Venezuela (introd.), Bolivia, Brazil (Amazonas, São Paulo, Pernambuco, Minas Gerais, Rio de Janeiro). Refs. — Myers, 1934 (biol.), Bartlett, 1941 (biol.); Parker, 1953 (immature stages).
myersi Townsend, 1939a:343. Type-locality: Brazil, Pará, Santarém.

Genus *NILEA* Robineau-Desvoidy

- NILEA* Robineau-Desvoidy, 1863:275. Type-species, *innoria* Robineau-Desvoidy (orig. des.).
Eutritochaeta Townsend, 1919a:580. Type-species, *carpocapsae* Townsend (orig. des.) = *noctuiiformis* (Smith).
AUREA (Blanchard), 1942b:99, fig. 4 (*Eutritochaeta*). Type-locality: Argentina, Chaco.
macromphaliae Blanchard in [Marchionatto], 1942:15 (*Phyllophorocera*). *Nomen nudum*.

Genus *OPHIROSTURMIA* Townsend

- OPHIROSTURMIA* Townsend, 1911b:133 (also, 1912a:335). Type-species, *cincta* Townsend (orig. des., as gen. n., sp. n.).
CINCTA Townsend, 1911b:133 (also, 1912a:335). Type-locality: Peru, Piura.



Genus PALPOZENILLIA Townsend

- PALPOZENILLIA Townsend, 1934:404. Type-species, *Zenillia palpalis* Aldrich (orig. des.).
DIATRAEAE Townsend, 1914b:340. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Ref. — Parker, 1953:56 (immature stages).
PALPALIS (Aldrich), 1932b:20, fig. 2 (*Zenillia*). Type-locality: Guyana, Wanaima. Distr. — Mexico, Venezuela, Guyana. Ref. — Box, 1952:485 (biology, immature stages).

Genus PAMMAERUS Aldrich

- PAMMAERUS Aldrich, 1927c:24. Type-species, *Sisyropa leptotrichopa* Brauer & Bergenstamm (orig. des.).
LEPTOTRICHOPUS (Brauer & Bergenstamm), 1891:347 (1891:43) (*Sisyropa*). Type-locality: Brazil. Ref. — Aldrich, 1927c:24.
QUADRUS (Wiedemann), 1830:328 (*Tachina*). Type-locality: "Brazil". Unrecognized.

Genus PATELLOA Townsend

- PATELLOA Townsend, 1916e:619. Type-species, *Phorocera leucaniae* Coquillett (orig. des.).
Yahuarphryno Townsend, 1927a:263. Type-species, *patelloides* Townsend (orig. des.). N. SYN.
Pateloa (error), Curran, 1934:444.
Patelloapsis Townsend, 1927a:263. Type-species, *similis* Townsend (orig. des.).
CONCOLOR (Townsend), 1929:378 (*Patelloapsis*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
FULVICEPS (Wulp), 1890:80 (*Phorocera*). Type-locality: Mexico, sev. places. N. COMB.
littoralis (Blanchard), 1963:209 (*Patelloapsis*). *Nomen nudum*.
OECETICOLA (Blanchard), 1963:209 (*Patelloapsis*). Type-locality: Argentina. San Juan.
NIGRIPALPIS Thompson, 1963b:268, fig. 62. Type-locality: Trinidad.
PATELLOIDES (Townsend), 1927a:368 (*Yahuarphryno*). Type-locality: Peru, Yahuar Mayo River.
PLURISERIATA (Aldrich & Webber), 1924:43 (*Phorocera*). Type-locality: U.S.A. Distr. — U.S.A., Mexico.
RUSTI (Aldrich), 1929b:3 (*Phorocera*). Type-locality: Argentina, Tucumán. Ref. — Blanchard, 1942:104.
SIMILIS (Townsend), 1927a:345 (*Patelloapsis*). Type-locality: Brazil, São Paulo, Itaquaquecetuba and Cantareira.
thyreyionis (Blanchard), 1963:209 (*Patelloapsis*). *Nomen nudum*.
TINCTA (Walker), 1852:287 (*Tachina*). Type-locality: "Brazil".
tucumana (Blanchard), 1963:209 (*Patelloapsis*). *Nomen nudum*.

Genus PATILLALIA Curran

- PATILLALIA Curran, 1934d:459. Type-species, *fasciata* Curran (orig. des.).
FASCIATA Curran, 1934d:469, figs. 158, 165. Type-locality: Panama, Canal Zone, Patilla Point (cf. Arnaud, 1963a:123).

Genus PETRARGYROPS Townsend

- PETRARGYROPS Townsend, 1927a:275. Type-species, *punctiger* Townsend (orig. des.).
- Sisyrohoughia* Townsend, 1927a:275. Type-species, *similis* Townsend (orig. des.). N. SYN.
- PUNCTIGER Townsend, 1927a:346. Type-locality: Brazil, São Paulo, Itaquaquecetuba. Ref. — Thompson, 1963:338.
- SIMILIS (Townsend), 1927a:357 (*Sisyrohoughia*). Type-locality: Brazil, São Paulo, Itaquaquecetuba. N. COMB.

Genus PHASMOFRONTINA Townsend

- PHASMOFRONTINA Townsend, 1931a:473. Types-pecies, *perarida* Townsend (orig. des.).
- PERARIDA Townsend, 1931a:473. Type-locality: Peru, Chosica, Canyon.

Genus PHRYNOTACHINA Townsend

- PHRYNOTACHINA Townsend, 1927a:263. Type-species, *minor* Townsend (orig. des.).
- MINOR Townsend, 1927a:347. Type-locality: Peru, Chaquimayo.

Genus PHRYXE Robineau-Desvoidy

- PHRYXE Robineau-Desvoidy, 1830:158. Type-species, *athaliae* Robineau-Desvoidy (Robineau-Desvoidy, 1863:329, 358) = *vulgaris* (Fallén).
- TOLUCANA Reinhard, 1956c:104. Type-locality: Mexico, Nevado Toluca.

Genus PLAGIMASICERA Townsend

- PLAGIMASICERA Townsend, 1915m:411. Type-species, *petiolata* Townsend (orig. des.).
- Aeolofrontina* Townsend, 1928c:160. Type-species, *arida* Townsend (orig. des.).
- PETIOLATA Townsend, 1915m:411. Type-locality: Peru, Chosica Canyon. Ref. — Townsend, 1931b:173 (descr. notes).
- arida* Townsend, 1928c:160 (*Aeolofrontina*). Type-locality: Peru, Chosica Canyon.

Genus PROMETOPLOPS Townsend

- PROMETOPLOPS Townsend, 1927a:268. Type-species, *politus* Townsend (orig. des.).
- POLITUS Townsend, 1927a:351. Type-locality: Brazil, São Paulo, Itaquaquecetuba.

Genus PROSPHERYSA Wulp

- PROSPHERYSA Wulp, 1890:116. Type-species, *aemulans* Wulp (Coquillett, 1910:595).



- Dexiophana* Brauer & Bergenstamm, 1891:374 (1891:70). Type-species, *Prosperrya aemulans* Wulp (mon.).
- Sarcolydella* Townsend, 1927a:272. Type-species, *analís* Townsend (orig. des.).
- AEMULANS Wulp, 1890:117, pl. 3, figs. 14-14a. Type-locality: Mexico, Veracruz and Tabasco.
- ANALIS (Townsend), 1927a:355 (*Sarcolydella*). Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- CASTANIFRONS (Bigot), 1889:261 (*Ceromasia*). Type-locality: "Mexico". Unrecognized.

Genus SERICODORIA Townsend

- SERICODORIA Townsend, 1928b:149. Type-species, *sericea* Townsend (orig. des.).
- SERICEA Townsend, 1928b:150. Type-locality: Paraguay, Angelito.

Genus TAPAJOHUGHIA Townsend

- TAPAJOHUGHIA Townsend, 1934a:405. Type-species, *tropica* Townsend (orig. des.).
- TROPICA Townsend, 1934a:405. Type-locality: Brazil, Bôa Vista, Tapajós River.

Genus TREFOPHRYX Townsend

- TREFOPHRYX Townsend, 1908:95. Type-species, *cinerea* Townsend (orig. des.).
- CINEREA Townsend, 1908:96. Type-locality: Mexico, Morelos, Cuernavaca.

Genus TRINITODORIA Townsend

- TRINITODORIA Townsend, 1935d:233. Type-species, *media* Townsend (orig. des.).
- MEDIA Townsend, 1935d:233. Type-locality: "Trinidad".

Genus TRIODONTOPYGA Townsend

- TRIODONTOPYGA Townsend, 1927a:268. Type-species, *tridens* Townsend (orig. des.).
- FLAVOLIMBATA (Bigot), 1889:258 (*Chaetolyga*). Type-locality: "Mexico".
- TRIDENS Townsend, 1927a:362. Type-locality: Brazil, São Paulo, Itaquaquecetuba.
- TRINITATIS Thompson, 1963c:992, figs. 2, 14, 28. Type-locality: "Trinidad".
- VORAX (Wiedemann), 1830:315 (*Tachina*). Type-locality: Mexico. N. SYN. Ref. — Aldrich, 1925d:460.

Genus UCAYALIMYIA Townsend

- UCAYALIMYIA Townsend, 1927a:378. Type-species, *antlerata* Townsend (orig. des.).



ANTLERATA Townsend, 1927a:382. Type-locality: Peru, Chanchamayo, Ucayali River.

Genus VERRUGOMYIA Townsend

VERRUGOMYIA Townsend, 1927a:217. Type-species, *orbitalis* Townsend (orig. des.).

ORBITALIS Townsend, 1927a:364. Type-locality: Peru, Quebrada de Verrugas.

Genus VERRUGOPHRYNO Townsend

VERRUGOPHRYNO Townsend, 1927a:262. Type-species, *exoristoides* Townsend (orig. des.).

EXORISTOIDES Townsend, 1927a:364. Type-locality: Peru, Verrugas Canyon.

Genus YPOPHAEMYIOPS Townsend

YPOPHAEMYIOPS Townsend, 1935d:233. Type-species, *Prophryno myersi* Aldrich (orig. des.).

MYERSI (Aldrich), 1933b:173 (*Prophryno*). Type-locality: Guyana, Pacaraima Mts., upper Ireng River.

Genus ZENILLIA Robineau-Desvoidy

ZENILLIA Robineau-Desvoidy, 1830:152. Type-species, *Musca libathrix* Panzer (Robineau-Desvoidy, 1863:471).

ADAMSONI Thompson, 1963b:293. Type-locality: "Trinidad".

ANGUSTATA (Wulp), 1890:70 (*Exorista*). Type-locality: Mexico, Guerrero, Chilpancingo. Distr. — U.S.A., Mexico.

GOWDEYI Curran, 1926:112. Type-locality: Jamaica, Hill Gardens and Blue Castle.

UNPLACED SPECIES OF ERYCINI

ARCUATIPENNIS Macquart, 1855:121 (1855:101) (*Masicera*). Type-locality: Ecuador, Quito.

ATYPICA Curran, 1927:11 (*Spathidexia*). Type-locality: Puerto Rico, Adjuntas.

BRASILIENSIS Moreira, 1915:227, fig. (*Masicera*). Type-locality: Rio de Janeiro.

CHILENSIS Cortés, 1950:7 (*Phorocera*). Type-locality: Chile, Coquimbo, Elqui, Gualiguaica.

CHRYSOCEPHALA Walker, 1837:351 (*Tachina*). Type-locality: "South America".

CUBENSIS Macquart, 1848:206 (1848:46), pl. 5, fig. 5 (*Masicera*). Type-locality: "Cuba".

DISGREGA Wulp, 1890:171 (*Telothyria*). Type-locality: Mexico, Guerrero.

DISPUTANS Walker, 1860:302 (*Masicera*). Type-locality: "Mexico".

FULIGINIPENNIS Wulp, 1890:164 (*Didyma*). Type-locality: Mexico, Guerrero, Omilteme.

HUMILIS Wulp, 1890:72 (*Exorista*). Type-locality: Mexico, Guerrero.

INFIMA Wulp, 1890:204 (*Anisia*). Type-locality: Mexico, Guerrero.

- INGLORIA Wulp, 1890:119 (*Prospberrya*). Type-locality: Mexico, Guerrero, Omilteme.
- MISELLA Wulp, 1890:204 (*Anisia*). Type-locality: Mexico, Guerrero.
- MUSCARIA Wulp, 1890:83 (*Phorocera*). Type-locality: Mexico, Tabasco, Teapa.
- PLAGIOIDES Wulp, 1890:125 (*Prospberrya*). Type-locality: Mexico, Guerrero and Morelos.
- PLUTO Curran, 1934c:513 (*Phorocera*). Type-locality: Guyana, Bartica, Kartabo.
- PULLULA Wulp, 1890:162 (*Didyma*). Type-locality: Mexico, Guerrero.
- SUBPILOSA Wulp, 1890:110 (*Masicera*). Type-locality: Mexico, Veracruz.
- TRIFILATA Wulp, 1890:192 (*Anisia*). Type-locality: Mexico, Guerrero.
- VITTATA Walker, 1852:273 (*Tachina*). Type-locality: "Colombia".

UNPLACED SPECIES OF GONHINAE.

- BERGI Brèthes, 1908:474 (*Exorista*). Type-locality: Argentina, Buenos Aires.
- BREVIVENTRIS Wiedemann, 1830:297 (*Tachina*). Type-locality: "Brazil".
- CESSATRIX Walker, 1860:305 (*Lydella*). Type-locality: "Mexico".
- CINCTA Giglio-Tos, 1893:4 (*Chaetogena*). Type-locality: Mexico, Orizaba.
- DESITA Walker, 1861:299 (*Eurigaster*). Type-locality: "Mexico".
- ELONGATA Rondani, 1848:75 (*Phorocera*). Type-locality: "Brazil".
- GRACILENS Giglio-Tos, 1893:5 (*Tricholyga*). Type-locality: "Mexico".
- INSITA Giglio-Tos, 1893:5 (*Tricholyga*). Type-locality: "Mexico".
- LEPIDA Robineau-Desvoidy, 1830:153 (*Zenillia*). Type-locality: "Cuba".
- LINEARIS Wulp, 1890:86 (*Phorocera*). Type-locality: Mexico, Guerrero, Venta de Zopilote.
- LONGA Rondani, 1850:364 (*Exorista*). Type-locality: "Cuba".
- MODESTUS Bigot, 1857:341 (*Eurigaster*). Type-locality: "Cuba".
- NIVEIFACIES Macquart, 1851:162 (1851:189), pl. 17, fig. 7 (*Exorista*). Type-locality: "Brazil".
- NOBILIS Williston, 1896:355, pl. 11, fig. 92 (*Exorista*). Type-locality: "St. Vincent", West Indies.
- OBSCURUS Bigot, 1857:341 (*Eurigaster*). Type-locality: Cuba.
- PORTERI Brèthes, 1910:67 (*Exorista*). Type-locality: "Chile".
- TESSELLATA Röder, 1885:345 (*Exorista*). Type-locality: "Puerto Rico".
- VAGABUNDA Wulp, 1890:161 (*Didyma*). Type-locality: Mexico, Guerrero, Omilteme.
- VOLUCRIS Wulp, 1890:165 (*Didyma*). Type-locality: Mexico, Guerrero.

UNRECOGNIZED SPECIES OF TACHINIDAE

- ALBOMACULATA Robineau-Desvoidy, 1863:622 (*Peleteria*). Type-locality: "Chile".
- BRASILIENSIS Schiner, 1868:329 (*Nemoraea*). Type-locality: "Brazil".
- CHRYSICEPS (Robineau-Desvoidy), 1830:37 (*Jurinia*). Type-locality: "Brazil".
- CILIPES Macquart, 1843:246 (1843:89) (*Gymnostylia*). Type-locality: "Surinam".
- DISCOLOR Robineau-Desvoidy, 1863:660 (*Jurinia*). Type-locality: "Brazil".
- DUBIA Brauer, 1897:357 (*Melanota*). Type-locality: Mexico.



- FULIGIPENNIS Macquart, 1835:77 (*Echinomyia*). Type-locality: "Brazil".
 GLAUCA Giglio-Tos, 1893:6 (*Masicera*). Type-locality: "Mexico".
 GYMNOPS Macquart, 1843:227 (1843:70), pl. 8, fig. 1 (*Tryphera*). Type-locality: "America".
 HISTRIX Rondani, 1863:18 (*Cryptopalpus*). Type-locality: "Bogotá".
 INCLINANS Walker, 1857:199 (*Masicera*). Type-locality: "South America".
 I'SEA Walker, 1849:776 (*Tachina*). Type-locality: "unknown". (possibly New World).
 INNOVATA Walker, 1861:296 (*Tachina*). Type-locality: "Mexico".
 LATERALIS Macquart, 1843:228 (1843:71), pl. 8, fig. 3 (*Scopolia*). Type-locality: "America".
 LATERALIS Robineau-Desvoidy, 1863:664 (*Faurella*). Type-locality: "Chile".
 NECOPINA Walker, 1860:303 (*Masicera*). Type-locality: "Mexico".
 NIGRIFERA Walker, 1852:303 (*Tachina*). Type-locality: "Brazil".
 OCCIDENTALIS Wiedemann, 1830:335 (*Tachina*). Type-locality: "St. Thomas".
 NIGRIFRONS Bigot, 1887:cxli (*Elachipalpus*). Type-locality: "Mexico".
 PARTITA Bigot, 1889:270 (*Dexiosoma*). Type-locality: Mexico.
 PROBATA Walker, 1861:318 (*Hylemyia*). Type-locality: "Mexico".
 PICTIPENNIS Macquart, 1851:156, pl. 16, fig. 7 (*Nemoraea*). Type-locality: "Colombia".
 PLANIVENTRIS Macquart, 1851:178 (1851:205), pl. 18, fig. 13 (*Tachina*). Type-locality: "South America".
porteri Reed, 1907:1045 (*Tachina*). Type-locality: "Chile". *Nomen nudum*.
 PUMILA Robineau-Desvoidy, 1863:622 (*Peleteria*). Type-locality: "America".
 SALTATRIX Wiedemann, 1830:300 (*Tachina*). Type-locality: "West Indies".
 SENSIFERA Walker, 1861:314 (*Musca*). Type-locality: "Mexico".
 SESQUIPLEX Giglio-Tos, 1893:6 (*Masicera*). Type-locality: "Mexico".
 SPINIPENNIS Wiedemann, 1830:303 (*Tachina*). Type-locality: "Brazil".
 TENEBRIFERA Walker, 1852:302 (*Tachina*). Type-locality: "Brazil".
 TRICHONEURA Wulp, 1890:111 (*Masicera*). Type-locality: Mexico, Guerrero.
 UNICOLOR Macquart, 1846:295 (1846:167), pl. 20, fig. 9 (*Miltogramma*). Type-locality: "Brazil".
 USTA Giglio-Tos, 1893:6 (*Masicera*). Type-locality: "Mexico".

UNPLACED GENERA OF TACHINIDAE.

Genus EGGONIA Brauer & Bergenstamm

- EGGONIA Brauer & Bergenstamm, 1893:104 (1893:16). Type-species, *wulphi* Townsend (sub. mon.; Townsend, 1927a:271).
 WULPHI Townsend, 1927a:271. Type-locality: "Mexico".

Genus HYADESIMYIA Bigot

- HYADESIMYIA Bigot, 1888b:26. Type-species, *clausa* (Bigot, 1891:cxxxv).
 CLAUSA Bigot, 1888b:27, pl. 3, fig. 7. Type-locality: Chile, Cape Horn. Ref. — Aldrich, 1935:85.

Genus OLIGOESTRUS Townsend

- OLIGOESTRUS Townsend, 1932:1. Type-species, *oestroideus* Townsend (orig. des.).
 OESTROIDEUS Townsend, 1932:4. Type-locality: Argentina, Bariloche, Lake Nahuel Huapi.



NOMINA NUDA

Hemiarchoytas Vimmer, 1940:101. *Nomen nudum*.

Gymnojurinella Vimmer, 1940:101, also Vimmer & Soukup, 1940:210 (key).
223 (as *Gimnojurinella*, pag. 223). *Nomen nudum*.

Parajurinia Vimmer, 1940:101. *Nomen nudum*.

pastranai Blanchard, in [Marchionatto], 1940:37 (*Xenosturmia*). *Nomen nudum*.

UNRECOGNIZED GENERA AND SPECIES OF TACHINIDAE

Genus CHAETOEPALPUS Vimmer & Soukup

CHAETOEPALPUS Vimmer & Soukup, 1940:218. Type-species, *coquilleti*
Vimmer & Soukup (mon.).

COQUILLETI Vimmer & Soukup, 1940:218. Type-locality: Peru, Puno.

Genus METOPIOPSIS Vimmer & Soukup

METOPIOPSIS Vimmer & Soukup, 1940:362. Type-species, *aurea* Vimmer &
Soukup (mon.).

AUREA Vimmer & Soukup, 1940:363 (as *Metopiops*, misspelling). Type-loca-
lity: Peru, Puno.

Genus OLINDA Robineau-Desvoidy

OLINDA Robineau-Desvoidy, 1830:116. Type-species, *brasiliensis* Robineau-
Desvoidy (mon.).

BRASILIENSIS Robineau-Desvoidy, 1830:116. Type-locality: "Brazil".

Genus PARABOTHRIA Vimmer & Soukup

PARABOTHRIA Vimmer & Soukup, 1940:365, 371. Type-species, *punoensis*
Vimmer & Soukup (mon.).

PUNOENSIS Vimmer & Soukup, 1940:365. Type-locality: Peru, Puno

LITERATURE CITED

D'AGUILAR, J.

1957. Revision des Voriini de l'ancien monde (Dipt. Tachinidae). *Ann. Epiphyt.* 3:235-270, 6 pls., 2 figs.

ALDRICH, J. M.

1921. The muscoid genera *Pseudeuantha* and *Uramyia*. *Ins. Ins. Menst.* 9:83-92.

1923. A new parasitic fly bred from bean beetle. *Proc. Ent. Soc. Washington* 25:95-96.

- 1924a. Notes on some types of American Muscoid Diptera in the collection of the Vienna Natural History Museum. *Ann. Ent. Soc. Amer.* 17:209-218.
- 1924b. The muscoid genus *Genea* in the North America (Dipt.). *Ent. News* 35:210-214.
- 1925a. New Diptera or two-winged flies in the United States National Museum. *Proc. U. S. Natl. Mus.* 66(18):1-36, 1 fig.
- 1925b. Two new species of the tachinid genus *Lixophaga*, with notes and key (Diptera). *Proc. Ent. Soc. Wash.* 27:132-136.
- 1925c. Notes on some types of American Muscoid Diptera in the collection of the Vienna Natural History Museum (cont.). *Ann. Ent. Soc. Amer.* 18:107-130, 2 figs.
- 1925d. Notes on some types of American Muscoid Diptera in the collection of the Vienna Natural History Museum (cont.). *Ann. Ent. Soc. Amer.* 18:456-469, 1 fig.
- 1926a. American two-winged flies of the genus *Microphthalma* Macquart with notes on related forms. *Proc. U. S. Natl. Mus.* 69(13):1-8.
- 1926b. Descriptions of new and little known Diptera or two-winged flies. *Proc. U. S. Natl. Mus.* 69(22):1-26.
- 1926c. Notes on muscoid flies (Diptera) with retracted hind crossvein, with key and several new genera and species. *Trans. Amer. Ent. Soc.* 52:7-28.
- 1926d. Notes on the metallic green tachinids allied to *Gymnochacta*, with keys and one new Chinese genus (Diptera). *Ins. Ins. Menst.* 14:51-58.
- 1926e. North American two-winged flies of the genus *Cylindromyia* Meigen (*Ocyptera*, of authors). *Proc. U. S. Natl. Mus.* 68(23):1-27, 1 pl.
- 1927a. Notes on muscoid synonymy. *Bull. Brooklyn Ent. Soc.* 22:18-25.
- 1927b. A new species of *Oedematocera* reared from the tropical migratory locust (Diptera). *Proc. Ent. Soc. Was.* 29:17-18.
- 1927c. Redescription of types of American muscoid flies in the collection of the Vienna Natural History Museum with incidental notes. *Proc. U. S. Natl. Mus.* 72(7):1-35.
- 1927d. Notes on the Dexiid genera *Cordyligaster* and *Eucordyligaster*. *Jour. Wash. Acad. Sci.* 17(4):84-86.
- 1928a. A revision of the American parasitic flies belonging to the genus *Belvosia*. *Proc. U. S. Natl. Mus.* 73(8):1-45.
- 1928b. New Diptera or two-winged flies from South America. *Proc. U. S. Natl. Mus.* 74(1):1-25.
- 1928c. Notes on the synonymy of Diptera. *Proc. Ent. Soc. Wash.* 30(8):142-145.
- 1928d. A new species of *Oedematocera* with notes on *Schistocercophaga* Townsend (Dipt. Tachinidae). *Ent. News* 39:301-304.
- 1929a. Further studies of types of American muscoid flies in the collection of the Vienna Natural History Museum. *Proc. U. S. Natl. Mus.* 74(19):1-34, 2 figs.
- 1929b. New genera and species of muscoid flies. *Proc. U. S. Natl. Mus.* 76(15):1-13.



- 1930a. Notes on the types of American two-winged flies of the genus *Sarcophaga* and a few related forms, described by the early authors. *Proc. U. S. Natl. Mus.* 78(12):1-39, 3 pls.
- 1930b. Notes on synonymy of Diptera, n° 4. *Proc. Ent. Soc. Wash.* 32(2):25-28.
- 1931a. Notes on the tachinid genus *Chetonodexodes*, with one new species. *Ann. Mag. Nat. Hist.* (10)8:205-207.
- 1931b. Notes on Diptera n° 5. *Proc. Ent. Soc. Wash.* 33:116-121.
- 1931c. Notes on Francis Walker's types of North American flies of the family Tachinidae. *Proc. U. S. Natl. Mus.* 80(10):1-16.
- 1931d. North American two-winged flies of the genus *Spathimeigenia*, with descriptions of five new species. *Proc. U. S. Natl. Mus.* 80(11):1-10.
- 1932a. Records of dipterous insects of the family Tachinidae reared by the late George Dimmock, with description of one new species and notes on the genus *Anetia* Robineau-Desvoidy. *Proc. U. S. Natl. Mus.* 80(20):1-8.
- 1932b. New diptera, or two winged flies from America, Asia, and Java with additional notes. *Proc. U. S. Natl. Mus.* 81(9):1-28, 1 pl.
- 1933a. A remarkable new genus of Tachinidae (Diptera) from Brazil. *Rev. de Ent.* 3:437-441, 2 figs.
- 1933b. Two reared species of Tachinidae from South America. *Proc. Ent. Soc. Wash.* 35:170-173, 1 fig.
1934. Tachinidae, in British Museum (Natural History), *Diptera of Patagonia and South Chile based mainly on material in the British Museum (Natural History)* 7(1):1-170, figs. 1-21. London.

ALDRICH, J. M. & R. T. WEBBER


1924. The North American species of parasitic two-winged flies belonging to the genus *Phorocera* and allied genera. *Proc. U. S. Natl. Mus.* 63(17):1-90.

ALLEN, H. W.

1929. An annotated list of the Tachinidae of Mississippi. *Ann. Ent. Soc. Amer.* 22:676-690, 1 fig.

ARNAUD, P. H., JR. (see also SABROSKY & ARNAUD)

1951. A study of the genus *Paradejeania* Brauer and Bergenstamm (Diptera: Tachinidae or Larvaevoridae). *Canad. Ent.* 83:317-329, 9 figs.
1952. *Reinhardiana* new name for *Hypenomyia* Townsend (Diptera: Tachinidae or Larvaevoridae). *Pan-Pacific Ent.* 28:58.
1958. The entomological publication of Charles Henry Tyler Townsend [1863-1944], with lists of his new generic and specific names. *Microentomology* 23:1-63, 1 fig.
1960. A review of the genus *Spathidexia* Townsend (Diptera: Tachinidae). *Wasman Journ. Biol.* 18:1-36, 33 figs.
- 1963a. Types of the Tachinidae (Diptera) in the American Museum of Natural History. *Bull. Amer. Mus. Nat. Hist.* 125:101-138.



- 1963b. Systematic studies in the tribe Neominthoini (Diptera, Tachinidae). *Amer. Mus. Novit.* 2135:1-55, 44 figs.
- 1963c. *Perumyia embiaphaga*, a new genus and species of Neotropical Tachinidae (Diptera) parasitic on Embioptera. *Amer. Mus. Novit.* 2143:1-9, 14 figs.
- 1963d. A revision of the genus *Borgmeiermyia* Townsend (Diptera, Tachinidae). *Amer. Mus. Novit.* 2133:1-18, 17 figs.
1966. A revision of the parasitic fly genus *Polistiopsis* Townsend (Diptera, Tachinidae). *Amer. Mus. Novit.* 2241:1-12, 20 figs.

AUSTEN, E. E.

1907. The synonymy and generic position of certain species of Muscidae (sens. lat.) in the Collection of the British Museum, described by the late Francis Walker. *Ann. Mag. Nat. Hist.* (7)19:326-347.

BARTLETT, K.

1941. The biology of *Metagonistylum minense* Tns. a parasite of the sugar cane borer. *Puerto Rico Exp. Stat., Bull.* 40:1-20, 10 figs.

BEARD, R. L.

1940. The biology of *Anasa tristis* De Geer with particular reference to the tachinid parasite, *Trichopoda pennipes* Fab. *Connect. Agric. Exp. Station, Bull.* 440:593-680, 3 pls., 18 figs.

BENEWAY, D. F.

1961. *Androeuryps*, a new genus of Tachinidae (Diptera) from Central America. *Journ. Kans. Ent. Soc.* 34:44-47, 6 figs.
1963. A revision of the flies of the genus *Lespesia* (= *Achaetoneura*) in North America (Diptera: Tachinidae). *Univ. Kansas Sci. Bull.* 44:627-686, 34 figs.

BERRY, P. A.

1951. Biology and habits of cotton stainers: (Hemiptera: *Dysdercus* spp.) their natural enemies in South America and two parasitic flies imported into Puerto Rico. *Rev. Ent.* 22(1-3):329-342, illustr.

BERRY, P. A. & H. PARKER


1949. Investigations on a South American *Epilachna* sp. and the importation of its parasites *Lydinolydella metallica* Tns. into the United States. *Proc. Ent. Soc. Washington* 51:93-104.

BERTHOLD, A. A.

1927. *Latreille's Natürliche Familien der Tierreichs. Aus dem Französischen. Mit Anmerkungen und Zusätzen*, 606 pp. Weimar.

BEZZI, M.

1906. Noch Namen für einige Dipteren-Gattungen. *Ztschr. f. System. Hymenopt. u. Dipt.* 2:190-192.



1923. Fissicorn Tachinidae, with description of new forms from Australia and South America. *Proc. Linn. Soc. N. S. Wales* 48: 647-659, 8 figs.

1925. On the tachinid genus *Euthera* (Diptera), with description of new species from Australia, Africa and South America. *Proc. Linn. Soc. N. S. Wales* 50:275-283, 3 figs.

BIEZANKO, J. L. M.

1961. Castniidae, Zygaenidae, Dalceridae, Eucleidae, Megalopygidae, Cossidae et Hepialidae. *Arq. Ent. (B)* 14 Inst. Agron. Sul, Pelotas, 12 pp.

BIGOT, J. M. F.

1857a. Diptères nouveaux provenant du Chili. *Ann. Soc. Ent. France* (3)5:277-308, 2 col. plates.

1857b. Dipteros. Pp. 328-349, pl. 20 of insects and crustaceans, bound in vol. 8. In Sagra, R. de la, *Historia física, política y natural de la isla de Cuba*, vol. 7, 371 pp. Paris, "1856" (Spanish edition); pags. 783-829 in French edition, 1857.

1859. Dipterorum aliquot genera. *Rev. et Mag. de Zool.* (2)11:305-315, pl. 11.

1876. Diptères nouveaux ou peu connus, 6^e partie, VIII: Curie des Phasides (Phasidae, mihl). Genres *Trichopoda* (Macq.) et *Bogusia* (Rond.). *Ann. Soc. Ent. France* (5)6:389-400.

1878. Diptères exotiques nouveaux ou peu connus, 9^e partie, XIII: Genres *Ocyptera* (Latr.), *Ocypterulla*, *Exogaster* (Rond.). *Ann. Soc. Ent. France* (5)8:40-47.

1880. (Diagnoses d'un nouveau genre de Diptères de la tribu des Tachinidae). *Ann. Soc. Ent. France* (5)10:liiii.

1884a. (Description d'un nouveau genre et d'une nouvelle espèce de Diptères de la famille des Dexidae). *Ann. Soc. Ent. France* (6)4 (Bul.):xxxvii.

1884b. (Diagnose d'un nouveau genre de Diptères). *Ann. Soc. Ent. France* (6)4 (Bul.):lxix-lxx.

1884c. (Diagnose d'un genre et d'une espèce de Diptères) *Ann. Soc. Ent. France* (1883) (6)3 (Bull.):cviii-cix.

1885a. (Diagnoses de trois genres nouveaux de Diptères du groupe des Déxiaires). *Ann. Soc. Ent. France* (6)5 (Bul.):xi-xii.


1885b. (Diagnoses de deux genres nouveaux de Diptères du groupe des Déxiaires). *Ann. Soc. Ent. France*, ser. 6,5 (Bul.):xxv-xxvi; xxii-xxxiii.

1885c. (Diagnose de deux genres nouveaux de Diptère du groupe des Tachinides). *Ann. Soc. Ent. France*, ser. 6,5 (Bul.):xliv-xlvi; liv-lv.

1887. (Diagnoses de quelques espèces nouvelles de Diptères). *Ann. Soc. Ent. France*, ser. 6,7 (Bul.):cxxxix-cxlii.

1888a. Diptères nouveaux ou peu connus, 33^e partie, XLI: Tachinidae. *Ann. Soc. Ent. France*, ser. 6,8:77-101.

1888b. Diptères in *Mission scientifique du Cap. Horn*. 1882-1883. Tomo 6. Zoologie, Insectes:1-45, pls. 1-4, Paris.



1889. Diptères nouveaux on peu connus, 34^e partie, XLII. Diagnoses de nouvelles espèces. *Ann. Soc. Ent. France*, ser. 6,8:253-270.
 1891. (Notes). *Ann. Soc. Ent. France*, 60 (Bul.):cxxxv-cxxxvi.

BISCHOF, J.

1904. Beitrag zur Kenntniss der Muscaria schizometopa. *Verh. Zool. Bot. Gesell. Wien*, 54:79-101.

BLANCHARD, E.

1852. Orden IX. Dipteros. Pp. 327-468. In Gay, C. ed. *Historia fisica y política de Chile*. Zoologia, vol. 7, 471 pp. Paris.

BLANCHARD, E. E.

1935. Apuntes sobre Dípteros argentinos. *Rev. Argentina Ent.* 1:5-12, 3 figs.
 1937. Dípteros Argentinos nuevos o poco conocidos. *Rev. Soc. Ent. Argentina* 9:35-58, 8 figs.
 1940. Apuntes sobre muscoideos argentinos. *Ann. Soc. Cient. Argentina*, 129:224-246, 5 figs.
 1941. Los dípteros muscoideos del Museo de La Plata. Primera parte: Tachinidae. *Rev. Mus. La Plata* (n.s.), 2, Zoología: 341-379, 12 figs.
 1942a. Nuevos dípteros y himenópteros parásitos de la Republica Argentina. *Rev. Soc. Ent. Argentina*, 11:340-379, 17 figs.
 1942b. Parásitos de *Alabama argillacea* Hbn. en la Republica Argentina. Estudio preliminar (concl.). *An. Soc. Cient. Argentina*, 134: 94-128, 17 figs.
 1943a. Un díptero y seis himenópteros Argentinos, nuevos para la ciencia. *Rev. Soc. Ent. Argentina* 12:92-104, 7 figs.
 1943b. Los dípteros muscoideos del Museo de La Plata. Primera parte: Tachinidae (conclusion). *Rev. Mus. La Plata* (n.s.), 3:123-161, 29 figs.
 1943c. Un nuevo Dexiido *Ceracia subandina*, parásito de la tucura, *Dichroplus arrogans* Stål. *Rev. Soc. Ent. Argentina* 12(1):19-21, 1 fig.
 1943d. Un nuevo exoristido, importante parasito del gorgojo de las hortalizas (*Listroderes obliquus* Klug). *Rev. Soc. Ent. Argentina* 11:450-454, figs. a-e.
 1944. *Dallasimya bosqi*, nuevo genero y especie de triquiopodino (Dipt. Gymnosomatidae). *Rev. Argentina Ent.* 2:7-11, illus.
 1947a. Nuevos géneros y especies de insectos parásitos (Hymenoptera & Diptera) del Uruguay. *Com. Zool. Mus. Hist. Nat. Montevideo* 2(42):1-19, 7 figs.
 1947b. *Dasyuromyia lloydi* nueva especie de prosénido argentino. *Rev. Soc. Ent. Argentina* 13:258-262, figs. a-g.
 1950. Un nuevo parásito del gorgojo del tabaco. *Rev. Invest. Agropecuarias* 4:135-138, 1 fig.
 1954. Contribución al conocimiento de los Oestromuscarios entomófagos argentinos. I. Los Belvoslinos (Dipt. Exoristidae). *Min. de*



- Agr. y Ganad. Argentina, Publ. Inst. San. Veg.*, Año 10. ser. A (57):1-53, 17 figs.
1955. Un nuevo género de Actiini sin nervadura transversa posterior (Dipt. Exoristidae). *Rev. Soc. Ent. Argentina* 18:23-25, figs.
1957. *Parapoliops grioti*, nuevo Actiino útil argentino (Dipt.). *Rev. Soc. Ent. Argentina* 19:45-46 (1956).
1958. Tres Dípteros parásitos del bicho quemador (Dipt. Exoristidae). *An. Soc. Cient. Argentina* 166:35-40.
1959. Dípteros parásitos de orugas de Arctiidae ("Gatas peludas"). *Rev. Invest. Agr.* 13:157-182, 13 figs.
1962. Dos nuevos enemigos de oruguitas que viven sobre el "yuyo colorado" (Diptera, Exoristidae). *Rev. Invest. Agr.* 16:249-255, 2 figs.
1963. Dípteros parásitos de Noctuidae argentinos. *Rev. Invest. Agric.* 17:129-254, 38 figs.
- 1966a. Descripción de un díptero de la oruga o perro del naranjo, *Papilio thoas thoanthiades* Burm. (Lepidóptera). *Inst. Nac. de Tecn. Agropecuaria. Rev. de Invest. Agropecuarias*, Buenos Aires, ser. 5 (Pat. Veg.), 3(4):27-31, fig. 1.
- 1966b. Dípteros parásitos de escarabeoideos argentinos. *Rev. Invest. Agropecuarias*, Buenos Aires, ser. 5 (Pat. Veg.), vol. III, n° 11: 175-229, 19 figs.
- 1966c. Nuevos triquiopodinos argentinos, parásitos de hemípteros nocivos (Diptera, Gymnosomatidae). *Rev. Invest. Agropecuarias*, Buenos Aires, ser. 5 (Pat. Veg.), vol. III, n° 7: 59-95, 2 plates, 12 figs.

BOX, H. E.

1928. Observations upon *Lixophaga diatraeae* Townsend, a tachinid parasite of *Diatraea saccharalis* Fabr. in Porto Rico. *Bull. Ent. Res.* 19:1-6.
1952. *Palpozenillia palpalis* (Aldr.), a tachinid parasite of sugar cane moth borers (*Castnia* and *Diatraea*). *Proc. Hawaiian Ent. Soc.* 14(3):485-490, figs. 1-2.

BRAUER, F.

1895. Bemerkungen zu einigen neuen Gattungen der Muscarien und Deutung einiger Original-Exemplare. *Sitzber. K. Akad. der Wissen. Wien, Math.-Nat. Cl. Abt.* 1(109):582-604, 8 figs.
- 1897-1899. Beiträge zur Kenntniss der Muscaria Schizometopa. Bemerkungen zu den Original exemplaren der von Bigot, Macquart und Robineau-Desvoidy, beschreiben Muscaria Schizometopa aus der Sammlung des Herrn G. H. Verral, I-III. *Sitzber. K. Akad. der Wissen. Wien, Math. Nat. Cl. Abt.* 1(106):329-377, 1897 (107):493-546, 1898 (108):495-529, 1899.

BRAUER, F. & J. E. VON BERGENSTAMM

1889. Die Zweiflügler des Kaiserlichen Museums zu Wien. IV. Vorarbeiten zu einer Monographie der Muscaria Schizometopa (exclusive Antomyiidae). Pars I. *Denkschr. K. Akad. der Wissen.*



- Wien (Math-Nath. Cl.) 56 (1):69-180, 11 pls. Also printed separately in Wien, 1889, 112 pp.
1891. Die Zweiflügler des Kaiserlichen Museums zu Wien. VI. Vorarbeiten zu einer Monographie der Muscaria Schizometopa (exclusive Anthomyidae). Pars II. *Denkschr. K. Akad. der Wissen, Wien* (Math — Nat. Cl.) 60:89-240.
1893. Die zweiflügler des Kaiserlichen Museums zu Wien. VI. Vorarbeiten zu einer Monographie der Muscaria Schizometopa (exclusive Anthomyidae). Pars III. *Denkschr. K. Akad. der Wiss. Wien* (Math — Nat. Cl.) 60:89-240. Also printed separately in Wien, 1893, 152 pp.
1894. Die Zweiflügler des Kaiserlichen Museums zu Wien. VII. Vorarbeiten zu einer Monographie der Muscaria Schizometopa (exclusive Anthomyidae). Pars IV. Pt. 7, 88 pp. Wien.
- BRÊTHES, J. (see also MASSINI, P. C. & J. BRÊTHES)
1908. Sobre tres *Exorista* (Diptera), parásitas de la *Palustra tenuis* Berg. *An. Mus. Nac. Buenos Aires* 16:473-476.
1909. Dipteros e Himenópteros de Mendoza. *An. Mus. Nac. Buenos Aires* 12:85-105, 3 figs.
1910. Dos insectos nuevos chilenos. *Rev. Chil. Hist. Nat.* xiv:67-69, figs.
1917. Sur quelques Diptères de Lima (Perou). *An. Zool. Aplic.* 4:16-18.
- 1918a. Quelques diptères du Chile. *Rev. Chil. Hist. Nat.* 22:49-50.
- 1918b. Description d'une nouvelle "Dexiinae" argentine. *Physis* 4(16):115.
1920. Cueillette d'insectes au Rio Blanco. III. Diptère. *Rev. Chil. Hist. Nat.* 23:40-44, figs. 8-10 (1919).
- 1920a. Insectes du Pérou. *An. Soc. Cient. Argentina* 89:27-54.
- 1920b. Description d'un nouveau diptère chilien parasite de *Laora variabilis*. *An. Zool. Apl. Chile* 7:12-13, fig. 1.
1922. El bicho de cesto *Oeceticus kirbyi*, var. *platensis* Berg. Campaña 1920-1921. Dos nuevos parásitos. *Inst. Biol. Soc. Rural Argentina* 28 pp. illus.
1925. Coléoptères et Diptères chiliens. Diptère. *Rev. Chil. Hist. Nat.* 29:207-208.
1927. Parásitos y hiperparásito de "*Diatraea saccharalis*" en la caña de azúcar, en Tucumán. *Rev. Ind. y Agrícola de Tucumán* 17(7-8): 163-166.
- BROOKS, A. R.
1942. *Clistomorpha*, *Psaliadopteryx* and allies (Diptera, Tachinidae). *Canad. Ent.* 74:140-150, 4 figs.
- 1945a. New Canadian Diptera (Tachinidae). *Canad. Ent.* 77:78-96.
- 1945b. A revision of the North American species of the *Phasia* complex (Diptera, Tachinidae). *Sci. Agr.* 24:647-679, 4 pls.
1946. A revision of the North American species of the *Rhodogynce* complex (Diptera, Larvaevoridae). *Canad. Ent.* (1945) 77:218-230, 2 pls.



1949. New North American larvacvorine flies (Diptera, Larvaeovoridae). *Canad. Ent.* 81:21-24.

CALTAGIRONE, L.

1953. Observaciones sobre *Incamiya chilensis* Aldrich y su multiplication en laboratorio. *Rev. Chil. Ent.* 3:87-99, 3 figs.
1966. Una nueva especie chilena de *Opsophagus* Aldrich, 1926 (Diptera, Tachinidae). *Publ. Centro de Estudios Entomologicos, Fac. Filos. Education, Univ. Chile* n° 8:61-67, 2 figs.

CAMPOS, L.

1953. Notas sobre Taquinidos Chilenos. I. (Dipt., Tachinidae). *Agr. Tec. Chile* 13:24-31.

CLAUSEN, C. P.

1940. *Entomophagous insects*, 688 pp., 257 figs., New York.

COPPEL, H. C.

1960. Key to adults of dipterous parasites of spruce budworm, *Choristoneura fumiferana* (Clem.) (Lepidoptera: Tortricidae). *Ann. Ent. Soc. Amer.* 53:94-97, 2 pls.

COQUILLET, D. W.

1889. The corn worm or boll worm in California. *Insect Life* (U. S. Dept. Agr. Div. Ent. Bul.) 1:331-332.
1890. The dipterous parasites of *Diabrotica soror*. *Insect Life* (U. S. Dept. Agr., Div. Ent. Period. Bull.) 2:233-236, 1 fig.
1895a. New Tachinidae with a slender proboscis. *Canad. Ent.* 27:125-128.
1895b. Descriptions of new genera and species. Pp. 307-319. In Johnson C. W., Diptera of Florida. *Proc. Acad. Nat. Sci. Phila.* 303-340.
1895c. New genera and species of Tachinidae. *Journ. N. Y. Ent. Soc.* 3:97-107.
1895d. Notes and descriptions of Tachinidae. *Journ. N. York Ent. Soc.* 3:49-58.
1897. Revision of the Tachinidae of America north of Mexico. A family of parasitic two-winged insects. *Proc. U. S. Dept. Agr., Div. Ent. Tech. Ser.* 7:1-156.
1900a. Report on a collection of dipterous insects from Puerto Rico. *Proc. U. S. Natl. Mus.* 22:249-270.
1900b. Two new genera of Diptera. *Ent. News* 11:429-430, 1 fig.
1902a. New cyclorhaphous Diptera from Mexico and New Mexico. *Canad. Ent.* 34:195-202.
1902b. New Diptera from North America. *Proc. U. S. Natl. Mus.* 25:83-126.
1904a. New Diptera from Central America. *Proc. Ent. Soc. Wash.* 6:90-98.

104.225

- 1904b. New North American Diptera. *Proc. Ent. Soc. Wash.* 6: 166-192.
1905. A new dextiid parasite of a Cuban beetle. *Canad. Ent.* 37:362.
1910. The types-species of the North American genera of Diptera. *Proc. U. S. Natl. Mus.* 37:499-647.

CORTÉS, R.

- 1944a. Las especies chilenas del género *Archytas* Jaen. (Dipt. Tachinidae). *Bol. Dept. Sanid. Vegetal* 3(1943):139-141.
- 1944b. Taquinidos nuevos para Chile (Dipt. Tachinidae). *Bol. San. Veg. Chile* 3(1943):142.
- 1944c. Las especies chilcnas del género *Cylindromyia* Meigen (Dipt. Tachinidae). *Bol. San. Veg. Chile* 3(2):177-179 (1943).
- 1944d. *Thichodischia* Bigot y *Dischotrichia* n. gen. géneros nuevos de tachinidos para Chile (Dipt. Tachinidae). *Rev. Univ. Chile* 29: 49-58.
- 1944e. Las especies chilcnas del género *Sturmia* R. D. (Diptera, Tachinidae). *Bol. Mus. Nac. Hist. Nat.* 22:159-169.
- 1945a. Algunos géneros goniinos de taquinidos chilenos (Dipt., Tachinidae). *Agr. Téc. (Chile)* 4:115-124 (1944).
- 1945b. Nuevo nombre genérico para un taquinido de la Republica Argentina (Dipt., Tachinidae). *Acta Zool. Lill.* 2:255-257 (1944).
- 1945c. Taquinidos chilenos nuevos o poco conocidos (Dipt., Tachinidae). *Agr. Téc. (Chile)* 5:24-30.
- 1945d. Espécies chilenas de los géneros *Phorocera* R. D. y *Parasetigena* B. B. (Dipt., Tachinidae). *Act. Zool. Lill.* 3:157-164.
- 1945e. Nuevos géneros de taquinidos chilenos aliados a *Trichoprosopus* Macquart (Dipt. Tachinidae). *Rev. Chil. Hist. Nat.* (1944), 48:149-160.
1946. Tachinidae, in Stuardo, C., Catálogo de los Dipteros de Chile, pp. 172-185, Santiago.
1950. Nueva especie chilcna del género *Phorocera* R. D. (Dipt. Tachinidae). *Agr. Tec. (Chile)* 9:5-10 (1949).
- 1951a. Sobre tres especies de taquinidos chilenos (Dipt. Tachinidae). *Agr. Tec. (Chile)* 10:59-65 (1950).
- 1951b. Nuevos géneros de Tachinidae chilenos con cerdas facio-orbitalis (Dipt. Tachinidae). *Rev. Chil. Ent.* 1:249-262.
1952. Los insectos de las Islas Juan Fernández. Tachinidac (Diptera). *Rev. Chil. Ent.* 2:109-111.
1963. Tipos de Tachinidae (Diptera) chilenos en algunos museos extranjeros. *Rev. Univers. Chile, Santiago, Univ. Catolica* 47:241-252 (1952).
- 1967a. Primeros generos de taquinidos chilenos con la cuarta vena longitudinal evanescente. Diptera: Tachinidae. *Bol. Técn. Est. Exp. Agron. Univ. Chile* 26:3-9, 4 figs.
- 1967b. Taquinidos chilenos nuevos o poco conocidos (Diptera: Tachinidae). *Bol. Técn. Est. Exp. Agron. Univ. Chile* 26:10-29, 6 figs.



- 1968a. Taquinidos chilenos (Dipt. Tachinidae) parásitos de Phásmidos (Phasmoidea). *Bol. Soc. Biol. Concepción* 40:101-111 (1965-66).
- 1968b. Nueva especie chilena del genero *Incamyia* Townsend 1912 (Diptera: Tachinidae). *Rev. Chil. Entom.* 6:17-20.
- 1968c. El Taquinido más austral de Chile (Dipt., Tachinidae). *Rev. Chil. Ent.* 6:142.
1969. Rehabilitación de una especie bigotiana. *Trichodischia caerulea* Bigot 1885 diferente de *T. soror* Bigot, 1885 (Diptera: Tachinidae). *Rev. Universitaria* 58(31):97-99 (1968).
- CORTÉS, R. & N. HICHINS
1969. *Taquinidos de Chile. Distribución geográfica y huéspedes conocidos*. Ediciones de la Universidad de Chile, 99 pp., 7 pls., Santiago, Chile.
- CROSSKEY, R. W.
1965. The immature stages and affinities of the tachinid fly *Glaurocara flava*, a parasite of the African bush-cricket *Homocoryphus nitidulus vicinus*. *Proc. Zool. Soc. London* 144:203-217, 1 plate, 18 figs.
1971. The type material of Australasian, Oriental and Ethiopian Tachinidac (Diptera) described by Macquart and Bigot. *Bull. British Mus. (Nat. Hist.)*, Entomology, 25:251-305.
- CROUZEL, I. S. DE
1965. Parasitismo em "gusanos blanco" en la Republica Argentina (Col. Scarabeidac). *Rev. Soc. Ent. Argentina* 27:83-87.
- CURRAN, C. H.
- 1925a. Descriptions of four new Neotropical Diptera. *Trans. Amer. Ent. Soc.* 51:259-264.
- 1925b. New Tachinidae in the Canadian National Collection (Diptera). *Canad. Ent.* 57:150-156.
- 1925c. The American species of the tachinid genus *Pcletcra* Desv. (Diptera). *Proc. Trans. Roy. Soc. Canada, ser. 3*, vol. 19 (= Whole ser. 31) (5):225-257, 2 pls.
- 1925d. New American Diptera. II. *Ann. Mag. Nat. Hist. ser. 9*, vol. 16: 338-354.
- 1925e. New Diptera in the American Museum of Natural History. *Amer. Mus. Novitates* 200:1-10.
- 1926a. New Diptera from Jamaica. *Ent. Bull.* 4 (1). *Dept. Agr. Jamaica*. Appendix: 102-114.
- 1926b. The Canadian species of the tachinid genera *Cryptomeigenia* B. B. and *Tachinomyia* Tns. (Dipt.). *Proc. Trans. Roy. Soc. Canada, ser. 3*, 20 (= Whole ser. 32) (5):155-171, 2 pls.
- 1927a. A new tachinid parasitic on armyworms in Mexico. *Proc. Haw. Ent. Soc.* 6:497-498.
- 1927b. Some new North American Diptera. *Canad. Ent.* 59:290-303, figs. 1-10.
- 1927c. New West Indies Tachinidac. *Amer. Mus. Novitates* 260:1-15, 5 figs.

- 1928a. Records and descriptions of Diptera, mostly from Jamaica. *Ent. Bull. Dept. Agric. Jamaica* 4(3):29-45.
- 1928b. Revision of the American species of *Archytas* (Tachinidae, Diptera). *Canad. Ent.* 60:201-208, 218-226, 249-256, 275-282, 3 pls.
- 1928c. Insects of Porto Rico and Virgin Island. Diptera or two-winged flies. Pt. 1, pp. 1-118, 39 figs. In New York Academy of Sciences, Scientific Survey of Porto Rico and the Virgin Island, vol. 11, New York.
1929. New Syrphidae and Tachinidae. *Ann. Ent. Soc. Amer.* 22: 489-510.
- (1930). Report on Diptera collected at the Station for the Study of Insects, Harriman Interstate Park, N. Y. *Bull. Amer. Mus. Nat. Hist.* (1931) 61:21-115.
1931. First supplement to the "Diptera of Porto Rico and the Virgin Island". *Amer. Mus. Novitates* 456:1-23, 7 figs.
1932. New American Diptera. *Amer. Mus. Novitates* 534:1-15.
1933. The North American species of *Actia* in the American Museum of Natural History. *Amer. Mus. Novitates* 614:1-7.
- 1934a. The Templeton Crocker expedition of the California Academy of Sciences, 1932, n° 13 (Diptera). *Proc. Calif. Acad. Sci.* 21(13): 147-172, 4 figs.
- 1934b. Review of the tachinid genus *Calodexia* van der Wulp (Diptera). *Amer. Mus. Novitates* 685:1-21.
- 1934c. The Diptera of Kartabo, Bartica district, British Guiana with descriptions of new species from other British Guiana localities. *Bull. Amer. Mus. Nat. Hist.* 66:287-532, 54 figs.
- 1934d. *The families and genera of North American Diptera*. 512 pp. 235 figs. New York.
1935. New American Diptera. *Amer. Mus. Novitates* 812:1-24.
- 1937a. Threc new Neotropical Diptera. *Rev. Chilena Hist. Nat.* 40 (1936):331-335.
- 1937b. Three new Tachinidae (Diptera) parasitic on *Polybia* species (Hymenoptera). *Amer. Mus. Novitates* 923:1-4.
- 1939a. The species of *Macromyia* Desvoidy (Tachinidae, Diptera). *Amer. Mus. Novitates* 1020:1-3.
- 1939b. The dipterous genus *Chrysotachina* Brauer & Bergenstamm (Tachinidae). *Amer. Mus. Novitates* 1021:1-3.
1941. New Neotropical Tachinidae (Diptera). *Amer. Mus. Novitates* 1113:1-5.
1942. American Diptera. *Bull. Amer. Mus. Nat. Hist.* 80:51-84.
1947. New and little known American Tachinidae. *Bull. Amer. Mus. Nat. Hist.* 89:41-122, 134 figs.
1960. Review of the tachinid genus *Juriniopsis* Townsend (Diptera). *Amer. Mus. Novitates* 2014:1-7.

DENIER, P. C. L.

1939. Lista de los artrópodos dañinos o útiles a los algodones argentinos. *Physis* 17:553-567.

DRURY, D.

1770. *Illustrations of Natural History. Wherein are exhibited upwards of 240 figures of exotic insects, according to their different genera* vol. 1, 130 pp., 4 figs., 50 pls. London.
1773. *Illustrations of Natural History Wherein are exhibited upwards of 240 figures exotic insects, according to their different genera.* Vol. 2, 90 pp., 50 pls. and index to vols. 1-2 (4 unnumbered pages). London.

DUPUIS, C.

1963. Essai monographique sur les Phasiinae (Diptères Tachinaires parasites d'Hétéroptères). *Mem. Mus. Nat. d'Hist. Nat. n. scr.* (= ser. 3), ser. A: Zool. 26:1-461, 73 figs.

EMDEN, F. I. VAN

1949. The scientific name for the common tachinid parasite of *Diatraea* spp. (L. Pyral.) in Central and South America, with notes on related species (Dipt.). *Rev. de Ent.* 20:499-508, 6 figs.
1954. Tachinidae and Calliphoridae. Sect. a, 133 pp., 42 figs. In Royal Entomological Society of London, *Handbooks for the identifications of British Insects*. Vol. 10: *Cyclorrhapha*, Pt. 4. *Calyptrata* (1) London.

ENGEL, E. O.

1920. Studien über neotropische Histriciidac sensu B. et B. (Dipt.). *Zool. Jahrb. Abt. Syst. Bd.* 43:273-323, 27 figs.

FABRICIUS, J. G.

1775. *Systema entomologiae, sistens insectorum classes, ordines, genera, species adiectis synonymis, locis, descriptionibus, observationibus* 832 pp. Flensburgi et Lipsiae. [= Flensburg and Leipzig].
1781. *Species insectorum exhibentes eorum differentias specificas, synonyma, auctorum, loca natalia, metamorphosis*. Vol. 2, 517 pp. Hamburgi et Kiloni [= Hamburg and Kiel].
1805. *Systema antliatorum secundum ordines, genera, species*. 373 pp. + 30 pp. Brunsvigae.

FALLÉN, C. F.

1810. Försök att bestämma dei Sverige funne flugarter, som kunna föras till släktet *Tachina*. *K. Vetensk. Acad. Nya. Handl.* [ser. 2], 31:235-287.
1820. *Monographia Muscidum Sveciae*. Pp. 1-12, 13-24, 25-40 cont. Lundae.

FISCHER, C. R.

1923. Nota sobre *Actinochaeta carlos-albertoi* (Costa Lima, 1926) (Diptera, Dexiidae). *Rev. de Ent.* 3:194-198, fig. 1.

FONTENELLE, T. J. H. & J. H. GUIMARÃES

1966. Biological observations on some Masiphyini (Diptera, Tachinidae) parasites of Mantodea. *Pap. Avul. Dep. Zool.* 19:233-244, 6 figs.

GIGLIO-TOS, E.

1893. Diagnosi di nuovi generi e di nuove specie di Ditteri. VIII. IX. *Bol. Real. Univ. Torino. Mus. Zool. e Anat. Comp.* 8(147):1-11 (158):1-14.
1894. *Ditteri del Messico*. Pt. 3, 76 pp. 1 pl. Torino. Also published in *R. Accad. delle Sci. Torino, Mem.* 44 (Cl. di Sci. Fis., Mat. e Nat.): 473-546, 1 pl., 1894.

GISTEL, J.

1848. *Naturgeschichte des Thierreichs. Für höhere Schulen*. xvi + 216 pp. 32 pls. + 4 pp. of explanation, Stuttgart.

GREENE, C. T.

- 1921a. Two new species of Diptera. *Proc. Ent. Soc. Wash.* 23(6):125-127, fig. 1.
- 1921b. An illustrated synopsis of the puparia of 100 muscoid flies (Diptera). *Proc. U. S. Natl. Mus.* 60(10):1-39, 20 pls.
1927. The larva and puparium of *Oedematocera dampfi* Aldrich (Diptera). *Proc. Ent. Soc. Wash.* 29:18-19, figs. a-d.
1934. Tachinid flies with an evanescent fourth vein, including a new genus and five new species. *Proc. Ent. Soc. Wash.* 36:27-40, 4 pls.

GRIMSHAW, P. H.

1901. Part 1. Diptera. Pp. 1-17, pls. 1-3. In Sharp, D. ed., *Fauna Hawaiensis*, vol. 3, 704 pp., 19 pls. Cambridge, England.

GRIOT, M.

1944. Oruga que vacia las capsulas del lino. *Rev. Argentina Agro-nomia* 11(11):44-57, 6 figs., 2 pls.

GUÉRIN-MÉNEVILLE, F. E.

- (1831). pls. 20-21 (= livrs. 23-24, part). In Duperrey, L. I. ed., *Voyage autour du monde sur la corvette de sa majesté La Coquille Zoologie. Atlas, Insectes*, 21 pls. Paris.
- (1838). Première division. Crustacés, arachnides et insectes. Div. 1, pp. 1-216, 217-319 (= livre 25, 26). In Duperrey, L. I. ed., *Voyage autour du monde sur la corvette de sa majesté La Coquille. Zoologie*, vol. 2, pt. 2. Paris.

GUIMARÃES, J. H. (see also FONTENELLE & GUIMARÃES)

1960. Contribuição ao conhecimento do gênero *Archytas* Jacnicke 1867 (Diptera, Tachinidae). *Mem. Inst. Oswaldo Cruz* 58:115-124, 38 figs.

104.230

- 1961a. Segunda contribuição ao conhecimento do gênero *Archytas* Jaennicke, 1867 (Diptera, Tachinidae). *Mem. Inst. Oswaldo Cruz* 59:163-179, 45 figs.
- 1961b. Terceira contribuição ao conhecimento do gênero *Archytas* Jaennicke, 1867 (Diptera, Tachinidae). *Mem. Inst. Oswaldo Cruz* 59:355-396, 106 figs.
1962. Espécies brasileiras do gênero *Peleteria* Desvoidy, 1830 (Diptera, Tachinidae). *An. Academ. Bras. Cien.* 34:483-495, 55 figs.
- 1963a. Primeira contribuição ao conhecimento da tribo *Cuphoceratini* (Diptera, Tachinidae). *Mem. Inst. Oswaldo Cruz* 61:47-72, 121 figs.
- 1963b. Quarta contribuição ao conhecimento do gênero *Archytas* Jaennicke, 1867 (Diptera, Tachinidae). *Mem. Inst. Oswaldo Cruz* 61:153-164, 37 figs.
- 1963c. Primeira contribuição ao conhecimento da tribo *Juriniini* Townsend, gênero *Euempheremyia* Townsend, 1927 (Diptera, Tachinidae). *Mem. Inst. Oswaldo Cruz* 61:341-349, 29 figs.
- 1963d. Fifth contribution to the knowledge of the genus *Archytas* Jaennicke, 1867 (Diptera, Tachinidae). *Mem. Inst. Oswaldo Cruz* 61(2):329-340, 48 figs.
1964. Segunda contribuição ao conhecimento da tribo *Cuphoceratini* Townsend (Diptera, Tachinidae). *Papéis Avulsos do Dept. de Zool. Secr. Agr. S. Paulo* 16:173-189, 52 figs.
- 1966a. Taxonomic studies on some Masiphyini (Diptera, Tachinidae) reared from Mantodea. *Papéis Avulsos do Dept. Zool.* 19:205-232, 53 figs.
- 1966b. Revisão parcial do gênero *Adejeania* Townsend, 1913 (Diptera, Tachinidae) com especial referência às espécies brasileiras. *Arq. Zool. Secr. Agr. S. Paulo* 14:155-196, 73 figs.

HARRIS, T. W.

1835. VIII. Insects. Pp. 553-602. Hitchcock, E. *Report on the geology, mineralogy, botany and zoology of Massachusetts*, Ed. 2, 702 pp., 104 figs., + Atlas, 17 pls., Amherst, Mass.
1841. *A report on the insects of Massachusetts, injurious to vegetation*, 459 pp. Cambridge, Mass.

HENDEL, F.

1908. *Nouvelle Classification des mouches à deux ailes* (Diptera L.) d'après un plan tout nouveau par J. G. Meigen, Paris, an VIII (1800 v.s.). Mit. eincm Kommentar. *Verhandl. K. k. Zool. Bot. Gesell. Wien* 58:43-69.

HERTING, B.

1957. Das weibliche Postabdomen des calyptraten Fliegen (Diptera) und sein Merkmalwert für die Systematik der gruppe. *Z. Morph. Ökol. Tierc* 45:429-461, 21 figs.
1960. Biologie des westpaläartischen Raupenfliegen. Dipt., Tachinidae. *Mon. z. Angew. Ent.* 16:1-188, 12 figs.



INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE

1954. Opinion 205. Rejection of the generic name *Phoranthella* Townsend (Class Insecta, Order Diptera), as published in 1915, as nomen nudum. Pp. 309-318 (= pt. 241). In Hemming, F. ed., *Opinions and declarations rendered by the International Commission on Zoological Nomenclature*. Vol. 3, 448 pp., 2 pls., London.
1963. Opinion 678. The suppression under the Plenary Powers of the pamphlet published by Meigen, 1800. *Bull. Zool. Nomencl. Internt. Comm. Zool. Nomencl.* 20:339-342.

JACOBS, J. C.

1900. Diptères. Diagnoses d'insectes recueillis par l'expédition antarctique belge. *Ann. Soc. Ent. Belgique* 44:106-107.
1906. Diptère. Bibionidae, Chironomidae, Culicidae, Tipulidae, Syrphidae, Muscidae, Rhyphidae, Anthomyidae in *Expedition Antarctique Belge. Résultats du Voyage du S. Y. Belgica en 1897-1898-1899*. Zoologie: 67-71, pl. 3 (1906).

JAENNICKE, F.

1867. Neue exotische Dipteren. *Abhandl. Seckenb. Naturf. Gesell.* 6:311-408, pls. 43-44. Also printed separately under the title "*Neue exotische Dipteren aus den Museen zu Frankfurt a. M. und Darmstadt*" 100 pp., 2 pls. Frankfurt, 1868.

JAMES, M. T.

1945. A new larvaevorid parasite of the social butterfly *Eucheira socialis* Westwood. *Journ. Wash. Acad. Sci.* 35:328-330, fig. 1.
1947. A review of the larvaevorid flies of the tribe Leskiini with the scutose first vein (R₁). *Proc. U. S. Natl. Mus.* 97:91-115, 1 fig.

JOHNSON, C. W.

1919. A revised list of the Diptera of Jamaica. *Bull. Amer. Mus. Nat. Hist.* 41:421-449.

LANDIS, B. J. & H. F. HOWARD

1940. *Paraderodes epilachnae*, a tachinid parasite of the Mexican bean beetle. *U. S. Dept. Agr., Tech. Bull.* 721, 31 pp., 23 figs.

LATREILLE, P. A.

1809. *Genera crustaceorum et insectorum secundum ordinem naturalem in familias disposita, iconibus exemplisque plurimis explicata*. Vol. 4, 399 pp., 4 pls. Parisiis et Argentorat.
1825. *Families naturelles du règne animal*, 570 pp. Paris.
1829. Les Crustacés, les Arachnides et les Insectes. Tome second, vol. 2, vol. 5, 556 pp., 5 pls. In Cuvier, G. C. L. D., *Le règne animal...* Ed. 2. Paris.

LE BARON, W.

1872. Insects injurious to the apple tree. Noxious and Beneficial. *Insects of the State of Ill. (State Ent.) Rpt.* (3):99-133, 3 figs.



LIMA, A. M. DA COSTA

1926. Sobre o typo curioso de pernas anteriores, observados na mosca "*Thelairodes earlos albertoi*" n.sp. (Dipt. Dexiidae). *Sciencia Medica* 4(10):1-7, 2 figs.
1935. Um novo parasito endófago de *Mormidea poecila*. *O Campo* 6:21-22, 2 figs.
1947. Sôbre endoparasitos de *Thecla basilides* (Lep. Lycaenidae). *An. Acad. Bras. de Ciencias* 19(3):277-281, 1 pl.
1950. Mosea parasita das lagartas do eucalipto (Tachinidae). *Chacaras e Quintais* 82:167-169, 3 figs.

LINNAEUS, c.

1758. *Systema naturae per regna tria naturae*. Ed. 10, vol. 1, 824 pp. Holmiae = Stockholm.

LLOY, P.

1864. I ditteri distribuiti seeondo un nuovo metodo di classificazione naturale [cont.]. *Atti I. R. Ist. Veneto di Sci., Let. ed Arti*, ser. 3. 9:499-518, 569-604, 719-771, 879-910, 889-1027, 1087-1126, 1311-1352 cont.

LLOYD, D. c.

1951. A survey for grasshopper parasites in temperate South America. *Canad. Ent.* 83:213-230, 2 pls., 1 fig.

LOEW, H.

1863. *Diptera Americae septentrionalis indigena*. Centuria quarta. *Berlin. Ent. Ztschr.* 7:275-326.
1866. *Diptera Americae septentrionalis indigena*. Centuria septima. *Berlin Ent. Ztschr.* 10:1-54.
1869. *Diptera Americae septentrionalis indigena*. Centuria octava. *Berlin Ent. Ztschr.* 13:1-52.

MACQUART, J.

1835. *Histoire naturelle des Insectes Diptères*. Tome deuxième. *Diptera*, Vol. 2, 803 pp., 12 pls. In Roret, N. E. ed. (Collection des suites à Buffon). Paris.
1843. *Diptères exotiques nouveaux ou peu connus*. *Mém. Soc. Roy des Sci. de l'Agr. et Arts, Lilli*, 1842:162-460, 36 pls. Also printed separately as his "*Diptères exotiques nouveaux ou peu connus*". Vol. 2, pt. 3, pp. 5-304, 36 pls. Paris, 1843.
1846. *Diptères exotique nouveaux ou peu connus (I^{er} er) Supplément*. *Mem. Soc. Roy des Sci., de l'Agr. et des Arts, Lilli* (1845), 1844: 133-364, 20 pls. Also printed separately as his *Diptères exotiques nouveaux ou peu connus*. Supplément I, pp. 5-238, 20 pls. Paris, 1846.
1847. *Diptères exotiques nouveaux ou peu connus. 2^o supplément*. *Mém. Soc. Roy des Sci., de l'Agr., et des Arts, Lilli*, 1846:21-120, 6 pls. Also printed separately as his *Diptères exotiques nouveaux ou peu connus*. Supplément II, pp. 5-104, 6 pls., Paris, 1847.

1848. Diptères exotiques nouveaux ou peu connus. Suite de 2^e supplément i.c., 3 rd. supplément. *Mem. Soc. Roy. des Sci. de l'Agr. et des Arts, Lilli*, 1847 (2):161-237, 7 pls. Also printed separately as his *Diptères exotiques nouveaux ou peu connus*. Supplément III, pp. 1-77, 7 pls. Paris, 1848.
1851. Diptères exotiques nouveaux ou peu connus. Suite du 4^e supplément. *Mém. Soc. Natl. des Sci. de l'Agr. et des Arts, Lilli*, 1850: 134-282 (text), 283-294 (explanation of figs, index), pls. 15-28. Also published separately as his *Diptères exotiques nouveaux ou peu connus*. Supplément IV part., pp. 161-309 (text), 317-323 (expl. 27 figs.), 324-336 (combined index of the 2 parts of this supplément), pls. 15-28. Paris, 1851.
- MALLOCH, J. R.
1932. Some new species of the dipterous family Tachinidae. *Stylops* 1:197-203.
- [MARCHIONATTO, J. B.]
- (1938-1942). Labor de la direccion y sus dependencias. *Bol. Inform. Dir. Sanid. Veg., Min. Agr.* (Argentina), (mimeo), ano I (2):28-29, 1938; (4):44, 1938; 2 (5):37, 1938; 4 (13):21 1941; 4 (15):29, 1941; 5 (19):15, 1942.
- MARNEF, L.
1965. *Lafuentemyia yanezi* nov. gen. nov. sp. de Taquinido (Diptera) de Chile. *Bull. & Ann. Soc. Roy. d'Ent. de Belgique* 101:243-250, 2 figs.
- MASSINI, P. C. & J. BRÊTHES
1918. Metodo biologico contra las plagas, aplicado al *Oeceticus platanensis* (bicho del canastro). *An. Soc. Rural Argentina* 52:207-215, 11 figs., 1 pl.
- MEIGEN, J. W.
1800. *Nouvelle classification des mouches à deux ailes* (Diptera L.) d'après un plan tout nouveau. 40 pp. Paris.
1803. Versuch einer neuen Gattungseintheilung der europäischen zweiflügeligen Insekten. *Mag. f. Insektenkunde* 2:259-281.
1824. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten*. Vol. 4, xii + 428 pp., pls. 33-41. Hamm.
1838. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten*. Vol. 7, "oder Supplement band", xii + 434 pp., pls. 67-74. Hamm.
- MENDES, L. O. T.
1938. Lista dos inimigos naturais de *Dysdercus* spp. observados no Estado de São Paulo. *Rev. de Ent.* (Rio de Janeiro) 9(1-2):215-217.
- MESNIL, L. P.
- 1949a. Essai de révision des espèces du genre *Drino* Robineau-Desvoidy. Sturmiinae à oeufs macro-types. *Bull. Inst. Roy. des Sci. Nat. de Belg.* 25(42):1-38.



- 1949b. Larvaevorinac (Tachininae) Fam. 64 g., pp. 49-104, figs. 29-30 (= lfg. 161) cont. In Lindner, E., ed., *Die Fliegen der palaearktischen Region*. Vol. 8. Stuttgart.
1960. Larvaevorinae (Tachininae) cont. (Fam.) 64 g., pp. 561-608 (lfg. 210), 1960; pp. 609-656 (= fig. 212). 1960. In Lindner, E., ed., *Die Fliegen der palaearktischen Region*. Vol. 8. Stuttgart.
1966. Larvaevorinae (Tachininae) cont., 64 g. pp. 881-928 (= lfg. 263), 1966. In Lindner, E. ed., *Die Fliegen der palaearktischen Region*. Stuttgart.
- MIK, J.
1864. Dipterologische Beiträge. *Verhandl. K. k. Zool. Bot. Gesell. Wien*, 14 (Abhandl.): 785-798.
1894. Ueber *Echinomyia popelii* Portsch. *Wien Ent. Ztg.* 13:100.
- MYERS, J. G.
1934. The discovery and introduction of the Amazon fly, a new parasite for cane-borers (*Diatraea* spp.). *Trop. Agr.* 11(8):191-195.
- MOREIRA, C.
1951. Description d'une tachinaire nouvelle. *Bull. Soc. Ent. France* 14:227-229, e figs.
- NUTTING, W. L.
1953. The biology of *Euphasiopteryx brevicornis* (Townsend). (Diptera, Tachinidae) parasitic in the cone-headed grasshoppers. (Orthoptera, Copiphorinae). *Psyche* 60:69-81, 1 fig., pl. 4.
- OLIVIER, G. A.
- 1811-[1812]. Insectes [i.e., Arthropoda]. Pt. 5], vol. 8, pp. 1-360 (= livr.?), 1811; pp. 361-722 (= livr. 77), [1812]. In Société de Gens de Lettres, de Savans et d'Artistes. *Encyclopédie méthodique Histoire naturelle*. Paris.
- OSTEN-SACKEN, C. R.
1877. Western Diptera. Descriptions of new genera and species of Diptera from California. [U. S. Dept. Int.]. *Bull. U. S. Geol. and Geog. Survey of the Ter.*, 3:189-354.
1887. Some North American Tachinac. *Canad. Ent.* 19:161-166.
- PANTEL, J.
1910. Recherches sur les Diptères à larves entomobies. I. Caractères parasitiques aux points de vue biologique, éthologique et histologique. *Cellule* 26:25-216, 26 figs., 5 pls.
- PARROTT, P. J. & W. J. SCHÖENF
1912. The apple and cherry ermine moths. N. Y. (Geneva). *Agr. Exp. Sta. Tech. Bull.* 24:1-40, 10 figs., 9 pls.



PARKER, H. L.

1953. Miscellaneous notes on South American dipterous parasites. *Boll. Lab. Ent. Agr. "Fillippo Silvestri". Portici* 12:45-73.

PARKER, H., P. A. BERRY & A. SILVEIRA

1950. Vegetable weevils and their natural enemies in Argentina and Uruguay. *U. S. Dept. Agr. Tech. Bull.* 1016:1-28, 8 figs.

PAZOS, J. H.

1914. Catalogo de los Dipteros de la Isla de Cuba. *Anal. Acad. Ciencias Habana* 1:990-1003.

PERTY, M.

- 1830-1833. *Insecta brasiliensia*. Pp. 1-60, pls. 1-12, 1830; pp. 61-124, pls. 13-24, 1832; 125-224; pls. 25-40, 1833. *In his Delectus animalium articulorum quae in itinere per Brasiliam annis MDCCCVII-MDCCCXX jussu et auspiciis Maximiliani Josephi I. Bavariae regis augustissimi peracto collegerunt Dr. J. B. de Spix et Dr. C. F. Ph. de Martius*. 44 + 224 pp., 40 pls. Monachii, 1830-1834".

RAPP, W. E., JR.

1945. *Pseudolutzomyia*, new name for *Lutzomyia* Curran, 1934 (Diptera). *Proc. Ent. Soc. Washington* 47:278.

REED, C. S.

1907. Entomología económica. Biología de la *Laora variabilis* F. Ph., y noticias acerca de un díptero cuya larva vive como parásito en su oruga. *Bol. Soc. Agr. Sur., Chile* 7, 5:1008-1014; 7, 6:1045-1051 (1907).

REINHARD, H. J.

1923. New Tachinidac from Texas (Diptera). *Ent. New* 34:266-269.
1924. A new species of *Gonia* from Texas (Diptera). *Ent. New* 35:357-358.
1930a. On the genus *Viviania* with the description of two new species from Texas (Tachinidae, Diptera). *Bull. Brooklyn Ent. Soc.* 25:102-107.
1930b. Two new North American species of muscoid flies (Tachinidae, Diptera). *Bull. Brooklyn Ent. Soc.* 25:199-202.
1931a. Revision of the American parasitic flies belonging to the genus *Winthemia*. *Proc. U. S. Natl. Mus.* 79(20):1-45, pl. 1.
1931b. The two-winged flies belonging to *Siphosturmia* and allied genera, with descriptions of two new species. *Proc. U. S. Natl. Mus.* 79(11):1-11.
1934a. American muscoid flies of the genera *Ceratomyiella* and *Paradidyma*. *Proc. U. S. Natl. Mus.* 83:9-43.
1934b. Two new species of the tachinid genus *Siphosturmiopsis* with key and notes (Diptera). *Ent. News* 45:15-19.



- 1934c. Revision of the American two-winged flies belonging to the genus *Cuphocera*. *Proc. U. S. Natl. Mus.* 83:45-70.
- 1935a. New genera and species of American Muscoid flies (Tachinidae: Diptera). *Ann. Ent. Soc. Amer.* 28:160-173.
- 1935b. North American two-winged flies of the genus *Doryphorophaga* (Tachinidae, Diptera). *Journ. N. Y. Ent. Soc.* 43:387-394.
- 1942a. New North American Tachinidae belonging to the genera *Microchaetina* and *Hypenomyia* with key to the known species (Diptera). *Canad. Ent.* 74:88-91.
- 1942b. A new species of *Oedematocera* with notes and key (Tachinidae: Diptera). *Ent. New* 53:106-108.
- 1943a. New North American Muscoidea (Tachinidae, Diptera). *Jour. Kansas Ent. Soc.* 16:14-23.
- 1943b. New Tachinidae from northeastern United States. (Diptera). *Bull. Brooklyn Ent. Soc.* 38:78-90.
- 1943c. New genera of North America muscoid Diptera. *Canad. Ent.* 75:163-169.
1944. New muscoid Diptera from the United States. *Journ. Kans. Ent. Soc.* 17:57-72.
1945. New genera and species of North American Tachinidae (Diptera). *Canad. Ent.* 77:28-36.
1947. New genera and species of muscoid Diptera. *Jour. Kans. Ent. Soc.* 20:15-24.
1951. New American muscoid Diptera. *Bull. Brooklyn Ent. Soc.* 46:1-9.
- 1952a. New genera and species of muscoid Diptera. *Bull. Brooklyn Ent. Soc.* 47:1-2.
- 1952b. Muscoid flies of the genus *Chaetophlesia* (Diptera). *Jour. Kans. Ent. Soc.* 25:13-21.
- 1953a. New Mexican Tachinidae (Diptera). *Jour. Kans. Ent. Soc.* 26:95-102.
- 1953b. New species of Tachinidae from Mexico (Diptera). *Bull. Brooklyn Ent. Soc.* 48:89-96.
1954. Parasitic flies of the genus *Proscnoides* (Tachinidae, Diptera). *Canad. Ent.* 86:408-413.
- 1955a. North American muscoidca (Diptera: Tachinidae). *Ent. News* 66:233-238.
- 1955b. North American tachinid flies of the genus *Wagneria* (Diptera). *Jour. Kans. Ent. Soc.* 28:49-59.
- 1955c. New genera and species of North American Tachinidae (Diptera). *Jour. Kans. Ent. Soc.* 28:123-130.
- 1956a. New Tachinidae (Diptera). *Ent. New* 67:121-129.
- 1956b. A synopsis of the tachinid genus *Leucostoma* (Diptera). *Jour. Kansas Ent. Soc.* 29:155-168.
- 1956c. New muscoid Diptera mainly from California. *Pan-Pacific Ent.* 32:103-110.
1957. New American muscoid Diptera (Sarcophagidae, Tachinidae). *Ent. News* 68:99-111.

- 1958a. New American Tachinidae (Diptera). *Ent. News* 69:233-242.
- 1958b. New genera and species of North American Tachinidae (Diptera). *Jour. Kans. Ent. Soc.* 31:225-232.
- 1958c. Parasitic flies of the genus *Mochlosoma* (Tachinidae, Diptera). *Canad. Ent.* 90:98-110.
- 1958d. Notes on *Spathimeigenia* with descriptions of four new species (Diptera, Tachinidae). *Proc. Ent. Soc. Wash.* 60:207-212.
- 1958e. North American Tachinidae (Diptera). *Jour. Kans. Ent. Soc.* 31:277-284.
- 1959a. New Nearctic Tachinidae (Diptera). *Pan-Pacific Ent.* 35:157-163.
- 1959b. New North American Tachinidae (Diptera). *Ent. News* 70:225-234.
1961. New American Tachinidae and Sarcophagidae (Diptera). *Jour. Kans. Ent. Soc.* 34:204-213.
- 1962a. New North American Tachinidae (Diptera). *Pan-Pacific Ent.* 38:215-224.
- 1962b. North American Muscoid Diptera. *Ent. News* 73(7):169-178.
- 1964a. North American Tachinidae (Diptera). *Jour. Kans. Ent. Soc.* 37(1):35-51.
- 1964b. Parasitic flies of the genera *Helioprosopa* and *Euhelioprosopa* (Diptera, Tachinidae). *Pan-Pacific Ent.* 40:117-124.
- 1964c. Mexican Tachinidae (Diptera). *Act. Zool. Mexicana* 7:1-21.
- 1967a. New Nearctic and Neotropical Muscoidean Diptera. (Sarcophagidae and Tachinidae). *Jour. Kans. Ent. Soc.* 40:94-110.
- 1967b. Change of names in Tachinidae (Diptera). *Jour. Kans. Ent. Soc.* 40(4):600.
1968. New World species of *Erynnia* (Tachinidae, Diptera). *Jour. Kans. Ent. Soc.* 41:501-508.

RETTEMEYER, C. W.

1961. Observation on the biology and taxonomy of flies found over Swarm raids of army ants. (Diptera: Tachinidae, Conopidae). *Univ. Kans. Sci. Bull* 42:993-1066, 5 figs.

RILEY, C. V.

1870. Second annual report on the noxious beneficial and other insects of the state of Missouri. *Mo., State Bd. Agr. Ann. Rept.* (1869) 5 (spec. pagination): 1-135, 99 figs.
1871. Third annual report on the noxious, beneficial and other insects of the State of Missouri. *Mo. State Bd. Agr. Ann. Rept.* (1870) 6 (spec. pagination): 1-175, 73 figs.

ROBERTSON, C.

1901. Some new Diptera. *Canad. Ent.* 33:284-286.

ROBINEAU-DESVOIDY, J. B.

1830. *Essai sur les Myodaires*. Paris. *Inst. de France Cl. des Sci. Math. et Phys. Acad. Roy. des Sci., Mém. présentés par divers Savants ser. 2*, 2:1-813.



1863. *Histoire naturelle des Diptères des environs de Paris*. Vol. 1. xvi + 1143 pp.; vol. 2, 920 pp. Paris.

RÖDER, V. VON

1885. Dipteren von Insel Portorico, erhalten durch Herrn Consul Krug in Berlin. *Stett. Ent. Ztg.* 46:337-349.
1886. Dipteren von den Cordilleren in Columbien Gesammelt durch Herrn Alphons Stübel. *Stett. Ent. Ztg.* 47:257-270. (Also printed separately with addition of a plate as "*Dipteren gesammelt in den Jahren 1868-1877 auf einer Reise durch Süd Amerika*" von Alphons Stübel, 16 pp., 1 col., plate. Berlin, 1892).

RONDANI, C.


1848. Esame di varie specie d'insetti ditteri brasiliani. *Truqui, Studi. Ent.* 1:63-112, 1 pl. (Turin).
- 1850a. Dipteriorum species aliquae in America Aequatoriali collectae a Cajetano Osculati, observatae et distinctae novis breviter descriptis. *Nuov. Ann. Sc. Nat.* (Bologna), ser. 3, 2:357-372.
- 1850b. Osservazioni sopra alquante specie di esapodi ditteri del Museo Torinese. *Nuovi Ann. delle Sci. Nat.* ser. 3, 2:165-197, pl. 4.
1856. Dipterologiae Italiae prodromus. Vol. 1: Genera Italica ordinis dipterorum ordinatim disposita et distincta et in familias et stirpes aggregata, 228 pp. Parniae [= Parmal].
1863. *Diptera exotica revisa et annotata*, 99 pp., 1 pl. Modena. (Also published under the title "Dipteriorum species et genera aliqua exotica", in *Arch. per Zool. l'Anat. e Fis. Modena* (1863) 3(1):1-99, pl. 5, 1864.
1865. Diptera Italica non vel minus cognita descripta vel annotata observationibus nonnullis additis. Fasc. I. Oestridae-Syrphidae, Conopidae; Fasc. II. Muscidae. *Atti Soc. Ital. di Sci. Nat.* 8:193-231.
1868. Diptera aliqua in America meridionali lecta a Prof. P. Strobel annis 1866-67 distincta et annotata, novis aliquibus descriptis. *Ann. Soc. Nat. Modena* 3:24-40, 1 pl.

SABROSKY, C. W. (see also SABROSKY & ARNAUD)

1947. A synopsis of the larvaevorid flies of the genus *Eudejeania*. *Proc. U. S. Natl. Mus.* 97:141-156.
1950. Notes on Trichopodini (Diptera, Larvaevoridae), with description of a new parasite of cotton stainers in Puerto Rico. *Jour. Wash. Acad. Sci.* 40:361-371, 11 figs. (Erratum in *Proc. Ent. Soc. Washington* 53, 1951, pag. 210).
1952. A new larvaevorid fly parasitic on tortoise beetles in South America (Diptera). *Jour. Wash. Acad. Sci.* 42:325-327.
1953. Taxonomy and host relations of the tribe Ormiini in the Western Hemisphere. I, II: *Proc. Ent. Soc. Wash.* 55:167-183; 289-305.



1955. The taxonomic status of the armyworm parasite known as *Archytas piliventris* (van der Wulp) (Diptera: Larvaevoridae). *Fla. Ent.* 38:77-83, 6 figs.
1963. The identity of *Lutzomyia* Curran (Diptera). *Bull. Brooklyn Ent. Soc.* 58:14-17.
1965. Host of the tachinid tribe Eutherini (Diptera). *Proc. Ent. Soc. Wash.* 67:61.
1969. A review of the genus *Juriniopsis* Townsend (Diptera: Tachinidae). *Florida Ent.* 52:80-90, 10 figs.
- SABROSKY, C. W. & P. H. ARNAUD, JR.
1965. Tachinidae. In Stone, A. et al. A catalog of the Diptera of America north of Mexico, U.S.D.A., *Agriculture Handbook* 276. 1696 pp.
- SAY, T.
1829. Descriptions of North American dipterous insects. *Jour. Acad. Nat. Sci. Phila.* 6:149-178.
- SCHINER, I. R.
1868. Diptera. Art. 1, 388 pp., 4 pls. In Wüllerstorff-Urbair, B. von. In charge. *Reise der österreichischen Fregatte Novara*. Zool. Vol. 2. Abt. 1. Sect. B. Wien.
- SÉGUY, E.
1925. Étude sur quelques Calliphorinés testacés rares ou peu connus. *Bull. Mus. Nat. d'Hist. Nat.* 31:439-441.
- 1926a. Sur deux Calliphorines nouveaux. *Bull. Soc. Ent. France* 5: 62-64.
- 1926b. Calliphorines nouveaux. *Encycl. Ent. ser. B.* II, Dipt. 3:17-20. Paris.
- 1926c. Sur une forme nouvelle se rapportant aux "Oestridae dubiosc". *Encycl. Ent. ser. B.* II, Dipt. 3:1-10, 11 figs. Paris.
- 1927a. Étude sur les Calliphorines inférieurs. *C. R. Congrès Soc. Savants*, Paris, 421-427.
- 1927b. Un nouveau Calliphorine de la Guyane française. (Dipt.). *Ann. Soc. Ent. France* 96:262.
- SHIMER, H.
1871. Additional notes on the striped squash beetle. *Amer. Nat.* 5: 217-220, 2 figs.
- SMITH, H. E.
1915. New species of Tachinidae (Diptera) from New England. *Psyche* 22:98-102.
1916. New Tachinidae from North America. *Proc. Ent. Soc. Wash.* 18:94-98.
- SWEZEY, O.
1926. (Notes and exhibitions, March 5, 1925). *Proc. Hawaiian Ent. Soc.* 6:224 (footnote).



TAVARES, O.

1962. Contribuição ao conhecimento da tribo Ormiini I. Gênero *Ormia* Robineau-Desvoidy, 1830 (Diptera, Tachinidae). *Mem. Inst. Oswaldo Cruz* 60:347-363, 40 figs.
1964. Contribuição ao conhecimento da tribo Ormiini. II. Gênero *Ormiophasia* Townsend, 1919 (Diptera, Tachinidae). *Mem. Inst. Oswaldo Cruz* 62:37-52, 60 figs.
- 1965a. Contribuição ao conhecimento da tribo Ormiini. III. Gênero *Euphasiopteryx* Townsend, 1915 (Diptera, Tachinidae). *Mem. Inst. Oswaldo Cruz* 63:13-25, 40 figs.
- 1965b. Contribuição ao conhecimento da tribo Ormiini IV. Gênero *Ormia* Robineau-Desvoidy, 1830 (Diptera, Tachinidae). *Rev. Brasil. Biol.* 25:211-216, 12 figs.

THOMPSON, W. R.

1954. *Hyalomoydes triangulifera* Loew (Diptera, Tachinidae). *Canad. Ent.* 86:137-144.
1960. The larval morphology of some tachinid parasites of *Diatraea* (Diptera). *Trans. Amer. Entom. Soc.* 86:207-224, 42 figs.
1961. The Tachinids (Diptera) of Trinidad. I. The Voriines. *Trans. Amer. Ent. Soc.* 87:21-44, 25 figs.
- 1963a. The Tachinids of Trinidad. II. Echynomyiines, Dexiines and allies. *Canad. Jour. Zool.* 41:335-576, 174 figs.
- 1963b. The tachinids of Trinidad. III. The Goniines with microtype eggs (Diptera, Tachinidae). *Stud. Entom.* 6:257-404, 68 figs.
- 1963c. The Tachinids of Trinidad. IV. Wintheminiines. *Canad. Ent.* 95: 593-995, 38 figs.
- 1963d. The Tachinids of Trinidad. V. Siphosturmiines and Masyphyiines. *Canad. Ent.* 95:1292-1320, 29 figs.
1964. The Tachinids of Trinidad. VI. The larviparous Goniines of the Carcelline type (Diptera, Tachinidae). *Studia Ent.* 7:97-151, 46 figs.
1966. The Tachinids of Trinidad. VII. The larviparous Goniines with broad cheeks (Diptera, Tachinidae). *Studia Ent.* 8(1965):353-434, 46 figs.
1968. The Tachinids of Trinidad. VIII. Phorocerines. *Memoirs Ent. Soc. Canada* 56:1-207, 140 figs.

THOMSON, C. G.

1869. 6. Diptera. Species novas descripsit. Pp. 443-614, pl. 9 (= h. 12, n° 2). In *K. Svenska Vetenskaps-Akademien, Kongliga svenska fregatten Eugenie resa omkring jorden. Pt. 2. Zoologie (Sec. 7) 1: Insekter*, 617 pp., 9 pls. Stockholm, "1868".

TOTHILL, J. D.

1924. A revision of nearctic species in the genus *Fabriciella* (Tachinidae). *Canad. Ent.* 56:257-269.

104.241

TOWNSEND, C. H. T.

- 1891a. Notes on North American Tachinidae sens. lat., with descriptions of new species. Paper. I. *Proc. Ent. Soc. Wash.* 2:134-146.
- 1891b. Notes on North American Tachinidae sens. str. with descriptions of new genera and species. Paper II. *Trans. Amer. Ent. Soc.* 18:394-382.
- 1891c. An *Exorista* parasitic on *Lagoa opercularis*. *Ent. News* 2(8): 159-160.
- 1892a. Notes on North American Tachinidae, with descriptions of new genera and species. Paper III. *Trans. Amer. Ent. Soc.* 19: 88-132.
- 1892b. New Jamaica Tachinidae I. *Ent. News* 3:146-147.
- 1892c. Notes on North American Tachinidae, with descriptions of new genera and species. Paper V. Paper VI. *Canad. Ent.* 24:64-70; 77-82; 165-172.
- 1892d. New North American Tachinidae. *Ent. News* 3:80-81; 129-131.
- 1892e. Notes on North American Tachinidae, with descriptions of new species. Paper VII. *Trans. Amer. Ent. Soc.* 19:284-289.
- 1892f. The North American genera of Calyptrate Muscidae. Paper II. *Trans. Amer. Ent. Soc.* 19:133-144.
- 1892g. A Tachinid bred from larva of *Protoparce jamaicensis* (Butl.) in Jamaica. *Jour. Inst. Jamaica* 1:70-71.
1893. Comments on Mr. van der Wulp's recent diagnoses of new species of Mexican Phasiidae, Gymnosomatidae, Ocypteridae, and Phaniidae. *Canad. Ent.* 25:165-168.
1894. Some new Santo Domingo Tachinidae. *Jour. New York Ent. Soc.* 2:78-79.
1895. Contributions to the dipterology of North America. II. Tabanidae, Conopidae, Tachinidae, etc. *Trans. Amer. Ent. Soc.* 22: 55-80.
- 1897a. Contributions from the New Mexico Biological Stations n° 2. On a collection of Diptera from the lowlands of the Rio Nautla in the State of Veracruz. I. *Ann. Mag. Nat. Hist.* ser. 6, 19:16-34.
- 1897b. Contributions from the New Mexico Biological Station. N° IV. Diptera from the Sacramento and the White Mountains, in southern New Mexico. I. *Ann. Mag. Nat. Hist.* ser. 6, 19:138-149.
- 1897c. Contributions from the New Mexico Biological Station. N° 2 (cont.). On a collection of Diptera from the lowlands of the Rio Nautla, in the State of Veracruz. *Ann. Mag. Nat. Hist.* 6(20): 19-33; 272-291.
1908. The taxonomy of the muscoidean flies, including description of new genera and species. *Smith. Misc. Collect.* 51 (= pub. 1803): 1-138.
- 1911a. Review of work by Pantel and Portchinski on reproductive and early stage characters of muscoid flies. *Proc. Ent. Soc. Wash.* 13:151-170.
- 1911b. Annoucement of further secured results in the study of muscoid flies. *Ann. Ent. Soc. Amer.* 4:127-152.




- 1912a. Descriptions of new genera and species of muscoid flies from the Andean and Pacific Coast regions of South America. *Proc. U. S. Natl. Mus.* 43(1935):301-367.
- 1912b. Foundation of some new genera and species of muscoid flies mainly on reproductive and early-stage characters. *Jour. N. Y. Ent. Soc.* 20:107-119.
- 1912c. A readjustment of muscoid names. *Proc. Ent. Soc. Wash.* 14: 45-53.
- 1912d. Six new genera of nearctic Muscoidea. *Proc. Ent. Soc. Wash.* 14:163-166.
- 1913a. Inquiry into the relationships and taxonomy of the muscoid flies. *Canad. Ent.* 45:37-57.
- 1913b. Muscoid parasites of the cotton-stainer and other lygaeids. *Psyche* 20:91-94.
- 1913c. New muscoid flies, mainly Hystriciidae and Pyrrhosinae from the Andean Montanya. *Insec. Inscit. Menstr.* 1(11):144-148.
- 1913d. On *Trichiopoda* Latreille, *Polistomyia* Townsend and *Trichopodopsis* new genera. *Jour. N. Y. Ent. Soc.* 21:147-148 (313, a correction).
- 1913e. On the tribe Dejeaniini of the muscoid family Hystriciidae with five new genera. *Psyche* 20:102-106.
- 1913f. Two new generic names in muscoidea (Dipt.). *Ent. News* 24 (3):133.
1914. New muscoid flies mainly Hystriciidae and Pyrrhosinae from the Andean Montanya (cont.). *Insec. Inscit. Menstr.* 2:10-16; 29-32; 42-48; 81-96; 123-128; 133-144; 153-160; 169-176 (see also 1913c and 1915h).
- 1915a. Correction to paper on Andean muscoidea (Dipt.). *Ent. News* 26(1):28 (mailed December 31, 1914).
- 1915b. Proposal of new muscoid genera for old species. *Proc. Biol. Soc. Wash.* 28:19-24.
- 1915c. Revision of *Myophasia*. *Proc. Ent. Soc. Wash.* 17:107-114.
- 1915d. New Masiceratidae and Dexiidae from South America. *Jour. N. Y. Ent. Soc.* 23:61-68.
- 1915e. New Andean Spallanzanline Flies. *Insec. Inscit. Menstr.* 3(5-7): 63-69.
- 1915f. A new genus of Hystriciine flies with white maggots. *Insec. Inscit. Menstr.* 3(1-4):45-46.
- 1915g. New western and southwestern Muscoidea. *Jour. N. Y. Ent. Soc.* 23:216-234.
- 1915h. New muscoid flies, mainly Hystriciidae and Pyrrhosinae from the Andean Montanya (concl.). *Insec. Inscit. Menstr.* (1914)2:183-7.
- 1915i. An acalyptrate genus of muscoidea (Diptera). *Insec. Inscit. Menstr.* 3(1-4):41.
- 1915j. A polistiform genus of muscoid flies (Diptera). *Insec. Inscit. Menstr.* 3(1-4):43-44.
- 1915k. Nine new tropical American genera of Muscoidea. *Insec. Inscit. Menstr.* 3:91-97.



- 1915l. New muscoid flies mainly Hystricilidae and Pyrrhosiniac from the Andean Montanya. *Insec. Inscit. Menstr.* 2:183-187 (1914).
- 1915m. New Neotropical muscoid flies. *Proc. U. S. Natl. Mus.* 49 (2115):405-440.
- 1915n. New Peruvian hystricine flies. *Insec. Inscit. Menstr.* 3(5-7): 69-76.
- 1915o. *Phyllophilopsis*, new name. *Can. Ent.* 48:19.
- 1916a. *Andrina radialis* Townsend, new name. *Canad. Ent.* 27:178.
- 1916b. New Muscoidea genera (Dipt.). *Ent. News* 27:178.
- 1916c. Designations of muscoid genotypes, with new genera and species. *Insec. Inscit. Menstr.* 4:4-12.
- 1916d. Description of two new Tachinids (Dipt.). *Ent. News* 27:217.
- 1916e. Some new North America muscoid forms. *Insec. Inscit. Menstr.* 4:73-78
- 1916f. Diagnoses of new genera of muscoid flies founded on old species. *Proc. U. S. Natl. Mus.* 49(2128):617-633.
- 1916g. New genera and species of muscoid flies. *Proc. U. S. Natl. Mus.* 51(2152):299-323.
- 1916h. New and noteworthy Brazilian Muscoidea collected by Herbert H. Smith. *Bull. Amer. Mus. Nat. Hist.* 35:15-22.
- 1917a. New genera and species of American Muscoid Diptera. *Proc. Biol. Soc. Wash.* 30:43-50.
- 1917b. Second paper on Brazilian Muscoidea collected by Herbert H. Smith. *Bull. Amer. Mus. Nat. Hist.* 37(6):221-233.
- 1917c. Miscellaneous muscoid notes and descriptions. *Insec. Inscit. Menstr.* (1916) 4:121-128.
1918. Some muscoid synonymy, with one new genus. *Proc. Ent. Soc. Wash.* 20:19-21.
- 1919a. New genera and species of muscoid flies. *Proc. U. S. Natl. Mus.* 56(2301):541-592.
- 1919b. New muscoid genera, species and synonymy (Diptera) (concl.). *Insec. Inscit. Menstr.* (1918) 6:157-182.
1926. New holarctic Muscoidea (Diptera). *Insec. Inscit. Menstr.* 14: 24-41.
- 1927a. Synopse dos gêneros muscoideos da região humida tropical da America, com gêneros e espécies novas. *Rev. Mus. Paulista* 15: 203-385, 7 figs.
- 1927b. Errata a "Synopses dos gêneros muscoideos da região humida tropical da America, com gêneros e espécies novas". Four unnumbered pp. following p. 386. *Rev. Mus. Paulista* 15.
- 1928a. *Schistocercophaga*, new genus of locust parasites (Larvaevoridae) (Diptera, Tachinidae). *Ent. News* 39:152.
- 1928b. New Muscoidea from humid tropical South America. *Wiener Ent. Zeit.* 44:143-154.
- 1928c. New muscoid genera and species of the coast of Peru and Chile. *Rev. Chilena Hist. Nat.* 31:158-164 (1927).



1929. New species of humid tropical American Muscoidea (Sic). *Rev. Chilena Hist. Nat.* 32(1928):365-382.
- 1931a. New genera and species of American oestromuscoid flies. *Rev. Ent.* 1:313-354; 437-479.
- 1931b. Notes on American oestromuscoid types. *Rev. de Ent.* 1:65-104; 157-182.
- 1932a. A remarkable new genus and species of two-winged flies related to the Oestridae. *Proc. U. S. Natl. Mus.* 82(2942):1-4, figs. 1-2.
- 1932b. Five new Brazilian oestromuscoid genera. *Rev. de Ent.* 2: 105-107.
1933. Two new generic names in Oestroidea. *Rev. de Ent.* 3:527.
- 1934a. New neotropical oestromuscoid flies. *Rev. de Ent.* 4:201-212; 390-406.
- 1934b. *Manual of Myiology in twelve parts.* Pt. 1. Development structure, 280 pp., São Paulo.
- 1935a. Multifissicorn male oestroids. *Rev. de Ent.* 5:292-293, figs. 1, 2.
- 1935b. New muscoid genera, mainly from the Neotropical Region. *Rev. de Ent.* 5:68-74.
- 1935c. *Manual of Myiology, in twelve parts.* Part 2: Muscoid classification and habits, 296 pp., 9 pls. São Paulo.
- 1935d. New South America oestroidea (Dipt.). *Rev. de Ent.* 5(2): 216-233.
- 1936a. *Manual of Myiology, in twelve parts.* Pt. 3: Oestroid classification and habits (Gymnosomatidae to Tachinidae), 250 pp. São Paulo.
- 1936b. *Manual of Myiology, in twelve parts.* Pt. 4: Oestroid classification and habits (Dexiidae and Exoristidae), 303 pp., São Paulo.
- 1936c. A new phasiine parasite of *Dysdercus*. *Rev. de Ent.* 6:489.
- 1937a. New fly parasites of *Dysdercus*. *Rev. de Ent.* 7:316-318.
- 1937b. Two new South American oestromuscarian genera (Dipt.). *Rev. de Ent.* 7:115-117.
- 1938a. Further fly parasites of Heteroptera. *Rev. de Ent.* 8:204.
- 1938b. *Manual of Myiology, in twelve parts.* Pt. 7. Oestroid generic diagnoses and data (Gymnosomatini to Senostomatini), 434 pp., São Paulo.
- 1938c. Further fly parasites of *Dysdercus*. *Rev. de Ent.* 8:347-348.
- 1939a. *Manual of Myiology, in twelve parts.* Pt. 8. Oestroid generic diagnoses and data (Microtropesini to Voriini), 408 pp., São Paulo.
- 1939b. New striking melanophorid tribe and gymnosomatid genera from Rio de Janeiro (Diptera, Oestromuscaria). *Rev. de Ent.* 10: 249-254.
- 1939c. Seven new genera of Brazilian oestromuscarian flies. *Rev. de Ent.* 10:446-452.
- 1939d. The species of the supergenus *Paratheresia* (Sic!) (Dipt.). *Rev. de Ent.* 10:546-549.
- 1939e. *Manual of Myiology, in twelve parts.* Pt. 9: Oestroid generic diagnoses and data (Thelairini to Clythoini), 270 pp., São Paulo.



- 1940a. *Manual of Myiology, in twelve parts*. Pt. 10: Oestroid generic diagnose and data (Anacamptomyini to Frontinini), 335 pp., São Paulo.
- 1940b. New oestroid flies from Brazil. *Rev. de Ent.* 11:889-894.
- 1941a. *Manual of Myiology, in twelve parts*. Pt. 11: Oestroid generic diagnoses and data (Goniini to Trypherini), 342 pp., São Paulo.
- 1941b. New fly parasites of *Diatraea* in São Paulo. *Rev. de Ent.* 12: 339-341.
- 1942a. *Manual of Myiology, in twelve parts*. Pt. 12: General consideration of the Oestromuscaria, 365 pp., 82 pls., São Paulo.
- 1942b. Possible fly parasite of *Diatraea* in São Paulo. *Rev. de Ent.* 12:339-341.
- 1942c. New fly parasites of *Diatraea*. *Rev. de Ent.* 13(1-2):149-150.
- 1942d. Two new reared South American flies. *Rev. de Ent.* 13(3): 438-439.

TRAVASSOS Fº, L. & M. CARRERA

1941. *Xanthozona melanopyga* (Wiedemann, 1830) (Diptera, Tachinidae) predadora de *Brassolis astyra* Godart, 1824 (Lepidoptera, Brassolidae) praga das palmeiras. *Arq. Zool. Secr. Agr. S. Paulo* 3:43-74, 24 figs.

VERBEKE, J.

1962. Contribution à l'étude des Tachinidae Africains (Diptera) 1. Description et valeur taxonomique des genitalia males. 2. Imatomyini Palpostomatini et Ethillini nouveaux ou peu connus. *Expl. Hidr. Lacs Kivu Edouard et Albert* 3(4):79-187, 27 pls.
1963. The structure of the male genitalia in Tachinidae (Diptera) and their taxonomic value. *Stuttgart Beitr. Naturk.* (Dedic. E. Lindner) 114:1-5, 27 figs.

VILLENEUVE, J.

1932. Notes diptérologiques. *Bull. Soc. Ent. France* 37:271-272.
1933. Description de *Aplomyiopsis galerucellae*, n. gen., sp. (Tachinidae), parasite de *Galerucella luteola* (F. Müll.) en Amérique du Nord. *Bol. R. Ist. Sup. Agr. Portici* 27:125-126.
1934. Notes Diptérologiques. *Rev. Franc. d'Entomologie* 1:180-183.
1937. Descriptions de Myodaire supérieurs. *Rev. Zool. Bot. Afr.* 29: 205-212.

VILLERS, C. DE

1789. *Caroli Linnaei entomologia, faunae suecicae descriptionibus aucta*. Vol. 3, 657 pp., 4 pls. Lugduni [= Lyon, France].

VIMMER, A.

1940. Prehled rodu jihoamerických hystriciid (Tachinoidea, Dipt.). Übersicht der Gattungen der südamerikanischen hystriciiden (Tachinidae, Dipt.). *Casopis (Acta Soc. Ent. Cechoslovenie*, vol. 37, nº 3-4, p. 100-13 (in Czech and German).



VIMMER, A. & SOUKUP, S. S.

1940. Los dípteros de los alrededores de Puno. Tachinoidea. *Bol. Mus. Nat. "Javier Prado"* (Peru), 4(13):205-223; (14); 360-372.

WACHTL, F. A.

1894. Analytische Uebersicht der europäischen Gattungen aus dem Verwandtschaftskreise von *Echinomya* Duméril nebst Beschreibung einer neuen *Eudora*. *Wien Ent. Ztg.* 13:140-144.

WALKER, F.

1837. Descriptions, etc, of the insects collected by Cpt. P. P. King, R. N. F. R. S., in the Survey of the Straits of Magellan. Diptera. *Trans. Linn. Soc. London* 17:331-359.
1849. *List of the specimens of dipterous insects in the collection of the British Museum*, Vol. 4. 689-1172. London.
1852. Diptera. Vol. 1, pp. 157-252; 233-414, 4 pls. (cont.). In Saunders, W. W., ed, *Insecta Saundersiana*. London. "1856".
- 1857-58. Caracteres of undescribed Diptera in collection of W. W. Saunders. *Trans. Ent. Soc. London*, n. ser (= ser. 2) 4:119-158; 1857; 190-235, 1958.
1860. Characters of undescribed Diptera in the collection of W. W. Saunders. *Trans. Ent. Soc. London* (1858-1861) n. ser. (= ser. 2). 5:268-296 (cont.).
1861. Characters of undescribed Diptera in collection of W. W. Saunders (concl.). *Trans. Ent. Soc. London* (1858-1861), n. ser. = ser. 2, 5:297-334.

WALTON, W. R.

- 1913a. A new species of Tachinidae from Porto Rico. *Proc. Ent. Soc. Wash.* 14:198-200, pl. X.
- 1913b. New North American Tachinidae (Dipt.). *Ent. News* 24:49-52, pl. III.
- 1914a. A new tachinid parasite of *Diabrotica vittata*. *Proc. Ent. Soc. Wash.* 16:11-14, 1 pl.
- 1914b. Four new species of Tachinidae from North America. *Proc. Ent. Soc. Wash.* 16:90-95, 7 figs.
1916. The tachinid genus *Argyrophylax* B. & B. *Proc. Ent. Soc. Wash.* 18:189-192, pl. xiv.

WEBBER, R. T. (see also ALDRICH & WEBBER)

1930. A revision of the North American tachinid flies of the genus *Achaetuncura*. *Proc. U. S. Natl. Mus.* 78(10):1-37, 14 figs.
1931. A new parasite fly of the genus *Chaetophlepsis*. *Proc. U. S. Natl. Mus.* 78(20):1-4.

WEST, L. S.

1925. New Phasiinae and Tachinidae from New York State. *Jour. N. Y. Ent. Soc.* 33:121-135.



WESTWOOD, J. O.

1840. Order XIII. Diptera Aristole (Antliata Fabricius, Halteriptera Clairv.). Pp. 125-128 (= signature I, part), 129-144 (= signature K), 145-158 (= signature L.). In his *An introduction to the modern classification of insects. Synopsis of the genera of British insects*, 158 pp. London.

WIEDEMANN, C. R. W.

1819. Brasilianische Zweiflüger. *Zool. Mag.* (Wiedemann's) 1(3):40-56.
1824. *Munus rectoris in Academia Christiana Albertina aditurus analecta entomologica ex Museo Regio Havniensi*, 60 pp., 1 pl. Kiliae [= Kiel].
1830. *Aussereuropäische zweiflügelige Insekten*. Vol. 2, xii + 684 pp., 5 pls. Hamm.

WILLISTON, S. W.

1886. Dipterological notes and descriptions. *Trans. Amer. Ent. Soc. and Proc. Acad. Nat. Sci. Phila.* 13:287-307.
1887. North American Tachinidae. *Gonia. Canad. Ent.* 19:6-12.
1888. An interesting new genus of South American Tachinidae. *Ent. News* 3:151-153.
1889. The dipterous parasites of North American butterflies. Pp. 1912-1924, pl. 89 (= pt. 12, part). In Scudder, S. H. *Butterflies of the eastern United States and Canada*, vol. 3, pp. 1775-1958, pls. 1-89. Cambridge, Mass.
1896. On the Diptera of St. Vincent (West Indies). *Trans. Ent. Soc. London* 1896:253-446, pls. 8-14.
1908. *Manual of North American Diptera*. Ed. 3, 405 pp., 163 figs. New Haven, Conn.

WULP, F. M. VAN DER

1867. Eenige Noord-Americaansche Diptera. *Tidschr. v. Ent.* 10 = ser. 2, 2:125-164, pls. 3-5.
1882. Remarks on certain American Diptera in the Leyden Museum and descriptions of nine species. *Notes Leyden Mus.* 4:73-92.
1883. Amerikaansche Diptera concl. *Tidjschr. v. Ent.* 26:223-282 (1-60), pls. 1-2.
1884. Quelques Diptère Exotiques. *Ann. Soc. Ent. Belg.* 28 (Bul.): cclxxxviii-cxcvii, 3 figs.
1885. Langwerpige dexinen-vormen. *Tidschr. v. Ent.* 28:301-311 (189-200), 1 col., plate.
1886. Notes on *Dejeania* Robineau-Desvoidy. *Tidschr. v. Ent.* 29:xxix-xxxii.
1888-1896. Fam. Muscidae. Pp. 2-40, pls. 1-2, 1888; pp. 41-56, 57-88, 89-112, 113-144, 145-176, 177-200, 201-208, pls. 3-4, 1890:273-281, 1896 (cont.). In Godman, F. D. and Salvin, O., eds. *Biol. Centrali-Americana. Zoologia, Insecta-Diptera*. Vol. 2, 489 pp., 11 figs., 13 pls. London.



1891. Fam. Muscidae (cont.). Pp. 209-224, 225-248, 249-264, pls. 5-6 (cont.). In Godman, F. D. and Salvin, O. eds. *Biologia Centrali Americana. Zoologia*.
1892. Diagnoses of new Mexican Muscidae. *Tidschr. v. Ent.* 35: 183-195.

ZUSKA, J.

1963. The puparia of the European species of the family Larvaevoridae (Diptera). I. (Subfamily Salmaciinae, part. 1). *Acta Ent. Mus. Nat. Pragae* 35:333-372, 85 figs.

INDEX

abbreviata Bigot, 1890 (Ceromasia)	104.152
abdominalis (Townsend), 1934, Abolodoria	104.204
abdominalis Townsend, 1929, Eutrichopoda	104.7
abdominalis (Townsend), 1914 (Eujurinella), Jurinella	104.56
abdominalis Townsend, 1914a, Minthomyia	104.141
abdominalis (Wulp), 1890, (Masicera), Myiosturmiopsis	104.190
Abepalpus Townsend, 1931a	104.74
aberrans (Giglio-Tos), 1893 (Nemochaeta), Archytas	104.48
aberrans Thompson, 1966, Pseudosiphosturmia	104.191
aberrans (Townsend), 1920, (Ptilolydella), Actinotachina	104.124
aberrans Townsend, 1935d, Bonnaniops	104.127
aberrans Wulp, 1890, Anisia	104.125
Abolodoria Townsend, 1934	104.204
abrupta (Wiedemann), 1830 (Tachina), Bombyliopsis	104.40
abscondens (Townsend), 1914 (Neojurinia), Jurinella	104.56
Acaulona Wulp, 1884	104.6
accedens Wulp, 1890, Anisia	104.125
Acemeigenia Townsend, 1927a	104.123
Acemyini	104.123
Acemyiopsis Townsend, 1915m	104.123
Achaetoneura Brauer & Bergenstamm, 1891	104.208
Achaetoneuropsis Townsend, 1927a	104.208
aclista Reinhard, 1956c, Peleteria	104.44
Acroceronia Cortés, 1951b	104.74
Acroglossa Williston, 1889	104.173
Acromiodexia Townsend, 1931a	104.105
acromion (Wiedemann), 1824 (Musca), Acromiodexia	104.105
Acronacantha Wulp, 1891	104.23
Acronarista Townsend, 1908	104.164
Acronaristopsis Townsend, 1919b	104.164
Actiini	104.164
Actia Robineau-Desvoidy, 1830	104.164
Actinactia Townsend, 1927a	104.165
Actinochaeta Brauer & Bergenstamm, 1889	104.113
Actinochaetopsis Townsend, 1934a	104.113
Actinocrocota Townsend, 1935d	104.165
Actinodoria Townsend, 1927a	104.123



Actinominthella Townsend, 1928c	104.114
<i>Actinomintho</i> Townsend, 1928b	104.94
<i>Actinophryno</i> Townsend, 1928b	104.124
<i>Actinoplagia</i> Blanchard, 1940	104.86
<i>Actinoprosopa</i> Townsend, 1927a	104.205
<i>Actinotachina</i> Townsend, 1927a	104.124
<i>acuminata</i> (Bigot), 1885b (<i>Oxydextia</i>), <i>Uramyia</i>	104.104
<i>acuminata</i> Tothill, 1924, <i>Fabriciella</i>	104.43
<i>acuminata</i> (Wulp), 1890 (<i>Macquartia</i>), <i>Paraporia</i>	104.104
<i>Acuphoceropsis</i> Blanchard, 1943b	104.74
<i>ada</i> Curran, 1947, <i>Amicrotrichomma</i>	104.47
<i>adamsoni</i> Thompson, 1963b, <i>Zenillia</i>	104.214
<i>Adejeania</i> Townsend, 1913e	104.45
<i>Adercomyia</i> Arnaud, 1963b	104.156
<i>Adidyma</i> Townsend, 1935d	104.205
<i>adjuncta</i> (Wulp), 1890 (<i>Mystacella</i>), <i>Carcelia</i>	104.204
<i>Admontia</i> Brauer & Bergenstamm, 1889	104.171
<i>Admontiini</i>	104.171
<i>Adoryphorophaga</i> Townsend, 1931	104.124
<i>adscrita</i> Wulp, 1890 (<i>Telothyria</i>), <i>Sitophaga</i>	104.148
<i>adunata</i> Reinhard, 1961, <i>Acronarista</i>	104.164
<i>adusta</i> (Wulp), 1890 (<i>Jurinia</i>), <i>Juriniopsis</i>	104.58
<i>adusta</i> , authors, <i>Juriniopsis</i>	104.58
<i>adustum</i> Reinhard, 1958c, <i>Mochlosoma</i>	104.30
<i>adversa</i> Townsend, 1935d, <i>Adidyma</i>	104.202
<i>Aeglops</i> Aldrich, 1934	104.88
<i>aegrota</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>aelops</i> (Walker), 1849 (<i>Tachina</i>), <i>Beskia</i>	104.14
<i>aemulans</i> Reinhard, 1964e, <i>Sitellitergus</i>	104.35
<i>aemulans</i> Wulp, 1890, <i>Prospherysa</i>	104.213
<i>aenea</i> (Aldrich), 1934 (<i>Aeglops</i>), <i>Ollacheryphe</i>	104.88
<i>aenea</i> Bigot, 1889 (<i>Chaetolyga</i>)	104.152
<i>aenea</i> (Coquillett), 1895d, (<i>Hypostena</i>) <i>Elfia</i>	104.167
<i>aenea</i> Wiedemann, 1830 (<i>Tachina</i>), <i>Mylophasia</i>	104.27
<i>aenescens</i> Curran, 1927b, <i>Hesperophasia</i>	104.93
<i>Aelofrontina</i> Townsend, 1928c	104.212
<i>aequabilis</i> (Walker), 1849 (<i>Tachina</i>), <i>Cryptopalpus</i>	104.61
<i>affinis</i> Reinhard, 1934e, <i>Paradidyma</i>	104.115
<i>affinis</i> (Schiner), 1868 (<i>Saundersia</i>), <i>Exopalpus</i>	104.64
<i>affinis</i> (Townsend), 1927a (<i>Achaetoneuropsis</i>), <i>Lespesia</i>	104.208
<i>afra</i> (Wulp), 1890 (<i>Brachycoma</i>), <i>Lespesia</i>	104.209
<i>Agicuphocera</i> Townsend, 1915m	104.74
<i>agilis</i> Curran, 1934b, <i>Calodextia</i>	104.154
<i>Aglummyia</i> Townsend, 1912a	104.28
<i>Agrarialia</i> Curran, 1934d	104.161
<i>ajax</i> Curran, 1947, <i>Jurinella</i>	104.56
<i>alabamae</i> (Townsend), 1940b (<i>Hemimasipoda</i>), <i>Winthemia</i>	104.196
<i>alacris</i> Walker, 1860 (<i>Masicera</i>)	104.193
<i>alba</i> Townsend, 1917a, <i>Euphoroceroides</i>	104.158
<i>albescens</i> Aldrich, 1934, <i>Admontia</i>	104.171
<i>albicauda</i> Townsend, 1917b, <i>Thysanopsis</i>	104.187
<i>albiceps</i> (Walker), 1860 (<i>Echinomyia</i>), <i>Archytas</i>	104.48
<i>albiceps</i> (Wulp), 1822 (<i>Hystricia</i>), <i>Bombyliomyia</i>	104.39
<i>albicincta</i> Townsend, 1912a, <i>Jaenimyia</i>	104.185



<i>albicincta</i> Wulp, 1891, <i>Stenodexia</i>	104.108
<i>albidula</i> Wulp, 1890, (<i>Meigenia</i>)	104.152
<i>albifacies</i> (Bigot), 1888a (<i>Blepharipeza</i>), <i>Chaeteprosopa</i>	104.184
<i>albifacies</i> (Bigot), 1889 (<i>Oplisa</i>), <i>Pseudodexia</i>	104.108
<i>albifacies</i> (Wulp), 1892 (<i>Meigenia</i>), <i>Paralispe</i>	104.145
<i>albifacies</i> Wulp, 1890 (<i>Prospheysa</i>)	104.180
<i>albifrons</i> (Walker), 1837 (<i>Tachina</i>)	104.193
<i>albilineata</i> Blanchard, 1943b, <i>Vibrissomyia</i>	104.81
<i>albimacula</i> (Wiedemann), 1830 (<i>Tachina</i>), <i>Deopalpus</i>	104.76
<i>albimana</i> Curran, 1942, <i>Hystricia</i>	104.41
<i>albincisa</i> (Wiedemann), 1830 (<i>Tachina</i>), <i>Argyrophylax</i>	104.162
<i>albipes</i> (Townsend), 1931e (<i>Lasiopalpus</i>), <i>Eulasiopalpus</i>	101.54
<i>albipes</i> Townsend, 1914, <i>Melanepalpellus</i>	104.68
<i>albipila</i> Curran, 1941, <i>Eudejeania</i>	104.53
<i>albocalyptrata</i> Bigot 1888a, <i>Trichophora</i>	104.72
<i>albocingulata</i> Wulp, 1890 (<i>Hypostena</i>)	104.152
<i>albofenestrata</i> Curran, 1928a, <i>Jamaicara</i>	104.20
<i>albolineata</i> Wulp, 1891, <i>Bathydexia</i>	104.36
<i>albolineatus</i> (Macquart), 1854 (<i>Micropalpus</i>), <i>Epalpodes</i>	104.63
<i>albomaculata</i> Robineau-Desvoidy, 1863 (<i>Pcleteria</i>)	104.215
<i>albomaculatus</i> (Jaenicke), 1867 (<i>Micropalpus</i>), <i>Epalpus</i>	104.64
<i>albomarginata</i> Wulp, 1890 (<i>Hypostena</i>)	104.152
<i>albomicans</i> Wulp 1890, <i>Didyma</i>	104.131
<i>albopicta</i> (Bigot), 1889 (<i>Chaetolyga</i>), <i>Spathimeigenia</i>	104.148
<i>albopilosa</i> (Curran), 1926a (<i>Masicera</i>), <i>Blondella</i>	104.126
<i>albosignata</i> (Wulp), 1892 (<i>Hystricia</i>), <i>Calohystricia</i>	104.40
<i>albula</i> Arnaud 1963b, <i>Cerotachina</i>	104.155
<i>albuquerquei</i> Guimarães, 1963c, <i>Euempheremyia</i>	104.66
<i>albuquerquei</i> Guimarães, 1962, <i>Peleteria</i>	104.44
<i>alcedo</i> (Loew), 1869 (<i>Gymnochaeta</i>), <i>Chrysotachina</i>	104.81
<i>aldrichi</i> Blanchard, 1943, <i>Peleteria</i>	104.44
<i>aldrichi</i> Blanchard, 1966b, <i>Proxanthobasis</i>	104.110
<i>aldrichi</i> Cortés, 1944c, <i>Cylindromyia</i>	104.15
<i>aldrichi</i> Curran, 1934b, <i>Calodexia</i>	104.154
<i>aldrichi</i> Reinhard, 1934a, <i>Paradidyma</i>	104.115
<i>aldrichi</i> Reinhard, 1935a, <i>Uramya</i>	104.104
<i>aldrichi</i> Sabrosky, 1947, <i>Eudejeania</i>	104.53
<i>aldrichi</i> (Séguy), 1925 (<i>Ormia</i>), <i>Euphasiopteryx</i>	104.20
<i>aldrichi</i> (Townsend), 1931a (<i>Willistonina</i>), <i>Belvosia</i>	104.181
<i>aldrichi</i> (Townsend), 1931b (<i>Spanipalpus</i>), <i>Deopalpus</i>	104.76
<i>aldrichi</i> Townsend, 1891a, <i>Hyalomya</i>	104.13
<i>Aldrichiopa</i> Guimarães, nom. nov.	104.165
<i>aleurites</i> Townsend, 1940b, <i>Conactiodoria</i>	104.130
<i>Alexogloblinia</i> Cortés, 1945b	104.97
<i>alipes</i> Wulp, 1892, <i>Trichopoda</i> (<i>Trichopoda</i>)	104.9
<i>alligans</i> (Walker), 1849 (<i>Tachina</i>), <i>Epalpus</i>	104.64
<i>Allosturmia</i> Blanchard, 1958	104.187
<i>Almugmyia</i> Townsend, 1911b	104.28
<i>Alophorella</i> Townsend, 1912e	104.11
<i>Alophorellopsis</i> Townsend, 1927a	104.13
<i>alpina</i> Townsend, 1915e, <i>Chaetocnephalia</i>	104.174
<i>alpina</i> Townsend, 1914, <i>Echinopyrrhosia</i>	104.62
<i>alpina</i> Townsend 1913, <i>Eudejeania</i>	104.54
<i>alpina</i> Townsend, 1931a, <i>Engelomyia</i>	104.41

alpina (Townsend), 1912a (Salmacia), Gonia	104.175
alpina Townsend, 1931a, Mesembrierigone	104.84
alpina Townsend, 1931a, Opticopteryx	104.84
Alpinoplagia Townsend, 1931a	104.89
alpinum Townsend, 1912a, Dolichostoma	104.77
Alsopsyche Brauer & Bergenstamm, 1891	104.198
alternus (Walker), 1849 (Tachina), Epalpus	104.64
alypiae Sellers, 1943, Sisyropa	104.204
Amazohoughia Townsend, 1934	104.201
<i>Amazonella</i> Myers, 1934	104.210
amazonica (Townsend), 1934a (Actinochaetospsis), Actinochaeta	104.113
amazonica (Townsend), 1934a (Azulihoughia), Calocarcelia	104.202
amazonica Townsend, 1927a, Egameigenia	104.132
amazonica Townsend, 1934a, Helioplagia	104.87
ambulatorix Wulp, 1890, Didyma	104.132
ambigua (Macquart), 1851 (Hystricia), Jurinella	104.56
Amblychaeta Aldrich, 1934	104.87
amblycera Aldrich, 1934, Actia	104.164
ambulatorix Wulp, 1890, Didyma	104.131
americana Bigot, 1889 (Prosopaea)	104.152
americana Brooks, 1945, Thelaira	104.95
americana (Curran), 1927c (Anacamptomyia), Hemisturmia ...	104.194
<i>americana</i> Macquart, 1843, Melanophora	104.4
americana (Schiner), 1868 (Cnephallia), Chaetocnephallia	104.174
americana (Townsend), 1892c (Tryptocera), Actia	104.164
<i>americana</i> Wulp, 1890 (Plagia), Voria	104.93
Ametadoria Townsend, 1927a	104.205
Amicrotrichomma Townsend, 1927a	104.47
amisias (Walker), 1849 (Tachina), Macromya	104.43
amoena Macquart, 1843, Hystricia	104.41
amydriae Townsend, 1908, Drepanoglossa	104.167
<i>Anacamptomyini</i> , in part	104.122
<i>Anachaetopsis</i> , authors	104.173
Anadiscalia Curran, 1934d	104.201
Anadistichona Townsend, 1934	104.178
anale Giglio-Tos 1893, Mochlosoma	104.30
analis Aldrich, 1934, Telodytes	104.84
<i>analis</i> , authors, Archytas	104.48
analis Curran, 1937, Chaetogyne	104.36
analis (Fabricius), 1805 (Tachina), Archytas	104.48
analis (Macquart) 1843 (Dejeania), Adejeania	104.46
analis (Macquart) 1843 (Jurinia), Archytas	104.48
analis Macquart, 1846, Belvosia	104.181
analis (Macquart), 1851 (Megistogaster), Cordyligaster	104.111
analis Macquart, 1843 (Gymnostylia), Macromya	104.43
analis (Macquart), 1846 (Echinomyia), Peleteria	104.44
analis (Macquart), 1846 (Microtrichodes), Winthemia	104.196
analis Robineau-Desvoidy, 1830, Macromya	104.42
analis Schiner 1868, Trichophora	104.72
analis Townsend, 1931a, Anhangabahuia	104.205
analis (Townsend) 1927a (Anazygosturmia), Drino	104.188
analis Townsend, 1927a, Eubischofimyla	104.65
analis Townsend, 1914, Gymnochaetopsis	104.82
analis Townsend, 1927a, Itacnephallia	104.176



analís (Townsend), 1927a, (Sarcolydella), Prospheysa	104.213
analís Townsend, 1915n, Uruhuasiopsis	104.73
analís (Townsend), 1927a, Yahuar Mayoia	104.102
analísella Thompson, 1963c, Winthemia	104.196
Anametopochaeta Townsend, 1919a	104.124
anametopochaetoides Townsend, 1934, Cuparymyia	104.130
Anaphorinia Townsend, 1927a	104.124
anax Curran, 1947, Jurinella	104.56
Anazygosturmia Townsend, 1927a	104.188
anceps (Fabricius), 1805 (Musca), Trichodura	104.103
anceps (Wulp), 1812 (Xanthomelana), Vanderwulpella	104.18
ancilla (Walker), 1852 (Tachina), Pseudomyothyria	104.146
ancisa Reinhard, 1952b, Chaetophlepsis	104.89
Ancylogaster Bigot, 1884b	104.16
ancylostomiae Thompson, 1968, Carmodymyia	104.128
andana Robineau-Desvoidy, 1863 (Jurinia), Archytas	104.52
andeana Sabrosky, 1947, Eudejeania	104.53
Andesimya Brèthes, 1909	104.178
andicola Bigot, 1888b (Fabricia), Peleteria	104.44
andicola Townsend, 1914, Jurinella	104.56
andicula Townsend, 1915h, Archytas	104.49
andina Bigot, 1888a (Blepharipeza)	104.193
andina (Blanchard), 1943b (Cuphoceropsis), Peleteria	104.44
andina (Townsend), 1912a, (Dejeania) Adejeania	104.46
andina Townsend 1919a, Copecrypta	104.75
andina Townsend, 1912a, Chaetophorocera	104.206
andina Townsend, 1927a, Epicuphocera	104.77
andina (Townsend), 1914 (Jurinella), Exopalpus	104.55
andina Townsend, 1915e, Germariopsis	104.175
andina Townsend, 1927a, Metopactia	104.140
andina Townsend, 1931a, Neotrafoiopsis	104.82
andina (Townsend), 1929 (Gymnosturmia), Topomeigenia	104.193
andinensis Townsend, 1929, Chrysophryno	104.207
andinensis Townsend, 1927a, Olindopsis	104.142
andinum (Townsend), 1927a (Epidolichostoma), Dolichostoma	104.77
andinus Townsend, 1912a, Megaprosopus	104.24
andinus Townsend, 1914, Rhachoepalpus	104.70
Andinomyia Townsend, 1912a	104.75
Androeuryps Beneway, 1961	104.12
anduzei Curran, 1947, Adejeania	104.46
Anemorilla Townsend, 1915m	104.187
Anaepalpus Townsend, 1931a	104.60
Anetia, authors	104.126
Angiorrhina Brauer & Bergenstamm, 1889	104.23
angolensis Aldrich, 1934, Lypha	104.85
angrensis Guimarães, 1963b, Archytas	104.48
angulata Wulp, 1890 (Myobia)	104.120
angulata (Wulp), 1891 (Rhynchodexia), Ptilodexia	104.33
angusta Macquart, 1843, Gonia	104.175
angusta Perty, 1833, Diaugia	104.101
angusta Reinhard, 1967, Ziziphomyia	104.193
angusta Townsend, 1927a, Actinotachina	104.124
angusta Townsend, 1928b, (Actinophryno), Actinotachina	104.124
angusta Townsend, 1927a, Catajurinia	104.75
angusta Townsend, 1927a, Cataphorinia	104.128

<i>angusta</i> Townsend, 1916h, <i>Frontocnephalia</i>	104.175
<i>angusta</i> Townsend, 1927a, <i>Mayophorinia</i>	104.139
<i>angusta</i> Townsend, 1927a, <i>Metarrhinomyia</i>	104.140
<i>angusta</i> (Walker), 1852 (<i>Dexia</i>), <i>Myiobiopsis</i>	104.118
<i>angustata</i> (Wulp), 1890 (<i>Exorista</i>)	104.214
<i>angustata</i> , error, <i>Gonia</i>	104.175
<i>angustifrons</i> Aldrich, 1934, <i>Xanthobasis</i>	104.110
<i>angustifrons</i> Reinhard, 1930a (<i>Viviania</i>), <i>Sitophaga</i>	104.148
<i>angustifrons</i> Townsend, 1916g, <i>Chrysoexorista</i>	104.207
<i>angustifrons</i> Wulp, 1892 (<i>Micropalpus</i>)	104.83
<i>angustipennis</i> (Townsend), 1927a, (<i>Tinanemorilla</i>), <i>Nemorilla</i>	104.195
<i>Anhangabahuia</i> Townsend, 1931a	104.205
<i>Anisia</i> Wulp, 1890	104.125
<i>anomala</i> Aldrich, 1934, <i>Prosopochaeta</i>	104.99
<i>anomala</i> Townsend, 1939, <i>Urohypomyia</i>	104.109
<i>Anoxynops</i> Townsend, 1927a	104.125
<i>Anoxynopsoidea</i> Thompson, 1968	104.126
<i>Anoxynopsella</i> Townsend, 1935d	104.126
<i>ansata</i> Reinhard, 1951, <i>Belvosia</i>	104.181
<i>antactica</i> (Thomson), 1869, (<i>Degeeria</i>), <i>Austrostaurochaeta</i> ...	104.172
<i>antarctiae</i> Blanchard, in [Marchionatto], 1940	104.204
<i>antennalis</i> Aldrich, 1934, <i>Myiophasia</i>	104.27
<i>antennalis</i> Thompson, 1968, <i>Maracasimyia</i>	104.139
<i>antennalis</i> Townsend, 1927a, <i>Italispidea</i>	104.137
<i>antennalis</i> Townsend, 1927a, <i>Itaplectopsis</i>	104.137
<i>antennalis</i> Townsend, 1927a, <i>Phyllophryno</i>	104.145
<i>anthemon</i> (Walker), 1849 (<i>Tachina</i>), <i>Macromya</i>	104.42
<i>anthomyformis</i> Lynch-Arribáizaga in Brauer, 1898, <i>Siphona</i>	104.170
<i>anthracina</i> (Bigot), 1889 (<i>Rhamphininina</i>), <i>Ptilodexia</i>	104.33
<i>anthracina</i> (Wiedemann), 1830 (<i>Tachina</i>), <i>Euloewiopsis</i>	104.26
<i>anthracinus</i> (Bigot), 1889 (<i>Degeeria</i>), <i>Oxynops</i>	104.143
<i>antillensis</i> Arnaud, 1960, <i>Spathidexia</i>	104.95
<i>Antillicolla</i> Curran, 1927a	104.49
<i>antillicolla</i> Curran, 1927c, <i>Archytas</i>	104.49
<i>antilliana</i> Curran, 1927 (<i>Belvosia</i>), <i>Triachora</i>	104.184
<i>Antistaseopsis</i> Townsend, 1934a	104.205
<i>antlerata</i> Townsend, 1927a, <i>Ucayalimyia</i>	104.214
<i>aperta</i> Aldrich, 1934, <i>Dasyuromyia</i>	104.97
<i>aperta</i> Brauer & Bergenstamm, 1893 (<i>Wulpia</i>), <i>Catemophryx</i>	104.114
<i>aperta</i> (Brauer & Bergenstamm), 1889 (<i>Oestrophasia</i>), <i>Euoeostro-</i> <i>phasia</i>	104.20
<i>aperta</i> Brauer & Bergenstamm, 1893, <i>Paradidyma</i>	104.115
<i>Aphelogaster</i> Aldrich, 1934	104.165
<i>Aphriosphyria</i> Townsend, 1927a	104.43
<i>Aphriiini</i>	104.85
<i>Aphriosphyriopsis</i> Blanchard, 1943b	104.43
<i>apicalis</i> (Bigot), 1878, <i>Cylindromyia</i>	104.15
<i>apicalis</i> Brêthes, 1920a, <i>Didyma</i>	104.131
<i>apicalis</i> Curran, 1934b, <i>Calodexia</i>	104.154
<i>apicalis</i> (Jaennicke), 1867 (<i>Jurinia</i>), <i>Archytas</i>	104.50
<i>apicalis</i> Reinhard, 1934a, <i>Paradidyma</i>	104.115
<i>apicalis</i> Wiedemann, 1830, <i>Trichopoda</i> (<i>Trichopoda</i>)	104.9
<i>apicalis</i> (Wulp), 1890 (<i>Prospherysa</i>), <i>Lespesia</i>	104.209
<i>Apinocyptera</i> Townsend, 1915k	104.15



<i>apicata</i> Curran, 1947, <i>Jurinella</i>	104.56
<i>apicata</i> Curran, 1925c, <i>Peleteria</i>	104.44
<i>apicifer</i> (Walker), 1849, (<i>Tachina</i>) <i>Archytas</i>	104.48
<i>Aplomyia</i> Robineau-Desvoidy, 1830	104.205
<i>Aplomyodoria</i> Townsend, 1928c	104.161
<i>Aplomyiopsis</i> Villeneuve, 1933	104.126
<i>Aplomyiopsis</i> Townsend, 1927a	104.205
<i>Aporia</i> Macquart, 1846	104.164
<i>appendiculata</i> (Bigot), 1889 (<i>Phorostoma</i>), <i>Myiomima</i>	104.31
<i>appendiculata</i> Bigot of Coquillett, 1910, <i>Bathydextia</i>	104.36
<i>appendiculata</i> (Wulp), 1890 (<i>Phorocera</i>), <i>Euphorocera</i>	104.157
<i>approximata</i> Wulp, 1890 (<i>Anisia</i>)	104.164
<i>arabella</i> Townsend, 1912a, <i>Phaenopsis</i>	104.163
<i>araujoi</i> Guimarães, 1963d, <i>Archytas</i>	104.48
<i>Araucosimus</i> Aldrich, 1934	104.173
<i>archippivora</i> (Riley), 1871 (<i>Tachina</i>), <i>Lespesia</i>	104.209
<i>Archynactia</i> , misprint	104.94
<i>Archynemochaeta</i> Blanchard, 1941	104.48
<i>Archytas</i> Jaennicke, 1867	104.47
<i>Archytodejeania</i> Blanchard, 1941	104.48
<i>archytoides</i> Townsend, 1931a, <i>Abepalpus</i>	104.74
<i>Archytoepalpus</i> Townsend, 1927a	104.60
<i>Arctosoma</i> Aldrich, 1934	104.85
<i>arcuata</i> (Bigot), 1876 (<i>Trichopoda</i>), <i>Ecatophasiopsis</i>	104.12
<i>arcuata</i> Townsend, 1919, <i>Nephoplagia</i>	104.91
<i>arcuatipennis</i> Machuart, 1885 (<i>Masicera</i>)	104.214
<i>arcuatus</i> (Say), 1829 (<i>Ocyptera</i>), <i>Xanthomelanodes</i>	104.10
<i>arcuatus</i> Townsend, 1915m, <i>Euepalpodes</i>	104.66
<i>arequipae</i> (Townsend), 1912a, (<i>Erigonopsis</i>), <i>Dolichostoma</i>	104.77
<i>argentaurea</i> Townsend, 1939d, <i>Paratheresia</i>	104.37
<i>argentea</i> Thompson, 1964, <i>Actinodoria</i>	104.123
<i>argentea</i> Townsend, 1914, <i>Signosomopsis</i>	104.72
<i>argentescens</i> Townsend, 1955d, <i>Anoxynopsella</i>	104.126
<i>argentescens</i> Townsend, 1927a, <i>Neargyrophylax</i>	104.142
<i>argenteus</i> Thompson, 1968, <i>Heliodexodes</i>	104.135
<i>argenteus</i> Townsend, 1914, <i>Rhachioepalpus</i>	104.70
<i>argenteiceps</i> Aldrich, 1925a, <i>Atacta</i>	104.180
<i>argenteiceps</i> Aldrich, 1928b, <i>Chaetocraniopsis</i>	104.174
<i>argenteiceps</i> (Wulp), 1892 (<i>Hyalomyia</i>), <i>Heyneophasia</i>	104.12
<i>argenteiceps</i> Wulp, 1890 (<i>Myobia</i>)	104.164
<i>argentifer</i> Aldrich, 1934, <i>Ganaproctus</i>	104.78
<i>argentifrons</i> Aldrich, 1925a, <i>Atacta</i>	104.180
<i>argentifrons</i> (Brooks), 1945 (<i>Alophorellopsis</i>), <i>Hyalomyia</i>	104.13
<i>argentifrons</i> Townsend, 1934, <i>Amazohoughia</i>	104.201
<i>argentifrons</i> (Wulp), 1890 (<i>Telothyria</i>), <i>Actinodoria</i>	104.124
<i>argentina</i> (Bigot), 1889 (<i>Rhamphiniina</i>), <i>Ptilodextia</i>	104.33
<i>argentina</i> (Brèthes), 1922 (<i>Gymnostylia</i>), <i>Oxyaporia</i>	104.143
<i>argentina</i> Townsend, 1931a, <i>Epimeigenia</i>	104.132
<i>argentinensis</i> Blanchard, 1941, <i>Bombyliopsis</i>	104.40
<i>argentinensis</i> Blanchard, 1941, <i>Fabriciopsis</i>	104.55
<i>argentinensis</i> Blanchard, 1942a, <i>Hyalomyodes</i>	104.18
<i>argentinensis</i> (Blanchard), 1966e (<i>Trichopodopsis</i>), <i>Trichopoda</i> (<i>Galactomyia</i>)	104.110
<i>argentinensis</i> Blanchard, 1941, <i>Zonoepalpus</i>	104.74

<i>Argentoepalpus</i> Townsend, 1919	104.63
<i>Argyreomyia</i> Townsend, 1915m	104.137
<i>Argyrochaetona</i> Townsend, 1919b	104.161
<i>Argyrodoria</i> Townsend, 1927a	104.126
<i>Argyromima</i> Brauer & Bergenstamm, 1889	104.14
<i>Argyrophylax</i> Brauer & Bergenstamm, 1889	104.161
<i>argyropha</i> (Schiner), 1868 (Dejeania), Eudejeania	104.53
<i>arida</i> Townsend, 1928c, <i>Aplomydoria</i>	104.161
<i>arida</i> (Townsend), 1911b, (<i>Almugmyia</i>), <i>Microchaetina</i>	104.28
<i>arida</i> Townsend, 1928c, (<i>Aelofrontina</i>), <i>Plagimasicera</i>	104.212
<i>arida</i> Townsend, 1915m, <i>Protogoniopsis</i>	104.184
<i>Aridalia</i> Curran, 1934d	104.202
<i>ariel</i> Curran, 1947, <i>Jurinella</i>	104.56
<i>aristalis</i> (Townsend), 1927a (<i>Ptilolydella</i>), <i>Actinotachina</i>	104.124
<i>aristata</i> Townsend, 1934, <i>Icteriodexia</i>	104.106
<i>armata</i> Röder, 1886, <i>Hystrichodexia</i>	104.30
<i>armata</i> (Townsend), 1915m (<i>Lachnommopsis</i>), <i>Paradidyma</i>	104.115
<i>armata</i> (Wiedemann), 1830 (<i>Tachina</i>), <i>Adejeania</i>	104.46
<i>armatus</i> Bigot, 1884b (<i>Ancylogaster</i>), <i>Hemyda</i>	104.16
<i>armigera</i> (Coquillett), 1889 (<i>Tachina</i>), <i>Eucelatoria</i>	104.133
<i>arnaudi</i> Guimarães, 1963b, <i>Archytas</i>	104.48
<i>Arrhinactia</i> Townsend, 1927a	104.94
<i>Arthrochaeta</i> Brauer & Bergenstamm, 1889	104.55
<i>articulata</i> (Wulp), 1892 (<i>Xanthomelanodes</i>), <i>Xenophasia</i>	104.11
<i>ascita</i> Reinhard, 1953b, <i>Microphthalma</i>	104.24
<i>Asilidotachina</i> Townsend, 1931a	104.155
<i>asolis</i> Gundlach in Pazos, 1944, <i>Melanophora</i>	104.4
<i>assimilis</i> Reinhard, 1954, <i>Prosenoides</i>	104.33
<i>assimilis</i> (Townsend), 1919a (<i>Austrolydella</i>), <i>Doryphorophaga</i> ..	104.132
<i>assimilis</i> (Wulp), 1892 (<i>Jurinia</i>), <i>Eujurinoides</i>	104.131
<i>assimulata</i> Wulp, 1891 (<i>Telothyria</i>)	104.152
<i>Atacta</i> Schiner, 1868	104.180
<i>Atactomima</i> Townsend, 1916h	104.180
<i>Atactopsis</i> Townsend, 1917b	104.180
<i>Atactosturmia</i> Townsend, 1916	104.180
<i>ategulata</i> Townsend, 1915i (<i>Eucordylidexia</i>), <i>Cordilygaster</i>	104.111
<i>Ateloglutus</i> Aldrich, 1934	104.89
<i>aterrimus</i> (Robineau-Desvoidy), 1830 (<i>Jurinia</i>), <i>Archytas</i>	104.48
<i>aterrimum</i> (Villers), 1938a (<i>Parapsalida</i>), <i>Leucostoma</i>	104.17
<i>Athrycia</i> Robineau-Desvoidy, 1830	104.90
<i>atra</i> (Aldrich), 1934 (<i>Lispidea</i>), <i>Elfia</i>	104.167
<i>atra</i> Bischof, 1904, <i>Empheremyia</i>	104.63
<i>atra</i> Brauer & Bergenstamm, 1893, <i>Gaediophana</i>	104.179
<i>atra</i> (Brauer & Bergenstamm), 1893, (<i>Pseudoclista</i>) <i>Myophasia</i> ..	104.27
<i>atra</i> (Röder), 1885 (<i>Ocyptera</i>), <i>Cylindromyia</i>	104.15
<i>atra</i> Townsend 1927a, <i>Bischofimyia</i>	104.60
<i>atra</i> Townsend, 1927a, <i>Euhuascaraya</i>	104.66
<i>atra</i> Townsend, 1927a, <i>Opsoempheria</i>	104.69
<i>atra</i> Townsend, 1927a, <i>Thelyphaenopsis</i>	104.150
<i>atrata</i> (Guérin-Ménéville), 1844 (Dejeania), Eudejeania	104.53
<i>atrata</i> (Walker), 1852 (<i>Tachina</i>), <i>Belvosia</i>	104.181
<i>atrata</i> (Wulp), 1888 (Dejeania)	104.60
<i>atratura</i> (Walker), 1852 (<i>Tachina</i>), <i>Paralispe</i>	104.145
<i>atricauda</i> Aldrich, 1934, <i>Cylindromyia</i>	104.15
<i>atriceps</i> Wulp, 1890 (<i>Phorocera</i>), <i>Metadoria</i>	104.140


Atrichiopoda Townsend, 1931a	104.6
atrifrons (Bigot), 1889 (Macquartia), Pseudeuantha	104.104
atrifrons (Wiedemann), 1830 (Musca), Tromodesiopsis	
atrifrons (Wiedemann), 1824 (Melanophora), Xanthomelanodes	104.10
atripennis Coquillett, 1895d, Chaetoplusia	104.90
atripennis (Say), 1829 (Ocyptera), Xanthomelanodes	104.10
atriventris Aldrich, 1934, Psecacera	104.25
Atrophopodus Townsend, 1892a	104.114
Atrophopoda Townsend, 1891b	104.115
atrophopodella Townsend, 1928c, Actinominthella	104.114
atrophopodoides Townsend, 1891b, Vanderwulpia	104.115
atypica Curran, 1927 (Spathidexia)	104.214
atypica Townsend, 1914, Echinopyrrhosia	104.62
aucta (Wiedemann), 1830 (Dexia), Euantha	104.111
aurantiacus Townsend, 1931a, Erythroepalpus	104.65
aurata Blanchard, 1963, Chaetogaedia	104.178
aurata Robineau-Desvoidy, 1830, Hemyda	104.16
aurata Townsend, 1927a, Anaphorinia	104.124
aurata Townsend, 1934a, Erviopsis	104.101
aurata Townsend, 1927a, Heliolydella	104.135
aurata Townsend, 1927a, Leskiolydella	104.138
aurata Townsend, 1934a, Orohoughia	104.163
aurata Townsend, 1934a, Urumyobia	104.120
auratilis Reinhard, 1951, Belvosia	104.181
auratofrontalis (Brèthes), 1908 (Exorista), Eusisyropa	104.203
auratus Campos, 1953, Pollops	104.169
aurea Aldrich, 1934, Notodytes	104.99
aurea (Blanchard), 1942b (Eutritochaeta), Nilea	104.210
aurea (Giglio-Tos), 1893 (Dejeania), Adejeania	104.46
aurea (Giglio-Tos), 1893 (Hystriodexia), Sumichrastia	104.35
aurea Thompson, 1963, Dexodioides	104.131
aurea Thompson, 1963, Dexodopsis	104.131
aurea Thompson, 1963d, Ignotomyia	104.198
aurea Thompson, 1963a, Trochiloglossa	104.119
aurea Townsend, 1911b, Azygobothria	104.187
aurea Townsend, 1934a, Conactiodoria	104.130
aurea Townsend, 1917, Diaphanomyia	104.62
aurea Townsend, 1915d, Dolichogonia	104.175
aurea (Townsend), 1916h (Macromeigenia), Mystacella	104.190
aurea Townsend, 1929, Vibrissosoria	104.178
aurea Vimmer & Soukup, 1940, Metopiopsis	104.217
aureocauda Thompson, 1966, Drino	104.188
aureocephala Thompson, 1964, Calocarcelia	104.202
aureolatus (Townsend), 1914 (Eusignosoma), Epalpus	104.64
aureonigra Thompson, 1963c, Winthemia	104.196
aurescens Townsend, 1917b, Eucelatoria	104.133
aurescens Townsend, 1931a, Itavoria	104.91
aurescens Townsend, 1915m, Micronychiops	104.85
aurescens Townsend, 1934a, Uruleskia	104.120
aureus (Aldrich), 1926b (Cuphocera), Chiloepalpus	104.61
aureus (Giglio-Tos), 1893 (Saundersia), Epalpus	104.64
aureus (Townsend), 1914 (Eusignosoma), Epalpus	104.64
auricaudata Brooks, 1945b, Paraphoranthia	104.13
auricephala (Blanchard), 1963, (Gaediosturmia) Chaetogaedia ..	104.178

104.257



<i>auricephala</i> Townsend, 1927a, <i>Eumachaeraea</i>	104.134
<i>auriceps</i> Aldrich, 1925d, <i>Chaeteprosopa</i>	104.184
<i>auriceps</i> Coquillett, 1904, <i>Distichona</i>	104.174
<i>auriceps</i> (Curran), 1927c, (<i>Antillicolla</i>), <i>Neossarromyia</i>	104.70
<i>auriceps</i> (Macquart), 1843 (<i>Masicera</i>), <i>Lespesia</i>	104.209
<i>auriceps</i> Reinhard, 1952a, <i>Cyosoprocta</i>	104.184
<i>aurifacies</i> Guimarães, 1963a, <i>Neocuphocera</i>	104.79
<i>aurifacies</i> (Robineau-Desvoidy), 1830 (<i>Jurinia</i>), <i>Diaphanomyia</i>	104.62
<i>aurifacies</i> Townsend, 1927a, <i>Chiloepalpus</i>	104.61
<i>aurifacies</i> Walton, 1912, <i>Cryptomeigenia</i>	104.130
<i>auriferus</i> (Walker), 1849 (<i>Tachina</i>), <i>Epalpus</i>	104.64
<i>aurifodina</i> Reinhard, 1954b, <i>Helioprosopa</i>	104.78
<i>aurifrons</i> Brooks, 1949, <i>Juriniopsis</i>	104.58
<i>aurifrons</i> (Macquart), 1846 (<i>Myobia</i>), <i>Myobiopsis</i>	104.118
<i>aurifrons</i> Thompson, 1968, <i>Anoxynops</i>	104.125
<i>aurifrons</i> (Townsend), 1919f (<i>Neoarchytas</i>), <i>Archytas</i>	104.49
<i>aurifrons</i> Townsend, 1915k, <i>Hystrieciella</i>	104.56
<i>aurifrons</i> Townsend, 1919a, <i>Schistostephana</i>	104.8
<i>auripilosa</i> (Blanchard), 1954 (<i>Willistonina</i>), <i>Belvosia</i>	104.181
<i>auroanalis</i> (Townsend), 1931a, (<i>Oxapampoepalpus</i>) <i>Parepalpus</i>	104.69
<i>aurocaudata</i> (Bigot), 1888a (<i>Blepharipeza</i>), <i>Drino</i>	104.188
<i>aurometallica</i> Townsend, 1927a, <i>Bolohoughia</i>	104.206
<i>auronigra</i> Townsend, 1927a, <i>Sturmioactia</i>	104.192
<i>aurulans</i> (Townsend), 1927a (<i>Prophryno</i>), <i>Lespesia</i>	
<i>aurulenta</i> (Bigot), 188a, (<i>Frontina</i>), <i>Belvosia</i>	104.181
<i>Austeniops</i> Townsend, 1915h	104.60
<i>australis</i> Aldrich, 1928, <i>Belvosia</i>	104.181
<i>australis</i> Aldrich, 1934, <i>Embiomyia</i>	104.85
<i>australis</i> (Macquart) 1855 (<i>Jurinia</i>), <i>Archytas</i>	104.48
<i>australis</i> Reinhard, 1935d, <i>Doryphorophaga</i>	104.132
<i>australis</i> Townsend, 1911b, <i>Eucelatoria</i>	104.133
<i>australis</i> (Townsend), 1911b (<i>Phasiopteryx</i>), <i>Euphasiopteryx</i> ...	104.21
<i>australis</i> (Townsend), 1928c (<i>Spanipalpus</i>), <i>Deopalpus</i>	104.76
<i>australis</i> Townsend, 1916c, <i>Myiophasia</i>	104.27
<i>australis</i> Townsend, 1929, <i>Leskiomima</i>	104.117
<i>austrina</i> (Coquillett), 1920b (<i>Sturmia</i>), <i>Drino</i>	104.188
<i>autographae</i> (Sellers), 1943 (<i>Zenillia</i>), <i>Eusisyropa</i>	104.203
<i>Austrolydella</i> Townsend, 1919a	104.132
<i>Austrophytomomyia</i> Blanchard, 1962	104.165
<i>Austrostaurochaeta</i> Townsend, 1931a	104.172
<i>Avibrissosturmia</i> Townsend, 1927a	104.194
<i>avida</i> Townsend, 1927a, <i>Avibrissosturmia</i>	104.194
<i>ayerzai</i> Blanchard, 1943, <i>Voria</i>	104.93
<i>Azygobothria</i> Townsend, 1911b	104.187
<i>Azulihoughia</i> Townsend, 1934	104.202
<i>azurea</i> (Curran), 1934d (<i>Zuanalia</i>), <i>Daetaleus</i>	104.25
<i>Baculocaptus</i> Cortés, 1968	104.123
<i>badiiventris</i> Wulp, 1888 (<i>Jurinia</i>)	104.66
<i>bahamensis</i> Townsend, 1919b, <i>Acronaristopsis</i>	104.164
<i>bakeri</i> Aldrich, 1926e, <i>Cylindromyia</i>	104.15
<i>balloui</i> (Curran), 1935 (<i>Sturmia</i>), <i>Drino</i>	104.188
<i>balteata</i> (Wulp), 1890 (<i>Prospheysa</i>), <i>Euphorocera</i>	104.158
<i>barbata</i> Bigot, 1887, <i>Jurinia</i>	104.68
<i>barbata</i> (Bigot), 1889 (<i>Phorocera</i>), <i>Metadoria</i>	104.140


barbatula (Wulp), 1890 (Brachycoma), Lespesia	104.209
barbiellinii Bezzi, 1925, Euthera	104.18
barbiellinii Townsend, 1935d, Belvosiomimops	104.183
barbiellinii Townsend, 1935d, Melanoromintho	104.140
barrosi (Brèthes), 1919 (Camarona), Morphodexia	104.24
basalis Curran, 1934d, Anadiscalia	104.201
basalis (Giglio-Tos) 1893a (Thelairodes), Cholomyia	104.106
basalis (Walker), 1852 (Tachina), Belvosia	104.181
basalis Walker, 1852 (Dexia)	104.122
<i>basalis</i> Walker, 1849 (Tachina), Juriniopsis	104.58
basalis (Walker), 1837 (Tachina), Notomanes	104.99
basalis Wulp, 1890 (Degeeria)	104.152
basifulvus (Walker), 1849 (Tachina), Archytas	104.49
<i>basifulvus</i> authors, Archytas	104.48
basilaris (Wulp), 1890 (Didyma), Metadoria	104.140
bassleri Townsend, 1929, Bathydexia	104.37
<i>bassleri</i> Curran, 1939, Macromyia	104.43
Bathydexia Wulp, 1891	104.36
<i>Bathytheresia</i> Townsend, 1928	104.37
beatus Curran, 1947, Rhachioepalpus	104.70
beckeri Engel, 1920, Dejeniops	104.70
bella Curran, 1934b, Calodexia	104.154
bella Giglio-Tos, 1893, Belvosia	104.181
<i>Belvoisia</i> , error or emendation	104.181
Belvosia Robineau-Desvoidy, 1830	104.180
Belvosiella Curran, 1934	104.183
Belvosiini	104.180
<i>Belvosiomima</i> Townsend, 1915	104.181
Belvosiomimops Townsend, 1935d	104.183
<i>Belvosioipsis</i> Townsend, 1927a	104.181
bennetti Thompson, 1963a, Archytas	104.49
bequaerti Curran, 1934b, Calodexia	104.154
bequaerti (Curran), 1925d, (Ptilomyia), Ptilomyoides	104.121
bergi Brèthes, 1908 (Exorista)	104.215
Beskia Brauer & Bergenstamm, 1889	104.215
Beskiocephala Townsend, 1916h	104.75
Beskioleskia Townsend, 191a	104.116
bezzii Arnaud, 1963b, Cryptocladocera	104.155
Bezzimya Townsend, 1919g	104.112
bibens (Wiedemann), 1830 (Stomoxys), Eumyobia	104.116
Bibiomima Brauer & Bergenstamm, 1889	104.6
bicaudata Curran, 1947, Adejeania	104.46
bicincta Robineau-Desvoidy, 1830, Belvosia	104.181
bicincta Reinhard, 1934a, Ceratomyiella	104.114
bicincta Robineau-Desvoidy, 1830, Belvosia	104.181
bicolor (Bigot), 1876 (Trichopoda), Homogenia	104.7
bicolor Coquillett, 1902, Euthera	104.18
bicolor (Giglio-Tos) 1893 (Hystrisiphona), Jurinodexia	104.30
<i>bicolor</i> Jaennicke, 1867, Archytas	104.49
bicolor Macquart, 1851. Exopalpus	104.54
bicolor (Macquart), 1846 (Blepharipeza), Parachaeta	104.186
<i>bicolor</i> Reinhard, 1931a, Winthemia	104.197
bicolor Townsend, 1931a, Chiloclista	104.97
bicolor Townsend, 1912a, Ecuadorana	104.83
bicolor Townsend, 1912a, Vibrissomyia	104.80



<i>bicolor</i> Townsend, 1931a, <i>Xanthotheresia</i>	104.38
<i>bicolor</i> (Wiedemann), 1830 (<i>Tachina</i>), <i>Jurinella</i>	104.56
<i>bicolor</i> (Williston), 1886 (<i>Saundersia</i>), <i>Xanthoepalpus</i>	104.73
<i>Bicruciostrumia</i> Townsend, 1932b	104.187
<i>bicrucis</i> Townsend, 1932b, <i>Bicruciostrumia</i>	104.187
<i>biezankoi</i> (Blanchard) <i>in</i> Biezanko, 1961 (<i>Williston</i>), <i>Belvosia</i>	104.181
<i>bifasciata</i> (Fabricius), 1775 (<i>Musca</i>), <i>Belvosia</i>	104.181
<i>bifuscata</i> , error, <i>Belvosia</i>	104.181
<i>bigeminata</i> (Curran), 1927c (<i>Frontina</i>), <i>Lespesia</i>	104.209
<i>bigeminata</i> Curran, 1927a, (<i>Lydella</i>)	104.152
<i>bilimeckii</i> Brauer & Bergenstamm, of Tavares, 1962, <i>Ormia</i>	104.22
<i>bilimeckii</i> (Brauer & Bergenstamm), 1889 (<i>Phasiapteryx</i>), <i>Ormia</i>	104.22
<i>bilineata</i> Wulp, 1890 (<i>Masicera</i>)	104.152
<i>Binghamimyia</i> Townsend, 1919b	104.105
<i>Biohypostena</i> Townsend, 1927a	104.124
<i>biornatus</i> Curran, 1947, <i>Rhachioepalpus</i>	104.70
<i>bipartitus</i> Wulp, 1888 (<i>Saundersia</i>), <i>Xanthoepalpus</i>	104.73
<i>birabeni</i> (Blanchard), 1941 (<i>Eudejeniops</i>), <i>Eudejeania</i>	104.53
<i>bischofi</i> Townsend, 1913a, <i>Xanthoepalpus</i>	104.73
<i>Bischofimyia</i> Townsend, 1927a	104.60
<i>biserialis</i> (Schiner), 1868 (<i>Phorocera</i>), <i>Euphorocera</i>	104.157
<i>biseriata</i> (Wulp), 1890, (<i>Brachycoma</i>), <i>Masiphya</i>	104.198
<i>biseta</i> (Arnaud), 1963b (<i>Adercomyia</i>), <i>Pelecotheca</i>	104.156
<i>bistrigata</i> (Wulp), 1890 (<i>Masicera</i>), <i>Eumasicera</i>	104.162
<i>bivittata</i> Townsend, 1919a, <i>Myothyriopsis</i>	104.190
<i>blanchardi</i> Guimarães, 1971, <i>Peleteria</i>	104.44
<i>blandita</i> (Wulp), 1890, <i>Calolydella</i>	104.127
<i>blandoides</i> Thompson, 1963b, <i>Eusisyropa</i>	104.203
<i>blandus</i> Curran, 1947, <i>Rhachioepalpus</i>	104.70
<i>Blephariatacta</i> Townsend, 1913a	104.183
<i>Blepharipeza</i> Robineau-Desvoidy, 1830	104.185
<i>blepharipus</i> Brauer & Bergenstamm, 1891 (<i>Ctenophorocera</i>) ...	104.160
<i>Blepharopoda</i> Rondani, 1850	104.36
<i>Blondelia</i> Robineau-Desvoidy, 1830	104.126
<i>Blondeliini</i>	104.123
<i>Bolbochaeta</i> Bigot, 1885	104.165
<i>boliviana</i> Townsend, 1931a, <i>Alpinoplagia</i>	104.89
<i>boliviana</i> Townsend, 1931a, <i>Rhachosaundersia</i>	104.71
<i>bolivianus</i> Bischof, 1904, <i>Epalpus</i>	104.64
<i>Bolodoria</i> Townsend, 1927a	104.127
<i>Bolohoughia</i> Townsend, 1927a	104.206
<i>Bolomyia</i> Brauer & Bergenstamm, 1891	104.187
<i>Bombyliomyia</i> Brauer & Bergenstamm, 1889	104.34
<i>Bombyliojeania</i> Townsend, 1931a	104.53
<i>Bombyliopsis</i> Townsend, 1915b	104.40
<i>bonasus</i> Reinhard, 1964c, <i>Ergolabus</i>	104.90
<i>Bonnaniops</i> Townsend, 1935d	104.127
<i>Bonnatia</i> Robineau-Desvoidy, 1830	104.81
<i>Borgmeiermyia</i> Townsend, 1935a	104.165
<i>borgmeieri</i> Guimarães, 1964, <i>Itacuphocera</i>	104.78
<i>bosqi</i> (Blanchard), 1954 (<i>Neobelvosioptis</i>), <i>Belvosia</i>	104.181
<i>bosqi</i> Blanchard, 1937, <i>Celatoria</i>	104.128
<i>bosqi</i> Blanchard, 1944, <i>Dallusimyia</i>	104.6
<i>bosqi</i> (Blanchard), 1968c (<i>Trichopodopsis</i>), <i>Trichopoda</i> (<i>Galactomyia</i>)	104.10



<i>bosqi</i> Blanchard in [Marchionatto], 1940 (<i>Achaetoneura</i>), <i>Lespesia</i>	104.209
<i>botyvora</i> Robineau-Desvoidy, 1830 (<i>Phorocera</i>)	104.160
<i>Bourquinia</i> Blanchard, 1935	104.206
<i>Brachybelvosia</i> Townsend, 1927a	104.183
<i>Brachynephalia</i> Townsend, 1927a	104.173
<i>Brachymasicera</i> Townsend, 1911b	104.202
<i>brachyptera</i> (Thomson), 1869 (<i>Myobia</i>), <i>Ceracia</i>	104.123
<i>brasiliانا</i> Brauer & Bergenstamm, 1891 (<i>Myiomima</i>), <i>Chaetotheresia</i>	104.36
<i>brasiliانا</i> Brauer & Bergenstamm, <i>Masiphya</i>	104.194
<i>brasiliانا</i> Brauer & Bergenstamm, 1891 <i>Paralispe</i>	104.145
<i>brasiliانا</i> Brauer & Bergenstamm, of Coquillett, 1891, <i>Promasiphya</i>	104.200
<i>brasiliانا</i> Reinhard, 1934z, <i>Paradidyma</i>	104.115
<i>brasiliانا</i> Townsend, 1937a, <i>Acaulona</i>	104.6
<i>brasiliانا</i> (Townsend), 1927a (<i>Biohypostena</i>), <i>Actinotachina</i> ..	104.124
<i>brasiliانا</i> Townsend, 1929, <i>Alophorella</i>	104.11
<i>brasiliانا</i> Townsend, 1931a, <i>Blephariatacta</i>	104.183
<i>brasiliانا</i> Townsend, 1935d, <i>Borgmeiermyia</i>	104.166
<i>brasiliانا</i> Townsend, 1927a, <i>Catocypthera</i>	104.15
<i>brasiliانا</i> Townsend, 1937a, <i>Paraphoranthra</i>	104.13
<i>brasiliانا</i> Townsend, 1929, <i>Voria</i>	104.93
<i>brasilianum</i> (Townsend), 1938a (<i>Parapsalida</i>), <i>Leucostoma</i>	104.17
<i>brasiliensis</i> Guimarães, 1966a, <i>Manteomasiphya</i>	104.198
<i>brasiliensis</i> Moreira, 1915, (<i>Masicera</i>)	104.214
<i>brasiliensis</i> (Robineau-Desvoidy), 1830 (<i>Dejeania</i>), <i>Adejeania</i> ..	104.46
<i>brasiliensis</i> Robineau-Desvoidy, 1830, <i>Harrisia</i>	104.185
<i>brasiliensis</i> (Robineau-Desvoidy), 1830 (<i>Jurinia</i>), <i>Macrojurinia</i>	104.59
<i>brasiliensis</i> Robineau-Desvoidy, 1863 (<i>Illigeria</i>), <i>Melanophora</i> ...	104.4
<i>brasiliensis</i> Robineau-Desvoidy, 1830, <i>Olinda</i>	104.217
<i>brasiliensis</i> Schiner, 1868, <i>Atacta</i>	104.180
<i>brasiliensis</i> Schiner, 1868 (<i>Nemoraea</i>)	104.215
<i>brasiliensis</i> (Schiner), 1868 (<i>Redtenbacheria</i>), <i>Pseudoredtenbacheria</i>	104.147
<i>brasiliensis</i> (Townsend), 1927a (<i>Neacroglossa</i>), <i>Acroglossa</i>	104.173
<i>brasiliensis</i> (Townsend), 1929 (<i>Trichodejeania</i>), <i>Adejeania</i>	104.46
<i>brasiliensis</i> (Townsend), 1938c (<i>Alophoropsis</i>), <i>Alophorella</i>	104.11
<i>brasiliensis</i> (Townsend), 1934a, <i>Antistaseopsis</i>	104.205
<i>brasiliensis</i> Townsend, 1934a, <i>Antistaseopsis</i>	104.205
<i>brasiliensis</i> Townsend, 1927a, (<i>Pseudoarchytopsis</i>), <i>Archytas</i> ...	104.50
<i>brasiliensis</i> Townsend, 1927a, (<i>Belvosioptis</i>), <i>Belvosia</i>	104.183
<i>brasiliensis</i> Townsend, 1927a, <i>Brachybelvosia</i>	104.183
<i>brasiliensis</i> Townsend, 1927a, <i>Bracnephalia</i>	104.174
<i>brasiliensis</i> Townsend, 1929, <i>Cclatoria</i>	104.128
<i>brasiliensis</i> Townsend, 1927a, <i>Ceromasiopsis</i>	104.206
<i>brasiliensis</i> Townsend, 1929, <i>Hyalomyodes</i>	104.18
<i>brasiliensis</i> (Townsend), 1917 (<i>Ypophaeomyia</i>), <i>Lespesia</i>	104.209
<i>brasiliensis</i> Townsend, 1929, <i>Leskiopsis</i>	104.121
<i>brasiliensis</i> Townsend, 1927a, (<i>Euthelairodopsis</i>), <i>Neomintho</i> ...	104.156
<i>brasiliensis</i> Townsend, 1927a, <i>Ophirionopsis</i>	104.122
<i>brasiliensis</i> Townsend, 1927a, <i>Oxyepalpus</i>	104.69
<i>brasiliensis</i> Townsend, 1927a, <i>Oxynops</i>	104.144
<i>brasiliensis</i> Townsend, 1939d, <i>Parathercsia</i>	104.37




<i>brasiliensis</i> Townsend, 1927a, <i>Procarcelia</i>	104.195
<i>brasiliensis</i> Townsend, 1917b, <i>Polygaster</i>	104.94
<i>brasiliensis</i> Townsend, error, <i>Voria</i>	104.93
<i>brasiliensis</i> (Townsend), 1927a, (<i>Hemimasipoda</i>), <i>Winthemia</i> ..	104.196
<i>brasiliensis</i> Townsend, 1929, <i>Xanthomelanodes</i>	104.10
<i>brasiliensis</i> Townsend, 1917b, <i>Xanthomelanopsis</i>	104.11
<i>brasiliensis</i> Townsend, 1927a, <i>Zonoepalpus</i>	104.74
<i>braueri</i> Townsend, 1931a, <i>Chrysotachina</i>	104.81
<i>braueri</i> Williston, 1896, <i>Diaphoropeza</i>	104.114
<i>Brauerimyia</i> Townsend, 1908	104.114
<i>Braziliomyia</i> Thompson 1968	104.128
<i>brevicauda</i> Curran, 1934c, <i>Uramya</i>	104.104
<i>brevicornis</i> Aldrich, 1934, <i>Clastoneura</i>	104.171
<i>brevicornis</i> Brauer & Bergenstamm, 1891 (<i>Selenomyia</i>), <i>Da-</i> <i>syuromyia</i>	104.98
<i>brevicornis</i> Blanchard, 1955, <i>Haywardimyia</i>	104.168
<i>brevicornis</i> (Macquart), 1851 (<i>Prosenia</i>), <i>Hystriochodexia</i>	104.30
<i>brevifrons</i> Townsend, 1927a, <i>Caeniopsis</i>	104.206
<i>brevihirta</i> Curran, 1947, <i>Adejeania</i>	104.46
<i>brevipennis</i> (Schiner), 1868 (<i>Miltogramma</i>), <i>Telothyria</i>	104.121
<i>brevirostria</i> Curran, 1947, <i>Adejeania</i>	104.46
<i>brevis</i> (Wulp), 1890 (<i>Exorista</i>), <i>Carcelia</i>	104.204
<i>breviventris</i> Wiedemann, 1830 (<i>Tachina</i>)	104.215
<i>brownae</i> Curran, 1947, <i>Saundersiopsis</i>	104.71
<i>browni</i> Aldrich, 1934, <i>Planomyia</i>	104.169
<i>browni</i> Curran, 1947, <i>Adejeania</i>	104.46
<i>browni</i> Curran, 1941, <i>Eudejeania</i>	104.53
<i>browni</i> Curran, 1942, <i>Hystricia</i>	104.41
<i>browni</i> Curran, 1947, <i>Lindigia</i>	104.58
<i>bruchi</i> (Blanchard), 1941 (<i>Archytodejeania</i>), <i>Archytas</i>	104.49
<i>bruchi</i> (Blanchard), 1941 (<i>Parafabricia</i>), <i>Archytas</i>	104.49
<i>bruchi</i> (Blanchard), 1954 (<i>Belvosiomima</i>), <i>Belvosia</i>	104.181
<i>bruchi</i> Blanchard, 1966c, <i>Eutrichopodopsis</i>	104.7
<i>bruchi</i> Blanchard, 1943a, <i>Lydinolydella</i>	104.139
<i>brunisquammis</i> Robineau-Desvoidy, 1830 (<i>Elaphoria</i>)	104.160
<i>bruneipennis</i> Bischof, 1904, <i>Epalpus</i>	104.64
<i>buccatus</i> (Reinhard), 1934c (<i>Cuphocera</i>), <i>Deopalpus</i>	104.76
<i>bullocki</i> Aldrich, 1934, <i>Araucosimus</i>	104.173
<i>buscki</i> Townsend, 1919a, <i>Beskioleskia</i>	104.116
<i>buscki</i> Townsend, 1919a, <i>Bezzimyia</i>	104.112
<i>busckii</i> (Townsend), 1915m (<i>Argyreomyia</i>), <i>Leptostylum</i>	104.137
<i>caelibata</i> Cortés, 1944d, <i>Discotrichia</i>	104.98
<i>Caeniopsis</i> Townsend, 1927a	104.206
<i>Caeniosoma</i> Townsend, 1927a	104.127
<i>Caenisomopsis</i> Townsend, 1934a	104.127
<i>caerulea</i> Bigot, 1855c, <i>Trichodischia</i>	104.100
<i>caerulea</i> Bigot of Cortés, 1944, <i>Trichodischia</i>	104.100
<i>caerulea</i> Macquart, 1851, <i>Gnadochaeta</i>	104.25
<i>caeruleiventris</i> (Wulp), 1890 (<i>Exorista</i>), <i>Chrysoexorista</i>	104.207
<i>caeruleonigra</i> (Macquart), 1846 (<i>Jurinia</i>), <i>Jurinella</i>	104.56
<i>caeruleonigra</i> Wulp, 1891, <i>Camaronia</i>	104.36
<i>calcarata</i> (Wulp), 1890 (<i>Masicera</i>), <i>Eumasicera</i>	104.162
<i>calceata</i> (Wulp), 1891 (<i>Calodexia</i>), <i>Cholomyia</i>	104.106



californicus Townsend, 1908, Xanthomelanodes	104.10
caliginosa (Walker), 1852 (Tachina), Hystricia	104.41
Calohystricia Townsend, 1931	104.40
Calolydella Townsend, 1927a	104.127
callani Thompson, 1968, Calodexia	104.154
callanganus Bischof, 1904, Epalpus	104.64
<i>Callesthes</i> Aldrich, 1928b	104.106
<i>callida</i> Townsend, 1919a (Charapomyia), Trafoia	104.83
callitris Reinhard, 1964c, Jalapamyia	104.87
Callotroxis Aldrich, 1929b	104.28
callipygus (Bigot), 1857a (Jurinia), Chiloepalpus	104.60
Calocarcelia Townsend, 1927a	104.202
Calodexia Wulp, 1891	104.153
Calodexiini	104.153
calogaster (Bigot), 1889 (Microphthalma), Irengia	104.23
<i>caloptera</i> Bigot, 1889 (Tricoliga), Pseudeuantha	104.104
Calothelaira Townsend, 1927a	104.105
Calpodomyia Townsend, 1915m	104.127
calyptrata (Aldrich), 1934 (Phorocera)	104.152
calyptrata Williston, 1896, Didyma	104.131
calva Coquillett, 1902, Oestrophasia	104.20
Camaronia Wulp, 1891	104.36
Camposiana Townsend, 1915k	104.60
Campsodes Cortès, 1967a	104.166
Campylochetini	104.89
Camptophryno Townsend, 1927a	104.206
Canalia Curran, 1934d	104.128
canalia Aldrich, 1928a, Belvosia	104.181
candicans Wulp, 1890, Anisia	104.125
canescens (Macquart), 1846 (Dejeania), Eudejeania	104.53
canus (Wulp), 1888 (Saundersia), Epalpus	104.64
capitata (Townsend, 1927a (Alophorellopsis), Hyalomya	104.13
capitata Townsend, 1927a, Gymnodoria	104.135
capitis (Townsend), 1915 (Zygofrontina), Lespesia	104.209
carbonaria Giglio-Tos, 1893 (Chaetogena)	104.122
carbonarius (Wulp), 1890 (Phorocera), Myiopharus	104.141
Carcelia Robineau-Desvoidy, 1830	104.202
Carcelliini	104.201
Carceliocephala Townsend, 1934	104.203
Carceliodoria Townsend, 1928b	104.203
carcelioides Townsend, 1927a, Hemisturmia	104.194
caridei (Brèthes) in Massini & Brèthes, 1918 (Parexorista), Euphorocera	104.158
carinata Curran, 1926a, Lixinia	104.138
carinata (Townsend), 1927a (Melanocyptera), Cylindromyia	104.15
carinata (Townsend), 1927a (Tachinophytopsis), Eucelatoriopsis	104.134
carinata Townsend, 1919a, Machairomasicera	104.134
carinata (Wulp), 1890 (Telothyria), Schizocerophaga	104.148
carlosalbertoi (Lima), 1926 (Thelairoides), Actinochaeta	104.113
Carmodymyia Thompson, 1968	104.128
carnata Reinhard, 1953b, Peleteria	104.44
caroniensis Thompson, 1963a, Archytas	104.49
carreirai Tavares, 1965b, Ormia	104.21
casanuevai Cortés, 1945, Phorocera	104.161
carrerai Guimarães, 1961b, Archytas	104.49

carrerei Guimarães, 1964, Itacuphocera	104.78
Casahüiria Townsend, 1919a	104.206
castanifrons (Bigot), 1889 (Ceromasia), Prospherysa	104.213
Catajurinia Townsend, 1927a	104.75
Catalinovoria Townsend, 1926	104.90
catamarcensis (Blanchard), 1954 (Belvosiomima), Belvosia ...	104.182
Cataphorinia Townsend, 1927a	104.128
Cataphoriniopsis Thompson, 1968	104.128
Catemophrys Townsend, 1908	104.114
Catocyptera Townsend, 1927a	104.15
caudata Curran, 1934b, Calodexia	104.154
caudata (Schiner), 1868 (Aporia), Uraporia	104.105
caudata Townsend, 1939c (Maracajuia), Neophyllophila	104.107
caudata (Townsend), 1927a (Urophylophila), Neophyllophila ..	104.107
causeyi Tavares, 1964, Ormiophasia	104.22
cauta Townsend, 1926, Catalinivoria	104.90
cayensis Townsend, 1914, Saundersiops	104.71
Celatoria Coquillett, 1890	104.128
Celatoriopsis Blanchard, 1963	104.129
chaetosus Townsend, 1928b, Chaetoxynops	104.129
Cenosoma Wulp, 1890	104.19
Ceracia Rondani, 1865	104.123
Ceratometopa Townsend, 1931a	104.23
Ceratomyiella Townsend, 1891	104.114
ceres (Curran), 1932 (Bucentes), Siphona	104.170
Ceromasiopsis Townsend, 1927a	104.206
Ceromya Robineau-Desvoidy, 1830	104.166
Cerotachina Arnaud, 1963b	104.155
cessatrix Walker, 1860 (Lydella)	104.215
chachapoyana Townsend 1929, Italydella	104.137
chacoensis Blanchard in Denier, 1939 (Tinanemorilla), Nemorilla	104.195
Chaeteprosopa Macquart, 1891	104.183
Chaetoandinomylia Vimmer & Soukup, 1940	104.75
Chaetocalirrhoe Townsend, 1935d	104.28
Chaetocnephalla Townsend, 1915e	104.174
Chaetocraniopsis Townsend, 1915e	104.174
Chaetodemoticus Brauer & Bergenstamm, 1891	104.87
Chaetodoria Townsend, 1927a	104.129
Chaetogaedia Brauer & Bergenstamm, 1891	104.178
Chaetogaedina	104.178
Chaetoglossa Townsend, 1892a	104.179
Chaetogyne Brauer & Bergenstamm, 1889	104.36
Chaetolixophaga Blanchard, 1940	104.129
Chaetona Wulp, 1891	104.105
Chaetonalia Curran, 1934d	104.105
Chaetonodexodes Townsend, 1916g	104.129
Chaetopalmus Vimmer & Soukup, 1940	104.217
chaetophora Brauer & Bergenstamm, 1891, Rhinomacquartia ..	104.100
Chaetophorocera Townsend, 1912a	104.206
Chaetoplagia Coquillett, 1895d	104.90
Chaetoprocta Brauer & Bergenstamm, 1891	104.186
Chaetoprosopa, emend.	104.183
chaetosa Aldrich, 1934, Lypha	104.85
chaetosa (Blanchard), 1954 (Latreillimyia), Belvosia	104.182
chaetosa Thompson, 1963d, Masiphyoidea	104.149



<i>chaetosa</i> Townsend, 1935d, <i>Actinocrocota</i>	104.165
<i>chaetosa</i> (Townsend), 1927a, (<i>Minthomima</i>), <i>Arrhinactia</i>	104.94
<i>chaetosa</i> Townsend 1912a (<i>Salmacia</i>), <i>Gonia</i>	104.175
<i>chaetosa</i> Townsend, 1915m, <i>Hypochaetopsis</i>	104.89
<i>chaetosa</i> Townsend, 1929, <i>Iteuthelaira</i>	104.156
<i>chaetosa</i> Townsend, 1911b, <i>Oomasicera</i>	104.204
<i>chaetosula</i> Townsend, 1892, <i>Muscopteryx</i>	104.99
<i>Chaetosisyrops</i> Townsend, 1912a	104.203
<i>Chaetotheresia</i> Townscnd, 1931a	104.36
<i>Chaetoxynops</i> Townsend, 1928b	104.129
<i>chalcos</i> (Townsend), 1934 (<i>Chrysodoria</i>), <i>Chrysoexorista</i>	104.207
<i>Chaquimayoia</i> Townsend, 1927a	104.129
<i>Charapemyia</i> Townsend, 1919a	104.83
<i>charapemyoides</i> Townsend, 1931a, <i>Neobrachelia</i>	104.86
<i>Chaetophlepsis</i> Townsend, 1915m	104.89
<i>charapense</i> Townsend, 1927a, <i>Caenisoma</i>	104.127
<i>charapensis</i> Townsend, 1927a, <i>Epiprospherysa</i>	104.133
<i>charapensis</i> Townsend, 1927a, <i>Siphoactia</i>	104.170
<i>charapensis</i> Townsend, 1919, <i>Urodexodes</i>	104.151
<i>Charapemyia</i> Townsend, 1919a	104.83
<i>Charapozelia</i> Townsend, 1927a	104.37
<i>charlini</i> Cortés, 1968, <i>Incamyia</i>	104.136
<i>chihuahuensis</i> Townsend, 1892 (<i>Brachycoma</i>)	104.193
<i>chilensis</i> Aldrich, 1928b, <i>Incamyia</i>	104.136
<i>chilensis</i> Bigot, 1880, <i>Psecacera</i>	104.25
<i>chilensis</i> (Brèthes), 1920b (<i>Phorichaeta</i>), <i>Ateloglutus</i>	104.89
<i>chilensis</i> Cortés, 1950 (<i>Phorocera</i>)	104.214
<i>chilensis</i> Curran, 1928b, <i>Archytas</i>	104.52
<i>chilensis</i> Macquart, 1851, <i>Hyalomyia</i>	104.13
<i>chilensis</i> Macquart, 1843, <i>Gonia</i>	104.176
<i>chilensis</i> (Schiner), 1868 (<i>Demoticus</i>), <i>Chaetodemoticus</i>	104.87
<i>chilensis</i> Townsend, 1915e, <i>Chaetocraniopsis</i>	104.174
<i>chilensis</i> Townsend, 1928c (<i>Valpogonia</i>), <i>Chaetocraniops</i>	104.174
<i>chilensis</i> Townsend, 1915m, <i>Ectophasiopsis</i>	104.12
<i>chiliensis</i> Bigot, 1880, <i>Psecacera</i>	104.25
<i>chillanensis</i> Cortés, 1951, <i>Epalpodes</i>	104.63
<i>Chiloclista</i> Townsend, 1931a	104.97
<i>Chiloepalpus</i> Townsend, 1927a	104.60
<i>chlorescens</i> Townsend, 1935d, <i>Chrysohoughia</i>	104.207
<i>Chlorohystricia</i> Townsend, 1927a	104.40
<i>Cholomyia</i> Bigot, 1884a	104.105
<i>chosica</i> Townsend, 1915m, <i>Oomeigenia</i>	104.143
<i>Chromoepalpus</i> Townsend, 1914	104.61
<i>chrysiceps</i> (Robineau-Desvoidy), 1830 (<i>Jurinia</i>)	104.215
<i>chrysocephala</i> Bigot, 1889 (<i>Ceromasia</i>)	104.152
<i>chrysocephala</i> Walker, 1837 (<i>Tachina</i>)	104.214
<i>Chrysodoria</i> Townsend, 1934	104.207
<i>Chrysoerigone</i> Townsend, 1927a	104.81
<i>Chrysoexorista</i> Townsend, 1915m	104.206
<i>Chrysohoughia</i> Townsend, 1935d	104.207
<i>Chrysometopiops</i> Townsend, 1919h	104.207
<i>chrysophora</i> (Wiedemann), 1830 (<i>Tachina</i>), <i>Hypotachina</i>	104.39
<i>Chrysophryno</i> 1927a	104.207
<i>Chrysophryna</i> , error	104.207

Chrysophryxe Sellers, 1943	104.207
chrysoprocta (Wiedemann), 1830 (Tachina), Mystacedia	104.191
chrysopyga (Bigot), 1887 (Frontina), Triachora	104.184
Chrysoprospherysa Townsend, 1928b	104.129
Chrysosturmia Townsend, 1916h	104.188
Chrysotachina Brauer & Bergenstamm, 1889	104.81
chrysotelus Walker, 1852 (Tachina)	104.161
Chrysotryphera Townsend, 1935d	104.207
christenseni (Blanchard), 1966c (Trichopodopsis), Trichopoda, (Galactomyia)	104.10
<i>Crangonimomma</i> , misprint	104.76
cidaris Reinhard, 1953d, Stolatosa	104.149
ciliata Aldrich, 1928, Belvosia	104.182
ciliata (Fabricius), 1805 (Ocyptera), Trichopoda (Trichopoda)	104.9
ciliata (Macquart), 1848 (Phorocera), Leschenaultia	104.185
ciliata Townsend, 1935d, Sturmimasiphya	104.200
ciliata Wulp, 1890, Anisia	104.125
ciliatum (Townsend), 1919c (Pterinoptera), Tyreomma	104.38
ciliger Robineau-Desvoidy, 1830, Winthemia	104.196
cilipes Macquart, 1843 (Gymnostylia)	104.215
<i>cilipes</i> Robineau-Desvoidy, 1830, Leschenaultia	104.186
cincrescens (Bigot), 1889 (Echinomyia), Peleteria	104.44
cincta Giglio-Tos, 1893 (Chaetogena)	104.215
cincta Giglio-Tos, 1893, Myiscotiptera	104.31
cincta Townsend, 1927a, Chroesoactia	104.130
cincta Townsend, 1911b, Ophiosturmia	104.210
cincta Walker, 1852 (Tachina)	104.152
cineracea (Wulp), 1890 (Brachycoma), Drino	104.188
cineraria Wulp, 1890, Anisia	104.125
cinerea Brèthes, 1909, Anisia	104.125
cinerea (Coquillett), 1895d (Paraplagia), Athrycia	104.90
cinerea Townsend, 1919a, Meigenelloides	104.139
cinerea Townsend, 1919a, Neoerigone	104.82
cinerea Townsend, 1908, Trepophrys	104.213
cinerea (Wulp) 1890 (Phorocera), Euphorocera	104.158
cinerea Wulp, 1891, Microchaetina	104.28
cinereicollis (Wulp), 1891 (Thelairodes), Spathidexia	104.195
cinereus Townsend, 1914, Rhachioepalpus	104.70
cingulata (Fabricius), 1805 (Musca), Calocarcelia	104.202
cingulatus (Schiner), 1868 (Meigenia), Urodexodes	104.151
cingulipes Blanchard, 1966, Ptilodexia	104.33
<i>cingulipes</i> Blanchard in Crouzel, 1965, Ptilodexia	104.33
ciniscula Reinhard, 1968, Macromyia	104.42
circularis Curran, 1947, Jurinella	104.56
cirphis Curran, 1927a, Archytas	104.49
<i>cirphisae</i> Curran, in Swezey, 1926, Archytas	104.49
cirrata Robineau-Desvoidy, 1830 (Phorocera)	104.161
Cistogaster Latreille, 1829	104.5
<i>citrina</i> Bigot, 1889 (Viviania), Chaetona	104.105
citrinus Blanchard, Blanchard, 1941, Ochroepalpus	104.69
claripalpis Thompson, 1968, Dexodiopsis	104.131
claripalpis Thompson, 1968, Neoxynopsoidea	104.142
claripalpis Thompson, 1968, Pseudoprospherysa	104.147
claripalpis Thompson, 1968, Stenochaeta	104.149




claripalpis (Wulp), 1896 (Sarcophaga), Paratheresia	104.37
claripennis Macquart, 1846, Ebenia	104.109
claripennis (Macquart), 1840 (Phorocera), Euphorocera	104.158
claripennis Thompson, 1963, Santacruziopsis	104.164
Clairvillia Robineau-Desvoidy, 1830	104.17
Clastoneura Aldrich, 1934	104.171
clarus Townsend, 1915e, Pictoepalpus	104.70
clausa Bigot, 1868a, Hyadesimyia	104.216
clausa (Townsend), 1927a, (Lydellactia), Actinotachina	104.124
clausa Townsend, 1927a, Euthelyconychia	104.135
clausa Townsend, 1934, Morphodexia	104.24
Clausicella Rondani, 1856	104.166
Clausicellana Curran, 1927a	104.166
clavipalpis Thompson, 1966m, Lespesia	104.209
Clinogaster Wulp, 1892	104.15
Clinoneura Brauer & Bergenstamm, 1889	104.33
Clythopsis Townsend, 1927a	104.123
Clythoxynops Townsend, 1927a	104.129
Cnephalodes Townsend, 1911b	104.174
Cnephalodopsis Townsend, 1912a	104.174
coerulea Macquart	
coerulea Townsend, 1927a, Cyanogymnomma	104.176
collaris Townsend, 1919a, Lydellothelaira	104.138
colombiana Townsend, 1929, Cordyllerodexia	104.29
colombiensis Arnaud, 1951, Paradejeania	104.59
colombiensis Curran, 1947, Saundersiopsis	104.71
columbiae Brauer & Bergenstamm, 1889, Actinochaeta	104.113
Colurus Reinhard, 1953a	104.113
comata Wulp, 1890 (Telothyria)	104.152
Comatacta Coquillett, 1902c	104.120
commetans (Walker), 1860 (Eurigaster), Bolomyia	104.187
commixta (Wulp), 1890 (Didyma), Mesochaeta	104.140
communis Aldrich, 1934, Admontia	104.171
communis Thompson, 1963a, Winthemia	104.196
communis Townsend, 1927a, (Aphriosphyria), Peleteria	104.45
Comops Aldrich, 1934	104.75
comosa (Wulp), 1890 (Prospheysa), Eucelatoria	104.133
compacta Curran, 1926a (Linnaemyia)	104.83
compacta (Walker), 1852 (Tachina), Drino	104.188
compascua (Wulp), 1892 (Echinomyia), Peleteria	104.44
complanata (Vimmer & Soukup), 1940 (Chaetoandinomyia), Andinomyia	104.75
compressa (Wulp), 1890 (Degeeria), Celatoria	104.128
Compsilurini	104.123
comta (Fallén), 1810 (Tachina), Bonnetia	104.81
Comyops Wulp, 1891	104.109
Comyopsis Townsend, 1919b	104.109
Conactia Townsend, 1927a	104.109
Conactiodoria Townsend, 1934	104.130
concina (Wulp), 1890 (Hyspostena), Calolydella	104.127
conclusa Curran, 1947, Adejeania	104.46
concolor (Townsend), 1929 (Patelloopsis), Patelloa	104.211
condor Curran, 1942, Hystricia	104.41

confluens Townsend, 1914, Saundersiops	104.71
confundens Townsend, 1927a, (Clythopsis), Ccracia	104.123
confusa (Aldrich), 1925c (Masiphya), Promasiphya	104.200
congerens Wulp, 1890 Anisia	104.125
congrua Wulp, 1891, Chaetona	104.105
congruens Wulp, 1892 (Jurinia), Archytas	104.50
conica Townsend, 1927a, Chaetodoria	104.129
conica Townsend, 1935d, Chysotryphera	104.207
conicus Townsend, 1927a, Anoxynops	104.125
conjuncta (Wulp), 1891 (Rhynchodexia), Ptilodexia	104.33
connectans (Townsend), 1912a (Tropidopsis), Macromya	104.42
connexus (Wulp), 1890 (Tclothyrina), Myiopharus	104.141
connota Curran, 1947, Jurinella	104.56
consanguineus (Wulp), 1892 (Saundersia), Epalpus	104.64
consistens (Curran), 1927 (Erycia), Sturmia	104.192
consobrina (Wulp), 1890 (Exorista), Winthemia	104.196
conspersa Wulp, 1890, Anisia	104.125
constans (Walker), 1849 (Tachina), Parepalpus	104.69
constrictans (Wulp), 1891 (Hystrichodexia), Ptilodexia	104.33
contermina (Walker), 1852 (Tachina), Belvosia	104.182
contigua (Wulp), 1890, (Prospheyrina) Paralipse	104.145
contiguus (Reinhard), 1934c (Cuphocera), Deopalpus	104.76
continua Curran, 1934c, Calodexia	104.154
contrarius (Walker), 1849 (Jurinia), Epalpus	104.64
convexa Curran, 1925c, Peleteria	104.44
convexa Walker, 1852 (Dexia)	104.122
convexinervis Wulp, 1892, Trichophora	104.72
Copecrypta Townsend, 1908	104.75
copulata (Wiedemann), 1830 (Tachina), Hystriciopsis	104.42
coquilletti Vimmer & Soukup, 1920, Chaetopalpus	104.217
cora (Bigot), 1889 (Degeeria), Eucelatoria	104.133
cora (Bigot), 1888a, (Echinomyia), Peleteria	104.44
Coracomia Aldrich, 1934	104.166
coracella (Aldrich), 1934 (Aphelogaster), Aldrichiopa	104.165
coracina (Wulp), 1891 (Hystrichodexia), Tropidodexia	104.35
corax Aldrich, 1934, Lypha	104.85
Cordilerodexia Townsend, 1927a	104.29
cordubensis (Blanchard), 1934b (Gymnocuphocera), Gymnom- mopsis	104.78
Cordylidexia Giglio-Tos, 1893	104.111
Cordyligaster Macquart, 1843	104.111
<i>Cordyligaster</i> , authors	104.111
cornifera Townsend, 1931a, Ceratometopa	104.23
cornuale Reinhard, 1947, Crocinosoma	104.166
cornuta Aldrich, 1934, Actia	104.165
cornuta Brauer & Bergestamm, 1889, Beskia	104.15
cornuta Townsend, 1934, Actia	104.165
cornuta Townsend, 1919, Casahuiria	104.206
cornutum Townsend, 1914, Corpulentosoma	104.61
Corozalia Curran, 1934	104.130
corpulenta (Townsend), 1927a (Tachinosoma), Jurinella	104.57
corpulenta (Wiedemann), 1830 (Tachina), Adejeania	104.46
Corpulentoepalpus Townsend, 1927a	104.61
Corpulentosoma Townsend, 1914	104.61



corpulentus Townsend, 1914, Eulasiopalpus	104.54
corpulentus Townsend, 1914 Epalpellus	104.63
corpulentus Townsend, 1927a, Melanoepaulpus	104.68
cortesi Catalgirona, 1966, Opsophagus	104.92
costalimai Tavares, 1964, Ormiophasia	104.22
costalis Townsend, 1917b, Cyanopsis	104.76
costalis Townsend, 1915m, Epiphanocera	104.133
costalis Townsend, 1927a, Melanorlopteryx	104.140
costalis Townsend, 1927a, Oedemamedina	104.142
costalis Walker, 1852 (Musca)	104.152
costalis Wulp, 1890 (Telothyría)	104.152
costata Wulp, 1888, Acaulona	104.6
cothurnata (Wiedemann), 1830 (Stomoxys), Stomatodexia	104.119
Cowania Reinhard, 1952a	104.90
cranei Thompson, 1966, Heliconiophaga	104.189
crassa (Wiedemann), 1830 (Musca), Chaetotheresia	104.36
crassicornis Aldrich, 1934, Coracomylia	104.166
crassicornis (Fabricius), 1774 (Musca), Gonia	104.175
crassisetia Cortés, 1945, Stuardomyia	104.26
crassisetia Reinhard, 1934a, Paradidyma	104.115
crebra (Wulp), 1890 (Prospheerisa), Chaetogaedia	104.178
crescentis (Reinhard), 1944 (Sturmia), Drino	104.188
crescentis (Townsend), 1916g (Atactomima), Atacta	104.180
crocata Reinhard, 1968, Macromya	104.42
<i>Crocota</i> Meigen, 1800	104.170
<i>Crocotini</i>	104.164
Crocinosoma Reinhard, 1947	104.166
Croesoactia Townsend, 1927a	104.130
croesus Townsend, 1928b, Chrysoprospherysa	104.129
crossi (Blanchard), 1942 (Pseudoprospherysa), Jurinella	104.57
crouzeli Blanchard, 1966b, Microphthalma	104.24
crouzeli Blanchard, 1966b, Plexiodexilla	104.27
cruciata Townsend, 1912a, Andimomyia	104.75
cruciata Townsend, 1914, Uruhuasia	104.73
cruciatus Townsend, 1914, Saundersiops	104.71
crucius (Giglio-Tos), 1893 (Nemochaeta), Archytas	104.49
crudelis (Wiedemann), 1830 (Tachina), Angiorrhina	104.23
cruenta Giglio-Tos, 1893, Chaetona	104.105
cruralis Giglio-Tos, 1893 (Degeeria)	104.152
cruzi Tavares, 1964, Ormiophasia	104.22
crypta Townsend, 1934a, Carceliocephala	104.203
Cryptocladocera Bezzi, 1923	104.203
Cryptomeigenia Brauer & Bergenstamm, 1891	104.130
Cryptomeigenioidea Thompson, 1968	104.130
Cryptopalpus Rondani, 1850	104.61
cubaecola (Jaennicke), 1867 (Tachina), Drino	104.188
<i>Cubaemyia</i> Townsend, 1931a	104.188
Cubaemyiopsis Thompson, 1963b	104.188
cubana Townsend, 1919b, Argyrochaetona	104.161
cubensis (Curran), 1927 (Distichona), Olenochaeta	104.177
cubensis Macquart, 1848 (Masicera)	104.214
cubensis Malloch, 1932, Echinodexia	104.29
cuestae Engel, 1920, Engelomyia	104.41



<i>cumutoensis</i> Thompson, 1968, <i>Euphorocera</i>	104.158
<i>Cuparimya</i> Townsend, 1934	104.130
<i>Cuphocera</i> , authors	104.76
<i>Cuphocerini</i>	104.74
<i>Cuphoceropsis</i> Townsend, 1935d	104.43
<i>Cuphoceromyia</i> Blanchard, 1943b	104.43
<i>cuprea</i> Townsend, 1927a, <i>Actinodoria</i>	104.124
<i>cupreiventris</i> Wulp, 1890, <i>Telothyria</i>	104.121
<i>cuprescens</i> (Walker) 1857 (<i>Eurigaster</i>), <i>Bolomyia</i>	104.187
<i>curepei</i> Thompson, 1966, <i>Drino</i>	104.188
<i>curepei</i> Thompson, 1964, <i>Pseudochaeta</i>	104.163
<i>curialis</i> Reinhard, 1958a, <i>Clairvillia</i>	104.17
<i>curta</i> Townsend, 1934a, <i>Macromasiphya</i>	104.199
<i>curta</i> (Wulp), 1890 (<i>Masicera</i>), <i>Drino</i>	104.188
<i>curupeinsis</i> Thompson, 1968, <i>Paranoxylops</i>	104.145
<i>curva</i> (Wulp), 1890 (<i>Telothyria</i>), <i>Spathimeigenia</i>	104.148
<i>curvipes</i> Aldrich, 1934, <i>Hyalomya</i>	104.13
<i>curvirostris</i> (Bigot), 1889 (<i>Prosenia</i>), <i>Prosenoides</i>	104.32
<i>curvirostris</i> , authors, <i>Prosenoides</i>	104.32
<i>cuspidata</i> Townsend, 1939b, <i>Shannoniella</i>	104.112
<i>cussirilis</i> Reinhard, 1953, <i>Euhysticia</i>	104.41
<i>cuyana</i> (Blanchard), 1963 (<i>Neowinthemia</i>), <i>Winthemia</i>	104.196
<i>cuzcana</i> (Townsend), 1912a (Perua), <i>Microphthalma</i>	104.24
<i>cuzcensis</i> Townsend, 1912a, <i>Incamiya</i>	104.136
<i>cyanea</i> Giglio-Tos, 1893, <i>Scotiptera</i>	104.34
<i>cyanea</i> Macquart, 1851, <i>Chaeteprosopa</i>	104.184
<i>cyaneiventris</i> (Macquart), 1846 (<i>Blepharipeza</i>), <i>Chaeteprosopa</i>	104.184
<i>cyaneiventris</i> (Wulp), 1884 (<i>Hysticia</i>), <i>Chlorohysticia</i>	104.40
<i>Cyanogymnomma</i> Townsend, 1927a	104.76
<i>Cyanopsis</i> Townsend, 1917b	104.76
<i>cylindrata</i> (Wulp), 1892 (<i>Phorocera</i>), <i>Eupelcotheca</i>	104.156
<i>cylindrica</i> Townsend, 1927a, <i>Arrhinactia</i>	104.94
<i>cylindrica</i> (Townsend), 1891a (<i>Minthohoughia</i>), <i>Spathidexla</i> ...	104.95
<i>cylindriventris</i> Wulp, 1890 (<i>Hypostena</i>)	104.152
<i>Cylindromasicera</i> Townsend, 1915d	104.162
<i>Cylindromyia</i> Meigen, 1803	104.15
<i>Cylindromyiini</i>	104.14
<i>Cylindrophasia</i> Townsend, 1916g	104.6
<i>Cyosoprocta</i> Reinhard, 1952a	104.184
<i>Cyphocera</i> , error	104.76
<i>Cyrtosoma</i> Brauer & Bergenstamm, 1893	104.37
 <i>Dactilodidyma</i> Townsend, 1927a	
<i>daemon</i> (Wiedemann), 1830 (<i>Tachina</i>), <i>Archytas</i>	104.49
<i>Daetaleus</i> Aldrich, 1928	104.25
<i>Dallasimya</i> Blanchard, 1944	104.6
<i>dammipus</i> (Walker), 1849 (<i>Tachina</i>), <i>Archytas</i>	104.49
<i>dampfii</i> Aldrich, 1927b, <i>Oedematocera</i>	104.172
<i>danai</i> (Townsend), 1940b, (<i>Zygofrontina</i>), <i>Lespesia</i>	104.209
<i>darwini</i> Aldrich, 1934, <i>Peliceps</i>	104.99
<i>Dasyuromyia</i> Bigot, 1885c	104.97
<i>datanae</i> (Townsend), 1892e (<i>Exorista</i>), <i>Winthemia</i>	104.196
<i>dcaurata</i> Blanchard, 1935, <i>Bourquinia</i>	104.206
<i>debilis</i> Aldrich, 1934, <i>Admontia</i>	104.71




debithrix (Walker), 1860 (Jurinia), Jurinella	104.57
decisa Walker, 1852, Trichopoda (Trichopoda)	104.9
decisus (Walker), 1849 (Tachina), Pararchytas	104.54
decora Aldrich, 1926e, Cylindromyia	104.15
decoratus (Townsend), 1931a (Saundersia), Echinopyrrhosioops	104.62
decoratus (Rondani), 1850a (Cuphocera), Deopalpus	104.76
decumana Reinhard, 1958a, Hemyda	104.16
degeerioides (Wulp), 1890 (Myothyria)	104.152
dellephilae (Osten-Sacken), 1887 (Tachina), Winthemia	104.196
Dejeania Robineau-Desvoidy, 1830	104.45
Dejeaniini	104.45
dejecta (Wulp), 1890 (Masicera), Doryphorophaga	104.132
Dejeniopalpus Townsend, 1916f	104.116
Dejaniops Townsend, 1913c	104.52
Dcloblepharis Aldrich, 1934	104.87
delta Townsend, 1931a, Deltoceromyia	104.167
delta Townsend, 1914, Uruhuasia	104.73
Deltoceromyia Townsend, 1931a	104.167
delusa Reinhard, 1964c, Mastigiomyia	104.46
demissum Reinhard, 1958c, Mochlosoma	104.30
demoticoideus (Brauer & Bergenstamm), 1889 (Arthrochaeta), Exopalpus	104.55
dentata (Coquillett), 1895b (Acemyia), Ceracia	104.123
denudata Townsend, 1931a, Pseudoxanthosona	104.70
denudatus Bischof, 1904, Epalpus	104.64
Deopalpus Townsend, 1908	104.76
deplanata Wulp, 1890 (Hypostena)	104.152
depleta (Wiedemann), 1830 (Tachina), Euphasiopteryx	104.21
depressa Robineau-Desvoidy, 1830, Macromyia	104.42
derelicta Reinhard, 1934, Paradidyma	104.115
desita Walker, 1861 (Eurigaster)	104.215
desvoidy Townsend, 1931a, Sophia	104.112
devia Reinhard, 1929a, Ginglymyia	104.167
devia Townsend, 1917b, Copecrypta	104.75
Dexiinae	104.93
Dexiini	104.96
Dexiini of authors	104.22
Dexillini	104.22
dexina Townsend, 1914, Neaphria	104.85
Dexiophana Brauer & Bergenstamm, 1891	104.213
Dexodimyia Thompson, 1968	104.131
Dexodioidea Thompson, 1968	104.131
Dexodiopsis Thompson, 1968	104.131
Dexodomintho Townsend, 1935d	104.131
destrella Reinhard, 1935d, Lasineura	104.168
diabrotica (Schiner), 1871 (Tachina), Celatoria	104.128
diacrita Reinhard, 1954, Prosenoides	104.32
diadema (Wiedemann), 1830 (Dexia), Myiobiopsis	104.118
diaphana (Brauer & Bergenstamm), 1891 (Paragymnomma), Diaphanomyia	104.62
diaphana (Fabricius), 1805 (Tachina), Archytas	104.49
diaphanus (Fabricius), 1805 (Ocyptera), Xanthomelanodes	104.10
diaphanoides Curran, 1925e, Gymnomma	104.67
Diaphanomyia Townsend, 1917b	104.62
Diaphoropeza Townsend, 1908	104.114



<i>diatraea</i> Brèthes, 1927a, (Sarcophaga), Paratheresia	104.37
<i>diatraea</i> Townsend, 1914b, Palpozenillia	104.211
<i>diatraea</i> (Townsend), 1915b (Euzenilliopsis), Lixophaga	104.138
<i>Diaghaia</i> Perty, 1833	104.101
<i>dicax</i> Giglio-Tos, 1893 (Degeeria)	104.152
<i>dichroma</i> (Wulp), 1888 (Jurinia), Protodejeania	104.59
<i>dicta</i> (Giglio-Tos), 1843 (Plagia), Voria	104.93
<i>Didyma</i> Wulp, 1890	104.131
<i>Didymops</i> Townsend, 1927a	104.132
<i>dieloceri</i> (Townsend), 1942, (Hylotomomyia) <i>Spathimeigenia</i> ...	104.148
<i>differens</i> (Wulp), 1890 (Trixa), <i>Microphthalma</i>	104.24
<i>difficilis</i> Aldrich, 1934, <i>Myiophasia</i>	104.27
<i>dileta</i> (Wiedemann), 1830 (Musca), <i>Medinophyto</i>	104.106
<i>diluta</i> Wulp, 1890, <i>Siphona</i>	104.170
<i>Dimasicera</i> Townsend, 1915d	104.163
<i>dimidiata</i> Townsend, 1937a, <i>Paraphasiana</i>	104.13
<i>dimmocki</i> (Aldrich), 1932a (Anetia), <i>Eucelatoriopsis</i>	104.134
<i>Diotrephes</i> Reinhard, 1964c	104.40
<i>dirphiae</i> Townsend, 1928, <i>Podosturmia</i>	104.191
<i>discalis</i> Aldrich, 1934, <i>Admontia</i>	104.171
<i>discalis</i> (Brèthes), 1909 (Saundersia), <i>Cryptopalpus</i>	104.61
<i>discalis</i> Thompson, 1968, <i>Dexodomyia</i>	104.131
<i>discalis</i> Townsend, 1927a, <i>Empheremyiops</i>	104.63
<i>discalis</i> (Townsend), 1927a (Myodoria), <i>Lydinolydella</i>	104.138
<i>discalis</i> Townsend, 1915m, <i>Neominthopsis</i>	104.142
<i>discalis</i> (Townsend), 1919a, <i>Opsozella</i>	104.102
<i>discalis</i> (Townsend), 1915m, (Paramyocera), <i>Rhamphimina</i> ...	104.34
<i>discalis</i> (Townsend), 1914 (Epalpus), <i>Saundersiops</i>	104.71
<i>Dischotrichia</i> Cortés, 1944d	104.98
<i>discolor</i> (Wulp), 1891 (Rhynchodexia), <i>Ptilodexia</i>	104.33
<i>discors</i> (Wulp), 1892 (Gymnomma), <i>Parepalpus</i>	104.69
<i>discrepans</i> (Wulp), 1890 (Baumhaueria), <i>Distichona</i>	104.174
<i>disgrega</i> Wulp, 1890 (Telothyria)	104.214
<i>disjuncta</i> (Wiedemann), 1824 (Tachina), <i>Microphthalma</i>	104.24
<i>dispar</i> Wulp, 1890 (Echinomyia), <i>Archytas</i>	104.50
<i>disparata</i> Brauer & Bergenstamm, 1891, <i>Hypotachina</i>	104.39
<i>disputans</i> Walker, 1860 (Masicera)	104.214
<i>dissepta</i> (Wulp), 1890 (Telothyria), <i>Eucelatoriopsis</i>	104.134
<i>dissimilis</i> (Wulp), 1888 (Nemochaeta), <i>Archytas</i>	104.49
<i>dissimiloides</i> Thompson, 1963a, <i>Archytas</i>	104.49
<i>dissitus</i> (Reinhard), 1967a (Eurylochus), <i>Eurythemymia</i>	104.162
<i>Distichona</i> Wulp, 1890	104.174
<i>distincta</i> (Townsend), 1929 (Anazygosturmia), <i>Drino</i>	104.188
<i>distincta</i> , authors, <i>Drino</i>	104.189
<i>distincta</i> error, Aldrich, 1926a, <i>Microphthalma</i>	104.24
<i>divaricata</i> Townsend, 1935d, <i>Dolichodinera</i>	104.29
<i>divergens</i> Aldrich, 1934, <i>Ganopleuron</i>	104.98
<i>diversus</i> (Walker), 1849 (Tachina), <i>Cryptopalpus</i>	104.61
<i>dives</i> Curran, 1934b, <i>Calodexia</i>	104.154
<i>divisus</i> (Walker), 1852 (Tachina), <i>Archytas</i>	104.49
<i>Dolichostoma</i> Townsend, 1912a	104.77
<i>Dolichocnephala</i> Townsend, 1915d	104.174
<i>Dolichocodia</i> Townsend, 1935d	104.29
<i>Dolichocryptera</i> Townsend, 1921a	104.15


Dolichodinera Townsend, 1935d	104.29
Dolichogonia Townsend, 1915d	104.175
Dolichopalpellus Townsend, 1927a	104.167
dolichotarsis (Curran), 1934c (Phillophilopsis), Microchaetona	104.107
dominica Curran, 1927c, Cistogaster	104.5
dominica (Townsend), 1919a (Ormia), Euphasiopsteryx	104.21
Doriella Townsend, 1931a	104.208
dorsalis (Walker), 1852 (Dexia), Trichodura	104.103
dorsalis (Wiedemann), 1830 (Ocyptera), Cyldromyia	104.15
dorsalis (Wulp), 1888 (Hystiricia), Macromyia	104.42
dorsalis (Wulp), 1892 (Xanthomelana), Xanthomelanodes	104.10
dorsopunctata (Macquart), 1843 (Micropalpus), Trichosaundersia	104.73
doryphorae (Riley), 1869 (Lydella), Doryphorophaga	104.132
Doryphorophaga Townsend, 1912d	104.132
dotata (Walker), 1849 (Ocyptera), Cyldromyia	104.15
downsi Curran, 1947, Protodejeania	104.59
downsi Reinhard, 1953a, Colurus	104.185
dreisbachi Reinhard, 1956a, Eudexia	104.29
Drepanoglossa Townsend, 1891	104.167
Drino Robineau-Desvoidy, 1863	104.188
dubia (Bigot), 1969a (Chaetolyga), Drino	104.188
dubia Bigot, 1885a, Rhamphiniina	104.34
dubia Brauer, 1887 (Melanota)	104.215
dubia (Townsend), 1927a (Dactylodidyma), Paradoria	104.144
dubiosa Thompson, 1968, Santacruzia	104.148
dubiosa Thompson, 1963c, Winthemia	104.196
dubius (Giglio-Tos), 1893 (Nemochaeta) Archytas	104.49
dubius (Walker), 1893 (Nemochaeta), Archytas	104.49
ducalis Reinhard, 1958, Admontia	104.171
ducke Guimaraes, 1961b, Archytas	104.49
dumalis Reinhard, 1964, Phobetromyia	104.43
Dumerillia Robineau-Desvoidy, 1830	104.53
dunningii (Coquillett), 1895e (Tryptocera), Spathimeigenia	104.95
duplare Reinhard, 1958c, Mochlosoma	104.30
durvillei Macquart, 1843, Trichoprosopus	104.25
dux Curran, 1928b, Archytas	104.49
dysderci Townsend, 1940b, Paraphasiana	104.13
dysderci (Townsend), 1938c (Euphorantha), Alohorella	104.11
dysderci (Townsend), 1937a (Eumogenia), Homogenia	104.7
Ebenia Macquart, 1846	104.109
Ebeniini	104.109
Ecatocypterops Townsend, 1935d	104.15
Ecatocypteropsis Townsend, 1915m	104.12
echinata Wulp, 1891, Hystirichodexia	104.30
Echinodexia Brauer & Bergenstamm, 1893	104.29
Echinomasicera Townsend, 1915m	104.185
Echinomyiini	104.43
Echinopyrrhosia Townsend, 1914	104.62
Echinopyrrhosiops Townsend, 1931a	104.62
Echinotachina Townsend, 1931e	104.45
ecitonis (Townsend), 1896 (Hyalomyia), Androeurops	104.12
Eclausicella, misprint	104.134
Ectophasiopsis Townsend, 1915m	104.12




Ecuadorana Townsend, 1912a	104.
edessae (Townsend), 1942b (Xenophyxis), Neobrachelia	104.86
edwardsi Aldrich, 1929b, Callotroxis	104.28
Edwinyia Aldrich, 1930a	104.63
Egameigenia Townsend, 1927a	104.132
egensis Townsend, 1927a, Chrysophryno	104.207
egle Curran, 1947, Jurinella	104.57
Eggonia Brauer & Bergenstamm, 1893	104.216
egregia Wulp, 1890, Polygaster	104.94
elata Brauer & Bergenstamm, 1889, Myiomintho	104.141
elactilis Reinhard, 1964b, Helioprosopa	104.78
Elfia Robineau-Desvoidy, 1850	104.167
elegans (Bigot), 1857b (Tachina), Winthemia	104.196
elegans (Giglio-Tos), 1893 (Aporia), Paraporia	104.104
elegans Reinhard, 1964a, Minthodexiopsis	104.107
elegans Townsend, 1917c, Erythroargyrops	104.133
elegans (Townsend), 1927a (Microtrichommodes), Exopalpus ...	104.55
elegans Townsend, 1919a, Microaporia	104.96
elegans Townsend, 1929, Parachaetolyga	104.195
elegans Townsend, 1915m, Rhombothyriops	104.148
elegans (Wulp), 1890 (Hypostena), Cryptomeigenia	104.130
elegantula Arnaud, 1963b, Cerotachina	104.155
elisae Cortés, 1945 (Phorocera)	104.161
elodioides Townsend, 1935d, Succingulodes	104.140
elongata Rondani 1948, (Phorocera)	104.215
elongata Townsend, 1919a, Ollachea	104.143
elongata Townsend, 1911b, Phasiatacta	104.177
elongata (Wulp), 1890 (Exorista), Tinalydella	104.151
elquiensis Cortés, 1951b, Acroceromia	104.74
elusa Aldrich, 1926b, Belvosia	104.182
Eleuthromyia Reinhard, 1964c	104.82
elyowaldi Guimarães, 1963c, Euempheremyia	104.66
emarginata Townsend, 1915k, Camposiana	104.60
emarginatus Townsend, 1914, Trichoepalpus	104.72
embiaphaga Arnaud, 1963c, Perumyia	104.177
Embiomyia Aldrich, 1934	104.85
Empheremyia Bischof, 1904	104.62
Empheremyiops Townsend, 1927a	104.63
emulatus Reinhard, 1962b, Mimologus	104.190
Enchomyia Aldrich, 1934	104.175
Engelomyia Townsend, 1913b	104.41
Epaulophasia Townsend, 1934	104.12
Epalpellus Townsend, 1914	104.63
Epalpodes Townsend, 1912a	104.63
Epalpus Rondani 1850	104.63
Epicoronimyia Blanchard, 1940	104.167
Epicuphocera Townsend, 1927a	104.77
Epidexia Townsend, 1912b	104.132
Epidolichostoma Townsend, 1927a	104.77
Epigrimyia, authors	104.166
epilachnae (Aldrich), 1923 (Paradexodes), Lydinolydella	104.126
epileuca (Walker), 1849 (Tachina), Jurinella	104.57
Epimeigenia Townsend, 1931a	104.132
Epicoronimyia Blanchard, 1940	104.167
Epicuphocera Blanchard	104.



Epiphanocera Townsend, 1915m	104.133
<i>Epiphyllaphila</i> Townsend, 1927a	104.107
Epiplagiops Blanchard, 1943d	104.158
Epiprospherysa Townsend, 1927a	104.133
equatorialis (Macquart), 1854 (Micropalpus), Epalpodes	104.63
equatorialis (Townsend), 1912a (Eugymnochaeta), Chrysotachina	104.81
<i>equatorialis</i> Townsend, 1852, Epalpodes	104.63
equinoctialis Townsend, 1912, Triachora	104.184
eques (Wiedemann), 1830 (Dexia), Pseudodexia	104.108
crecta Aldrich, 1934, Theresia	104.38
Ergolabus Reinhard, 1964c	104.90
erigonopsidis Townsend, 1931a, Ernestiopsis	104.83
<i>Erigonopsis</i> Townsend, 1912a	104.77
Ernestiini	104.83
Ernestiopsis Townsend, 1931a	104.83
eronis Curran, 1929, Signosomopsis	104.72
errans Curran, 1927c, Prorhynchops	104.32
erratis Reinhard, 1958e, Dolichocodia	104.29
erro Townsend, 1931a, Itamintho	104.96
Ervia Robineau-Desvoidy, 1830	104.101
Erviopsis Townsend, 1934	104.101
Eryciini	104.204
Erynnia Robineau-Desvoidy, 1830	104.173
Erythroargyrops Townsend, 1917c	104.133
erythrocauda (Curran), 1934c (Frontina), Lespesia	104.209
erythrocauda (Bigot), 1888a (Gonia), Enchomyia	104.175
Erythrocerini	104.173
Erythroepalpus Townsend, 1931a	104.65
Erythromelana Townsend, 1919b	104.106
<i>Erythrophasia</i> Townsend, 1917c	104.10
erythropyga (Bigot), 1889 (Chaetolyga), Winthemia	104.196
erythropyga Sabrosky, 1950, Acaulona	104.6
erythropyga (Wulp), 1892 (Nemoraea), Xanthophyto	104.84
erythrostoma (Bigot), 1888a, (Epalpus), Vibrissomyia	104.80
csuriens (Fabricius), 1805 (Musca), Iteuthelaira	104.156
<i>esuriens</i> , authors, Belvosia	104.181
ethelius Curran, 1947, Rhachoeopalpus	104.70
etythrina Bigot, 1889 (Hystiricia)	104.60
Euacaulona Townsend, 1908	104.7
Euantha Wulp, 1885	104.111
Euanthoides Townsend, 1931a	104.111
<i>Eubelvosiopsis</i> Blanchard, 1954	104.181
Eubischofimyia Townsend, 1927a	104.65
<i>Eublepharipeza</i> Townsend, 1914	104.39
Eucelatoria Townsend, 1908	104.133
Eucelatoriodea Thompson, 1968	104.133
cucelatorioides Blanchard, 1963, Celatoriopsis	104.129
Eucelatoriopsis Townsend, 1927a	104.134
Euchaetogyne Townsend, 1908	104.37
Euchcirophaga James, 1945	104.83
Euclausicella Townsend, 1927a	104.134
<i>Eucordylidexia</i> Townsend, 1915i	104.111
Eucordyligaster Townsend, 1927c	104.111
Eucorpulentosoma Townsend, 1914	104.65




<i>Eucyrtophloe</i> Townsend, 1916f	104.90
<i>Eudejeania</i> Townsend, 1912a	104.53
<i>Eudexia</i> Brauer & Bergenstamm, 1889	104.29
<i>Euemphercmyia</i> Townsend, 1927a	104.66
<i>Euepalpodes</i> Townsend, 1915m	104.66
<i>Euepalpus</i> Townsend, 1908	104.66
<i>Eufabricia</i> Townsend, 1908	104.48
<i>Eufabriciopsis</i> Townsend, 1915b	104.66
<i>Eugaedia</i> Townsend, 1916f	104.179
<i>Eugaediopsis</i> Townsend, 1916c	104.179
<i>Eugymnochaeta</i> Townsend, 1915b	104.81
<i>Euhemiargyra</i> Townsend, 1927a	104.144
<i>Euhelioprosopa</i> Reinhard, 1964b	104.77
<i>Euhuascaraya</i> Townsend, 1927a	104.66
<i>Euhystricia</i> Macquart, 1843	104.41
<i>Eujurinia</i> Townsend, 1908	104.56
<i>Eujurinella</i> Townsend, 1914	104.56
<i>eulalia</i> Townsend, 1927a, <i>Euloewiodoria</i>	104.162
<i>Eulasiopampus</i> Townsend, 1913e	104.54
<i>Euloewiodoria</i> Townsend, 1927a	104.162
<i>Euloewiopsis</i> Townsend, 1917b	104.26
<i>Eumachaeraea</i> Townsend, 1927a	104.134
<i>Eumasicera</i> Townsend, 1909	104.162
<i>Eumasicerini</i>	104.161
<i>Eumelanepampus</i> Townsend, 1915m	104.66
<i>Eumogenia</i> Townsend, 1908	104.7
<i>Eumyobia</i> Townsend, 1911b	104.116
<i>Eunemorilla</i> Townsend, 1919b	104.189
<i>Euolestrogaster</i> Townsend, 1935d	104.155
<i>Euolestrogastrodes</i> Townsend, 1935d	104.156
<i>Euolestrophasia</i> Townsend, 1892f	104.20
<i>Euolestropsis</i> Townsend, 1913	104.25
<i>Euomogenia</i> Townsend, 1908	104.7
<i>Euparafrofrontina</i> Brèthes, 1917	104.208
<i>Eupelecotheca</i> Townsend, 1919b	104.156
<i>Euphasiopteryx</i> Townsend, 1915b	104.20
<i>Euphorocera</i> Townsend, 1892a	104.157
<i>Euphoroceropsis</i> Townsend, 1917a	104.158
<i>Euprospherysa</i> Townsend, 1928b	104.134
<i>Euptilodegeeria</i> Townsend, 1931a	104.134
<i>Euptililomyia</i> Townsend, 1939c	104.120
<i>Euquadrosoma</i> Townsend, 1915m	104.61
<i>eurotae</i> Blanchard, 1937, <i>Euphorocera</i>	104.158
<i>Eurylochus</i> Reinhard, 1927a	104.162
<i>Eurythemys</i> Reinhard, 1967b	104.162
<i>Eurythiopsis</i> Townsend, 1914	104.55
<i>Eusaundersiops</i> Townsend, 1915m	104.67
<i>Euscopliopterygini</i>	104.19
<i>Eusisyropa</i> Townsend, 1908	104.203
<i>Eusignosoma</i> Townsend, 1914	104.63
<i>Eustomatodexia</i> Townsend, 1892c	104.119
<i>Euthelothyria</i> Townsend, 1931a	104.120
<i>Euthelaira</i> Townsend, 1912a	104.156
<i>Euthelairopsis</i> Townsend, 1927a	104.156
<i>Euthelyconychia</i> Townsend, 1927a	104.135



<i>Euthera</i> Loew, 1866	104.18
<i>Eutheresiops</i> Townsend, 1917a	104.38
<i>Eutherini</i>	104.18
<i>Euthyprosopiella</i> Blanchard, 1963	104.98
<i>Eutrichopoda</i> Townsend, 1908	104.7
<i>Eutrichopodopsis</i> Blanchard, 1966c	104.7
<i>Eutritochaeta</i> Townsend, 1919a	104.210
<i>Eutrixoides</i> Walton, 1913b	104.23
<i>Euwinthemia</i> Blanchard, 1963	104.194
<i>Euzenilltops</i> Townsend, 1916	104.138
<i>eva</i> Townsend, 1935d, <i>Eujurinioides</i>	104.54
<i>evanescens</i> Cortés, 1967a, <i>Campsodes</i>	104.166
<i>exigua</i> Wulp, 1890, <i>Didyma</i>	104.131
<i>exilis</i> (Townsend), 1892 (<i>Pseudohystricia</i>), <i>Hystriciella</i>	104.56
<i>exilis</i> (Wulp), 1890 (<i>Exorista</i>) <i>Thelairodoria</i>	104.150
<i>Exodexia</i> Townsend, 1927a	104.29
<i>Exoernestia</i> Townsend, 1927a	104.83
<i>Exopalpus</i> Macquart, 1851	104.54
<i>Exoristini</i>	104.157
<i>Exoristinae</i>	104.122
<i>exoristoides</i> Townsend, 1927a, <i>Verrugophryno</i>	104.214
<i>Exoristopsis</i> Townsend, 1915n	104.87
<i>expergita</i> Walker, 1860, (<i>Masicera</i>)	104.152
<i>externa</i> (Fabricius), 1805 (<i>Dictya</i>), <i>Euscopoliopteryx</i>	104.19
<i>extrema</i> Walker, 1857 (<i>Dexia</i>), <i>Xanthodexia</i>	104.96
<i>Fabricia</i> Latreille, 1829	104.43
<i>Fabricia</i> Meigen, 1838	104.148
<i>Fabricia</i> Robineau-Desvoidy, 1830	104.43
<i>Fabriciella</i> Bezzi, 1906	104.43
<i>Fabriciopsis</i> Townsend, 1914	104.55
<i>faceta</i> Reinhard, 1957, <i>Plagiomima</i>	104.92
<i>facialis</i> (Aldrich), 1928b (<i>Selenomyia</i>), <i>Morphodexia</i>	104.24
<i>facialis</i> (Sellers), 1943 (<i>Zenillia</i>), <i>Chrysoexorista</i>	104.207
<i>facialis</i> Townsend, 1917b, <i>Atactopsis</i>	104.180
<i>facialis</i> Townsend, 1927a, <i>Actinoprosopa</i>	104.205
<i>facialis</i> Townsend, 1927a, <i>Helioprosopa</i>	104.78
<i>facialis</i> Townsend, 1928a, <i>Mayodistichona</i>	104.176
<i>facialis</i> Townsend, 1927a, <i>Nepophasmophaga</i>	104.172
<i>facialis</i> Townsend, 1927a, <i>Ollacheryphe</i>	104.88
<i>facialis</i> Townsend, 1935d, (<i>Cuphoceropsis</i>), <i>Peleteria</i>	104.45
<i>facialis</i> Townsend, 1931a, <i>Plaxactia</i>	104.145
<i>facialis</i> Townsend, 1933, <i>Trixodopsis</i>	104.26
<i>fallaciosus</i> Engel, 1920, <i>Dejeaniops</i>	104.53
<i>famelica</i> (Wiedemann), 1830 (<i>Stomoxys</i>), <i>Myobiopsis</i>	104.118
<i>famelica</i> (Wiedemann), 1830, (<i>Stomoxys</i>), <i>Myobiopsis</i>	104.118
<i>fantastica</i> Townsend, 1931a, <i>Fasslomyia</i>	104.84
<i>fasciata</i> Curran, 1934b, <i>Calodexia</i>	104.154
<i>fasciata</i> Curran, 1934d, <i>Canalia</i>	104.128
<i>fasciata</i> Curran, 1934d, <i>Patillalia</i>	104.221
<i>fasciata</i> Curran, 1934d, <i>Thclairalia</i>	104.164
<i>fasciata</i> (Macquart), 1848 (<i>Gymnostylia</i>), <i>Gymnaporia</i>	104.103
<i>fasciata</i> Macquart, 1842, <i>Macromya</i>	104.42
<i>fasciata</i> Townsend, 1927a, <i>Calocarcelia</i>	104.202

<i>fasciata</i> Townsend, 1927a, <i>Hcmilydella</i>	104.135
<i>fasciata</i> Wulp, 1890 (<i>Telothyria</i>)	104.152
<i>fraudulenta</i> Aldrich, 1928b, <i>Piriona</i>	104.99
<i>Fasslomyia</i> Townsend, 1931a	104.84
<i>fatua</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>feminea</i> Curran, 1947, <i>Jurinella</i>	104.57
<i>femoralis</i> Curran, 1941, <i>Eudejeania</i>	104.53
<i>femoratus</i> (Wulp), 1892 (<i>Saundersia</i>), <i>Epalpus</i>	104.64
<i>ferox</i> Townsend, 1912a, <i>Spathimya</i>	104.149
<i>ferruginea</i> Townsend, 1934 (<i>Procistogaster</i>), <i>Gymnoclytia</i>	104.5
<i>ferruginea</i> Townsend, 1929, <i>Jurinella</i>	104.57
<i>ferruginosa</i> Townsend, 1895, <i>Belvosia</i>	104.182
<i>ferruginea</i> Townsend, 1934 (<i>Procistogaster</i>), <i>Gymnoclytia</i>	104.5
<i>fertoria</i> Walker, 1861 (<i>Eurigaster</i>)	104.152
<i>festiva</i> Cortés, 1944a, <i>Sturmia</i>	104.192
<i>fidelis</i> (Reinhard), 1967 (<i>Punaclista</i>), <i>Prosopochaeta</i>	104.99
<i>fiebrigi</i> Townsend, 1931a, <i>Phyllaristomyia</i>	104.184
<i>filipalpis</i> (Blanchard) <i>in</i> Denier, 1939 (<i>Plagiprosphercysa</i>)	
<i>filipalpis</i> Rondani of Curran, 1925c (<i>Peleteria</i>)	104.45
<i>filipalpis</i> (Rondani), 1863 (<i>Echinomyia</i>), <i>Peleteria</i>	104.40
<i>filipalpis</i> authors, <i>Bombyliomyia</i>	104.40
<i>filipalpis</i> Townsend, 1927a, <i>Metamyobia</i>	104.117
<i>filipes</i> Robineau-Desvoidy, 1830, <i>Sophia</i>	104.112
<i>filipes</i> (Walker), 1857 (<i>Dexia</i>), <i>Cholomyia</i>	104.106
<i>fimbriata</i> (Wulp), 1890 (<i>Brachycoma</i>), <i>Thysanomyia</i>	104.192
<i>fimbricata</i> Wulp, 1890 (<i>Telothyria</i>)	104.152
<i>finita</i> Reinhard, 1964b, <i>Helioprosopa</i>	104.78
<i>fitzgeraldi</i> Curran, 1937, <i>Polybiophila</i>	104.122
<i>flava</i> (Aldrich), 1929 (<i>Euantha</i>), <i>Leptidosophia</i>	104.112
<i>flava</i> Röder, 1885, <i>Trichopoda</i> (<i>Trichopoda</i>)	104.9
<i>flava</i> Townsend, 1917b, <i>Trochiloleskia</i>	104.119
<i>flava</i> Townsend, 1911b, <i>Eumyobia</i>	104.116
<i>flava</i> Townsend, 1916h, <i>Beskioccephala</i>	104.75
<i>flava</i> Townsend, 1919a, <i>Opsoleskia</i>	104.108
<i>flavescens</i> Robineau-Desvoidy, 1830, <i>Icelia</i>	104.101
<i>flavescens</i> Townsend, 1935d, <i>Euolestrogastroides</i>	104.155
<i>flavescens</i> (Townsend), 1929 (<i>Leskiopalpus</i>), <i>Myobiopsis</i>	104.118
<i>flavescens</i> (Wulp), 1890 (<i>Masicera</i>)	104.152
<i>flavibasis</i> Aldrich, 1934, <i>Admontia</i>	104.171
<i>flavicans</i> (Macquart), 1846 (<i>Micropalpus</i>), <i>Epalpus</i>	104.64
<i>flavicans</i> Townsend, 1908 (<i>Eufabricia</i>), <i>Archytas</i>	104.49
<i>flavicans</i> (Wulp), 1890 (<i>Exorista</i>), <i>Lespesia</i>	104.209
<i>flavicauda</i> Townsend, 1908, <i>Euepalpus</i>	104.66
<i>flavicauda</i> (Wulp), 1890 (<i>Phorocera</i>), <i>Gaediopsis</i>	104.179
<i>flavicornis</i> (Brauer & Bergenstamm), 1891 (<i>Minthodexia</i>), <i>Minthodexiopsis</i>	104.107
<i>flavicornis</i> (Wulp), 1890, (<i>Myobia</i>) <i>Calothelaira</i>	104.105
<i>flavicosta</i> Wulp, 1891, <i>Rhombothyria</i>	104.94
<i>ssp. flavida</i> Townsend, 1915d, <i>Aglummyia</i>	104.28
<i>flavifacies</i> Bigot, 1889 (<i>Masicera</i>)	104.193
<i>flavifacies</i> (Macquart), 1851 (<i>Jurinia</i>), <i>Archytas</i>	104.49
<i>flavifrons</i> (Jaenicke), 1867 (<i>Jurinia</i>), <i>Archytas</i>	104.49
<i>flavifrons</i> Rondani, 1863, <i>Spathipalpus</i>	104.118
<i>flavifrons</i> Wulp, 1890, <i>Mystacella</i>	104.191

<i>flavipalpis</i> (Macquart), 1846 (Hystricia), Bombyliomyia	104.39
<i>flavipalpis</i> authors, Bombyliomyia	104.40
<i>flavipalpis</i> Thompson, 1964, Pseudochaeta	104.163
<i>flavipennis</i> (Wiedemann), 1830 (Dexia), Myoblopsis	104.118
<i>flavipes</i> Aldrich not Schiner, Calodexia	104.155
<i>flavipes</i> (Bigot), 1889 (Homodexia), Eucalodexia	104.106
<i>flavipes</i> Coquillett, 1895, Prosenoides	104.32
<i>flavipes</i> (Schiner), 1868 (Meigenia), Calodexia	104.154
<i>flavipes</i> Thompson, 1964, Amazohoughia	104.201
<i>flavipes</i> Thompson, 1968, Pelecotheca	104.157
<i>flavipes</i> Thompson, 1968, Synorbitalia	104.173
<i>flavipes</i> Townsend, 1915n, Vibrissoepalpus	104.64
<i>flavipes</i> Vimmer & Soukup, 1940, Epalpus	104.202
<i>flavirostris</i> (Wulp), 1890 (Exorista), Carcelia	104.202
<i>flavirostris</i> , authors, Carcelia	104.39
<i>flavitaris</i> (Macquart), 1846 (Hystricia), Bombyliomyia	104.42
<i>flavitaris</i> Macquart, 1847, Lasiopalpus	104.70
<i>flavitaris</i> (Macquart), 1843 (Micropalpus), Rhachoeopalpus	104.41
<i>flavitibia</i> Curran, 1942, Hystricia	104.126
<i>flaviventris</i> (Macquart), 1843 (Lydella), Blondelia	104.44
<i>flaviventris</i> (Wulp), 1888 (Echinomyia), Peleteria	104.19
<i>flavocalyptrata</i> Bigot, 1889 (Stevcnia)	104.126
<i>flavocalyptratus</i> (Wulp), 1890 (Hypostena), Anoxynops	104.213
<i>flavolimbata</i> (Bigot), 1889 (Chaetolyga), Triodontopyga	104.69
<i>flavoscutellatus</i> (Bischof), 1904 (Epalpus), Oxyepalpus	104.110
<i>flavotegulata</i> Townsend, 1927a, Myiophasiopsis	104.121
<i>Floradalia</i> Thompson, 1963a	104.150
<i>floridensis</i> Townsend, 1892, Tachynophyto	104.158
<i>floridensis</i> Townsend, 1916d, Euphorocera	104.58
<i>floridensis</i> Townsend, 1916e, Juriniopsis	104.43
<i>florum</i> (Walker), 1849 (Tachina), Fabriciella	104.7
<i>Forcipophasia</i> Townsend, 1935d	104.27
<i>Formicomylia</i> Townsend, 1916h	104.29
<i>formidabilis</i> (Bigot), 1889 (Rhamphina), Eudexia	104.182
<i>formosa</i> Aldrich, 1928a, Belvosia	104.182
<i>formosana</i> (Blanchard), 1954 (Eubelvosioipsis), Belvosia	104.40
<i>formosus</i> Reinhard, 1964c, Diotrephes	104.40
<i>forreri</i> (Wulp), 1964c, Diotrephes	104.152
<i>forticula</i> Wulp, 1890 (Telothyria)	104.182
<i>fosteri</i> (Townsend), 1915m (Belvosiomima), Belvosia	104.193
<i>foveata</i> Wulp, 1890 (Brachycoma)	104.108
<i>foxii</i> Johnson, 1919, Stenodexia	104.167
<i>fracida</i> Reinhard, 1962a, Ginglymyia	104.33
<i>fraterna</i> Reinhard, (Wulp), 1891 (Rhynchodexia), Ptilodexia ...	104.168
<i>fraudentula</i> (Wulp), 1890 (Masicerca), Drino	104.50
<i>frenguelli</i> (Blanchard), 1941 (Archynemochaeta), Archytas ...	104.13
<i>freyreisi</i> (Wiedemann), 1830 (Phasia), Hyalomylia	104.182
<i>frontalis</i> Aldrich, 1928a, Belvosia	104.167
<i>frontalis</i> (Aldrich), 1934 (Lispidea), Elfia	104.120
<i>frontalis</i> Townsend, 1939c, Euptilomyia	104.135
<i>frontalis</i> Townsend, 1927a, Hemiargyropsis	104.118
<i>frontalis</i> Townsend, 1934, Proleskiomima	



frontalis (Wulp), 1892 (Nemochaeta), Archytas	104.50
Frontiniella Townsend, 1918	104.208
Frontocnephalia Townsend, 1916h	104.175
frontosa Cortés, 1951, Ruiziella	104.45
<i>Frontina</i> , authors	104.208
frugale (Curran), 1934e (Lydella), Blondelia	104.126
fucatus (Wulp), 1892 (Trichopoda), Deopalpus	104.76
fulgens Reinhard, 1967, Tarpessita	104.88
fuliginipennis Bigot, 1888a, Jurinia	104.68
fuliginipennis Wulp, 1890 (Didyma)	104.214
fuliginosum Robineau-Desvoidy, 1830, Gymnosomma	104.5
fulva Townsend, 1927a, Nepocarcelia	104.204
fulvescens Townsend, 1927a, Itabiomyia	104.137
fulvibasis Curran, 1934a, Calodexia	104.154
fulvidapex (Bigot), 1889 (Tricoliga), Winthemia	104.196
fulvicauda (Walton), 1914 (Linnaemyia), Gymnochaetopsis	104.82
fulviceps (Wulp), 1890 (Phorocera), Patelloa	104.211
fulvipennis Wulp, 1890, Anisia	104.125
<i>fulvipes</i> Townsend, 1927a (Oligolydella), Pseudoredtenbacheria	104.147
fulvipes (Wulp), 1890 (Phorocera), Patelloa	104.211
fulviventris (Robineau-Desvoidy), 1830 (Jurinia), Archytas	104.50
<i>fulviventris</i> Townsend, 1935 (Makasinoceropsis), Archytas	104.52
fulviventris Townsend, 1927a, Charapozelia	104.37
fulvus Townsend, 1914m, Melanepalpellus	104.68
<i>fulvus</i> Townsend, 1927a, (Phosococephalops), Gonia	104.176
fumata Wulp, 1895 (Morinia)	104.110
fumata Townsend, 1919b, Comyopsis	104.109
fumifera Townsend, 1919a, Phasiophyto	104.110
fumipennis (Bigot), 1889, (Dexiosoma), Ptilodexia	104.33
fumipennis Thompson, 1968, Cataphoriniopsis	104.128
fumipennis Thompson, 1968, Santacruziniopsis	104.164
fumipennis Engel, 1920, Engelomyia	104.41
fumipennis Townsend, 1935d, Dexodomintho	104.131
fumipennis Townsend, 1928b, Euprospherysa	104.134
fumipennis Townsend, 1927a, Paraprospherysa	104.145
fumosa (Townsend), 1912a (Oestrogaster), Calodexia	104.154
fumosa Townsend, 1917b, Ommaleskia	104.107
funditor Curran, 1934, Belvosiella	104.183
funebri Reinhard, 1952a, Cyosoprocta	104.184
funebri Reinhard, 1966c, Eutrichopodopsis	104.7
funebri Reinhard, 1964c, Gigamyiopsis	104.29
furcata (Wulp), 1890 (Labidigaster), Clairvillia	104.17
furtum Reinhard, 1958a, Mochlosoma	104.31
fusca Townsend, 1935d, Forcipophasia	104.7
fuscus Townsend, 1935d, Euoestrogaster	104.155
fuscanipennis Bischof, 1904, Epalpus	104.64
fuscanipennis Macquart, 1846 (Dexia)	104.122
fuscicornis Curran, 1925e, Jurinella	104.57
fuscicostallis (Wulp), 1890 (Mystacella), Paramesochaeta	104.159
fuscifacies Bigot, 1888a, Cordyligaster	104.111
<i>fuscipennis</i> Macquart, 1851 (Megistogaster), Cordyligaster	104.111
fuscisquamma Curran, 1929c, Peleteria	104.44
fuscisquamula (Bigot), 1954 (Willistonella), Belvosia	104.182

<i>fuscosa</i> Townsend, 1912a, Chaetophorocera	104.206
<i>fusca</i> Townsend, 1935d, Forcipophasia	104.7
<i>fusicornis</i> Curran, error, Jurinella	104.57
<i>futilis</i> Wulp, 1890, Siphona	104.170
<i>gabana</i> (Townsend), 1919 (Eublepharipeza), Bombyliomyia	104.40
<i>Gabanimyia</i> Townsend, 1914	104.72
<i>gabanus</i> Townsend, 1914, Xanthoepalpus	104.73
<i>Gacdiophana</i> Brauer & Bergenstamm, 1893	104.179
<i>Gaediophanopsis</i> Blanchard, 1954	104.179
<i>Gaediopsis</i> Brauer & Bergenstamm, 1891	104.179
<i>Gaediosturmia</i> Blanchard, 1963	104.178
<i>gagatea</i> (Robineau-Desvoidy), 1830 (Sophia), Scotiptera	104.35
<i>gagatea</i> Robineau-Desvoidy, 1830, Jurinia	104.68
<i>gagatea</i> Townsend, 1927a, Gymnommopsis	104.78
<i>gagateum</i> Townsend, 1927a, Juriniosoma	104.68
<i>gagatina</i> Wulp, 1890, Anisia	104.125
<i>Galactomyia</i> Townsend, 1908	104.9
<i>Galapagosia</i> Curran, 1934a	104.116
<i>galbae</i> Thompson, 1968, Cataphoriniopsis	104.128
<i>Ganopleuron</i> Aldrich, 1934	104.98
<i>Ganoproctus</i> Aldrich, 1934	104.78
<i>geminata</i> (Brauer & Bergenstamm), 1889 (Masipoda), Winthemia	104.196
<i>geminata</i> Curran, 1926a (Atacta), Drino	104.189
<i>geminata</i> Townsend, 1927a, Calolydella	104.127
<i>geminata</i> Townsend, 1927a, Italydella	104.137
<i>genalis</i> Aldrich, 1934, Setolestes	104.100
<i>genalis</i> Townsend, 1927a, Neocampylochaeta	104.89
<i>Genea</i> Rondani, 1850b	104.116
<i>genei</i> Rondani, 1863, Gonias	104.175
<i>Geneodes</i> Townsend, 1934	104.117
<i>Geneoglossa</i> Townsend, 1935d	104.116
<i>Geneopsis</i> Townsend, 1927a	104.116
<i>generosa</i> (Wulp), 1892 (Echinomyia), Peleteria	104.44
<i>gentica</i> Walker, 1860 (Masicera)	104.152
<i>genuina</i> Wulp, 1891:243 (Dexia)	104.122
<i>Germariini</i>	104.86
<i>Germariopsis</i> Townsend, 1915e	104.175
<i>gerstchi</i> Curran, 1942, Calohystricia	104.40
<i>gerstchi</i> Curran, 1947, Eulasiopalpus	104.54
<i>gertschi</i> Curran, 1947, Jurinella	104.57
<i>giacomellii</i> (Blanchard), 1941 (Proarchytoides), Archytas	104.50
<i>giacomellii</i> (Blanchard), 1966c (Trichopodopsis), Trichopoda, (Galactomyia)	104.10
<i>giacomelli</i> (Blanchard), 1943b (Aphriosphyriopsis), Peleteria ...	104.44
<i>Gigamyiopsis</i> Reinhard, 1964c	104.29
<i>Gigantachinosoma</i> Townsend, 1932b	104.56
<i>gigantea</i> (Townsend), 1932b (Gigantachinosoma), Jurinella	104.57
<i>Gigantoepalpus</i> Townsend, 1913a	104.67
<i>Ginglymia</i> Townsend, 1892a	104.167
<i>Ginglymyia</i> , emend.	104.167
<i>glabricula</i> Wulp, 1890 (Phorocera), Winthemia	104.198
<i>gladiatrix</i> Townsend, 1917c, Xiphomyia	104.151



glaucia Aldrich, 1934, Hyalomya	104.13
glaucia Giglio-Tos, 1893 (Masicera)	104.216
<i>Glaurocarini</i> , in part	104.19
globosa (Reinhard), 1953a (Nicetria), Calodexia	104.154
globosa (Townsend), 1892d (Loewia), Myiophasia	104.27
glossata (Townsend), 1935d (Geneoglossa), Genea	104.116
Glossidionophora Bigot, 1885c	104.16
Gnadochaeta Macquart, 1851	104.25
gollath (Brauer & Bergenstamm), 1889, Eudexia	104.29
Gonatorrhina Röder, 1886	104.98
goncalvesi Guimarães, 1963b, Archytas	104.50
Gonia Meigen, 1803	104.175
Goniina	104.173
Goniinae	104.122
Goniini	104.173
<i>gonioides</i> (Bigot), 1888a (Jurinia), Archytas	104.50
gonioides Cortés, 1967b, Gonzalezodoria	104.26
<i>Goniomima</i> Townsend, 1908	104.184
Gonistylum Macquart, 1851	104.176
<i>Gonistylum</i> error	104.176
Gonzalezodoria Cortés, 1967b	104.26
gortys Walker, 1949 (Dexia)	104.122
Gopaulia Thompson, 1968	104.158
gowdeyi Curran, 1928a, Argyrophyllax	104.162
gowdeyi Curran, 1926, Zenillia	104.214
gracilens Giglio-Tos, 1893 (Tricholyga)	104.215
gracilentus (Wulp), 1892 (Xanthomelana), Xanthomelanoides ...	104.10
gracilis Giglio-Tos, 1893 (Chaetogena)	104.122
gracilis James, 1947, Genea	104.116
gracilis Townsend, 1919a, Leptodexia	104.30
gracilis Townsend, 1927a, Medinophyto	104.106
gracilis Townsend, 1919a, Microchaetona	104.107
gracilis Townsend, 1927a, Xanthophyllophilla	104.109
<i>gracilis</i> Wiedemann, 1830 (Dexia), Zelia	104.103
gradata Wiedemann, 1830, Trichopoda, (Trichopoda)	104.9
grandis Guimarães, 1966b, Adejeania	104.46
grandis Reinhard, 1954, Prosenoides	104.32
grandis Townsend, 1935d, Chaetocalirrhoe	104.28
grandis Townsend, 1927a, Itaxanthomelana	104.8
grandis (Townsend), 1940b (Xenopyxis), Neobrachelia	104.86
Graphogastrini	104.171
grata (Wulp), 1890 (Myobia), Calodexia	104.154
gratiosa Wulp, 1890 (Meigenia)	104.152
<i>gravipes</i> Brauer & Bergenstamm, 1891 (Minthodexia), Xantho- dexia	104.96
gravipes Wulp, 1890, Leucostoma	104.18
grioti (Blanchard), 1963 (Ypophaeomyia), Lespesia	104.209
grioti (Blanchard), 1963 (Winthemiopsis), Winthemia	104.196
grioti Blanchard, 1957, Parapoliops	104.169
grioti (Blanchard), 1943b (Aphriosphyria), Peleteria	104.44
griphus Reinhard, 1964c, Hegesinus	104.55
grisea (Curran), 1926a (Hypostena), Lixophaga	104.138
grisea Thompson, 1968, Anoxynopsoidea	104.126
grisea Thompson, 1968, Vivianoidea	104.137


<i>grisea</i> (Townsend), 1927a (<i>Gymnosturmia</i>), <i>Topomeigenia</i>	104.193
<i>griseomicans</i> (Wulp), 1890 (<i>Exorista</i>), <i>Carcelia</i>	104.202
<i>griseus</i> Townsend, 1934, <i>Geneodes</i>	104.117
<i>guianensis</i> Townsend, 1935d, <i>Irengia</i>	104.23
<i>guianensis</i> (Townsend), 1919 (<i>Trichoduropsis</i>), <i>Trichodura</i>	104.103
<i>guianica</i> (Curran), 1934c (<i>Ormia</i>), <i>Euphasiopsteryx</i>	104.21
<i>guianica</i> Curran, 1934c, (<i>Phorocera</i>)	104.161
<i>Gymnaporia</i> Townsend, 1919b	104.103
<i>Gymnocarcelia</i> Townsend, 1919a	104.203
<i>Gymnochaetopsis</i> Townsend, 1914	104.82
<i>Gymnoclytia</i> Brauer & Bergenstamm, 1893	104.5
<i>Gymnocuphocera</i> Blanchard, 1943b	104.78
<i>Gymnodoria</i> Townsend, 1927a	104.135
<i>Gymnojurinella</i> Vimmer, 1940	104.217
<i>Gymnomma</i> Wulp, 1888	104.67
<i>Gymnommopsis</i> Townsend, 1927a	104.78
<i>Gymnopalpus</i> Townsend, 1919b	104.94
<i>gymnops</i> Macquart, 1843 (<i>Tryphera</i>)	104.216
<i>Gymnosomma</i> Meigen, 1803	104.5
<i>Gymnosomatinae</i>	104.4
<i>Gymnosomatini</i>	104.4
<i>Gymnosturmia</i> Townsend, 1927a	104.193
<i>Gymnostylia</i> Macquart, of Brauer & Bergenstamm	104.143
<i>haemorrhoidalis</i> Bigot, 1889 (<i>Tromodesia</i>), <i>Tromodesiopsis</i>	104.102
<i>haenschi</i> Townsend, 1931a, <i>Homosaundersiops</i>	104.67
<i>haitensis</i> Robineau-Desvoidy, 1830 <i>Trichopoda</i> , (<i>Trichopoda</i>) ...	104.9
<i>halliana</i> Cortés, 1945c, <i>Carcelia</i>	104.202
<i>hamata</i> Wulp, 1890 (<i>Telothyria</i>), <i>Velocia</i>	104.151
<i>handlirschi</i> Brauer & Bergenstamm, 1889, <i>Bibliomima</i>	104.6
<i>hariola</i> Reinhard, 1961, <i>Uramya</i>	104.104
<i>harpeus</i> Reinhard, 1952a, <i>Saundersiops</i>	104.71
<i>Harrisia</i> Robineau-Desvoidy, 1830	104.185
<i>Harrisiini</i>	104.185
<i>harrisinae</i> Coquillett, 1897, <i>Sturmia</i>	104.192
<i>Harrisopsis</i> Townsend, 1927a	104.185
<i>haustellata</i> Bigot, 1885c, <i>Bolbochaeta</i>	104.165
<i>haustellata</i> (Townsend) 1927a, (<i>Neoprosena</i>), <i>Prosenoides</i>	104.32
<i>haywardi</i> (Blanchard), 1927a (<i>Plagiotachina</i>), <i>Euphorocera</i>	104.158
<i>haywardi</i> (Blanchard), 1927a, (<i>Metagymnopsis</i>) <i>Gymnommopsis</i>	104.78
<i>haywardi</i> (Blanchard), 1942a (<i>Ypophamyia</i>), <i>Prottypophamyia</i>	104.195
<i>Haywardiamyia</i> Blanchard, 1955	104.168
<i>hebes</i> (Fallén), 1820 (<i>Tachina</i>), <i>Spallanzania</i>	104.177
<i>hebes</i> (Wulp), 1892 (<i>Hyalomyia</i>), <i>Heyncophasia</i>	104.12
<i>hedemannni</i> (Brauer & Bergenstamm), 1891 (<i>Paragaedia</i>), <i>Chaeteprosopa</i>	104.184
<i>Hegesinus</i> Reinhard, 1964c	104.55
<i>heinrichi</i> (Lima), 1947 (<i>Zygosturmia</i>), <i>Drino</i>	104.188
<i>Heliaea</i> Curran, 1934c	104.90
<i>heliconiinarum</i> (Townsend), 1929 (<i>Plagiotachina</i>), <i>Euphorocera</i>	104.158
<i>Heliconiophaga</i> Thompson, 1966	104.189
<i>Heliodes</i> Thompson, 1968	104.135
<i>Helioplaga</i> Townsend, 1934a	104.87
<i>Heliolydella</i> Townsend, 1927a	104.135



<i>Heliolydellops</i> Townsend, 1934a	104.87
<i>Heliolydella</i> Townsend, 1934a	104.87
<i>hemiargyroides</i> Townsend, 1927a, <i>Argyrodoria</i>	104.126
<i>Helioprosopa</i> Townsend, 1927a	104.78
<i>Hemiarchytas</i> Vimmer, 1940	104.217
<i>Hemiargyra</i> Townsend, 1908	104.141
<i>Hemiargyrophylax</i> Townsend, 1927a	104.135
<i>Hemiargyropsis</i> Townsend, 1927a	104.135
<i>Hemilydella</i> Townsend, 1927a	104.135
<i>Hemimasipoda</i> Townsend, 1927a	104.196
<i>Hemisturmia</i> Townsend, 1927a	104.194
<i>Hemyda</i> Robineau-Desvoidy, 1830	104.16
<i>heros</i> (Schiner), 1868 (<i>Phorocera</i>), <i>Palpexorista</i>	104.159
<i>heros</i> Townsend, 1931a, <i>Gigantoepalpus</i>	104.67
<i>hesperidarum</i> Williston, 1889, <i>Acroglossa</i>	104.173
<i>Hesperophasia</i> Townsend, 1915g	104.93
<i>Hesperophasiini</i>	104.93
<i>heynei</i> Townsend, 1934, <i>Heyneophasia</i>	104.12
<i>Heyneophasia</i> Townsend, 1934	104.12
<i>heyrovsky</i> Vimmer & Soukup, 1940, <i>Neocyrtophoeba</i>	104.91
<i>hichinsi</i> Cortés, 1967b, <i>Parasetigena</i>	104.159
<i>hiemalis</i> Thompson, 1939b, <i>Archytas</i>	104.50
<i>hirsutus</i> Townsend, 1908, <i>Deopalpus</i>	104.76
<i>hirta</i> Townsend, 1927a, <i>Proleskia</i>	104.86
<i>hirticollis</i> (Wulp), 1892 (<i>Cistogaster</i>), <i>Gymnoclytia</i>	104.5
<i>hispidula</i> Wulp, 1890 (<i>Exorista</i>), <i>Trafoia</i>	104.83
<i>histrion</i> (Aldrich), 1928b, (<i>Callesthes</i>), <i>Medinophyto</i>	104.106
<i>homeonychioides</i> (Townsend), 1934a, (<i>Heliolydella</i>), <i>Heliolydellops</i>	104.87
<i>Homogenia</i> Wulp, 1892	104.7
<i>Homohypochaeta</i> Townsend, 1927a	104.89
<i>Homosaundersia</i> Townsend, 1931a	104.67
<i>Homosaunderslops</i> Townsend, 1931a	104.67
<i>honesta</i> (Rondani), 1850 (<i>Dejeania</i>), <i>Adejeania</i>	104.46
<i>horrida</i> (Giglio-Tos), 1893 (<i>Cyrtophloeoba</i>), <i>Eucyrtophloeoba</i>	104.90
<i>hospita</i> Reinhard, 1952a, <i>Leschenaultia</i>	104.186
<i>Huascaraya</i> Townsend, 1914	104.72
<i>huascarayana</i> Townsend, 1914, <i>Eudejeania</i>	104.53
<i>Huascarayopsis</i> Townsend, 1927a	104.67
<i>huascarayus</i> (Townsend), 1917c (<i>Hemiargyra</i>), <i>Myiopharus</i>	104.141
<i>Huascarodexia</i> Townsend, 1919b	104.101
<i>humeralis</i> Curran, 1942, <i>Hystricia</i>	104.41
<i>humeralis</i> (Wulp), 1890 (<i>Telothyria</i>), <i>Lydinolydella</i>	104.138
<i>huntingtoni</i> Curran, 1947, <i>Jurinella</i>	104.57
<i>hyalinipennis</i> (Macquart), 1835 (<i>Echinomyia</i>), <i>Jurinella</i>	104.68
<i>hyalinipennis</i> (Wulp), 1890 (<i>Degeeria</i>), <i>Doryphorophaga</i>	104.132
<i>Hyadesimyia</i> Bigot, 1888b	104.216
<i>Hyalomya</i> Robineau-Desvoidy, 1830	104.13
<i>Hyalomyia</i> , emend.	104.13
<i>Hyalomyodes</i> Townsend, 1893	104.18
<i>Hyasoma</i> Aldrich, 1934	104.98
<i>hyphena</i> (Townsend), 1927a, (<i>Hypophorinia</i>), <i>Argyrodoria</i>	104.126
<i>Hypenomyia</i> Townsend, 1919a	104.28
<i>Hyperecteinini</i> , in part	104.172



Hypertrophocera Townsend, 1891b	104.114
Hypertrophomma Townsend, 1915k	104.163
Hypochaetopsis Townsend, 1915m	104.89
hypodermica Townscnd, 1927a, Hypomyothyria	104.136
Hypodoria Townsend, 1927a	104.135
Hypohoughia Townsend, 1927a	104.136
Hypomyothyria Townsend, 1927a	104.136
Hypophorinia Townsend, 1927a	104.126
Hypophylax Townsend, 1935d	104.136
Hypoproxynops Townsend, 1927a	104.136
Hypotachina Brauer & Bergenstamm, 1891	104.39
Hypotachinini	104.39
Hystrichodexia Röder, 1886	104.30
Hystricia Macquart, 1843	104.41
<i>Hystricia</i> , authors	104.40
<i>Hystriciella</i> Engel, 1920	104.40
<i>Hystriciella</i> Townsend, 1915k	104.55
Hystriciopsis Townsend, 1914	104.42
<i>hystricosa</i> Townsend, 1914 (Gabanimyia), Trichophora	104.72
<i>hystricosa</i> (Williston), 1886 (Dejeania), Protodejeania	104.59
<i>Hystrisyphona</i> , emend.	104.30
<i>hystrix</i> Brauer & Bergenstamm, 1891 (Paragymnoma), Trichophora	104.72
<i>hystrix</i> Fabricius, 1775 (Musca), <i>Hystriciella</i>	104.56
<i>hystrix</i> Rondani, 1865, <i>Cryptopalpus</i>	104.61
<i>hystrix</i> (Schiner) in Brauer & Bergenstamm, 1891 (Saundersia), <i>Cryptopalpus</i>	104.61
<i>hystrix</i> Townsend, 1931a, <i>Anepalpus</i>	104.60
<i>hystrix</i> Townsend, 1914 (Eublepharipeza), <i>Bombyliomyia</i>	104.39
<i>hystrix</i> Townsend, 1915m, <i>Echinomasicera</i>	104.185
<i>hystrix</i> Townsend, 1914, <i>Fabriciopsis</i>	104.55
<i>hystrix</i> , authors, <i>Archytas</i>	104.51
Icelia Robineau-Desvoidy, 1830	104.101
Iceliini	104.100
Iconofrontina Townsend, 1931a	104.208
icterica (Wiedemann), 1830 (Tachina), Chaetona	104.105
Ictericodexia Townsend, 1934a	104.106
ignobilis (Wulp), 1890 (Exorista), Winthemia	104.196
Ignotomyia Reinhard, 1961	104.198
illucens Wulp, 1890, Telothyria	104.121
<i>imaculata</i> Macquart, 1851 (Echinomyia), Archytas	104.49
<i>imaculata</i> (Macquart), 1843 (Cistogaster), <i>Gymnoclytia</i>	104.5
<i>immaculata</i> Wulp, 1890, (Phorocera)	104.161
<i>immaculatus</i> (Macquart), 1846 (Hystricia), <i>Rhachoepalpus</i>	104.70
<i>Imaguncula</i> Reinhard, 1958b	104.176
<i>imitans</i> Blanchard, 1966c, <i>Eutrichopodopsis</i>	104.7
<i>imitator</i> (Townsend), 1929 (Argentoepalpus), <i>Epalpus</i>	104.64
<i>imitatrix</i> Townsend, 1919a, <i>Incamyiopsis</i>	104.137
<i>imitatrix</i> (Wulp), 1891 (Rhynchodexia), <i>Ptilodexia</i>	104.33
<i>immissa</i> (Reinhard), 1959b (Lydella), <i>Oswaldia</i>	104.143
<i>immista</i> Reinhard, 1955c, <i>Lydina</i>	104.87
<i>immunda</i> (Wulp), 1890 (Hypostena), <i>Ptilodegeeria</i>	104.147
<i>impedita</i> (Wulp), 1890 (Masicera), <i>Eumasicera</i>	104.152



<i>imperialis</i> Curran, 1926, (<i>Sturmia</i>)	104.193
<i>impressum</i> Townsend, 1914, <i>Signosoma</i>	104.71
<i>inaequipes</i> Bigot, 1884, <i>Cholomyia</i>	104.106
<i>inambarica</i> Townsend, 1912a, <i>Euthelaira</i>	104.156
<i>inambaricus</i> (Townsend) 1915m (<i>Neoarchytas</i>), <i>Archytas</i>	104.50
<i>inca</i> Curran, 1925c, <i>Peleteria</i>	104.44
<i>inca</i> Townsend, 1927a, <i>Acemeigenia</i>	104.123
<i>inca</i> (Townsend), 1911b, (<i>Zygosturmia</i>), <i>Drino</i>	104.188
<i>Incamyia</i> Townsend, 1912a	104.136
<i>Incamiopsis</i> Townsend, 1919a	104.137
<i>inacaram</i> (Townsend), 1912a (<i>Neotrafoia</i>), <i>Trafoia</i>	104.83
<i>incasanus</i> Townsend, 1912a, <i>Archytas</i>	104.49
<i>incertus</i> , authors, <i>Archytas</i>	104.51
<i>incertus</i> (Macquart), 1851 (<i>Gonia</i>), <i>Archytas</i>	104.50
<i>incertus</i> (Giglio-Tos), 1893 (<i>Nemochaeta</i>), <i>Archytas</i>	104.50
<i>inclinans</i> Walker, 1857 (<i>Masicera</i>)	104.216
<i>incognita</i> (Blanchard), 1966c:62 (<i>Trichopodopsis</i>), <i>Trichopoda</i> ..	104.9
<i>incognita</i> (Wulp), 1890 (<i>Plagia</i>), <i>Plagiomima</i>	104.92
<i>incompleta</i> (Curran), 1928a, (<i>Lydella</i>), <i>Blondelia</i>	104.126
<i>incompta</i> (Wulp), 1890 (<i>Brachycoma</i>), <i>Drino</i>	104.189
<i>inconspicua</i> Wulp, 1890, <i>Didyma</i>	104.131
<i>inconstans</i> (Wiedemann), 1830 (<i>Trichopoda</i>), <i>Homogenia</i>	104.8
<i>incontestata</i> Curran, 1926b, <i>Peleteria</i>	104.44
<i>indulta</i> (Wiedemann), 1830 (<i>Tachina</i>), <i>Carcelia</i>	104.202
<i>indita</i> (Walker), 1860 (<i>Lydella</i>), <i>Pseudeuantha</i>	104.104
<i>indivisa</i> Townsend, 1897c, <i>Trichopoda</i> (<i>Trichopoda</i>)	104.9
<i>inepta</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>infernalis</i> (Townsend), 1919a, (<i>Stomatolydella</i>), <i>Paralispe</i>	104.145
<i>infima</i> Wulp, 1890 (<i>Anisia</i>)	104.214
<i>infirma</i> (Walker), 1849 (<i>Tachina</i>), <i>Archytas</i>	104.52
<i>inflata</i> (Séguy), 1927b, (<i>Pseudormia</i>), <i>Ormiophasia</i>	104.22
<i>inflaxa</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>infumatus</i> (Bigot), 1887, (<i>Fabricia</i>), <i>Archytas</i>	104.50
<i>infuscata</i> (Wulp), 1888 (<i>Hystricia</i>), <i>Exopalpus</i>	104.55
<i>infuscatus</i> (Wulp), 1887 (<i>Nemochaeta</i>), <i>Archytas</i>	104.50
<i>ingloria</i> Wulp, 1890 (<i>Prospheysa</i>)	104.215
<i>inornata</i> (Walker), 1837 (<i>Tachina</i>), <i>Dasyuromyia</i>	104.97
<i>innovata</i> (Walker), 1860 (<i>Jurinia</i>), <i>Archytas</i>	104.50
<i>innovata</i> Walker, 1861 (<i>Tachina</i>)	104.216
<i>innupta</i> Cortés, 1945a, <i>Chaetocnephalia</i>	104.174
<i>inornatus</i> (Schiner), 1868 (<i>Saundersia</i>), <i>Eusaundersiops</i>	104.67
<i>inquinata</i> (Wulp), 1890 (<i>Masicera</i>), <i>Drino</i>	104.189
<i>insecta</i> Giglio-Tos, 1893 (<i>Degeeria</i>)	104.152
<i>insignis</i> (Wulp), 1882 (<i>Masicera</i>), <i>Sturmia</i>	104.192
<i>insita</i> Giglio-Tos, 1893 (<i>Tricholyga</i>)	104.215
<i>insolita</i> Curran, 1934b, <i>Calodexia</i>	104.154
<i>insolita</i> (Walker), 1852 (<i>Dexia</i>), <i>Hystrichodexia</i>	104.30
<i>insularis</i> (Brauer & Bergenstamm), 1891 (<i>Prosopaea</i>), <i>Prosopaea</i>	104.160
<i>insularis</i> Curran, 1927 (<i>Belvosia</i>), <i>Triachora</i>	104.184
<i>insularis</i> Curran, 1960, <i>Juriniopsis</i>	104.58
<i>insularis</i> Curran, 1927c, <i>Comatacta</i>	104.120
<i>insularis</i> Williston, 1896, <i>Cistogaster</i>	104.5
<i>insulensis</i> (Townsend), 1892c (<i>Eustomatodexia</i>), <i>Stomatodexia</i> ..	104.119

<i>intermedia</i> Townsend, 1927a, <i>Itasturmia</i>	104.190
<i>intermedia</i> Townsend, 1927a, <i>Iteuthelaira</i>	104.156
<i>intermedia</i> (Wulp), 1890 (Nemoraea), <i>Exopalpus</i>	104.55
<i>interrupta</i> Aldrich, 1927c, <i>Euantha</i>	104.111
<i>interrupta</i> (Aldrich), 1934 (Lispidea), <i>Elfia</i>	104.167
<i>interrupta</i> Curran, 1933, <i>Actia</i>	104.165
<i>interrupta</i> Curran, 1934b, <i>Calodexia</i>	104.154
<i>interstincta</i> Wulp, 1890 (Exorista)	104.198
<i>intritus</i> (Walker), 1860 (Nemoraea), <i>Archytas</i>	104.50
<i>intrusa</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>inuitata</i> Reinhard, 1964c, <i>Eleuthromyia</i>	104.82
<i>ipsea</i> Walker, 1849 (Tachina)	104.216
<i>Irengia</i> Townsend, 1935	104.23
<i>irregularis</i> (Wulp), 1890 (Brachycoma), <i>Drino</i>	104.189
<i>Irwinia</i> Cortés, 1967a	104.168
<i>Ischyrophaga</i> Townsend, 1915b	104.106
<i>ischyri</i> (Coquillett), 1905 (Thelairoides), <i>Ischyrophaga</i>	104.106
<i>Isodorus</i> Reinhard, 1954, <i>Prosenoides</i>	104.32
<i>Itabiomyia</i> Townsend, 1927a	104.137
<i>Itacnephalia</i> Townsend, 1927a	104.176
<i>Itacuphocera</i> Townsend, 1927a	104.78
<i>Italispidea</i> Townsend, 1927a	104.137
<i>Italydella</i> Townsend, 1927a	104.137
<i>Itamintho</i> Townsend, 1931a	104.96
<i>Itaplectopsis</i> Townsend, 1927a	104.137
<i>itaguaquecetubae</i> Townsend, 1913a, <i>Euthelothyria</i>	104.121
<i>itaguaquecetubae</i> (Townsend), 1929 (Argyrcomyia), <i>Leptostylum</i>	104.137
<i>itaguaquecetubae</i> (Townsend), 1929 (Jicaltepecia), <i>Spathimeigenia</i>	104.149
<i>Itarchytas</i> Blanchard, 1940	104.48
<i>Itasaundersia</i> Townsend, 1927a	104.67
<i>Itasturmia</i> Townsend, 1927a	104.189
<i>Itavoria</i> Townsend, 1931a	104.91
<i>Itaxanthomelana</i> Townsend, 1927a	104.8
<i>iterans</i> (Walker), 1849 (Tachina), <i>Peleteria</i>	104.44
<i>Iteuthelaira</i> Townsend, 1927a	104.156
<i>ivu</i> (Townsend), 1928b (Actinomlintho), <i>Polygaster</i>	104.94
<i>jaena</i> Townsend, 1919b, <i>Erythromelana</i>	104.106
<i>Jaenimyia</i> Townsend, 1912a	104.185
<i>jaennicki</i> (Giglio-Tos), 1894 (Saundersia), <i>Epalpus</i>	104.64
<i>Jalapamyia</i> Reinhard, 1964c	104.87
<i>Jamaicaria</i> , error	104.20
<i>jamaicensis</i> Curran, 1926a, <i>Lixinia</i>	104.145
<i>Jamacaria</i> Curran, 1928a	104.20
<i>Jaynesleskia</i> Townsend, 1934	104.117
<i>jayncsi</i> (Aldrich), 1924, (Leskiomima), <i>Jaynesleskia</i>	104.117
<i>Jicaltepecia</i> Townsend, 1917	104.148
<i>jicaltepecia</i> (Townsend), 1931a, (Eufurinia), <i>Jurinella</i>	104.57
<i>johnsoni</i> Coquillett, 1895d, <i>Lasionoura</i>	104.168
<i>jonesii</i> Walton, 1913b, <i>Eutrixoides</i>	104.23
<i>juyensis</i> (Blanchard), 1941 (Tachinosomopsis), <i>Jurinella</i>	104.57
<i>Jurinella</i> Brauer & Bergenstamm, 1889	104.56
<i>Jurinia</i> Robineau-Desvoidy, 1830	104.68



Juriniini	104.60
jurinioides (Townsend), 1895 (Blepharipeza), Leschenaultia ...	104.186
Juriniopsis Townsend, 1916d	104.58
juriniodes (Giglio-Tos), 1893 (Nemochaeta), Archytas	104.50
Jurinosoma Townsend, 1927a	104.68
Jurinodexia Townsend, 1915b	104.30
koehleri Blanchard, 1954, Gaediophanopsis	104.179
koheleri (Blanchard), 1941 (Pseudohystricia), Jurinella	104.57
koehleri Blanchard, 1940, Actinoplaga	104.86
kreibohmi (Blanchard), 1942 (Pronemorilla), Winthemia	104.194
kuscheli Cortés, 1952, Phantasiosiphona	104.169
labeosus Reinhard, 1957, Parepalpus	104.69
lacertosum (Wulp), 1891 (Prosenia), Mochlosoma	104.31
<i>Lachnommopsis</i> Townsend, 1915m	104.115
lacteata (Townsend), 1908, (Euomogenia), Homogenia	104.8
laetus (Wulp), 1892 (Saundersia), Xanthoepalpus	104.74
Lafuentemyia Marnet, 1965	104.98
laevigata (Wulp), 1890 (Rhinophora), Periscepsia	104.92
lagoae (Townsend), 1891c, (Exorista), Carcelia	104.202
lalandii Robineau-Desvoidy, 1830, Peleteria	104.44
lampuris Reinhard, 1953b, Juriniopsis	104.58
lanei Guimarães, 1961, Archytas	104.50
lanei Tavares, 1964, Ormiophasia	104.22
laniferae (Webber), 1930 (Achaetoneura), Lespesia	104.209
lanipes (Fabricius), 1805 (Thereva), Trichopoda, (Trichopoda)	104.9
Larvaevoridae	104.1
<i>Larvaevorinae</i>	104.39
<i>Larvaevorini</i>	104.43
Lasionera Coquillett, 1895d	104.168
Lasiopalpus Macquart, 1847	104.42
Lasiona Wulp, 1830	104.42
laspeyresiae Blanchard, 1940, Chaetolixophaga	104.129
lata Aldrich, 1928a, Belvosia	104.182
lata Curran, 1947, Jurinella	104.57
lata (Wiedemann), 1830 (Tachina), Lespesia	104.209
lateralis Curran, 1934d, Aridalia	104.202
lateralis Curran, 1934d, Chaetonalia	104.105
lateralis (Fabricius), 1805 (Musca), Opsozelia	104.102
lateralis (Macquart), 1843 (Jurinia), Archytas	104.50
lateralis Macquart, 1843 (Scopelia)	104.216
lateralis (Walker), 1837 (Sarcophaga), Ptilodexia	104.33
lateralis (Walker), 1849 (Trichopoda), Cylindrophasia	104.6
latevittata (Wulp), 1890 (Exorista), Winthemia	104.197
latevittata (Wulp), 1890 (Exorista), Winthemia	104.197
laticeps (Wulp), 1890 (Brachycoma), Atacta	104.180
laticornis Macquart, 1846, Jurinia	104.68
laticornis (Wulp), 1888, (Saundersia) Epalpus	104.64
<i>latifasciata</i> Blanchard in [Marchionatto], 1941, Myothyriopsis	104.190
latiforceps Aldrich, 1934, Psecacera	104.25
latifrons (Walker), 1852 (Tachina), Leschenaultia	104.186
latimana (Wulp), 1890 (Exorista), Winthemia	104.197



latipennis (Brauer), 1898 (Hyalomyia), Alophorella	104.11
latitarsus Thompson, 1964, Pseudochaeta	104.163
lativentris (Curran), 1934 (Microphthalma), Irengia	104.23
lativittus (Walker), 1852 (Tachina), Epalpus	104.64
<i>Latreillia</i> Robineau-Desvoidy, 1830	104.181
<i>Latreillimyia</i> Townsend, 1908	104.181
laudatum Reinhard, 1958c, Mochlosoma	104.31
lavinia Curran, 1934c, Thelairodes	104.95
laxa Curran, 1942, Hystricia	104.42
lenis Reinhard, 1955c, Tesseracephalus	104.35
lenkoi Guimarães, 1961a, Archytas	104.50
lenkoi Guimarães, 1966a, Neomasiphyia	104.191
Ienkoi Tavares, 1965b, Ormia	104.21
Ienkoi Tavares, 1965b, Ormia	104.21
leopoldiensis (Brauer), 1897 (Micropalpus), Empheremyia	104.63
lepida Robineau-Desvoidy, 1830 (Zenillia)	104.215
Leptidosophia Townsend, 1931a	104.112
<i>Leptoda</i> Wulp, 1885	104.102
Leptodexia Townsend, 1919a	104.30
Leptomacquartia Townsend, 1919a	104.106
Leptostylum Macquart, 1851	104.137
leptotrichopus (Brauer & Bergenstamm), 1891 (Sisyropa), Pam- maerus	104.211
leschenaulti (Robineau-Desvoidy), 1830 (Peleteria), Archytas ...	104.50
Leschenaultia Robineau-Desvoidy, 1830	104.185
Leskiolydella Townsend, 1927a	104.137
<i>Leskiopalpus</i> Townsend, 1916e	104.117
Leskiomima Brauer & Bergenstamm, 1891	104.117
Leskiini	104.116
Leskiopsis Townsend, 1916e	104.121
Lespesia Robineau-Desvoidy, 1863	104.208
leucocyla Wulp, 1890, (Degeeria)	104.152
leucomelanus (Walker), 1849 (Tachina), Epalpus	104.64
leuconotum (Wulp), 1892 (Exorista), Leptostylum	104.137
leucophaca (Wulp), 1890 (Hypostena), Ptilodegeeria	104.147
leucophrys (Wiedemann), 1830 (Tachina), Leschenaultia	104.186
leucopyga Wulp, 1882, Belvosia	104.182
Leucostoma Meigen, 1803	104.17
Leucostomatini	104.17
liciata Reinhard, 1964b, Helioprosopa	104.78
liebermanni Blanchard, 1940, Prosenactia	104.169
lilloi (Blanchard), 1954 (Willistonina), Belvosia	104.182
limbata (Blanchard), 1966c (Trichopodopsis), Trichopoda (Galactomyia)	104.10
limbata (Wiedemann), 1830 (Dexia), Zelia	104.102
limbisquamum Aldrich, 1934, Hyasoma	104.98
Lindigepalpus Townsend, 1931a	104.68
lindigi Townsend, 1931a, Ptilodeeria	104.147
Lindigia Townsend, 1931a	104.58
lindigii Bischof, 1904, Epalpus	104.64
linearis Townsend, 1915m, Calpodomyia	104.127
linearis Wulp, 1890 (Phorocera)	104.215
lineata (Blanchard), 1943b, (Prosteatosoma), Peleteria	104.44
lineata Macquart, 1851, Gonia	104.175



<i>lineata</i> Townsend, 1914, <i>Trichosaundersia</i>	104.73
<i>lineata</i> Townsend, 1912a, <i>Vibrissomyia</i>	104.81
<i>lineata</i> Townsend, 1934, <i>Trichodura</i>	104.103
<i>lineata</i> (Wulp), 1890 (<i>Mystacella</i>), <i>Chrysoexorista</i>	104.207
<i>lineatus</i> Townsend, 1914, <i>Epalpus</i>	104.64
<i>lineifrons</i> Sabrosky, 1953, <i>Ormia</i>	104.21
<i>linellii</i> Townsend, 1915m, <i>Pseudeuantha</i>	104.104
<i>linellii</i> Townsend, 1931, <i>Pseudobombyliomyia</i>	104.43
<i>lineolata</i> (Bigot), 1888a (<i>Epalpus</i>), <i>Vibrissomyia</i>	104.81
<i>Linnaemyini</i>	104.81
<i>lipernis</i> Reinhard, 1952b, <i>Chaetophlepsis</i>	104.89
<i>Lispideosoma</i> Reinhard, 1934c	104.168
<i>Littoralis</i> (Blanchard), 1963 (<i>Patellopsis</i>), <i>Patelloa</i>	104.211
<i>littoralis</i> Blanchard, 1943d, <i>Epiplagiops</i>	104.158
<i>litturata</i> (Olivier), 1812 (<i>Ocyptera</i>), <i>Euantha</i>	104.111
<i>liturata</i> Aldrich, 1934, <i>Lypha</i>	104.86
<i>liturata</i> , emed., <i>Euantha</i>	104.111
<i>Lixophaga</i> Townsend, 1908	104.138
<i>Lixinia</i> Curran, 1926a	104.138
<i>lloydii</i> Blanchard, 1947b, <i>Dasyuromyia</i>	104.97
<i>lloydii</i> Blanchard, 1966b, <i>Myiophasiomima</i>	104.22
<i>lluyi</i> Townsend, 1929, <i>Exoernestia</i>	104.84
<i>lobulatus</i> Curran, 1928b, <i>Archytas</i>	104.50
<i>longa</i> Rondani, 1850 (<i>Exorista</i>)	104.215
<i>longa</i> (Walker), 1852 (<i>Dexia</i>), <i>Orthaporia</i>	104.103
<i>longicornis</i> Aldrich, 1934, <i>Ganaproctus</i>	104.78
<i>longicornis</i> Aldrich, 1934, <i>Lypha</i>	104.86
<i>longicornis</i> (Bigot), 1889 (<i>Homodexia</i>), <i>Pseudodexia</i>	104.108
<i>longicornis</i> Macquart, 1843 (<i>Myiobia</i>)	104.120
<i>longicornis</i> Wulp, 1890 (<i>Brachycoma</i>)	104.204
<i>longipalpis</i> Wulp, 1883 (<i>Prosenia</i>)	104.39
<i>longipalpis</i> (Wulp), 1890 (<i>Myobia</i>), <i>Dejeniopalpus</i>	104.116
<i>longipes</i> Fabricius, 1805 (<i>Musca</i>), <i>Cholomyia</i>	104.105
<i>longipes</i> Thompson, 1968, <i>Cryptomeigenia</i>	104.130
<i>longipes</i> Townsend, 1919a, <i>Urochaetona</i>	104.109
<i>longipes</i> Wulp, 1890 (<i>Degeeria</i>)	104.152
<i>longipes</i> , authors, <i>Cholomyia</i>	104.106
<i>longipulvilli</i> Tothill, 1924, <i>Gonia</i>	104.175
<i>longiseta</i> (Wiedemann), 1830 (<i>Dexia</i>), <i>Chaetona</i>	104.105
<i>longitarsis</i> (Macquart), 1843 (<i>Prosenia</i>), <i>Stomatodexia</i>	104.119
<i>longitarsis</i> Wulp, 1893 (<i>Morinia</i>)	104.109
<i>longiuscula</i> (Walker), 1857 (<i>Masicera</i>), <i>Palpexorista</i>	104.159
<i>longula</i> Curran, 1934, <i>Corozalia</i>	104.130
<i>lopesi</i> Guimarães, 1966b, <i>Adejeania</i>	104.46
<i>lopesi</i> Guimarães, 1966a, <i>Prophasiopsis</i>	104.200
<i>lopesi</i> Guimarães, 1961b <i>Archytas</i>	104.50
<i>lopesi</i> Tavares, 1962, <i>Ormia</i>	104.21
<i>loriola</i> Reinhard, 1955c, <i>Sipholeskia</i>	104.118
<i>loxostegi</i> Blanchard, 1962, <i>Frontiniella</i>	104.208
<i>lucens</i> Reinhard, 1963, <i>Macromya</i>	104.42
<i>luctuosa</i> Cortés, 1951, <i>Ruiziella</i>	104.45
<i>luctuosa</i> (Wulp), 1890 (<i>Masicera</i>), <i>Xiphomyia</i>	104.152
<i>ludens</i> Walker, 1860 (<i>Echinomyia</i>), <i>Diaphanomyia</i>	104.62
<i>lugens</i> Wulp, 1890 (<i>Telothyria</i>)	104.152


lugubris (Blanchard), 1954 (Pseudobelvosia), Belvosia	104.182
lugubris James, 1945, Eucheirophaga	104.83
lugubris (Wulp), 1890 (Mystacella), Gaediophana	104.179
lutea Townsend, 1927a, Actinactia	104.165
lutea Townsend, 1935b, Trypetidomima	104.112
luteipennis Curran, 1934d, Schwarzalla	104.108
luteipennis Wiedemann, 1830, Trichopoda (Trichopoda)	104.9
luteola (Coquillett), 1900 (Belvosia), Triachora	104.185
lutescens Townsend, 1931a, Leptidosophia	104.112
lutzi Curran, 1947, Jurinella	104.57
lutzi Townsend, 1916g, Muscinotheralaira	104.141
lutzi Townsend, 1915c, Tropidodexia	104.35
Lydella Robineau-Desvoidy, 1830	104.210
<i>Lydella</i> , authors	104.126
<i>Lydellactia</i> Townsend, 1927a	104.124
<i>Lydellini</i>	104.204
Lydellohoughia Townsend, 1927a	104.138
Lydellothelaira Townsend, 1919a	104.138
Lydina Robineau-Desvoidy, 1830	104.87
Lydinolydella Townsend, 1927a	104.138
Lygaeomyia Aldrich, 1934	104.168
Lypha Robineau-Desvoidy, 1830	104.85
Lyphini	104.85
macarensis (Townsend), 1928c (Anazygosturmia), Drino	104.189
macella Reinhard, 1935b, Doryphorophaga	104.132
Machairomasicera Townsend, 1919a	104.139
macilenta (Wiedemann), 1830 (Tachina), Neomintho	104.156
macilenta (Wulp), 1890 (Prospheysa), Pelecotheca	104.157
macquarti Townsend, 1916, Pachymyia	104.37
Macquartiini	104.97
macra (Wulp), 1890 (Phorocera), Pelecotheca	104.157
macrocera (Bigot), 1888a (Echinomyia), Peleteria	104.44
macrocera Reinhard, 1964b, Helioprosopa	104.78
macrocera Townsend, 1927a, Mellanactia	104.139
<i>macrocera</i> , authors, Deopalpus	104.76
<i>macrocera</i> Wiedemann, of Townsend, Neocuphocera	104.79
macrocerus (Wiedemann), 1830 (Tachina), Deopalpus	104.76
Macrohoughia Townsend, 1927a	104.210
Macrohoughiopsis Townsend, 1927a	104.159
Macrojurinia Townsend, 1916h	104.59
<i>Macromeigenia</i> Brauer & Bergenstamm, 1891	104.190
Macrometopa Brauer & Bergenstamm, 1889	104.23
<i>macrophallae</i> Blanchard, in [Marchionatto], 1942 (Phyllopho- rocera), Nilea	104.210
Macromya Robineau-Desvoidy, 1830	104.42
Macromyini	104.39
macroptera Wulp, 1890, Anisia	104.125
macroptera (Wulp), 1891 (Rhynchodexia), Ptilodexia	104.33
maculata Vimmer & Soukup, 1940, Jurinella	104.57
maculifera (Bigot), 1889 (Prosenia), Stomatodexia	104.119
maculiventris (Brèthes), 1909 (Trichophora), Saundersiops	104.71
<i>maculiventris</i> Rondani, 1850b, Genea	104.117



maculus (Macquart), 1843 (Micropalpus), Exopalpus	104.64
madera (Townsend), 1915 (Ennyomma), Myiophasia	104.27
Madremyia Townsend, 1916g	104.210
madrensis (Townsend), 1915e (Euloewia), Myiophasia	104.27
magalhaesi Guimarães, 1966b, Adejeania	104.46
megellana Townsend, 1916g (Mesembriophyto), Dasyuromyia ...	104.97
maignicornis Wulp, 1890 (Degeeria), Doryphorophaga	104.132
major Aldrich, 1934, Notodytes	104.99
major (Bigot), 1889 (Rhamphinina), Ptilodexia	104.33
major Curran, 1934b, Calodexia	104.154
major (Curran), 1925e, (Jurinella), Exopalpus	104.55
major Curran, 1925e, Jurinella	104.57
major Curran, 1947, Protodejeania	104.60
major Thompson, 1963a, Floradalia	104.121
major Thompson, 1968, Opsoleskia	104.108
major (Townsend), 1927a (Geneopsis), Genea	104.116
major Townsend, 1911a (Almugmyia), Nomen nudum	104.28
majorina (Wulp), 1891 (Myothyria), Ceracia	104.123
majuscula Wulp, 1891, Calodexia	104.154
maldonadoi Blanchard, in Lloyd, 1951, Ceracia	104.123
Makasinocera Townsend, 1915m	104.48
Makasinocerops Townsend, 1935d	104.48
malleola (Bigot), 1884c (Sphyromyia), Peleteria	104.44
malloi Blanchard, 1962, Austrophytomyptera	104.165
manni Aldrich, 1928a, Belvosia	104.182
Manteomasiphya Guimarães, 1966a	104.198
manteophaga Guimarães, 1966a, Phasiopsis	104.200
Maracafula Townsend, 1939c	104.107
maracasi Thompson, 1968, Celatoria	104.128
maracasi Thompson, 1968, Metaphorocera	104.159
maracasi Thompson, 1968, Thelairodopsis	104.150
Maracasimya Thompson, 1968	104.139
marginalis (Brèthes), 1909 (Saundersia), Cryptopalpus	104.61
marginalis Curran, 1947, Adejeania	104.46
marginalis Townsend, 1935d, Myiodoriops	104.141
maris Townsend, 1929, Actinoprosopa	104.205
marmorata Townsend, 1927a, Macrohoughia	104.210
marmorata Townsend, 1915h, Peleteria	104.45
marmoratus (Townsend), 1915h (Pseudoarchytas), Archytas ...	104.51
marnefi Cortés, 1967b, Myiochaeta	104.91
marshalli Aldrich, 1931a, Chaetonodexodes	104.129
martinezi (Brèthes), 1917 (Euparafrontina), Lespesia	104.209
Masicerini	104.204
Masiphya Brauer & Bergenstamm, 1891	104.198
Masiphyini	104.198
Masiphyoidca Thompson, 1963d	104.199
Masipoda Brauer & Bergenstamm, 1889	104.196
Mastigiomyia Reinhard, 1964e	104.96
matamorosa Reinhard, 1951, Belvosia	104.182
mattoensis Townsend, 1939c, Miamimyops	104.172
mattoensis Townsend, 1939c, Monoestrogaster	104.107
Matucania Townsend, 1919a	104.98
matutina Townsend, 1919a, Topomeigenia	104.193
maura Townsend, 1932, Moreiria	104.186



<i>maura</i> Walker, 1837 (Tachina), Notomanes	104.99
<i>maura</i> Wulp, 1890 (Exorista)	104.193
<i>mayensis</i> Townsend, 1929, Diaphoropeza	104.114
<i>Mayodistichona</i> Townsend, 1928a	104.176
<i>Mayophorinia</i> Townsend, 1927a	104.139
<i>Medinophyto</i> Townsend, 1927a	104.106
<i>medeola</i> Reinhard, 1961, Thematheca	104.151
<i>media</i> Curran, 1947, Euhuascaraya	104.66
<i>media</i> Townsend, 1935d, Trinitodoria	104.213
<i>Meigenielloides</i> Townsend, 1919a	104.139
<i>medinops</i> (Townsend), 1934a (Microchaetonops), Microchaetona	104.107
<i>Megaparia</i> Wulp, 1891	104.23
<i>Megaprosopus</i> Macquart, 1843	104.23
<i>Megistogaster</i> Macquart, 1851	104.111
<i>Meigenelloides</i> Townsend, 1919a	104.139
<i>melaena</i> (Wulp), 1891, (Prosenia), Microchaetogyne	104.30
<i>melaleuca</i> Wiedemann, 1830 (Dexia), Scotiptera	104.35
<i>melaleuca</i> (Wiedemann), 1830 (Tachina), Sisyropa	104.204
<i>Melaleuca</i> Wulp, 1891	104.102
<i>melaleuca</i> Wulp, 1890 (Hypostena)	104.153
<i>melanaria</i> Wulp, 1896 (Calliphora)	104.39
<i>melanax</i> (Walker), 1849 (Tachina), Eudcjeania	104.53
<i>melanica</i> Townsend, 1914, Echinopyrrhosia	104.62
<i>melanoccps</i> Bigot, 1889 (Phorocera), Metadoria	104.140
<i>melanaria</i> Wulp, 1896 (Calliphora)	104.39
<i>melanax</i> (Walker), 1849 (Tachina), Eudejeania	104.53
<i>melanica</i> Townsend, 1914, Echinopyrrhosia	104.62
<i>melanoceps</i> Bigot, 1889 (Phorocera), Metadoria	104.140
<i>Melanocyptera</i> Townsend, 1927a	104.15
<i>Melanodoria</i> Townsend, 1927a	104.139
<i>Melanepalpellus</i> Townsend, 1914	104.68
<i>Melanepalpus</i> Townsend, 1927a	104.68
<i>Melanophora</i> Meigen, 1803	104.4
<i>Melanopforini</i>	104.4
<i>melanops</i> (Robineau-Desvoidy), 1830 (Trichopoda), Eutrichopoda	104.7
<i>Melanoroptcryx</i> Townsend, 1927a	104.140
<i>Melanoromintho</i> Townsend, 1935d	104.140
<i>Melanorophasia</i> Townsend, 1934	104.8
<i>melanosoma</i> (Wulp), 1892 (Cistogaster), Gymnoclytia	104.5
<i>melas</i> (Bigot), 1885 (Siphonimyia), Trichophora	104.72
<i>Meleterus</i> Aldrich, 1926e	104.91
<i>Melisoneurini</i>	104.28
<i>Mellachnus</i> Aldrich, 1934	104.88
<i>Mellanactia</i> Townsend, 1927a	104.139
<i>mellca</i> Giglio-Tos, 1893, Hystrichodexia	104.30
<i>mellea</i> (Macquart) in Brauer, 1898 (Hystricia), Epalpus	104.64
<i>mellicornis</i> Townsend, 1917b, Paraphasiopsis	104.200
<i>mellisquamma</i> Townsend, 1919a, Matucania	104.98
<i>mendes</i> (Townsend), 1940b (Zygofrontina), Lespesia	104.209
<i>mendes</i> Tavares, 1965b, Ormia	104.21
<i>mendocina</i> Blanchard in [Marchionatto], 1941, Bolohoughia	104.206
<i>mendocina</i> Blanchard, 1963, Euthyprosopiella	104.98
<i>Mericina</i> Curran, 1927c	104.82
<i>meridionale</i> (Townsend), 1915m (Siphopsalida), Leucostoma ...	104.18




meridionalis Thompson, 1968, Palpexorista	104.159
meridionalis (Townsend), 1912a (Eumyothyria), Cryptomeigenia	104.130
meridionalis (Townsend), 1917b (Zygosturmia), Drino	104.189
meridionalis (Townsend), 1912a (Dexodes), Eucelatoriopsis	104.134
meridionalis Townsend, 1929, Mesochaeta	104.140
meridionalis (Townsend), 1912a (Plagiops), Stomatomyia	104.160
merista Reinhard, 1953a, Paradidyma	104.115
mesacarrioni Blanchard, 1947a, Euanisia	104.133
Mesembrierigone Townsend, 1931a	104.84
Mesembrinormia Townsend, 1931a	104.20
Mesembriophyto Townsend, 1916g	104.97
Mesochaeta Brauer & Bergenstamm, 1891	104.140
Metadexia Coquillett, 1899	104.101
Metadoria Brauer & Bergenstamm, 1893	104.139
Metagonistylum Townsend, 1927a	104.210
Metagymnopsis Blanchard, 1943	104.78
metallica (Aldrich), 1934 (Hyalomya), Alophorella	104.12
metallica, authors, Juriniopsis	104.58
metallica Townsend, 1927a, Lydinolydella	104.139
metallica Townsend, 1908, Phosocephala	104.80
metallicus Curran, 1947, Rhachoepalpus	104.70
metallicus Curran, 1947, Saundersiops	104.71
metallicus Robineau-Desvoidy, 1830, Archytas	104.51
Metamyobia Townsend, 1927a	104.117
Metamyophasia Blanchard, 1966b	104.27
Metaphorocera Thompson, 1968	104.159
Metarrhinomyia Townsend, 1927a	104.140
metopia Bischof, 1904, Parachaetolyga	104.195
metopia Brauer & Bergenstamm, 1889, Myiopharus	104.142
Metopiops Townsend, 1912	104.163
Metopiopsis Vimmer & Soukup, 1940	104.217
Metopoaetia Townsend, 1927a	104.140
mexicana Aldrich, 1928a, Belvosia	104.182
mexicana Aldrich, 1931d, Spathimeigenia	104.149
mexicana Brauer & Bergenstamm, 1891, Gaediopsis	104.179
mexicana Brauer & Bergenstamm, 1889, Macrometopa	104.140
mexicana Brauer & Bergenstamm, 1893, Metadoria	104.140
mexicana (Brauer & Bergenstamm), 1893 (Microchira), Paradi- dyma	104.115
mexicana Curran, 1947, Jurinella	104.57
mexicana Giglio-Tos, 1893 (Plagia), Voria	104.93
mexicana Giglio-Tos, 1893 (Degeeria)	104.153
mexicana (Giglio-Tos), 1893 (Blepharipoda), Drino	104.189
mexicana Macquart, 1846, Trichopoda (Trichopòda)	104.9
mexicana (Robineau-Desvoidy), 1863 (Dejeania), Eudejeania ...	104.54
mexicana (Townsend), 1915m (Oestrogastropsis), Calodexia ...	104.154
mexicana Wulp, 1890, Gonia	104.175
mexicanum (Macquart), 1851 (Prosenia), Mochlosoma	104.31
Miamimyops Townsend, 1939c	104.172
micans (Wulp), 1888 (Hystricia), Bombyliopsis	104.40
micans (Wulp), 1883 (Alophora), Hyalomya	104.13
Microaporia Townsend, 1919a	104.96
Microchaetina Wulp, 1891	104.28



<i>Microchaetogyne</i> Townsend, 1931a	104.30
<i>Microchaetona</i> Townsend, 1919a	104.107
<i>Microchaetonops</i> Townsend, 1934a	104.107
<i>Microchira</i> Brauer & Bergenstamm, 1893	104.115
<i>Microgymnomma</i> Townsend, 1916h	104.79
<i>Microleskia</i> Thompson, 1968	104.117
<i>Micromasiphya</i> Townsend, 1934a	104.199
<i>Micronychiops</i> Townsend, 1915m	104.85
<i>micropalpus</i> (Curran), 1925e (<i>Ptilomyia</i>), <i>Comatacta</i>	104.120
<i>Microphthalma</i> Macquart, 1843	104.24
<i>Microphthalmia</i> , error	104.24
<i>microphthalmoides</i> Townsend, 1931a, <i>Morphodexia</i>	104.24
<i>Microsillus</i> Aldrich, 1926b	104.201
<i>Microplagia</i> Townsend, 1915	104.91
<i>Microtownsendia</i> Curran, 1934d	104.140
<i>Microtrichodes</i> Macquart, 1846	104.196
<i>Microtrichomma</i> Giglio-Tos, 1893	104.55
<i>Microtrichommodes</i> Townsend, 1927a	104.55
<i>milleri</i> Curran, 1947, <i>Jurinella</i>	104.57
<i>Milonius</i> Reinhard, 1955c	104.190
<i>mima</i> Reinhard, 1931a, <i>Winthemia</i>	104.197
<i>mima</i> Townsend, 1919b, <i>Nephochaetona</i>	104.94
<i>mima</i> Townsend, 1917b, <i>Opsophasiopteryx</i>	104.21
<i>mima</i> Townsend, 1915j, <i>Polistiopsis</i>	104.17
<i>mima</i> Townsend, 1935d, <i>Pronemorilla</i>	104.195
<i>mima</i> Townsend, 1931a, <i>Tipuloleskia</i>	104.119
<i>mimela</i> Reinhard, 1953a, <i>Epidexia</i>	104.132
<i>Mimologus</i> Reinhard, 1955c	104.190
<i>minense</i> Townsend, 1927a, <i>Metagonistylum</i>	104.210
<i>minima</i> Thompson, 1964, <i>Calocarcelia</i>	104.202
<i>minima</i> Wulp, 1890 (<i>Hypostena</i>)	104.153
<i>minor</i> (Curran), 1925e (<i>Jurinella</i>), <i>Exopalpus</i>	104.55
<i>minor</i> (Röder), 1885 (<i>Ocyptera</i>), <i>Cylindromyia</i>	104.16
<i>minor</i> Thompson, 1963b, <i>Eumacrohoughia</i>	104.208
<i>minor</i> Thompson, 1963a, <i>Floradalia</i>	104.121
<i>minor</i> Thompson, 1968, <i>Palpexorista</i>	104.159
<i>minor</i> Townsend, 1912a, <i>Euphorocera</i>	104.158
<i>Minthodexia</i> Brauer & Bergenstamm, 1891	104.96
<i>Minthodexiopsis</i> Townsend, 1927a	104.107
<i>Minthohoughia</i> Townsend, 1919a	104.95
<i>minthoidea</i> Townsend, 1931a, <i>Iconofrontina</i>	104.208
<i>Minthoini</i>	104.113
<i>Mintholeskia</i> Townsend, 1934	104.117
<i>Minthomima</i> Townsend, 1927a	104.94
<i>Minthomyia</i> Townsend, 1919a	104.41
<i>Minthopsis</i> Townsend, 1915m	104.141
<i>Minthoplagia</i> Townsend, 1915h	104.91
<i>Minthotachina</i> Townsend, 1935d	104.141
<i>minuscula</i> Townsend, 1934, <i>Melanorophasia</i>	104.8
<i>minuscula</i> (Wulp), 1891 (<i>Cordyligaster</i>), <i>Eucordyligaster</i>	104.112
<i>minuta</i> Curran, 1934a, <i>Galapagosis</i>	104.116
<i>minuta</i> Curran, 1947, <i>Jurinella</i>	104.57
<i>minuta</i> Wulp, 1890 (<i>Prospheerysa</i>)	104.64
<i>mira</i> Reinhard, 1958b, <i>Belvosia</i>	104.182

mirabile Townsend, 1911b, Ophirion	104.122
mirabilis Brauer & Bergenstamm, 1889, Argyromima	104.14
mirabilis Curran, 1934c, Heliæa	104.90
mirabilis Thompson, 1963d, Mystacomyoidea	104.109
mirabilis Townsend, 1912a, Metopiops	104.163
mirabilis (Townsend), 1940b (Xenopyxis), Neobrachelia	104.86
mirabilis Townsend, 1927a, Dolichopalpellus	104.167
miscella Townsend, 1935d, Minthotachina	104.141
misella Wulp, 1890 (Anisia)	104.215
misionensis (Blanchard), 1943b (Metagymnopsis), Gymnom- mopsis	104.78
misionensis (Blanchard), 1941 (Parafabricia), Archytas	104.51
mitis Curran, 1927a, Clausicellana	104.166
mitis Schiner in Brauer & Bergenstamm, 1891 (Trichophora)	104.72
mixta Townsend, 1935d, Myersimyia	104.159
mixta Townsend, 1927a, Myiosturmia	104.190
Mochlosoma Brauer & Bergenstamm, 1889	104.30
modesta Wulp, 1890, Didyma	104.131
modestus Bigot, 1857 (Eurigaster)	104.215
moerens (Wulp), 1892 (Hyalomyia), Alophorella	104.12
moestus (Wulp), 1890 (Didyma), Myiopharus	
montagna (Townsend), 1912a, (Blepharipeza), Leschenaultia ...	104.186
montana Townsend, 1929, Eucelatoria	104.133
Montserratia Thompson, 1964	104.203
mojingensis Arnaud, 1963b, Cryptocladocera	104.155
molinari Blanchard, in [Marchionatto], 1941, Podosturmia	104.191
monensis Curran, 1926a, (Phorocera)	104.193
monnula Reinhard, 1951, Gaediophana	104.179
monochroma (Wulp), 1890 (Degeeria), Oxynops	104.143
Monoestrogaster Townsend, 1939c	104.107
montana Reinhard, 1931a, Winthemia	104.197
montana (Wulp), 1891, (Dejeania), Protodejeania	104.60
montanus Aldrich, 1926c, Meleterus	104.91
montanus Townsend, 1912a, Chaetosisyrops	104.203
montivagus (Wulp), 1892 (Saundersia), Epalpus	104.64
morardi (Séguy), 1926b, Ormiophasia	104.22
Moreiria Townsend, 1932b	104.186
morionella Wulp, 1890, Anisia	104.125
Morphodexia Townsend, 1931a	104.24
moyobambensis Townsend, 1929, Platyrhinodexia	104.32
mucronata Townsend, 1927a, Ollachactia	104.143
mucorea Wulp, 1890, Anisia	104.125
multisetosa Wulp, 1890, Lasiona	104.42
multispinosa Thompson, 1963a, Peleteria	104.45
munda (Wulp), 1892 (Hyalomya), Heyneophasia	104.12
mundelli (Blanchard), 1935 (Epigrimyia), Epicoronimyia	104.167
murinus (Wulp), 1890 (Telothyria), Myiopharus	104.142
muscaeformis (Wulp), 1890 (Clista), Myiophasia	104.27
muscaria (Walker), 1852 (Dexia), Ptilodexia	104.33
muscaria Wulp, 1890, (Phorocera)	104.215
Muscinothelaira Townsend, 1916g	104.141
muscinum Wulp, 1890, Tyreomma	104.38
Muscopteryx Townsend, 1892	104.98
mutatus (Wiedemann), 1830 (Tachina), Xanthomelanodes	104.10



<i>myersi</i> (Aldrich), 1933b (Prophryno), <i>Ypophaemyiops</i>	104.214
<i>myersi</i> Townsend, 1939a, <i>Metagonistylum</i>	104.210
<i>Myersimyia</i> Townsend, 1935d	104.159
<i>Myobiopsis</i> Townsend, 1916e	104.117
<i>Myiochaeta</i> Cortés, 1967b	104.141
<i>Myiodoria</i> Townsend, 1927a	104.138
<i>Myiodoriopsis</i> Townsend, 1935d	104.141
<i>Myiomima</i> Brauer & Bergenstamm, 1889	104.31
<i>Myiomintho</i> Erauer & Bergenstamm, 1889	104.141
<i>Myiopharus</i> Brauer & Bergenstamm, 1889	104.141
<i>Myiophasia</i> Brauer & Bergenstamm, 1891	104.27
<i>Myiophasiini</i>	104.26
<i>Myiophasiomima</i> Blanchard, 1966b	104.27
<i>Myiophasiopsis</i> Townsend, 1927a	104.110
<i>Myioscotiptera</i> Giglio-Tos, 1893	104.31
<i>Myiosturmia</i> Townsend, 1927a	104.190
<i>Myiosturmiopsis</i> Thompson, 1963b	104.190
<i>Myioxynops</i> Townsend, 1927a	104.142
<i>Myothyria</i> Wulp, 1890	104.123
<i>Myothyriopsis</i> Townsend, 1919a	104.190
<i>Mystacomyia</i> Giglio-Tos, 1893	104.201
<i>Mystacomyoidea</i> Thompson, 1963d	104.199
<i>Myocuphocera</i> Townsend, 1931b	104.79
<i>Mystacella</i> Wulp, 1890	104.190
<i>mystacina</i> Aldrich, 1934, <i>Phaeodema</i>	104.110
<i>nana</i> (Townsend), 1915g (Odontocypthera), <i>Cylindromyia</i>	104.16
<i>nana</i> Townsend, 1927a, <i>Lydellohoughia</i>	104.138
<i>nana</i> (Townsend), 1934a, (Neoxynops), <i>Tachinophyto</i>	104.150
<i>nasuta</i> Townsend, 1927a, <i>Neosolieria</i>	104.85
<i>nattereri</i> Bischof, 1904, <i>Epalpus</i>	104.64
<i>nautiana</i> Townsend, 1908, <i>Comatacta</i>	104.120
<i>Neacroglossa</i> Townsend, 1927a	104.173
<i>Neaphria</i> Townsend, 1914	104.85
<i>Neaporia</i> Townsend, 1908	104.104
<i>nebulosa</i> Townsend, 1917b, <i>Euscopiopteryx</i>	104.19
<i>necopina</i> Walker, 1860 (Masicera)	104.216
<i>neglecta</i> Curran, 1925, <i>Peleteria</i>	104.44
<i>neglecta</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>negrensis</i> Aldrich, 1934 (Phorocera)	104.153
<i>nemo</i> (Curran), 1947 (Euhuascaraya), <i>Euempheremyia</i>	104.66
<i>Nemochaeta</i> Wulp, 1888	104.47
<i>nemochaetoides</i> Blanchard, 1943b, <i>Peleteria</i>	104.45
<i>memoralis</i> Brauer & Bergenstamm, <i>Alsopsyche</i>	104.198
<i>Nemorilla</i> Rondani, 1856	104.194
<i>memorina</i> Schiner in Brauer & Bergenstamm, 1889, <i>Eudexia</i> ...	104.29
<i>Neoarhytas</i> Townsend, 1915m	104.48
<i>Neoargyrophylax</i> Townsend, 1927a	104.142
<i>Neobelvosiopsis</i> Blanchard, 1954	104.181
<i>Neobrachelia</i> Townsend, 1931a	104.86
<i>Neobracheiliini</i>	104.86
<i>Neocampylochaeta</i> Townsend, 1927a	104.89
<i>Neochaetoplagia</i> Blanchard, 1963	104.91
<i>Neocraspedothrix</i> Townsend, 1927a	104.168



Neocuphocera Townsend, 1927a	104.79
Neocyrtophoeba Vimmer & Soukup, 1904	104.91
Neerigone Townsend, 1919a	104.82
Neoeuantha Townsend, 1931a	104.111
Neogymnomma Townsend, 1915e	104.69
Neojurinia Townsend, 1914	104.56
Neolophosia Townsend, 1939b	104.16
Neomasiphya Guimarães, 1966a	104.199
Neometachaeta Townsend, 1915m	104.115
neomexicana (Townsend), 1915e, (Ennyomma), Myiophasia ...	104.27
Neomintho Brauer & Bergenstamm, 1891	104.156
Neominthoidea Thompson, 1968	104.156
Neominthoini	104.155
Neominthopsis Townsend, 1915m	104.142
Neommasicera Townsend, 1927a	104.147
Neomyostoma Townsend, 1935d	104.31
Neopaedarium Blanchard, 1943b	104.91
Neophyllophila Townsend, 1927a	104.107
Neopodomyia Townsend, 1927a	104.191
neopolinosa Blanchard, 1966b, Xanthobasis	104.110
Neoprosena Townsend, 1927a	104.32
Neopsalida Townsend, 1916e	104.17
Neoptera Wulp, 1890	104.21
Neosolieria Townsend, 1927a	104.85
Neossarromyia Townsend, 1927a	104.79
Neotrafoia Townsend, 1912a	104.83
Neotrafoiopsis Townsend, 1931a	104.82
neotropica Bischof 1904, Ptilogonia	104.177
neotropica Townsend, 1927a, Neophyllophila	104.107
neotropica Townsend, 1927a, Neossarromyia	104.79
neotropicum (Curran), 1926a, (Voria), Paedarium	104.92
Neoxanthobasis Blanchard, 1966b	104.110
Neoxynops Townsend, 1934a	104.150
Neoxynopsoidca Thompson, 1968	104.142
Nephochaetona Townsend, 1919b	104.94
Nephoplaga Townsend, 1919	104.91
Nepocarcelia Townsend, 1927a	104.204
Nepophasmophaga Townsend, 1927a	104.172
nepos of Guimarães, Neocuphocera	104.79
nepos Townsend, 1927a, Neocuphocera	104.79
neptilucus (Wulp), 1892 (Jurinia), Archytas	104.51
nervosa (Walker), 1837 (Tachina), Dasyuromyia	104.97
nexa Reinhard, 1953b, Spatidexia	104.95
Nicetria Reinhard, 1953a	104.154
Nilea Robineau-Desvoidy, 1863	104.210
nitidiuscula (Wulp), 1890 (Degeeria), Doryphorophaga	104.132
nitidiventris Bigot, 1889 (Chaetolyga), Winthemia	104.198
nitidiventris Curran, 1933, Actia	104.165
nitidiventris Curran, 1928a, Melanophora	104.4
nigrifera Walker, 1852 (Tachina)	104.216
nigella Wulp, 1890, Anisia	104.125
niger (Townsend), 1914 (Eusignosoma), Epalpus	104.65
niger (Townsend), 1908 (Hemiargyra), Myiopharus	104.142
nigra Aldrich, 1934, Deloblepharis	104.87



nigra Aldrich, 1934, <i>Morphodexia</i>	104.24
nigra Bigot, 1889 (<i>Cestonia</i>)	104.153
nigra Bigot, 1885c, <i>Glossidionophora</i>	104.16
nigra Bigot, 1859, <i>Hystriisiphona</i>	104.30
nigra (Bigot), 1857 (<i>Scopolia</i>), <i>Periscepsia</i>	104.92
nigra Blanchard, 1966b, <i>Neoxanthobasis</i>	104.110
nigra Brauer & Bergenstamm, 1891, <i>Paradoria</i>	104.144
nigra Macquart, of Aldrich, 1926a, <i>Microphthalma</i>	104.24
nigra Macquart, 1843, <i>Trichophora</i>	104.72
nigra Townsend, 1915m, <i>Agicuphocera</i>	104.74
nigra (Townsend), 1914 (<i>Eublepharipeza</i>), <i>Bombyliomyia</i>	104.40
nigra Townsend, 1931a (<i>Sophia</i>), <i>Euantha</i>	104.111
nigra Townsend, 1912a, <i>Eudejeania</i>	104.54
nigra Townsend, 1914, <i>Euhystricia</i>	104.41
nigra Townsend, 1908, <i>Eutrichopoda</i>	104.7
nigra Vimmer & Soukup, 1940, <i>Andinomyia</i>	104.75
nigrens (Wulp), 1892 (<i>Hyalomya</i>), <i>Alophorella</i>	104.12
nigricalyptata (Macquart), 1885, (<i>Masicera</i>), <i>Parachaeta</i>	104.186
nigricans (Wulp), 1890 (<i>Degeeria</i>), <i>Celatoria</i>	104.128
nigricauda Bigot, 1876, <i>Trichopoda</i> (<i>Trichopoda</i>)	104.9
nigricaudum Blanchard, 1963, <i>Dolichostoma</i>	104.77
nigricauda Blanchard, 1966b, <i>Metamyiophasia</i>	104.27
nigricauda Thompson, 1966, <i>Drino</i>	104.189
nigricauda (Wulp), 1890 (<i>Exorista</i>), <i>Chrysoexorista</i>	104.207
nigriceps Aldrich, 1934, <i>Dasyuromyia</i>	104.97
nigripes Townsend, 1928c, <i>Brachymasicera</i>	104.202
nigripes Williston, 1908 (<i>Thelairoides</i>), <i>Cholomyia</i>	104.106
nigricolor Wulp, 1890, <i>Didyma</i>	104.131
nigricornis Blanchard, 1943b, <i>Acuphoceropsis</i>	104.74
nigricornis (Townsend), 1914, (<i>Epalpus</i>) <i>Saundersiops</i>	104.71
nigrifacies Bigot, 1889 (<i>Oplisa</i>), <i>Metadoria</i>	104.140
nigrifera Walker, 1852 (<i>Tachina</i>)	104.216
nigrifrons Aldrich, 1928a, <i>Belvosia</i>	104.182
nigrifrons Bigot, 1887 (<i>Elachipalpus</i>)	104.216
nigrifrons Bigot, 1889 (<i>Micropalpus</i>)	104.83
nigrifrons Thompson, 1968, <i>Gopaulia</i>	104.158
nigrifrons Wulp, 1890 (<i>Phorocera</i>)	104.161
nigrifrontalis (Blanchard), 1966c (<i>Trichopodopsis</i>), <i>Trichopoda</i> (<i>Galactomyia</i>)	104.10
nigrina (Wulp), 1883, (<i>Ocyptera</i>), <i>Cylindromyia</i>	104.16
nigripalpis Aldrich, 1934, <i>Aretosoma</i>	104.85
nigripalpis (Aldrich), 1932b (<i>Achaetoneura</i>), <i>Lespesia</i>	104.190
nigripalpis Aldrich, 1926c, <i>Opsothagus</i>	104.92
nigripalpis (Bigot), 1889 (<i>Chaetolyga</i>), <i>Eucelatoria</i>	104.133
nigripalpis Thompson, 1966, <i>Drino</i>	104.189
nigripalpis Thompson, 1968, <i>Eucelatorioidea</i>	104.133
nigripalpis Wulp, 1890 (<i>Brachycoma</i>)	104.194
nigripennis (Bigot), 1876 (<i>Trichopoda</i>), <i>Homogenia</i>	104.8
nigripennis (Wiedemann), 1830 (<i>Tachina</i>), <i>Talarocera</i>	104.80
nigripennis Wulp, 1891, <i>Comyops</i>	104.109
nigripes Thompson, 1968, <i>Calodexia</i>	104.154
nigripes Townsend, 1928c <i>Brachymasicera</i>	104.202
nigripes Wulp, 1892, <i>Trichopoda</i> , (<i>Trichopoda</i>)	104.9
nigripilosus Wulp, 1888 (<i>Saundersia</i>), <i>Parepalpus</i>	104.69




<i>nigrisquammis</i> (Townsend), 1892 (Blepharipeza), <i>Leschenaultia</i>	104.186
<i>nigrisquammis</i> Townsend, 1927a, <i>Melanodoria</i>	104.171
<i>nigrita</i> Thompson, 1968, <i>Admontia</i>	104.171
<i>nigrita</i> (Wulp), 1890 (Phorocera), <i>Paralispe</i>	104.145
<i>nigrithorax</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>nigriventris</i> Aldrich, 1934, <i>Steatosoma</i>	104.45
<i>nigriventris</i> (Macquart), 1843 (Hystricia), <i>Cryptopalpus</i>	104.61
<i>nigriventris</i> Robineau-Desvoidy, 1863, <i>Jurinia</i>	104.68
<i>nigriventris</i> Williston, 1896 (Degeeria)	104.153
<i>nigriventris</i> (Wulp), 1882, (Jurinia), <i>Archytas</i>	104.51
<i>nigrocalyptrata</i> (Macquart), 1855 (Masicera), <i>Parachaeta</i>	104.186
<i>nigrocalyptratus</i> (Macquart), 1846 (Jurinia), <i>Archytas</i>	104.51
<i>nigrocalyptratus</i> (Wulp), 1888 (Echinomyia), <i>Archytas</i>	104.51
<i>nigrocincta</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>nigrocostalis</i> Wulp, 1890 (Degeerla)	104.153
<i>nigroscutata</i> Rondani, 1863, <i>Hystricia</i>	104.41
<i>nigroscutellata</i> Wulp, 1892, <i>Homogenia</i>	104.8
<i>nigrothoracica</i> (Vimmer & Soukup), 1940 (Dejeania), <i>Adejeania</i>	104.46
<i>nigrotibiata</i> Curran, 1942, <i>Hystricia</i>	104.41
<i>Nimioglossa</i> Reinhard, 1945	104.31
<i>nimirum</i> Reinhard, 1956b, <i>Leucostoma</i>	104.18
<i>nitens</i> Aldrich, 1934, <i>Atelolglutus</i>	104.90
<i>nitens</i> Blanchard, 1966c, <i>Eutrichopodopsis</i>	104.7
<i>nitens</i> Curran, 1934d, <i>Microtownsendia</i>	104.140
<i>nitens</i> Curran, 1934d, <i>Zonalia</i>	104.104
<i>nitens</i> (Schiner), 1863 (Aporia), <i>Uramya</i>	104.104
<i>nitens</i> Thompson, 1964, <i>Pseudochaeta</i>	104.163
<i>nitens</i> Townsend, 1915, <i>Microplagia</i>	104.91
<i>nitens</i> Townsend, 1927a, <i>Prophaenopsis</i>	104.146
<i>nitens</i> Townsend, 1914, <i>Trichophoropsis</i>	104.80
<i>nitens</i> (Wiedemann), 1830 (Tachina), <i>Copecrypta</i>	104.75
<i>nitens</i> , authors, <i>Copecrypta</i>	104.76
<i>nitida</i> Curran, 1934c, <i>Oxynopsalia</i>	104.144
<i>nitidus</i> Townsend, 1914, <i>Rhachioepalpus</i>	104.70
<i>nitidifrons</i> Wulp, 1888, (Trichophora), <i>Copecrypta</i>	104.75
<i>nitidiventris</i> Bigot, 1889 (<i>Chaetolyga</i>)	104.198
<i>nitidiventris</i> Curran, 1933, <i>Actia</i>	104.165
<i>nitidiventris</i> (Curran), 1928b (Jurinia), <i>Juriniopsis</i>	104.58
<i>nitidiventris</i> Curran, 1928a, <i>Melanophora</i>	104.4
<i>nitidiventris</i> Macquart, 1851, <i>Prosopochaeta</i>	104.100
<i>nitidiventris</i> (Wulp), 1892 (Trichopoda), <i>Eutrichopoda</i>	104.7
<i>nitidiventris</i> Wulp, 1888, <i>Gymnomma</i>	104.67
<i>nitida</i> (Townsend), 1915d (Dimasicera), <i>Phaenopsis</i>	104.163
<i>nitidula</i> (Wulp), 1892, (Jurinia), <i>Juriniopsis</i>	104.58
<i>nitidula</i> Wulp, 1890, <i>Sphaerina</i>	104.149
<i>nitidus</i> (Macquart), 1851 (Micropalpus), <i>Epalpus</i>	104.65
<i>nitidus</i> (Wulp), 1892, (Nemochaeta), <i>Archytas</i>	104.51
<i>nitidiuscula</i> (Wulp), 1890 (Degeeria), <i>Doryphorophaga</i>	104.131
<i>niveifacies</i> Macquart, 1851 (Exorista)	104.215
<i>niveisquamma</i> Engel, 1920, <i>Jurinella</i>	104.57
<i>niveomarginata</i> (Wulp), 1890 (Anisia), <i>Spathidexia</i>	104.95
<i>niveus</i> Townsend, 1914, <i>Epalpus</i>	104.65
<i>niveus</i> Townsend, 1914, <i>Eulasiopalpus</i>	104.54
<i>nobilis</i> Williston, 1896 (Exorista)	104.215

<i>noctifer</i> Aldrich, 1934, <i>Lypha</i>	104.86
<i>nocturna</i> (Curran), 1934c (<i>Ormia</i>), <i>Euphasiapteryx</i>	104.21
<i>noera</i> (Reinhard), 1957 (<i>Phorocera</i>), <i>Euphorocera</i>	104.158
<i>nora</i> Curran, 1947, <i>Trichosaundersia</i>	104.73
<i>normula</i> Wulp, 1890 (<i>Masicera</i>)	104.204
<i>notabilis</i> Wulp, 1892, <i>Clinogaster</i>	104.15
<i>notata</i> (Bigot), 1888a (<i>Echinomyia</i>), <i>Exopalpus</i>	104.55
<i>notata</i> Cortés, 1967b, <i>Vibrissomyia</i>	104.81
<i>notata</i> Townsend, 1915m, <i>Eusaundersiops</i>	104.67
<i>notatus</i> Curran, 1947, <i>Rhachocpalpus</i>	104.70
<i>Notodytes</i> Aldrich, 1934	104.99
<i>Notomanes</i> Aldrich, 1934	104.99
<i>nova</i> Curran, 1947, <i>Rhachoeopalpus</i>	104.70
<i>nova</i> Townsend, 1927a, <i>Neocraspedothrix</i>	104.168
<i>novum</i> Giglio-Tos, 1893, <i>Gymnomma</i>	104.67
<i>nubecula</i> (Wulp), 1890 (<i>Telothyria</i>), <i>Sitophaga</i>	104.148
<i>nubilipennis</i> Wulp, 1891, <i>Acronacantha</i>	104.23
<i>nubilis</i> (Townsend) 1929 (<i>Pseudeuanthia</i>), <i>Uramya</i>	104.104
<i>nubilosa</i> Wulp, 1890 (<i>Hypostena</i>)	104.153
<i>nuda</i> Aldrich, 1934, <i>Incamyia</i>	104.136
<i>nuda</i> Thompson, 1963b, <i>Leschenaultia</i>	104.186
<i>nuda</i> Townsend, 1927a, <i>Eumacrohoughia</i>	104.207
<i>nudigaena</i> (Brauer), 1898 (<i>Jurinia</i>), <i>Archytas</i>	104.51
<i>nuditibia</i> Sabrosky, 1947, <i>Eudejeania</i>	104.54
<i>nyomula</i> (Townsend), 1914 (<i>Cordyligaster</i>), <i>Eucordyligaster</i> ...	104.112
<i>Oberonomyia</i> Reinhard, 1964	104.31
<i>obesa</i> Engel, 1920, <i>Hystricia</i>	104.41
<i>obesa</i> Townsend, 1928b, <i>Parajurinia</i>	104.159
<i>obesa</i> (Wiedemann), 1830 (<i>Tachina</i>), <i>Jurinella</i>	104.57
<i>obesula</i> (Wulp), 1890 (<i>Cnephalia</i>), <i>Belvosia</i>	104.182
<i>obesus</i> Coquillett, 1902, <i>Trixodes</i>	104.22
<i>obliteratus</i> Cortés, 1945a, <i>Chaetocraniopsis</i>	104.194
<i>obscura</i> Aldrich, 1931a, <i>Spathimeigenia</i>	104.149
<i>obscura</i> (Bigot), 1876 (<i>Trichopoda</i>), <i>Cylindromyia</i>	104.6
<i>obscura</i> (Bigot), 1876 (<i>Trichopoda</i>), <i>Cylindrophasia</i>	104.6
<i>obscura</i> (Bigot), 1889 (<i>Prosenia</i>), <i>Eudexia</i>	104.29
<i>obscura</i> Curran, 1947, <i>Euhuascaraya</i>	104.66
<i>obscura</i> (Séguy), 1926b, (<i>Plagiotormia</i>), <i>Ormiophasia</i>	104.22
<i>obscura</i> Thompson, 1968, <i>Braziliomyia</i>	104.128
<i>obscura</i> Thompson, 1965, <i>Sturmiopsoides</i>	104.192
<i>obscura</i> Townsend, 1914, <i>Hystriociopsis</i>	104.42
<i>obscura</i> (Walker), 1852 (<i>Dexia</i>), <i>Stomatodexia</i>	104.119
<i>obscurata</i> (Wulp), 1890 (<i>Exorista</i>), <i>Heliolydella</i>	104.135
<i>obscurella</i> (Wulp), 1890 (<i>Nemoraëa</i>), <i>Winthemia</i>	104.197
<i>obscurifrons</i> Wulp, 1890, <i>Anisla</i>	104.125
<i>obscuripes</i> Wulp, 1891, <i>Calodexia</i>	104.154
<i>obscurus</i> Bigot, 1857 (<i>Eurigaster</i>)	104.215
<i>obscurus</i> Townsend, 1914, <i>Eulasopalpus</i>	104.54
<i>obsoleta</i> Curran, 1925c, <i>Peleteria</i>	104.45
<i>obsoleta</i> (Wulp), 1890 (<i>Trixa</i>), <i>Microphthalma</i>	104.24
<i>obumbrata</i> Wulp of Brauer & Bergenstamm, <i>Ptilodegeeria</i> ...	104.147
<i>obumbrata</i> (Wulp), 1890 (<i>Hypostena</i>), <i>Euptilodegceria</i>	104.134
<i>occidentalis</i> Bigot, 1889, (<i>Chaetolyga</i>)	104.198


<i>occidentalis</i> (Coquillett), 1895a (<i>Drepanoglossa</i>), <i>Sipholeskia</i> ...	104.118
<i>occidentalis</i> Wiedemann, 1830, (<i>Tachina</i>)	104.216
<i>occidentalis</i> Wulp, 1892, <i>Admontia</i>	104.171
<i>occidentalis</i> Reinhard, 1931a, <i>Winthemia</i>	104.197
<i>occidua</i> (Walker), 1849 (<i>Gymnosoma</i>), <i>Gymnoclytia</i>	104.5
<i>occulta</i> (Wulp), 1890, (<i>Telothyria</i>), <i>Eucelatoriopsis</i>	104.134
<i>ocellaris</i> Townsend, 1927a, <i>Itacuphocera</i>	104.78
<i>ocellaris</i> Townsend, 1927a, <i>Orophrocera</i>	104.143
<i>ocellaris</i> (Townsend), 1912a, (<i>Protogonia</i>), <i>Protogoniops</i>	104.177
<i>ochracea</i> (Townsend), 1929 (<i>Eublepharipeza</i>), <i>Bombyliomyia</i> ...	104.39
<i>ochracea</i> Townsend, 1927a (<i>Tuberculocera</i>), <i>Dumerillia</i>	104.53
<i>ochracea</i> (Townsend), 1914, (<i>Eurythiopsis</i>), <i>Exopalpus</i>	104.55
<i>ochracea</i> (Wulp), 1890 (<i>Exorista</i>), <i>Chrysoexorista</i>	104.207
<i>ochraceus</i> (Townsend), 1914, <i>Epalpus</i>	104.65
<i>ochraceus</i> (Townsend), 1927a, <i>Ochroepalpus</i>	104.69
<i>ochricauda</i> (Wulp), 1890 (<i>Prosphegysa</i>), <i>Chaetogaedia</i>	104.178
<i>ochrescens</i> (Townsend), 1931a, (<i>Ocyptera</i>), <i>Cylindromyia</i>	104.16
<i>ochriceps</i> (Wulp), 1892 (<i>Cnephalia</i>), <i>Chaetogaedia</i>	104.178
<i>ochriceps</i> (Wulp), 1892 (<i>Hyalomya</i>), <i>Heyneophasia</i>	104.12
<i>ochricornis</i> (Bigot), 1888a, (<i>Epalpus</i>), <i>Deopalpus</i>	104.76
<i>ochricornis</i> (Townsend), 1916h, (<i>Euloewia</i>), <i>Myiophasia</i>	104.27
<i>ochrifrons</i> (Wulp), 1890 (<i>Telothyria</i>), <i>Adoryphorophaga</i>	104.124
<i>ochriceps</i> (Wulp), 1892 (<i>Cnephalia</i>), <i>Chaetogaedia</i>	104.178
<i>ochriceps</i> (Wulp), 1892 (<i>Hyalomya</i>), <i>Heyneophasia</i>	104.12
<i>Ochroepalpus</i> Townsend, 1927a	104.69
<i>octomaculata</i> Townsend, 1919a, <i>Pseudeuantha</i>	104.104
<i>oculata</i> Townsend, 1927a, <i>Xylocamptomima</i>	104.84
<i>oculatus</i> Curran, 1947, <i>Saundersiops</i>	104.71
<i>Ocyptera</i> , authors	104.15
<i>Ocypterosipha</i> Townsend, 1894	104.14
<i>Ocypteryx</i> Townsend, 1931a	104.15
<i>Ocyrtosoma</i> Townsend, 1912	104.37
<i>oeceticola</i> (Blanchard), 1963 (<i>Yahuarphryno</i>), <i>Patelloa</i>	104.211
<i>Oedemamedina</i> Townsend, 1927a	104.142
<i>Oedemapeza</i> Townsend, 1916f	104.172
<i>Oedematocera</i> Townsend, 1916f	104.172
<i>Oedematocerini</i>	104.172
<i>Oestrogaster</i> Townsend, 1912a	104.153
<i>Oestrogastroides</i> Townsend, 1915m	104.154
<i>Oestrogastropsis</i> Townsend, 1915m	104.153
<i>Oestrohystricia</i> Townsend, 1912	104.69
<i>oestroideus</i> Townsend, 1932, <i>Oligoestrus</i>	104.210
<i>Oestrophasia</i> Brauer & Bergenstamm, 1889	104.20
<i>Oestrophasiana</i> Townsend, 1931a	104.20
<i>Oestrophasiini</i>	104.19
<i>officialis</i> Townsend 1934, <i>Epaulophasia</i>	104.12
<i>okefenokeensis</i> Smith, 1916, <i>Winthemia</i>	104.197
<i>Okeopsis</i> Townsend, 1927a	104.196
<i>Ollachactia</i> Townsend, 1927a	104.143
<i>Ollachea</i> Townsend, 1919a	104.143
<i>Ollacheryphe</i> Townsend, 1927a	104.88
<i>Olenochaeta</i> Townsend, 1892a	104.176
<i>Oligoestrus</i> Townsend, 1932	104.216
<i>Olinda</i> Robineau-Desvoidy, 1830	104.217

olindoides Townsend, 1919a, Anametopochaeta	104.124
Olindopsis Townsend, 1927a	104.142
olivaceus Townsend, 1908, Rhachoepalpus	104.71
olivaurea Townsend, 1914, Jurinia	104.68
ollacheus Townsend, 1913e, Dejeniops	104.53
Oomasicera Townsend, 1916b	104.204
Oomeigenia Townsend, 1915n	104.143
omissa (Wulp), 1891 (Rhynchodexia), Ptilodexia	104.33
Ommaleskia Townsend, 1917b	104.107
onusta (Wulp), 1890 (Cnephalia), Spallanzania	104.178
opaca Wulp, 1890, Anisia	104.125
Ophirodexia Townsend, 1911b	104.101
Ophirion Townsend, 1911b	104.122
Ophirionini	104.122
Ophirionopsis Townsend, 1927a	104.122
Ophiosturmia Townsend, 1911b	104.210
ophthalmica Wulp, 1890, Anisia	104.125
opima Wulp, 1890 (Myobia)	104.120
opipare Reinhard, 1958c, Mochlosoma	104.31
Opsoempheria Townsend, 1927a	104.69
Oestropsis Townsend, 1912a	104.25
Opsoleskia Townsend, 1919a	104.108
Opsophagus Aldrich, 1926a	104.92
Opsophasiopteryx Townsend, 1917b	104.21
Opsosturmia Townsend, 1927a	104.191
Opsozelia Townsend, 1919a	104.101
Opticopteryx Townsend, 1931a	104.84
oppugnator Walton, 1914b (Compsilura), Eucelatoria	104.133
optica (Schiner), 1968 (Exorista), Eumasicera	104.162
oralis Townsend, 1927a, Neopodomyia	104.191
orbitalis Aldrich, 1934, Lypha	104.86
orbitalis (Aldrich), 1929a, (Copecryta), Neocuphocera	104.79
orbitalis Coquillett, 1904, Paradidyma	104.115
orbitalis (Curran), 1934c, (Sturmia), Drino	104.189
orbitalis Townsend, 1927a, Amicrotrichomma	104.47
orbitalis Townsend, 1927a, Camptophryno	104.206
orbitalis Townsend, 1961h, Chrysosturmia	104.188
orbitalis Townsend, 1927a, Clythoxynops	104.129
orbitalis Townsend, 1927a, Hypodoria	104.135
orbitalis Townsend, 1916h, Microgymnomma	104.79
orbitalis Townsend, 1927a, Verrugomyia	104.214
ordinaris (Wulp), 1890 (Exorista), Tinalydella	104.151
orellana Townsend, 1929, Calocarcelia	104.202
orientalis Townsend, 1927a, Cordillerodexia	104.29
orientalis (Townsend), 1929 (Plagiotachina), Euphorocera	104.158
Ormia Robineau-Desvoidy, 1830	104.21
Ormiini	104.20
Ormiophasia Townsend, 1919	104.22
ornata Aldrich, 1934, Lypha	104.86
ornatus Aldrich, 1926c, Opsophagus	104.92
ornata (Brauer & Bergenstamm), 1889 (Gymnostylia), Oxyaporía	104.143
ornata (Townsend), 1927a, (Chrysoerigone), Chrysotachina	104.81
ornata Townsend, 1927a, Oromasiphya	104.199
ornatipennis Engel, 1920, Engelomyia	104.41



ornatus (Macquart), 1843 (Micropalpus), Cryptopalpus	104.61
Orohoughia Townsend, 1934a	104.163
Oromasiphya Townsend, 1927a	104.199
Orophocera Townsend, 1927a	104.143
oroya Townsend, 1915f, Sorochemya	104.80
oroyensis Townsend, 1919a, Punamyocera	104.34
oroyensis Townsend, 1914, Vibrissomyia	104.81
ortalidoptera Townsend, 1939b, Shannonomyiella	104.19
Orthaporia Townsend, 1919b	104.103
Oswaldia Robineau-Desvoidy, 1863a	104.143
otiosa Wulp, 1890, Didyma	104.131
ovata Townsend, 1916h, Formicomya	104.77
ovata (Wulp), 1890 (Telothyria), Adoryphorophaga	104.124
ovipara Thompson, 1964, Montserratia	104.203
oviventris Townsend, 1931a, Atrichiopoda	104.6
Oxapampoepalpus Townsend, 1931a	104.69
Oxsnopsalia, misprint	104.144
Oxyaporia Townsend, 1919b	104.143
Oxydemia Bigot, 1885b	104.109
Oxydosphyria Townsend, 1926	104.43
Oxyepalpus Townsend, 1927a	104.69
Oxynops Townsend, 1912b	104.143
Oxynopsalia Curran, 1934c	104.144
Oxynopsis Townsend, 1927a	104.144
Oxyophirion Townsend, 1927a	104.144
pacata Wulp, 1890 (Telothyria)	104.153
pachecoi Curran, 1941, Eudejeania	104.54
pachecoi Curran, 1939, Macromya	104.39
pachecoi Curran, 1947, Protodejeania	104.60
pachecoi Curran, 1947, Saundersiops	104.71
Pachymyia Macquart, 1843	104.47
Pachynocera Townsend, 1919a	104.92
pacifica (Townsend), 1912a (Salmacia), Gonias	104.176
pactilis Reinhard, 1964b, Euhelioprosopa	104.77
Paedarium Aldrich, 1926c	104.92
pallens Reinhard, 1945, Trypheromyia	104.151
pallens authors, Gonias	104.175
pallens Wiedemann, 1830, Gonias	104.176
palliatum Curran, 1928b, Archytas	104.51
palliatum Wulp, 1891, Thelairodes	104.95
palliceps Bigot, 1888a, (Cryptopalpus), Rhynchoepalpus	104.71
pallida (Robineau-Desvoidy), 1863 (Dejeania), Eudejeania	104.54
pallidicornis (Bigot), 1889 (Anthracomyia), Pseudodexia	104.108
pallidipalpis Wulp, 1890, Anisia	104.125
pallidiventris Bigot, 1889 (Stevenia)	104.19
pallidula Wulp, 1890, (Brachycoma), Comatacta	104.120
pallipes (Macquart), 1843 (Dejeania), Eudejeania	104.54
pallitarsis Rondani, 1850b, Epalpus	104.65
pallmarum Townsend, 1934a, Prolypha	104.88
palpalis Aldrich, 1934, Morphodexia	104.24
palpalis (Aldrich), 1932b, (Zenillia), Palpozenillia	104.211
palpalis Curran, 1947, Adejeania	104.47

<i>palpalis</i> Curran, 1947, <i>Jurinella</i>	104.57
<i>palpalis</i> Reinhard, 1964, <i>Oberonomyia</i>	104.31
<i>palpalis</i> Townsend, 1928b, <i>Carceliodoria</i>	104.203
<i>palpalis</i> Townsend, 1927a, <i>Myioxynops</i>	104.142
<i>palpalis</i> Townsend, 1927a, <i>Paralispe</i>	104.145
<i>palpalis</i> Townsend, 1927a, <i>Plectopsis</i>	104.169
<i>palpalis</i> Townsend, 1927a, (<i>Okeopsis</i>), <i>Winthemia</i>	104.197
<i>Palpexorista</i> Townsend, 1926	104.159
<i>palpina</i> Rondani, 1850a, <i>Hystricia</i>	104.41
<i>Palpolinnaemyia</i> Townsend, 1927a	104.79
<i>palposa</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>Palpotachina</i> Townsend, 1915g	104.59
<i>Palpozenillia</i> Townsend, 1934	104.211
<i>palustrae</i> (Brèthes), 1908 (<i>Exorista</i>), <i>Nepocarcelia</i>	104.204
<i>Pammaerus</i> Aldrich, 1927c	104.211
<i>Panacemyia</i> Townsend, 1919b	104.144
<i>panamena</i> Curran, 1947, <i>Jurinella</i>	104.57
<i>panamensis</i> Curran, 1933, <i>Atacta</i>	104.165
<i>panamensis</i> Curran, 1939b, <i>Chrysotachina</i>	104.81
<i>panamensis</i> Townsend, 1934b, <i>Calodexia</i>	104.154
<i>panamensis</i> Townsend, 1919b, <i>Panacemyia</i>	104.144
<i>panamensis</i> Townsend, 1919a, <i>Parathelaira</i>	104.145
<i>panamensis</i> Townsend, 1919b, <i>Pelecotheca</i>	104.157
<i>papillo</i> Brauer & Bergenstamm, 1891, <i>Prosenoides</i>	104.32
<i>Parabelvosia</i> Blanchard, 1954	104.181
<i>Parabillaea</i> Blanchard, 1937	104.37
<i>Parabothria</i> Vimmer & Soukup, 1940	104.217
<i>Parabrachycoma</i> Blanchard, 1940	104.82
<i>Parachaeta</i> Coquillett, 1897	104.186
<i>Parachaetolyga</i> Bischof, 1904	104.195
<i>parancilla</i> Townsend, 1918, <i>Frontiniella</i>	104.208
<i>Paractinochaeta</i> Townsend, 1934a	104.113
<i>Paradejeania</i> Brauer & Bergenstamm, 1893	104.59
<i>Paradidyma</i> Brauer & Bergenstamm, 1891	104.115
<i>Paradionaea</i> Townsend, 1916e	104.17
<i>Paradoria</i> Brauer & Bergenstamm, 1891	104.144
<i>Paradomyops</i> Townsend, 1935d	104.94
<i>paradoxa</i> Townsend, 1931a, <i>Pirionimyia</i>	104.32
<i>paradoxicus</i> (Townsend) 1917b (<i>Paradomyia</i>), <i>Thelairodes</i>	104.95
<i>Paradrino</i> Mesnil, 1949b	104.191
<i>paracnsis</i> Townsend, 1934a, <i>Caenisomopsis</i>	104.127
<i>Parafabricia</i> Brauer & Bergenstamm, 1894	104.48
<i>Paragaedia</i> Brauer & Bergenstamm, 1891	104.183
<i>paraguayana</i> Townsend, 1928b, (<i>Anaphorinia</i>), <i>Chaetoxynops</i> ..	104.129
<i>paraguayensis</i> Townsend, 1928b, <i>Proparachaeta</i>	104.186
<i>paraguayensis</i> Townsend, 1928b, <i>Prospheysodoria</i>	104.146
<i>paraguayensis</i> (Townsend), 1928c (<i>Prowinthemia</i>), <i>Winthemia</i>	104.197
<i>Paragymnochaeta</i> Townsend, 1915g	104.81
<i>Paragymnomma</i> Brauer & Bergenstamm, 1891	104.72
<i>Parajurinia</i> Townsend, 1928b	104.59
<i>Parajurinia</i> Vimmer, 1940	104.217
<i>Paralispc</i> Brauer & Bergenstamm, 1891	104.144
<i>Paramesochaeta</i> Brauer & Bergenstamm, 1891	104.159



<i>paramonensis</i> Röder, 1886, <i>Gonatorrhina</i>	104.98
<i>Paramyocera</i> Townsend, 1915m	104.34
<i>Paranetia</i> Townsend, 1927a	104.145
<i>Paranoxylops</i> Thompson, 1968	104.145
<i>Paraphasiana</i> Townsend, 1940b	104.13
<i>Paraphasiopsis</i> Townsend, 1917b	104.199
<i>Paraphoranthia</i> Townsend, 1915b	104.13
<i>Parapoliops</i> Blanchard, 1957	104.168
<i>Paraporis</i> Townsend, 1912c	104.104
<i>Paraprospherysa</i> Townsend, 1927a	104.145
<i>Pararchytas</i> Brauer & Bergenstamm, 1894	104.59
<i>Pararrhinactia</i> Townsend, 1935d	104.163
<i>Parasetigena</i> Brauer & Bergenstamm, 1891	104.159
<i>Paratactopsis</i> Townsend, 1917b	104.184
<i>Parathelaira</i> Townsend, 1919a	104.145
<i>Paratheresia</i> Townsend, 1915e	104.37
<i>Paraxanthobasis</i> Blanchard, 1966b	104.110
<i>Parazelia</i> Townsend, 1919a	104.96
<i>Parcipromus</i> Reinhard, 1958b	104.85
<i>Parechinotachina</i> Townsend, 1931a	104.59
<i>Parepalpodes</i> Townsend, 1915m	104.72
<i>Parepalpus</i> Coquillett, 1902	104.69
<i>Parkerella</i> Townsend, 1942c	104.88
<i>parkeri</i> Sabrosky, 1952, <i>Eucelatoriopsis</i>	104.134
<i>parkeri</i> Townsend, 1941b, <i>Parthenoleskia</i>	104.118
<i>Parodomyia</i> Townsend, 1917b	104.95
<i>Parthenoleskia</i> Townsend, 1941b	104.118
<i>partita</i> Bigot, 1889 (<i>Dexiosoma</i>)	104.216
<i>parva</i> (Johnson), 1920, (<i>Plagia</i>), <i>Voria</i>	104.93
<i>parva</i> Townsend, 1916g, <i>Madremyia</i>	104.210
<i>parva</i> (Townsend), 1927a, (<i>Euhemilargyra</i>), <i>Paradoria</i>	104.144
<i>parva</i> Townsend, 1935d, <i>Pararrhinactia</i>	104.163
<i>parva</i> Townsend, 1942c, <i>Parkerella</i>	104.88
<i>parvicornis</i> Wulp, 1883 (<i>Dexia</i>)	104.122
<i>parvipalpis</i> (Wulp), 1890, (<i>Prospheerysa</i>), <i>Stomatomyia</i>	104.160
<i>parvipes</i> Townsend, 1891b, <i>Hypertrophocera</i>	104.115
<i>parvula</i> (Wulp), 1890, (<i>Phorocera</i>), <i>Metadoria</i>	104.140
<i>parvum</i> Aldrich, 1926e, <i>Paedarium</i>	104.92
<i>pastranai</i> Blanchard, 1963, <i>Neochaetoplagia</i>	104.91
<i>pastranai</i> Blanchard, in [Marchionatto], 1940, (<i>Xenosturmia</i>) ..	104.217
<i>patagonica</i> (Blanchard), 1963c (<i>Neowinthemia</i>), <i>Winthemia</i>	104.197
<i>Patelloa</i> Townsend, 1916e	104.211
<i>Patelloapsis</i> Townsend, 1927a	104.211
<i>patelloides</i> (Townsend), 1927a, (<i>Yahuarphryno</i>), <i>Patelloa</i>	104.211
<i>Pateloa</i> , Curran, 1934	104.211
<i>Patillalia</i> Curran, 1934d	104.211
<i>patula</i> (Walker), 1849 (<i>Tachina</i>), <i>Bombyliomyia</i>	104.40
<i>paulensis</i> (Townsend), 1927a, (<i>Proepalpus</i>), <i>Jurinia</i>	104.68
<i>paulensis</i> Townsend, 1927a, <i>Euempheremyia</i>	104.66
<i>paulensis</i> Townsend, 1927a, <i>Huascarayopsis</i>	104.67
<i>paulensis</i> (Townsend), 1927a (<i>Proepalpus</i>), <i>Jurinia</i>	104.68
<i>paulensis</i> Townsend, 1929, <i>Microgymnomma</i>	104.79
<i>paulensis</i> Townsend, 1929, <i>Pelecotheca</i>	104.157



paulista Townsend, 1929, <i>Gymnoclytia</i>	104.5
paulista Townsend, 1929, <i>Mesochaeta</i>	104.140
paulista Townsend, 1929, <i>Thysanomyia</i>	104.192
pavida Wulp, 1890, <i>Didyma</i>	104.131
pearsoni Guimarães, 1963d, <i>Archytas</i>	104.51
peccator Reinhard, 1956b, <i>Leucostoma</i>	104.18
pedalis Reinhard, 1953, <i>Microphthalma</i>	104.24
Pelecotheca Townsend, 1919b	104.156
Peleteria Robineau-Desvoidy, 1830	104.43
<i>Peleteropsis</i> Townsend, 1916c	104.43
<i>Peleteria</i> , emend	104.43
<i>Pelixia</i> Curran, 1934c	104.194
pellucens (Curran), 1925a (<i>Leskia</i>), <i>Genea</i>	104.116
pellucens Guimarães, 1966b, <i>Adejeania</i>	104.47
pellucida Robineau-Desvoidy, 1830 (<i>Sophia</i>), <i>Scotiptera</i>	104.35
pellucida (Séguy), 1925a, (<i>Ormia</i>), <i>Euphasiopteryx</i>	104.21
Pelycops Aldrich, 1934	104.99
penai Cortés, 1967, (<i>Enchomyia</i>), <i>Comops</i>	104.75
penaltis (Curran), 1934c, (<i>Myobia</i>), <i>Myobiopsis</i>	104.118
pendula Townsend, 1927a, <i>Perioptichochaeta</i>	104.79
<i>penicillata</i> Bigot, 1885c, <i>Dasyuromyia</i>	104.97
Pennapoda Wulp, 1892	104.3
pennipes (Fabricius), 1781 (<i>Musca</i>), <i>Trichopoda</i> , (<i>Trichopoda</i>)	104.9
Penthosia Wulp, 1892	104.16
peponis (Reinhard), 1953a (<i>Nicetria</i>), <i>Calodexia</i>	104.154
perarida Townsend, 1931a, <i>Phasmofrontina</i>	104.215
Peracroglossa Townsend, 1931a	104.177
percinerea Townsend, 1912a, <i>Aglummyia</i>	104.28
peregrina Wulp, 1890, <i>Anisia</i>	104.125
Perioptichochaeta Townsend, 1927a	104.79
Periscepsia Gistel, 1948	104.92
<i>pernambucensis</i> Townsend, 1939d, <i>Parasetigena</i>	104.38
pernox (Giglio-Tos), 1893 (<i>Nemochaeta</i>), <i>Archytas</i>	104.51
perorbitalis Townsend, 1927a, <i>Palpolinnaemyia</i>	104.79
perplexa (Townsend), 1931a, (<i>Parafabricia</i>), <i>Archytas</i>	104.51
perplexa Townsend, 1911b, <i>Pseudomyothyria</i>	104.147
perfecta (Walker), 1860 (<i>Dexia</i>), <i>Myobiopsis</i>	104.118
pertinax (Curran), 1934a (<i>Myobia</i>), <i>Myobiopsis</i>	104.118
pertyi Townsend, 1931a, <i>Mesembrinormia</i>	104.20
<i>Perua</i> Townsend, 1912a	104.24
peruana Arnaud, 1963d, <i>Borgmeiermyia</i>	104.106
peruana Curran, 1960, <i>Jurinopsis</i>	104.58
peruana Townsend, 1928c, <i>Argyrochaetona</i>	104.161
peruana Townsend, 1911b, <i>Diaphoropeza</i>	104.114
peruana (Townsend), 1928c (<i>Atrophopoda</i>), <i>Paradidyma</i>	104.115
peruana Townsend, 1929, <i>Stenostumia</i>	104.192
peruana Townsend, 1929, <i>Thysanomyia</i>	104.192
peruana (Townsend), 1911, (<i>Xanthomelana</i>), <i>Xanthomelanopsis</i>	104.11
peruanus Curran, 1928b, <i>Archytas</i>	104.51
Perumyia Arnaud, 1963c	104.177
peruviana (Brèthes), 1920 (<i>Melaleuca</i>), <i>Zelia</i>	104.102
peruviana Curran, 1939b, <i>Chrysotachina</i>	104.81
peruviana Townsend, 1931a, <i>Acaulona</i>	104.6

peruviana Townsend, 1911b, Diaphoropeza	104.114
peruviana Townsend, 1919b, Eunemorilla	104.189
peruviana Townsend, 1912 (Euphorocera)	104.161
peruviana (Townsend), 1927a, (Plagiotachina), Euphorocera ...	104.158
peruviana Townsend, 1912a (Salmacia), Gonia	104.176
peruviana (Townsend), 1927a, (Prophrynopsis), Incamyia	104.136
peruviana Townsend, 1928c, Olenochaeta	104.177
peruviana Townsend, 1928c, Paradidyma	104.115
peruviana Townsend, 1936e, Paraphoranthra	104.13
peruviana Townsend, 1931a, Peracroglossa	104.177
peruviana Townsend, 1927a, Pygophorinia	104.147
peruviana Townsend, 1929, Schizotachina	104.170
peruviana (Townsend), 1928c (Okea), Winthemia	104.192
peruvianum Townsend, 1928c, Leucostoma	104.18
peruvianum (Macquart), 1848 (Micropalpus), Epalpus	104.65
peruvianus Townsend, 1931a, Prospanipalpus	104.80
petentis Reinhard, 1958c, Muscopteryx	104.99
petiolata Thompson, 1963d Pseudomasiphya	104.200
petiolata Townsend, 1931a, Euanthoides	104.111
petiolata Townsend, 1919a, Pachynocera	104.92
petiolata Townsend, 1928c, Plagimasicera	104.212
petiolata Townsend, 1931a, Promegaparia	104.24
petiolata Townsend, 1931a, Pyrrhoernestia	104.82
petiolata (Townsend), 1919a, (Hypenomyia), Reinhardiana	104.28
petiolata Townsend, 1919a, Vibriosoovia	104.178
petiolata (Wiedemann), 1830 (Dexia), Cordyligaster	104.111
Petrargyrops Townsend, 1927a	104.212
<i>pfeifferi</i> Schiner in Brauer & Bergenstamm, 1897, Belvosia	104.182
Phaenopsis Townsend, 1912a	104.163
<i>Phaeodema</i> Aldrich, 1934	104.110
phaeoptera (Wiedemann), 1830 (Dexia), Yahuarmayoia	104.102
Phantasiosiphona Townsend, 1915h	104.169
phasiana (Townsend), 1897c, (Trichopoda), Pennapoda	104.8
Phasiatacta Townsend, 1911b	104.177
Phasiinae	104.4
Phasiini	104.11
Phasiocypthera Townsend, 1927a	104.16
Phasiophyto Townsend, 1919a	104.110
Phasiopsis Townsend, 1912b	104.200
<i>Phasiopteryx</i> Brauer & Bergenstamm, 1889	104.21
Phasiostoma Townsend, 1915g	104.169
Phasmofrontina Townsend, 1931a	104.212
phasmophagae Cortés, 1968a, Phasmovora	104.179
Phasmovora Cortés, 1968a	104.172
phillipii Rondani, 1863, Spathipalpus	104.113
<i>Phillophilopsis</i> , error	104.108
Phobetromyia Reinhard, 1964	104.43
Phoranthella Brooks, 1945b	104.14
<i>Phoranthella</i> Townsend, 1915b	104.14
<i>Phoranthella</i> Townsend, 1936a	104.14
<i>Phoriniini</i>	104.157
<i>Phorocerini</i>	104.157
Phosocephala Townsend, 1908	104.79
<i>Phosococephala</i> Townsend, 1936b	104.80




<i>Phosococephalops</i> Townsend, 1927a	104.175
<i>Phrissopolia</i> Townsend, 1908	104.178
<i>Phrynoiini</i>	104.204
<i>Phrynotachina</i> Townsend, 1927a	104.212
<i>Phryxe</i> Robineau-Desvoidy, 1830	104.212
<i>Phyllaristomyia</i> Townsend, 1931a	104.184
<i>Phyllolabella</i> Townsend, 1919b	104.39
<i>Phyllophila</i> Townsend, 1915b	104.108
<i>Phyllophilopsis</i> Townsend, 1915o	104.108
<i>Phyllophryno</i> Townsend, 1927a	104.145
<i>physonotae</i> Thompson, 1968, Eucelatoriodea	104.133
<i>piarconensis</i> Thompson, 1963a, Archytas	104.51
<i>pica</i> (Fabricius), 1805 (Musca), Rhamphinia	104.34
<i>pica</i> (Walker), 1852 (Tachina), Winthemia	104.197
<i>piceipes</i> (Wulp), 1892 (Hyalomyia), Alophorella	104.12
<i>piceiventris</i> (Walker), 1837 (Tachina), Drino	104.189
<i>piceus</i> (Giglio-Tos), 1893 (Saundersia), Epalpus	104.65
<i>pichilinguensis</i> Arnaud, 1963b, Cryptocladocera	104.155
<i>pickeli</i> Townsend, 1935d, Taperamyia	104.102
<i>picta</i> (Bigot), 1889 (Rhamphinia), Zelia	104.103
<i>picta</i> (Wulp), 1890 (Masicera), Myothyriopsis	104.190
<i>picticornis</i> Aldrich, 1934, Amblychaeta	104.87
<i>picticornis</i> Townsend, 1892, Chaetoglossa	104.179
<i>pictigaster</i> Bigot, 1889 (Ceromasia)	104.97
<i>pictipennis</i> Bigot, 1876, Trichopoda, (Trichopoda)	104.9
<i>pictipennis</i> Curran, 1941, Echinopyrrhosia	104.62
<i>pictipennis</i> Macquart, 1851 (Nemoraea)	104.216
<i>pictipennis</i> (Macquart), in Brauer & Bergenstamm, 1891, (Saundersia)	104.61
<i>pictipennis</i> (Townsend), 1934 (Procyanopsis), Deopalpus	104.76
<i>pictipennis</i> Townsend, 1935d, Thelairomima	104.96
<i>pictipennis</i> Wulp, 1891, Pseudomorinia	104.100
<i>pictiventris</i> Aldrich, 1934, Admontia	104.17
<i>Pictoepalpus</i> Townsend, 1915e	104.70
<i>pictus</i> (Schiner), 1868 (Saundersia), Epalpus	104.65
<i>piligera</i> Townsend, 1919a, Proroglutea	104.146
<i>pilipes</i> (Fabricius), 1805, (Thereva), Trichopoda, (Trichopoda)	104.9
<i>piliseta</i> (Wulp), 1890, (Masicera), Chaetona	104.105
<i>pilitarsis</i> Rondani, 1850 (Blepharipoda)	104.194
<i>pilitarsis</i> Rondani, 1850, Blepharopoda	104.36
<i>piliventris</i> authors, Archytas	104.51
<i>piliventris</i> Wulp, 1883 (Echinomyia), Archytas	104.50
<i>piliventris</i> Reinhard, 1934a, Paradidyma	104.115
<i>pilosa</i> Curran, 1941, Eudejeania	104.54
<i>pilosa</i> (Drury), 1770, (Musca), Hystriciclla	104.55
<i>pilosa</i> Drury of Curran, Archytas	104.51
<i>pilosus</i> (Walker), 1852, (Tachina), Archytas	104.51
<i>pilosa</i> (Wulp), 1890, (Hypostena)	104.153
<i>pinguoides</i> (Townsend), 1934a (Promasipoda), Winthemia	104.197
<i>pinguis</i> (Fabricius), 1805 (Musca), Winthemia	104.197
<i>piogaster</i> Pazos, 1914, Trichopoda (Trichopoda)	104.9
<i>Piriona</i> Aldrich, 1928b	104.99
<i>pirioni</i> Aldrich, 1934, Myiopharus	104.142
<i>pirioni</i> (Townsend), 1931a, (Dolichocyptra), Cylindromyia	104.16


Pirionimyia Townsend, 1931	104.32
piste (Walker), 1849 (Tachina), Nemorilla	104.195
piurana Townsend, 1911b, Belvosia	104.182
Piximactia Townsend, 1927a	104.145
placida Wulp, 1890, (Telothyria)	104.121
Plagia Meigen, 1838	104.93
Plagiatormia Ségué, 1926b	104.22
Plagimasicera Townsend, 1915m	104.212
plagioides Wulp, 1890 (Prospherysa)	104.215
Plagiomima Brauer & Bergenstamm, 1891	104.92
Plagiotachina Townsend, 1927a	104.157
Plagiotormia, error	104.22
Plagiprospherysa Townsend, 1892	104.160
plana Walker, 1852, (Dexia)	104.120
plangens Curran, 1928b, Archytas	104.51
planifrons Townsend, 1919a, Leptomacquartia	104.106
planifrons (Wulp), 1891 (Rhynchodexia), Ptilodexia	104.33
planiventris Macquart, 1851 (Nemoraëa)	104.216
Planomyia Aldrich, 1934	104.169
platensis (Brèthes), 1922 (Parasetigena), Euphorocera	104.158
Platyphasia Townsend, 1935d	104.8
platypoda Brauer & Bergenstamm, 1891, Pseudoviviania	104.169
Platyrhinodexia Townsend, 1927a	104.32
Plaxactia Townsend, 1931a	104.146
Plectopsis Townsend, 1927a	104.169
plena Aldrich, 1928b (Selenomyia), Psecacera	104.25
Plaxidexilla Blanchard, 1966d	104.27
Flexidexilla Blanchard, in Crouzel	104.27
plumata (Curran), 1925e, (Ptilomyia), Ptilomyiopsis	104.121
plumbea Townsend, 1927a, Zygozenillia	104.193
plumitarsis (Wulp), 1886 (Dejeania), Parechinotachina	104.59
plumosa (Wiedemann), 1830 (Dexia), Zelia	104.103
plumosula Townsend, 1927a, Chaquimayoiia	104.129
pluriseriata (Aldrich & Webber), 1924 (Phorocera), Patelloa	104.211
plusiae Coquillett, 1895c, Siphona	104.170
pluto Curran, 1934c, (Phorocera)	104.215
podicera Rondani, 1864 (Dejeania), Eudejeania	104.54
Podosturmia Townsend, 1928b	104.191
Poliophrys Townsend, 1908	104.179
Poliops Aldrich, 1934	104.169
Polistiopsis Townsend, 1915j	104.17
polita (Townsend), 1919a (Phrynolydella), Blondelia	104.127
polita Townsend, 1911b, Brachymasicera	104.202
polita Townsend, 1915m, Neometacheata	104.115
polita Townsend, 1919a, Pseudocelatoria	104.108
polita Townsend, 1927a, Prophasiopsis	104.200
polita Townsend, 1919, Pseudochaetona	104.108
politana (Townsend), 1911b (Blepharipa), Atactosturmia	104.180
Polybiophila Curran, 1937	104.122
Polygaster Wulp, 1890	104.94
Polygastropsis Townsend, 1919b	104.94
Polystomyia Townsend, 1908	104.9
politana Townsend, 1938c, Paraphoranthia	104.13
politus Townsend, 1927a, Prometopiops	104.212



<i>pollens</i> (Wulp), 1890 (Telothyria), Eucelatoria	104.133
<i>pollinosa</i> Aldrich, 1934, Xanthobasis	104.110
<i>pollinosa</i> Cortés, 1967a, Irwinia	104.168
<i>pollinosa</i> Curran, 1916, Myiophasia	104.27
<i>pollinosa</i> Thompson, 1966, Lespesia	104.209
<i>pollinosa</i> Thompson, 1968, Talparamyia	104.150
<i>pollinosa</i> Thompson, 1963, Winthemia	104.197
<i>pollinosa</i> (Wulp), 1888 (Hystricia), Jurinella	104.57
<i>pollinosa</i> Wulp of Townsend, (Eujurinia), Jurinella	104.57
<i>pollinosus</i> Curran, 1928b, Archytas	104.51
<i>pollinosus</i> Townsend, 1911b, Cnephelodes	104.174
<i>pollinosus</i> Townsend, 1912a (Siphosturmia), Microsillus	104.201
<i>polybia</i> (Curran), 1937 (Telothyriosoma), Ophironopsis	104.122
<i>Polybiophila</i> Curran, 1937	104.122
<i>porteri</i> (Brèthes), 1925 (Ocyptera), Cylindromyia	104.16
<i>porteri</i> Brèthes, 1918, Epalpus	104.65
<i>porteri</i> Brèthes, 1910 (Exorista)	104.215
<i>porteri</i> (Brèthes), 1920 (Parasetigena), Stomatotachina	104.160
<i>porteri</i> Reed, 1907 (Tachina)	104.216
<i>postera</i> Wulp, 1890 (Mystacella), Gaediopsis	104.179
<i>postica</i> Walker, 1861 (Eurigaster)	104.194
<i>potens</i> (Wiedemann), 1830 (Tachina), Belvosia	104.182
<i>potens</i> (Wiedemann), 1830 (Dexia), Zelia	104.103
<i>praeusta</i> (Wulp), 1891 (Rhynchodexia), Ptilodexia	104.33
<i>pretiosus</i> (Curran), 1934c (Cuphocera), Deopalpus	104.76
<i>pretiosa</i> (Curran), 1934d, Leptodexia	104.30
<i>prima</i> Townsend, 1915d, Cylindromasicera	104.162
<i>proboscidea</i> Townsend, 1931a, Pyrrhotachina	104.80
<i>Proarchytas</i> Townsend, 1931a	104.48
<i>Proarchytoides</i> Blanchard, 1941	104.48
<i>proбата</i> Walker, 1861 (Hylemyia)	104.216
<i>Procarcelia</i> Townsend, 1927a	104.195
<i>Procarcella</i> , misprint	104.145
<i>Procistogaster</i> Townsend, 1934	104.5
<i>Procleonice</i> Townsend, 1935d	104.99
<i>procteri</i> Curran, 1947, Jurinella	104.57
<i>Procyanopsis</i> Townsend, 1934	104.76
<i>Prodexodes</i> Townsend, 1927a	104.146
<i>prodigiosa</i> Bezzi, 1923, Cryptocladocera	104.155
<i>producta</i> Curran, 1947, Jurinella	104.58
<i>producta</i> Robineau-Desvoidy, 1830, Uramya	104.104
<i>productus</i> Curran, 1938b, Archytas	104.51
<i>Proepalpus</i> Townsend, 1927a	104.68
<i>profusa</i> Curran, 1947, Jurinella	104.58
<i>Proleskia</i> Townsend, 1927a	104.86
<i>Proleskiomima</i> Townsend, 1934	104.118
<i>prolixa</i> Townsend, 1935d, Procleonice	104.99
<i>Prolypa</i> Townsend, 1934a	104.88
<i>Promasiphya</i> Townsend, 1937a	104.200
<i>Promasipoda</i> Townsend, 1934a	104.196
<i>Promegaparia</i> Townsend, 1931a	104.24
<i>Prometopiops</i> Townsend, 1927a	104.212
<i>Promicrosillus</i> Townsend, 1935d	104.201
<i>prompta</i> Wulp, 1890, Didyma	104.131




Pronemorilla Townsend, 1935d	104.195
Proparachaeta Townsend, 1928b	104.186
<i>Proparachaetopsis</i> Blanchard, 1942a	104.186
Prophaenopsis Townsend, 1927a	104.146
Prophasiopsis Townsend, 1927a	104.200
Prophorinia Townsend, 1927a	104.146
Prophorostoma Townsend, 1927a	104.32
<i>Prophryno</i> Townsend, 1927a	104.208
<i>Prophrynopsis</i> Townsend, 1927a	104.136
propinqua (Wulp), 1892 (Cistogaster), <i>Gymnoclytia</i>	104.5
propusilla Sabrosky & Arnaud, 1965, <i>Cylindromyia</i>	104.16
Proroglutea Townsend, 1919a, <i>Proroglutea</i>	104.146
Prorynchops Brauer & Bergenstamm, 1891	104.32
Prosenactia Blanchard, 1940	104.169
Proseninae	104.19
Prosenini	104.28
Prosenoides Brauer & Bergenstamm, 1891	104.32
prosopina Brauer & Bergenstamm, 1891, <i>Sisyropa</i>	104.204
Prosopochaeta Macquart, 1851	104.99
Prospalaea Aldrich, 1925c	104.160
Prospanipalpus Townsend, 1931a	104.80
prospberyx Townsend, 1935d, <i>Hypophylax</i>	104.136
Prospberyxa Wulp, 1890	104.212
Prospberyssodoria Townsend, 1928b	104.146
<i>Prosteatosoma</i> Blanchard, 1943b	104.44
Protodejeania Townsend, 1915b	104.59
protoginoi (Blanchard), 1966a (<i>Achaetoneuropsis</i>), <i>Lespesia</i>	104.209
<i>Protoponia</i> Townsend, 1912a	104.177
Protoponiops Townsend, 1913f	104.177
Protoponiopsis Townsend, 1915m	104.184
Protonotodytes Blanchard, 1966b	104.28
<i>protoparcis</i> Townsend, 1892c (<i>Masicera</i>), <i>Drino</i>	104.189
Protrichoprosopis Blanchard, 1966b	104.25
Protypophaemyia Blanchard, 1963	104.195
<i>Prowinthemia</i> Townsend, 1928b	104.196
Proxanthobasis Blanchard, 1966b	104.110
proxima (Walker), 1852 (<i>Tachina</i>), <i>Belvosia</i>	104.182
<i>proximus</i> Townsend, 1927a (<i>Proxynops</i>), <i>Sphaerina</i>	104.149
<i>Proxynops</i> Townsend, 1927a	104.149
prudens Curran, 1928b, <i>Archytas</i>	104.51
prudens (Curran), 1934c (<i>Lydella</i>), <i>Blondelia</i>	104.127
pruinosis (Rondani), 1863 (<i>Cyphocera</i>), <i>Deopalpus</i>	104.77
<i>Psammopia</i> Townsend, 1915b	104.99
<i>Psecacera</i> Bigot, 1880	104.25
<i>Pseudeauantha</i> Townsend, 1915m	104.104
<i>Pseudoarchytas</i> Townsend, 1915h	104.48
<i>Pseudoarchytopsis</i> Townsend, 1927a	104.48
<i>Pseudobelvosia</i> Blanchard, 1954	104.181
<i>Pseudobombyliomyia</i> Townsend, 1931	104.43
<i>Pseudocelatoria</i> Thompson, 1968	104.146
<i>Pseudochaeta</i> Coquillett, 1895	104.163
<i>Pseudochaetona</i> Townsend, 1919a	104.108
<i>Pseudoclista</i> Brauer & Bergenstamm, 1893	104.27
pseudodaemon (Blanchard), 1940 (<i>Itarchytas</i>), <i>Archytas</i>	104.51



<i>Pseudodexia</i> Brauer & Bergenstamm, 1891	104.108
<i>Pseudohystiricia</i> Brauer & Bergenstamm, 1889	104.56
<i>pseudohystiricia</i> (Brauer & Bergenstamm), 1889 (<i>Hystirisiphona</i>), <i>Echinodexia</i>	104.29
<i>pseudomaculata</i> Blanchard, 1963, <i>Siphona</i>	104.170
<i>Pseudomasiphya</i> Thompson, 1963d	104.200
<i>Pseudomorinia</i> Wulp, 1891	104.100
<i>Pseudomyothyria</i> Townsend, 1892d	104.146
<i>Pseudoneoptera</i> Séguy, 1926b	104.22
<i>Pseudoprospherysa</i> Thompson, 1968	104.147
<i>pseudopyrrhopoda</i> (Blanchard), 1941 (<i>Eudejeaniops</i>), <i>Eudejeania</i> <i>Pseudoredtenbacheria</i> Brauer & Bergenstamm, 1889	104.54
<i>Pseudormia</i> Séguy, 1926c	104.147
<i>Pseudorrrhinactia</i> Thompson, 1968	104.22
<i>Pseudosiphosturmia</i> Thompson, 1966	104.92
<i>Pseudosturmia</i> Thompson, 1966	104.191
<i>Pseudoviviania</i> Brauer & Bergenstamm, 1891	104.191
<i>Pseudoxanthozona</i> Townsend, 1931a	104.169
<i>Pseudoxanthozonella</i> Townsend, 1931a	104.70
<i>Pterinopterna</i> Townsend, 1919a	104.70
<i>Pteropeza</i> Townsend, 1908	104.38
<i>Ptilodegeeria</i> Brauer & Bergenstamm, 1891	104.186
<i>Ptilodexia</i> Brauer & Bergenstamm, 1889	104.147
<i>ptilodexioides</i> Townsend, 1935d, <i>Neomyostoma</i>	104.33
<i>Ptilogonia</i> Bischof, 1904	104.31
<i>Ptilolydella</i> Townsend, 1927a	104.177
<i>Ptilomyioides</i> Curran, error or cmend.	104.124
<i>Ptilomyoides</i> Curran, 1928c	104.121
<i>Ptilomyia</i> Curran, 1925e	104.121
<i>Ptilomyiopsis</i> Townsend, 1933	104.121
<i>puer</i> (Williston), 1896 (<i>Phorocera</i>), <i>Stomatomyia</i>	104.160
<i>pueyrredoni</i> Brèthes, 1918b, <i>Hystrichodexia</i>	104.30
<i>pulchella</i> (Curran), 1934c (<i>Lydella</i>), <i>Blondelia</i>	104.127
<i>pulchellum</i> Macquart, 1851, <i>Leptostylum</i>	104.137
<i>pulchra</i> Townsend, 1919b, <i>Huascarodexia</i>	104.101
<i>pulchra</i> Townsend, 1911b, <i>Ophirodexia</i>	104.101
<i>pulchra</i> Townsend, 1919a, <i>Parazelia</i>	104.96
<i>pulchra</i> Wulp, 1891, <i>Euantha</i>	104.111
<i>pulchripes</i> (Aldrich), 1934 (<i>Cuphocera</i>), <i>Deopalpus</i>	104.77
<i>pulchrum</i> Townsend, 1927a, <i>Prophorostoma</i>	104.32
<i>pullata</i> Cortés, 1951, <i>Vibrissomyia</i>	104.81
<i>pulicaria</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>pullicaria</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>pullula</i> Wulp, 1890 (<i>Didyma</i>)	104.125
<i>pulverulentus</i> (Schiner), 1868 (<i>Saundersia</i>), <i>Rhachoeopalpus</i> ...	104.71
<i>pumila</i> Macquart, 1851 (<i>Echinomyia</i>), <i>Archytas</i>	104.49
<i>pumila</i> Robineau-Desvoidy, 1863 (<i>Peleteria</i>)	104.216
<i>pumila</i> (Wulp), 1890 (<i>Masicera</i>), <i>Lespesia</i>	104.209
<i>puna</i> Townsend, 1915d, <i>Dolichocnephalia</i>	104.174
<i>puna</i> Townsend, 1914, <i>Trichophoropsis</i>	104.80
<i>Punaclista</i> Townsend, 1915m	104.20
<i>Punamyocera</i> Townsend, 1919a	104.34
<i>Punamyia</i> Townsend, 1915m	104.80
<i>punctata</i> Robineau-Desvoidy, 1830, <i>Ormia</i>	104.21



<i>punctata</i> Robineau-Desvoidy, 1830 (Sophia), Scotiptera	104.35
<i>punctata</i> Townsend, 1912a, Jaenimyia	104.185
<i>punctata</i> Townsend, 1927a, Phasiocyptera	104.17
<i>punctata</i> Wulp, 1892 (Jurinia), Pararchytas	104.59
<i>punctiger</i> Townsend, 1927a, Petrargyrops	104.212
<i>punctigerum</i> Townsend, 1927a, Oxyophirion	104.144
<i>punctilucis</i> Townsend, 1927a, Hemiargyrophylax	104.135
<i>punctipennis</i> (Walker), 1857 (Sarcophaga), Paedarium	104.92
<i>punctipennis</i> (Wulp), 1891 (Rhynchodexia), Ptilodexia	104.33
<i>punctulata</i> Townsend, 1927a, Platyrhinodexia	104.32
<i>punensis</i> (Townsend), 1915m (Acemyiopsis), Ceracia	104.123
<i>punensis</i> Townsend, 1913e, Eudejeania	104.54
<i>punensis</i> Townsend, 1915h, Eutrichophora	104.77
<i>punensis</i> Townsend, 1914, Saundersiops	104.71
<i>punoensis</i> Vimmer & Soukup, 1940, Gaediopsis	104.179
<i>punoensis</i> Vimmer & Soukup, 1940, Parabothria	104.217
<i>puntarenensis</i> Townsend, 1928c (Erigonopsis), Dolichostoma ...	104.77
<i>purseglovei</i> Thompson, 1963a, Archytas	104.51
<i>purpurescens</i> Townsend, 1929, Argyrophylax	104.162
<i>purpurescens</i> Townsend, 1891a, Hyalomya	104.13
<i>purpurea</i> Curran, 1939b, Chrysotachina	104.81
<i>purpurea</i> Thompson, 1963a, Bombyliomyia	104.40
<i>purpurea</i> Townsend, 1927a, Chlorohystricia	104.40
<i>purpureus</i> Aldrich, 1928d, Daetaleus	104.25
<i>pygmae</i> (Macquart), 1851 (Echinomyia), Peleteria	104.45
<i>pygmaea</i> Townsend, 1927a, Sisypomyia	104.170
<i>pygmaea</i> Wulp, 1890 (Brachycoma)	104.180
<i>Pygophorinia</i> Townsend, 1927a	104.147
<i>pyrrhaspis</i> (Wiedemann), 1830 (Tachina), Macromya	104.42
<i>Pyrrhodexia</i> Townsend, 1931a	104.25
<i>Pyrrhoernestia</i> Townsend, 1931a	104.82
<i>pyrrhogaster</i> (Wiedemann), 1830, (Trichopoda) Eutrichopoda ...	104.7
<i>pyrrhopoda</i> Engel, 1920, Eudejeania	104.54
<i>pyrrhoprocta</i> (Wiedemann), 1830 (Dexia), Pyrrhodexia	104.25
<i>pyrrhopyga</i> (Wiedemann), 1819 (Tachina), Winthemia	104.197
<i>Pyrrhotachina</i> Townsend, 1931a	104.80
<i>pyste</i> (Walker), 1849 (Tachina), Nemorilla	104.195
 Quadratosoma Townsend, 1914	104.70
<i>quadrimalculata</i> Macquart, 1855 (Gymnostylia)	104.100
<i>quadrimalculata</i> (Macquart), 1846 (Aporia), Paraporia	104.104
<i>quadrimalculata</i> (Walker), 1852 (Dexia), Stomatodexia	104.119
<i>quadrissetosa</i> (Coquillett), 1902 (Gymnomma), Eufabriciopsis ...	104.66
<i>quadristriata</i> Wulp, 1890, (Hypostena)	104.153
<i>quadrivittata</i> Bigot, 1889 (Ceromasia)	104.122
<i>quadrus</i> (Wiedemann), 1830 (Tachina), Pammaerus	104.211
<i>quatuornotatus</i> (Townsend), 1935d, (Rhachoealpodes), Rhachoe-	
palpus	104.71
<i>quinquevittata</i> Blanchard, 1942a, Proparachaeta	104.186
 rachoui Tavares, 1962, Ormia	104.21
<i>radicis</i> (Townsend), 1916g (Andrina), Lydella	104.210
<i>rafaela</i> (Townsend), 1915h (Siphosturmiopsis), Siphosturmia ...	104.201




rafaela (Townsend), 1917a (Jicaltepecia), Spathimeigenia	104.149
rafaeli Townsend, 1913g, Chaetonodexodes	104.129
rafaeli Townsend, 1915h, Minthoplagia	104.91
ramata Townsend, 1927a, Mayoschizocera	104.168
rasilis (Wulp), 1890 (Telothyria), Lydinolydella	104.139
ratzeburgi (Jaennicke), 1867 (Demoticus), Deopalpus	104.77
rava Wulp, 1891 (Myocera)	104.39
rava Wulp, 1890 (Telothyria)	104.153
ravida Reinhard, 1945, Nimiloglossa	104.31
recincta Reinhard, 1964, Paradidyma	104.115
reclinata (Aldrich & Webber), 1924 (Zenillia), Carcelia	104.203
reclinata Townsend, 1919b, Binghamimyia	104.105
reclinata Townsend, 1927a, Conactia	104.130
reclinata Townsend, 1927a, Homohypochaeta	104.89
reclinata Townsend, 1927a, Hypohoughia	104.136
recondita Wulp, 1890 (Telothyria)	104.153
recta Schiner, 1868, Trichodura	104.103
recticornis (Macquart), 1855 (Gonia), Belvosia	104.182
rectinervis (Wulp), 1890, (Prospherysa), Lespesia	104.209
reducta Curran, 1947, Jurinella	104.58
<i>Reedia</i> Aldrich, 1928b	104.62
refuga (Wulp), 1890 (Telothyria), Argyrodoria	104.126
reinhardi Guimarães, 1963, Deopalpus	104.77
Reinhardiana Arnaud, 1952	104.28
reinhardtii (Wiedemann), 1830, (Tachina), Chlorohystricia	104.40
<i>reinhardtii</i> Wiedemann, authors, Chrysotachina	104.81
relicta Wulp, 1890 Telothyria	104.121
remissa Wulp, 1890, Anisia	104.125
remora Reinhard, 1953, Lixophaga	104.138
remota (Wulp), 1890 (Telothyria), Spathimeigenia	104.149
retrorsa Townsend, 1931a, Rhamphopteryx	104.84
Rhachoeopalpus Townsend, 1908	104.70
<i>Rhachoeopalpodes</i> Townsend, 1935d	104.70
Rhachosaundersia Townsend, 1931a	104.71
<i>Rhamphinina</i> , authors	104.34
Rhamphinina Bigot, 1885	104.34
Rhamphopteryx Townsend, 1931a	104.84
Rhinomacquartia Brauer & Bergenstamm, 1891	104.100
Rhinophorinae	104.5
<i>Rhodogyne</i> Meigen, 1800	104.90
rhois Townsend, 1917f, Eucyrtophloeoba	104.94
Rhombothyria Wulp, 1891	104.148
Rhombothyriopsis Townsend, 1915m	104.38
<i>rynchophorae</i> Blanchard, 1937 (Parabillaea), Paratheresia	104.33
<i>Rhynchodexia</i> , authors	104.33
<i>Rhynchodexia</i> , emend.	104.33
ricinorum Townsend, 1919a, Gymnocarcella	104.203
Ricosia Curran, 1927a	104.177
rigidirostris (Wulp), 1890, (Plagia), Plagiomima	104.92
<i>rimacensis</i> Townsend, 1915, Epalpodes	104.63
<i>rimacensis</i> (Townsend), 1915m (Parepalpodes), Trichophora ...	104.72
rivalis Reinhard, 1953a, Xiphomyia	104.152
robusta (Aldrich), 1928b, (Reedia), Edwynia	104.63
robusta Aldrich, 1934, (Achaetoneura)	104.194



<i>robusta</i> Aldrich, 1934, <i>Psecacca</i>	104.25
<i>robusta</i> Curran, 1925a, <i>Scotipecta</i>	104.35
<i>robusta</i> Thompson, 1968, <i>Pseudocelatoria</i>	104.146
<i>robusta</i> Townsend, 1919b (<i>Phyllolabella</i>), <i>Hypotachina</i>	104.39
<i>robusta</i> Townsend, 1927a, <i>Itasaundersia</i>	104.67
<i>robusta</i> (Wiedemann), 1830 (<i>Tachina</i>), <i>Peleteria</i>	104.45
<i>robusta</i> (Wulp), 1890 (<i>Brachycoma</i>), <i>Sitophaga</i>	104.148
<i>roederi</i> (Williston), 1893, (<i>Hystrihodexia</i>), <i>Euchaetogyne</i>	104.37
<i>roralis</i> (Linnaeus), 1758 (<i>Musca</i>), <i>Melanophora</i>	104.4
<i>rosenoi</i> Tavares, 1965a, <i>Euphasiopteryx</i>	104.21
<i>rostratus</i> Rondani, 1868, <i>Epalpus</i>	104.65
<i>rotundata</i> (Aldrich), 1934 (<i>Lispidea</i>), <i>Elfia</i>	104.167
<i>rozeni</i> Arnaud, 1963d, <i>Borgmeiermyia</i>	104.166
<i>rubentis</i> Reinhard, 1961 <i>Eugaediopsis</i>	104.179
<i>ruber</i> (Townsend), 1915m, (<i>Eucadratosoma</i>), <i>Cryptopalpus</i> ..	104.61
<i>ruber</i> Townsend, 1915m, <i>Eumelanepalpus</i>	104.66
<i>rubescens</i> Townsend, 1919a, <i>Sturmioidexia</i>	104.35
<i>rubicundus</i> (Wulp), 1892 (<i>Xanthomelana</i>), <i>Xanthomelanodes</i> ...	104.10
<i>rubida</i> Robineau-Desvoidy, 1830, <i>Dumerillia</i>	104.53
<i>rubidapex</i> Brauer, 1895 (<i>Chaetolyga</i>), <i>Winthemia</i>	104.196
<i>rubra</i> Vimmer & Soukup, 1940, <i>Winthemia</i>	104.197
<i>rubella</i> Robineau-Desvoidy, 1830, <i>Carcelia</i>	104.203
<i>rubicornis</i> (Wulp), 1891 (<i>Rhynchodexia</i>), <i>Ptilodexia</i>	104.33
<i>rubicornis</i> Thompson, 1968, <i>Pseudorrhinactia</i>	104.92
<i>rubicornis</i> (Wulp), 1891 (<i>Rhynchodexia</i>), <i>Ptilodexio</i>	104.33
<i>rubicornis</i> (Wulp), 1890 (<i>Exorista</i>), <i>Winthemia</i>	104.197
<i>rubrifrons</i> (Bigot), 1888a (<i>Echinomyia</i>), <i>Peleteria</i>	104.45
<i>rubripes</i> (Schiner) in Brauer & Bergenstamm, 1891 (<i>Saundersia</i>)	104.62
<i>rubripes</i> Thompson, 1968, <i>Calodexia</i>	104.154
<i>rubripes</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>rubripilla</i> (Rondani), 1850b (<i>Epalpus</i>), <i>Trichosaundersia</i>	104.73
<i>rubripilosa</i> (Wulp), 1888 (<i>Saundersia</i>), <i>Trichosaundersia</i>	104.73
<i>rubriventris</i> (Macquart), 1846 (<i>Dexia</i>) <i>Ptilodexia</i>	104.33
<i>rubriventris</i> (Wulp), 1890 (<i>Mystacella</i>), <i>Mystacomyia</i>	104.201
<i>rufa</i> (Schiner), 1868 (<i>Saundersia</i>), <i>Homosaundersia</i>	104.67
<i>rufa</i> Townsend, 1915e, <i>Neogymnonima</i>	104.69
<i>rufa</i> (Wulp), 1890 (<i>Neoptera</i>), <i>Ormia</i>	104.21
<i>rufata</i> (Bigot), 1889 (<i>Exorista</i>), <i>Bolomyia</i>	104.187
<i>rufescens</i> Aldrich, 1929b, <i>Gaediopsis</i>	104.179
<i>rufescens</i> Blanchard, 1966b, <i>Proxanthobasis</i>	104.110
<i>rufescens</i> Townsend, 1915m, <i>Anemorilla</i>	104.187
<i>rufianallis</i> (Wulp), 1891 (<i>Rhynchodexia</i>), <i>Ptilodexia</i>	104.33
<i>ruficauda</i> Blanchard, 1940, <i>Parabrachycoma</i>	104.82
<i>ruficauda</i> Curran, 1927a, <i>Mericina</i>	104.82
<i>ruficauda</i> Thompson, 1966, <i>Drino</i>	104.189
<i>ruficauda</i> Thompson, 1963b, <i>Tunapunia</i>	104.193
<i>ruficauda</i> Wulp, 1890 (<i>Brachycoma</i>)	104.199
<i>ruficauda</i> (Wulp), 1867 (<i>Schineria</i>), <i>Copecrypta</i>	104.75
<i>ruficaudus</i> Thompson, 1963d, <i>Microsillus</i>	104.201
<i>ruficorne</i> Macquart, 1851, <i>Gonistylum</i>	104.176
<i>ruficornis</i> Aldrich, 1934, <i>Ateloglutus</i>	104.90
<i>ruficornis</i> Aldrich, 1928a, <i>Belvosia</i>	104.182
<i>ruficornis</i> Aldrich, 1934, <i>Comops</i>	104.75
<i>ruficornis</i> (Blanchard), 1942b (<i>Okeopsis</i>), <i>Winthemia</i>	104.197


<i>ruficornis</i> (Thompson), 1869 (Exorista), <i>Winthemia</i>	104.195
<i>ruficornis</i> (Walker), 1852 (Tachina), <i>Crysotachina</i>	104.81
<i>ruficornis</i> (Wulp), 1892 (Cistogaster), <i>Gymnosoma</i>	104.5
<i>ruficoxa</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>rufifrons</i> Blanchard, 1954, <i>Belvosia</i>	104.182
<i>rufifrons</i> (Röder), 1885 (Frontina), <i>Lespesia</i>	104.209
<i>rufifrons</i> (Wulp), 1890 (Prosphecris), <i>Chaetogacdia</i>	104.178
<i>rufilabris</i> (Wulp), 1890 (Phorocera), <i>Euthelaira</i>	104.156
<i>rufilatera</i> (Rondani), 1850b, (Exorista), <i>Winthemia</i>	104.197
<i>rufina</i> Wulp, 1888, <i>Trichophora</i>	104.72
<i>rufipalpis</i> (Bigot), 1889 (Exorista), <i>Trafoia</i>	104.83
<i>rufipalpis</i> (Macquart), 1843, (Dejeania), <i>Adejeania</i>	104.47
<i>rufipalpis</i> Macquart, 1843, <i>Jurinia</i>	104.68
<i>rufipalpis</i> Macquart, 1843 (Blepharipeza), <i>Leschenaultia</i>	104.186
<i>rufipennis</i> (Macquart), 1846, (Micropalpus), <i>Epalpus</i>	104.65
<i>rufipes</i> Blanchard, 1966b, <i>Proxanthobasis</i>	104.110
<i>rufipes</i> Coquillett, 1902a, <i>Mochlosoma</i>	104.31
<i>rufipes</i> Curran, 1926a, <i>Atrophopalpus</i>	104.114
<i>rufipes</i> Jaennicke, 1867, (Micropalpus), <i>Epalpus</i>	104.64
<i>rufipes</i> Macquart, 1851, <i>Hystriola</i>	104.41
<i>rufipes</i> Wulp, 1892, <i>Homogenia</i>	104.8
<i>rufitibia</i> (Wulp), 1888 (Saundersia), <i>Epalpus</i>	104.65
<i>rufitorax</i> (Pazos), 1914 (Plagia)	104.93
<i>rufiventris</i> Aldrich, 1934, <i>Steatosoma</i>	104.45
<i>rufiventris</i> Brauer & Bergenstamm, 1891, <i>Sisyropa</i>	104.204
<i>rufiventris</i> (Macquart), 1846 (Micropalpus), <i>Cryptopalpus</i> ...	104.62
<i>rufiventris</i> (Macquart), 1843, (Jurinia), <i>Epalpus</i>	104.65
<i>rufiventris</i> Macquart, 1843, <i>Megaprosopus</i>	104.24
<i>rufiventris</i> Townsend, 1927a, <i>Archytoepalpus</i>	104.60
<i>rufiventris</i> Townsend, 1927a, <i>Hypoproxylops</i>	104.136
<i>rufiventris</i> Townsend, 1929, <i>Paratheresia</i>	104.37
<i>rufiventris</i> Townsend, 1927a, <i>Paradexodes</i>	104.146
<i>rufiventris</i> Townsend, 1929, <i>Sarcoprosena</i>	104.38
<i>rufiventris</i> Vimmer & Soukup, 1940, <i>Juriniella</i>	104.58
<i>rufohirta</i> Curran, 1942, <i>Hystriola</i>	104.41
<i>rufohirta</i> (Engel), 1920 (Bombyliomyia), <i>Bombyliopsis</i>	104.40
<i>rufomaculata</i> (Blanchard), 1963 (Rillella), <i>Lespesia</i>	104.209
<i>rufomaculata</i> Vimmer & Soukup, 1940, <i>Andinomyia</i>	104.75
<i>rufopalpus</i> Curran, 1926a, <i>Atrophopalpus</i>	104.114
<i>rufopilosa</i> (Wulp), 1898 (Saundersia), <i>Trichosaundersia</i>	104.72
<i>rufopygata</i> Bigot, 1889 (Viviania)	104.120
<i>rufostriata</i> Wulp, 1890, <i>Telothyria</i>	104.121
<i>rufostylata</i> (Bigot), 1887, (Frontina), <i>Chaetogacdia</i>	104.178
<i>rufum</i> (Brauer & Bergenstamm), 1893 (Cyrtosoma), <i>Ocyrtosoma</i>	104.37
<i>rufum</i> Townsend, 1914, <i>Quadratosoma</i>	104.70
<i>rufus</i> Townsend, 1927a, <i>Corpulentoepalpus</i>	104.61
<i>Ruiziella</i> Cortés, 1951	104.45
<i>ruralis</i> (Fallén), 1810 (Tachina), <i>Voria</i>	104.93
<i>russatus</i> Reinhard, 1962b, <i>Archytas</i>	104.51
<i>russulum</i> Reinhard, 1958c, <i>Mochlosoma</i>	104.31
<i>rusti</i> (Aldrich), 1929 (Phorocera), <i>Palpexorista</i>	104.159
<i>rusti</i> (Aldrich), 1929b (Phorocera), <i>Patelloa</i>	104.211
<i>rutilans</i> (Wulp), 1891 (Rhynchodexia), <i>Ptilodexia</i>	104.33
<i>rutilioides</i> (Jaennicke), 1867 (Dejeania), <i>Paradejeania</i>	104.59



sabroskyi (Arnaud), 1963b (Adercomyia), Pelecotheca	104.57
sabroskyi Guimarães, 1966b, Adejeania	104.47
sabroskyi Guimarães, 1963b, Archytas	104.51
sabroskyi Reinhard, 1958c, Mochlosoma	104.31
sabroskyi Tavares, 1965a, Ormia	104.22
saetigera Guimarães, 1966b, Adejeania	104.47
saginata Walker, 1861 (Eurigaster), Copecrypta	104.75
salla Curran, 1947, Jurinella	104.58
saltatrix Wiedemann, 1830 (Tachina)	104.216
Salmacia Meigen, 1800	104.175
sanctaerucis Thompson, 1963a, Archytas	104.52
sanctaerucis (Thompson), 1963a, (Leskiopalpus), Myobiopsis ..	104.118
sanctipauli Guimarães, 1966b, Adejeania	104.47
santacruz Thompson, 1968, Pseudoprospherysa	104.147
Santacruzia Thompson, 1968	104.148
Santacruziosis Thompson, 1968	104.164
sarcinale Reinhard, 1958c, Mochlosoma	104.31
Sarcocalirrhoe Townsend, 1928b	104.34
Sarcolydella Townsend, 1927a	104.213
sarcophagidea (Bigot), 1888b (Hyadesimyia), Dasyuromyia ...	104.97
sarcophagina Brauer & Bergenstamm, 1889, Myiomima	104.31
sarcophagina Wulp, 1883 (Prosenia)	104.39
Sarcoprosena Townsend, 1927a	104.38
satanica (Bigot), 1889, (Scopolia) Penthosia	104.17
Saundersia Schiner, 1868	104.61
saundersii (Williston), 1889 (Phorocera), Madremyia	104.210
Saundersiopmima Townsend, 1927a	104.56
Saundersiops Townsend, 1914	104.71
saxatilis Reinhard, 1952a, Phasiostoma	104.169
Schistocercophaga Townsend, 1928a	104.172
Schistostephana Townsend, 1919a	104.38
Schizocerophaga Townsend, 1916	104.148
Schizotachina Walker, 1852	104.170
Schlingermiya Cortés, 1967b	104.100
Schwarzalia Curran, 1934d	104.108
schwarzi Curran, 1947, Jurinella	104.58
schwarzi Curran, 1947, Saundersiops	104.71
scintilla Reinhard, 1953, Lixophaga	104.138
scita Walker, 1852 (Tachina)	104.153
Scopolia Robineau-Desvoidy, 1830	104.92
scordalus Reinhard, 1955c, Milonius	104.190
Scotipectera Macquart, 1835	104.34
scurra (Wulp), 1890 (Myobia), Calodexia	104.154
scutellaris Aldrich, 1934, Xanthopelta	104.88
scutellaris Robineau-Desvoidy, 1830, Harrisia	104.185
scutellaris (Robineau-Desvoidy), 1830 (Jurinia), Xanthozona ...	104.74
scutellaris Townsend, 1927a, Thysanosturmia	104.193
scutellaris (Wulp), 1890, (Phorocera), Euphorocera	104.158
scutellatus, authors, Archytas	104.51
scutellatus (Macquart), 1843, (Jurinia), Archytas	104.52
scutellata Brêthes, 1909, Andesimyia	104.178
scutellata (Wulp), 1891, (Rhynchodexia), Ptilodexia	104.33
seabrai Guimarães, 1961b, Archytas	104.52
Selenomyia Brauer & Bergenstamm, 1891	104.97



sellersi (Thompson), 1966 (Myxexoristopsis), Aplomya	104.205
semiater Bischof, 1904, Epalpus	104.65
semiater (Schiner), 1968 (Saundersia), Cryptopalpus	104.62
semiflava Aldrich, 1928a, Belvosia	104.182
semiflavus (Bischof), 1904 (Epalpus), Xanthoepalpus	104.74
semiflavus Bischof, 1904, Epalpus	104.65
seminiger (Wiedemann), 1830 (Tachina), Archytas	104.52
semipicta (Walker), 1852 (Dexia), Xanthodexia	104.96
semirufa (Blanchard), 1943b, (Aphriosphyria) Peleteria	104.45
semirufa (Wulp), 1891, (Leptoda), Zelia	104.103
sensifera Walker, 1861, Musca	104.216
sequax Williston, 1887, Gonia	104.176
sequens (Townsend), 1892a (Vanderwulpia), Catemophryx	104.114
sericea Townsend, 1928b, Sericodoria	104.213
sericea Wiedemann, of Wulp, Xanthodexia	104.96
sericea (Wiedemann), 1830 (Tachina), Xanthodexia	104.96
sericeum Giglio-Tos, 1893, Mochlosoma	104.31
Sericodoria Townsend, 1928b	104.213
sermyla (Walker), 1849 (Dexia), Uramya	104.104
serotina (Wulp), 1890 (Prospheya), Doryphorophaga	104.132
serotinum Reinhard, 1958c, Mochlosoma	104.31
serrei Séguy, 1925, Ormia	104.22
sesquipleux Giglio-Tos, 1893, (Masicera)	104.216
setifacies Brauer & Bergenstamm, 1891, Cryptomeigenia	104.130
setifacies Curran, 1928b, Archytas	104.52
setifera Townsend, 1915m, Exoristopsis	104.37
setifrons (Wulp), 1890 (Plagia), Minthoplagia	104.91
setigena Curran, 1927a, Ricosia	104.177
setigera Wulp, 1890 (Phorocera)	104.161
setipennis Townsend, 1919b, Gymnopalpus	104.94
setiventris (Wulp), 1890 (Macquartia), Uramya	104.104
Setolestes Aldrich, 1934	104.100
setosa (Coquillett), 1897, (Gaediopsis), Eugaediopsis	104.179
setosa Curran, 1925c, Peleteria	104.45
setosa (Townsend), 1915m, (Punaclista), Prosopochaeta	104.100
setosa Townsend, 1917b (Paratactopsis), Chaeteprosopa	104.184
setosa Townsend, 1915g, Hesperophasia	104.93
setosa Townsend, 1914, Xenoplagia	104.93
setosula (Wulp), 1890, (Mystacella), Trafoia	104.83
setosa (Wiedemann), 1830 (Tachina), Euloewiopsis	104.26
sexualis Curran, 1927c, Winthemia	104.97
sexualis Curran, 1934d, Agrarialia	104.161
shannoni Aldrich, 1933a, Tarassus	104.11
shannoni (Aldrich), 1934 (Metopomuscopteryx), Alexogloblinia	104.97
shannoni Guimarães, 1966, Archytas	104.52
shannoni Townsend, 1939d, Neolophosia	104.16
Shannoniella Townsend, 1939b	104.112
Shannoniellini	104.112
Shannonomyiella Townsend, 1939b	104.19
Shermanalia Curran, 1934d	104.30
sibilans Curran, 1928b, Archytas	104.52
sierricola (Townsend), 1908 (Poliophrys), Gaediopsis	104.180
sierricola (Townsend), 1917 (Ennyomma), Myiophasia	104.27



<i>sicricola</i> Townsend, 1926, <i>Visayalydina</i>	104.88
<i>sicsta</i> Curran, 1947, <i>Euhuascaraya</i>	104.66
<i>siestua</i> Curran, 1947, <i>Saundersiops</i>	104.71
<i>signata</i> Reinhard, 1931a, <i>Winthemia</i>	104.197
<i>signata</i> (Townsend), 1915h (<i>Apinocyptera</i>), <i>Cylindromyia</i>	104.16
<i>signata</i> (Walker), 1861 (<i>Eurigaster</i>), <i>Voria</i>	104.93
<i>signata</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>signatipennis</i> (Wulp), 1892 (<i>Ocyptera</i>), <i>Cylindromyia</i>	104.16
<i>signifera</i> Townsend, 1915d, <i>Paratheresia</i>	104.37
<i>signiferum</i> Wulp, 1890, <i>Cenosoma</i>	104.20
<i>Signoepalpus</i> Townsend, 1931	104.71
<i>Signosoma</i> Townsend, 1914	104.71
<i>Signosomopsis</i> Townsend, 1914	104.72
<i>sila</i> (Reinhard), 1958b (<i>Parcipromus</i>), <i>Neosolieria</i>	104.85
<i>simile</i> Townsend, 1914, <i>Eucorpulentosoma</i>	104.65
<i>similigena</i> Wulp, 1891, <i>Stomatodexia</i>	104.119
<i>similis</i> Blanchard, 1966c, <i>Eutrichopodopsis</i>	104.7
<i>similis</i> Thompson, 1968, <i>Pseudoprospherysa</i>	104.147
<i>similis</i> (Townsend), 1915m (<i>Oestrogastrodes</i>), <i>Calodexia</i>	104.154
<i>similis</i> (Townsend), 1928c (<i>Valpogonia</i>), <i>Chaetocraniopsis</i>	104.174
<i>similis</i> Townsend, 1927a, <i>Macrohoughiopsis</i>	104.159
<i>similis</i> Townsend, 1919b, <i>Orthaporia</i>	104.103
<i>similis</i> Townsend, 1915g, <i>Palpotachina</i>	104.59
<i>similis</i> Townsend, 1914, <i>Parepalpus</i>	104.69
<i>similis</i> (Townsend), 1927a, (<i>Patelloopsis</i>), <i>Patelloa</i>	104.211
<i>similis</i> (Townsend), 1927a (<i>Sisyrohoughia</i>), <i>Petrargyrops</i>	104.212
<i>similis</i> Townsend, 1935d, <i>Platyphasia</i>	104.8
<i>similis</i> Townsend, 1931a, <i>Pseudoxanthozonella</i>	104.70
<i>similis</i> Townsend, 1934a, <i>Xanthochaetona</i>	104.109
<i>similis</i> (Walker), 1852 (<i>Tachina</i>), <i>Peletcia</i>	104.45
<i>similis</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>simillima</i> (Fabricius), 1805 (<i>Ocyptera</i>), <i>Cylindromyia</i>	104.9
<i>simillima</i> Thompson, 1963b, <i>Carceliocephala</i>	104.203
<i>simillima</i> Vimmer & Soukup, 1940, <i>Gonia</i>	104.176
<i>simillimus</i> Townsend, 1914, <i>Saundersiops</i>	104.71
<i>simiolus</i> Reinhard, 1964c, <i>Sitellitergus</i>	104.35
<i>simmondsi</i> Thompson, 1963b, <i>Eugaediopsis</i>	104.179
<i>simplex</i> (Bigot), 1878 (<i>Ocyptera</i>), <i>Cylindromyia</i>	104.16
<i>simplex</i> (Bigot), 1889, (<i>Myocera</i>), <i>Ptilodexia</i>	104.34
<i>simplex</i> (Fallén), 1820 (<i>Ocyptera</i>), <i>Leucostoma</i>	104.18
<i>simulans</i> (Wulp), 1891, (<i>Rhynchodexia</i>), <i>Ptilodexia</i>	104.34
<i>singularis</i> Reinhard, 1931a, <i>Winthemia</i>	104.197
<i>singularis</i> (Townsend), 1891b (<i>Atrophopoda</i>), <i>Paradidyma</i>	104.115
<i>singularis</i> (Wiedemann), 1830 (<i>Tachina</i>), <i>Siphona</i>	104.170
<i>Siphoactia</i> Townsend, 1927a	104.170
<i>Siphocrocota</i> Townsend, 1935d	104.170
<i>Sipholeskia</i> Townsend, 1916c	104.118
<i>Siphona</i> Meigen, 1803	104.170
<i>siphonina</i> (Villeneuve), 1937, (<i>Myobia</i>), <i>Myobiopsis</i>	104.118
<i>Siphonini</i>	104.164
<i>Siphoniomyia</i> Bigot, 1885a	104.72
<i>Siphopsalida</i> Townsend, 1915m	104.17
<i>Siphosturmia</i> Coquillett, 1897	104.201
<i>Siphosturmiini</i>	104.201




<i>Siphosturmiopsis</i> Townsend, 1915n	104.201
<i>Sisyrohoughia</i> Townsend, 1927a	104.212
<i>Sisypomyia</i> Townsend, 1927a	104.170
<i>Sisyropa</i> Brauer & Bergenstamm, 1889	104.204
<i>Sitellitergus</i> Reinhard, 1964c	104.35
<i>Sitophaga</i> Gistel, 1843	104.148
<i>smaragdinus</i> (Macquart), 1843 (Jurinia), <i>Archytas</i>	104.52
<i>smithi</i> Aldrich, 1928a, <i>Belvosia</i>	104.182
<i>smithi</i> (Wulp), 1891 (Nemoraea), <i>Exopalpus</i>	104.55
<i>smithii</i> Townsend, 1916h, <i>Chrysometopiops</i>	104.207
<i>smithii</i> Williston, 1888, <i>Talarocera</i>	104.80
<i>sobrina</i> Wulp, 1890 (Phorocera)	104.161
<i>socia</i> (Walker), 1852 (Tachina), <i>Belvosia</i>	104.182
<i>sociabilis</i> (Greene), 1912a, <i>Drino</i>	104.189
<i>sodalis</i> (Wulp), 1890 (Masicera), <i>Blondelia</i>	104.127
<i>solita</i> Wulp, 1890, <i>Mystacella</i>	104.191
<i>solitaria</i> Aldrich, 1934 (Myiophasia)	104.100
<i>solitaria</i> (Curran), 1926a, (Hypostena), <i>Lixophaga</i>	104.138
<i>solitaris</i> Thompson, 1966, <i>Paradrino</i>	104.191
<i>Sophia</i> Robineau-Desvoidy, 1830	104.112
<i>Sophiini</i>	104.111
<i>sordida</i> Aldrich, 1934, <i>Peleteria</i>	104.45
<i>sordida</i> Giglio-Tos, 1893, <i>Microphthalma</i>	104.24
<i>sordida</i> (Walker), 1852 (Tachina), <i>Sitophaga</i>	104.148
<i>Sorochemyia</i> Townsend, 1915f	104.80
<i>soror</i> (Bigot), 1878 (Ocyptera), <i>Cylindromyia</i>	104.16
<i>soror</i> Bigot, 1885c, <i>Trichodischia</i>	104.100
<i>soror</i> (Williston), 1886 (Hystricia), <i>Bombyliomyia</i>	104.40
<i>sororcula</i> (Wulp), 1890 (Exorista), <i>Winthemlia</i>	104.197
<i>Spallanzania</i> Robineau-Desvoidy, 1830	104.177
<i>Spanipalpus</i> Townsend, 1908	104.76
<i>Spathidexia</i> Townsend, 1912b	104.95
<i>Spathimeigenia</i> Townsend, 1915b	104.148
<i>Spathimyia</i> Townsend, 1912a	104.149
<i>Spathipalpus</i> Rondani, 1863	104.118
<i>spatulata</i> (Townsend), 1928c (Stenaulacodoria), <i>Spathidexia</i> ...	104.95
<i>spectabilis</i> (Wulp), 1891 (Melaleuca), <i>Zelia</i>	104.103
<i>Sphaerina</i> Wulp, 1890	104.149
<i>Sphalloglandulus</i> Townsend, 1915m	104.136
<i>Sphyromyia</i> Bigot, 1884	104.43
<i>spinicosta</i> Aldrich, 1928b, <i>Incamiya</i>	104.136
<i>spinicoxa</i> Aldrich, 1928a, <i>Belvosia</i>	104.182
<i>spinigera</i> Wulp, 1892 (Jurinia), <i>Jurinella</i>	104.56
<i>spinipennis</i> Wiedemann, 1830, (Tachina)	104.216
<i>spinipes</i> Bigot, 1889 (Ceromasia)	104.153
<i>spiniventris</i> Guimarães, 1966b, <i>Adejeania</i>	104.47
<i>spinosa</i> (Bigot), 1889 (Rhynchodexia), <i>Ptilodexia</i>	104.34
<i>spinosa</i> Guimarães, 1966b, <i>Adejeania</i>	104.47
<i>spinosa</i> Guimarães, 1966a, <i>Mystacomyoidea</i>	104.199
<i>spinosa</i> Guimarães, 1966b, <i>Adejeania</i>	104.47
<i>spinosa</i> (Townsend), 1927a (Saundersiopmima), <i>Jurinella</i>	104.58
<i>spinosa</i> Townsend, 1927a, (Harrisioipsis), <i>Leschenaultia</i>	104.186
<i>spinosula</i> authors, <i>Athrycia</i>	104.90
<i>spinus</i> (Bigot), 1889, (Homodexia), <i>Thelairoides</i>	104.95

spinosus (Townsend), 1931a (Signoepalpus), Saundersiops	104.71
splendida Townsend, 1931a, Stomatotachina	104.160
spuria Wulp, 1890 (Brachycoma)	104.194
squamata Townsend, 1934a, Squamomedina	104.108
squamipes Wulp, 1892, Trichopoda (Trichopoda)	104.9
Squamomedina Townsend, 1934a	104.108
Steatosoma Aldrich, 1934	104.45
Stenaulacodoria Townsend, 1928c	104.95
Stenochaeta Thompson, 1968	104.149
Stenodexia Wulp, 1891	104.108
stenomae Curran, 1935, Lixophaga	104.138
Stenosturmia Townsend, 1927a	104.191
sternalis Aldrich, 1934, Dasyuromyia	104.95
sternalis Reinhard, 1955a, Cyldromyia	104.16
stipatus (Reinhard), 1958b (Dolichotarsus), Panacemyia	104.144
Stolatosoma Reinhard, 1953b	104.149
stolida Wulp, 1890, Anisia	104.125
Stomatodexia Brauer & Bergenstamm, 1889	104.119
Stomatomyia Brauer & Bergenstamm, 1889	104.160
Stomatolydella Townsend, 1919a	104.144
Stomatotachina Townsend, 1931a	104.160
strenua (Robineau-Desvoidy), 1830 (Zelia), Ptilodexia	104.34
striata Aldrich, 1928b, Incamyia	104.136
striata Aldrich, 1927a, Oedematocera	104.172
striata (Wulp), 1891 (Rhynchodexia), Ptilodexia	104.34
striatella Wulp, 1890 (Brachycoma)	104.194
striaticollis Wulp, 1891, Comyops	104.109
striatus Aldrich, 1934, Poliops	104.169
stricta Townsend, 1927a, Stenosturmia	104.192
strigata Wulp, 1890 (Masicera)	104.153
strigilata (Wulp), 1891 (Rhynchodexia), Ptilodexia	104.34
strigosa (Wulp), 1891 (Hypostena), Calothelaira	104.105
striolata (Wulp), 1890, (Telothyria), Eucelatoriopsis	104.134
Strongygastrini	104.18
Stuardomyia Cortés, 1945e	104.26
Sturmia, authors	104.188
Sturmia, Robineau-Desvoidy, 1830	104.192
Sturmiellina Thompson, 1963b	104.192
Sturmiini	104.187
Sturmioactia Townsend, 1926a	104.192
Sturmiodes Townsend, 1919	104.35
sturmioides Townsend, 1934, Sturmiomima	104.192
Sturmiomima Townsend, 1934	104.192
Sturmiopsoidea Thompson, 1966	104.192
Sturrimasiphya Townsend, 1935d	104.200
suavis Wulp, 1883, (Dexia)	104.122
subaenea Aldrich, 1934, Morphodexia	104.24
subaeneus Aldrich, 1934, Myiopharus	104.142
subalipes Townsend, 1897a, Trichopoda (Trichopoda)	104.9
subalpina Townsend, 1912, Eudejeania	104.54
subalpina Townsend, 1912, Oestrohystricia	104.69
subalpinus Townsend, 1912a, Eulasiopopus	104.54
subandina Blanchard, 1943c, Ceracia	104.123
subandina (Blanchard), 1943c (Cuphoceropsis), Peleteria	104.45



subauratum Blanchard, 1943b, Neopaedarium	104.91
subcilipes Macquart, 1843, Trichopoda (Trichopoda)	104.9
subcilipes Townsend, 1894, Trichopoda (Trichopoda)	104.9
subcinerea (Lynch Arribáizaga) in Brauer, 1898, Hyalomya ...	104.12
subcompressus Gundlach in Pazos, 1914 (Gonia)	104.176
subcyanea (Wulp), 1890 (Mystacella), Chrysotachina	104.81
subita Reinhard, 1958, Hypertrophomma	104.163
sublineata (Wulp), 1890 (Telothyria), Spathimeigenia	104.149
sublucens (Wulp), 1890 (Brachycoma), Sitophaga	104.148
subopaca (Aldrich), 1934, (Actia), Ceromya	104.166
subpetiolata (Wulp), 1892 (Cistogaster), Gymnoclytia	104.5
subpicea (Walker), 1852, (Tachina), Winthemia	104.197
subpilosa Wulp, 1890, (Masicera)	104.215
subpolita Townsend, 1912a, Brachymasicera	104.202
subtilipalpis Wulp, 1890 (Brachycoma), Promasiphya	104.200
subtilis Wulp, 1890, (Hypostena)	104.153
subvaria Walker, 1852 (Tachina)	104.194
subviridis (Wulp), 1892 (Gymnochaeta), Chrysotachina	104.81
Succingulodes Townsend, 1935d	104.149
succincta (Wulp), 1890 (Myobia), Pseudoredtenbacheria	104.147
suffusa Walker, 1852 (Dexia), Xanthodexia	104.96
sumichrasti Townsend, 1908, Euacaulona	104.7
Sumichrastia Townsend, 1916b	104.35
superpus Cortés, 1945a, Araucosimus	104.173
surinamensis Macquart, 1843, Jurinia	104.68
Synaplomyia Villeneuve, 1934	104.126
syngamiae Thompson, 1964, Pseudochaeta	104.163
Synorbitalia Thompson, 1968	104.157
Syringosoma Townsend, 1917b	104.8
tabida Reinhard, 1958b, Imaguncula	104.176
Tachininae	104.39
Tachinini	104.43
Tachinodes Brauer & Bergenstamm, 1889	104.47
tachinomoides Townsend, 1892, Euphorocera	104.158
Tachinomyia Townsend, 1892	104.160
Tachinophasia Townsend, 1931a	104.17
Tachinophyto Townsend, 1892d	104.150
Tachinophytopsis Townsend, 1927a	104.134
Tachinosoma Townsend, 1927a	104.56
Tachinosomopsis Blanchard, 1941	104.56
taglinoi (Sellers), 1945 (Zenillia), Chrysoexorista	104.207
Talacrocera Townsend, 1936b	104.80
Talarocera Williston, 1888	104.80
Talparomyia Thompson, 1963	104.150
Tamanamyia Thompson, 1963a	104.173
tantilla (Wulp), 1890 (Masicera), Blondelia	104.127
tanumeana Townsend, 1931, (Macropatelloa)	104.161
Tapajohoughia Townsend, 1934a	104.213
Tapajoleskia Townsend, 1934	104.119
Tapajosia Townsend, 1934	104.8
Tapajomintho Townsend, 1934a	104.113
Taperamyia Townsend, 1935d	104.102



Tarassus Aldrich, 1933a	104.11
Tarassini	104.11
Tarpessita Reinhard, 1967	104.88
tarsalis Aldrich, 1934, Dasyuromyia	104.98
tarsalis (Schiner), 1868 (Saundersia), Epalpus	104.65
tarsalis (Schiner), 1868 (Blepharipeza), Pteropeza	104.186
tarsalis (Séguy), 1925a, (Ormia), Euphasiopteryx	104.21
tarsalis Townsend, 1915m, Chaetophleps	104.89
tarsalis Townsend, 1927a, Opsosturmia	104.191
tarsata Schiner of Engel, 1920 (Hystriciella), Engelomyia	104.41
tarsata (Schiner), 1863 (Hystricia), Engelomyia	104.41
tatei Curran, 1939b, Chrysotachina	104.82
tatei Curran, 1947, Eulasiopalpus	104.54
tatei Curran, 1947, Saunderslops	104.71
taurea Townsend, 1934, Tapajoleskia	104.119
teffeensis Townsend, 1927a, Eucelatoriopsis	104.134
tegulata (Townsend), 1897a, (Trichopoda), Eutrichopoda	104.7
tegulata (Townsend), 1925c, (Peleteriosis), Peleteria	104.45
tegulata (Townsend), 1914, (Huascaraya), Trichophora	104.72
tehuantepeca (Townsend), 1908 (Acaulona), Cylindrophasia	104.6
Telodytes Aldrich, 1934	104.84
Telothyria Wulp, 1890	104.121
Telothyriini	104.120
Telothyriosoma Townsend, 1919a	104.150
tenebricosa Wulp, 1890 (Phorocera), Gaediophana	104.179
tenebrifera Walker, 1852 (Tachina)	104.216
tenuicornis Wulp, 1883 (Dexia)	104.122
tenuifrons Curran, 1934c, Phyllophilopsis	104.108
tenuipalpis Wulp, 1890 (Exorista)	104.198
tenuirostris James, 1947, Dejeniopalpus	104.116
tenuis Blanchard, 1959, Actinotachina	104.124
tenuiseta (Macquart), 1846 (Masicera), Eumasicera	104.162
tenuiseta Macquart, 1846 (Phorocera)	104.161
tersum Townsend, 1919a, Telothyriosoma	104.150
tcsselans Wulp, 1891, Mochlosoma	104.31
tessellata Brauer & Bergenstamm, 1891, Tetragrapha	104.160
tessellata (Giglio-Tos), 1893 (Acroglossa), Chaetogaedia	104.178
tessellata Röder, 1885 (Exorista)	104.215
tessellata (Wulp), 1890 (Mystacella), Winthemia	104.197
Tesseracephalus Reinhard, 1955c	104.35
testacea (Macquart), 1843 (Hystricia), Bombyliopsis	104.40
testaceiventris (Wulp), 1892, (Hystricia), Hystriciopsis	104.42
testaceus (Macquart), 1846 (Hystricia), Lindigepalpus	104.68
testaceus (Robineau-Desvoidy), 1830 (Jurinia), Zonoepalpus	104.74
testaceus (Wulp), 1888 (Saundersia), Epalpus	104.65
Tetragrapha Brauer & Bergenstamm, 1891	104.160
texana (Webber), 1930 (Achaetoneura), Lespesia	104.209
texensis Aldrich, 1931, Spathimeigenia	104.149
texensis Curran, 1925c, Peleteria	104.45
texensis Reinhard, 1924, Gonia	104.176
Thelaira Robineau-Desvoidy, 1830	104.95
Thelairalia Curran, 1934d	104.164
Thelairini	104.93
Thelairochaetona Townsend, 1919a	104.157



<i>Thelairoides</i> Wulp, 1891	104.95
<i>thelairodops</i> Townsend, 1935d, <i>Parodomyiops</i>	104.94
<i>Thelairodoria</i> Townsend, 1927a	104.150
<i>Thelairodoriopsis</i> Thompson, 1968	104.150
<i>Thelairomima</i> Townsend, 1936d	104.96
<i>Thelairophasia</i> Townsend, 1919b	104.200
<i>Thelyoxynops</i> Townsend, 1927a	104.150
<i>Thelyphaenopsis</i> Townsend, 1927a	104.150
<i>Thematheca</i> Reinhard, 1961	104.151
<i>Theresia</i> , authors	104.37
<i>Theresia</i> Robineau-Desvoidy, 1830	104.38
<i>Theresiini</i>	104.36
<i>Thereuops</i> Brauer & Bergenstamm, 1891	104.121
<i>thermithana</i> Cortés, 1945e, <i>Trichoceronia</i>	104.25
<i>thomae</i> (Wiedemann), 1830 (<i>Dexia</i>), <i>Tromodesiana</i>	104.102
<i>thompsoni</i> Guimarães, 1963b, <i>Adejeania</i>	104.47
<i>thompsoni</i> Guimarães, 1966a, <i>Neomasiphya</i>	104.199
<i>thoracica</i> Curran, 1925c, <i>Jurinella</i>	104.58
<i>thrix</i> Townsend, 1919a, <i>Thelairochaetona</i>	104.157
<i>thrix</i> Townsend, 1927a, <i>Thelairodoria</i>	104.150
<i>thyreionis</i> (Blanchard), 1963, <i>Patelloopsis</i>	104.211
<i>Thysanomyia</i> Brauer & Bergenstamm, 1891	104.192
<i>Thysanopsis</i> Townsend, 1917b	104.187
<i>Thysanosturmia</i> Townsend, 1927a	104.193
<i>tibialis</i> Aldrich, 1934, <i>Psecacera</i>	104.25
<i>tibialis</i> (Blanchard), 1954 (<i>Parabelvosia</i>), <i>Belvosia</i>	104.183
<i>tibialis</i> Blanchard, 1966b, <i>Paraxanthobasis</i>	104.110
<i>tibialis</i> Sellers, 1943, <i>Chrysophryxc</i>	104.207
<i>timida</i> Wulp, 1890, <i>Didyma</i>	104.131
<i>Tinalydella</i> Townsend, 1927a	104.151
<i>Tinanemorilla</i> Townsend, 1927a	104.195
<i>tincta</i> (Walker), 1852 (<i>Tachina</i>), <i>Patelloa</i>	104.211
<i>tinctipennis</i> (Curran), 1934c (<i>Rhynchodexia</i>), <i>Ptilodexia</i>	104.34
<i>tinctipennis</i> Curran, 1934c (<i>Rhynchodexia</i>), <i>Ptilodexia</i>	104.34
<i>tinctisquammae</i> Curran, 1926a, <i>Stomatodexia</i>	104.119
<i>tincensis</i> Townsend, 1927a, <i>Tinalydella</i>	104.151
<i>tipuliformis</i> Walker, 1857, <i>Cordyligaster</i>	104.111
<i>Tipuloleskia</i> Townsend, 1931a	104.119
<i>tolucana</i> Reinhard, 1956c, <i>Phryxc</i>	104.212
<i>Topomeigenia</i> Townsend, 1919a	104.193
<i>tortricis</i> (Coquillett), 1895 (<i>Pseudomyothyria</i>), <i>Erynnia</i>	104.173
<i>tortricis</i> (Coquillett), 1897 (<i>Phorocera</i>), <i>Hemisturmia</i>	104.194
<i>townsendi</i> (Aldrich), 1925 (<i>Masiphya</i>), <i>Promasiphya</i>	104.200
<i>townsendi</i> (Blanchard), in Griot, 1944 (<i>Ypophaeomyia</i>), <i>Lespesia</i>	104.209
<i>townsendi</i> Blanchard, 1963, <i>Protypophaemyia</i>	104.195
<i>townsendi</i> Curran, 1928b, <i>Archytas</i>	104.52
<i>townsendi</i> Curran, 1947, <i>Adejeania</i>	104.47
<i>townsendi</i> Curran, 1934b, <i>Calodexia</i>	104.155
<i>townsendi</i> (Curran), 1931 (<i>Epigrimyia</i>), <i>Clausicella</i>	104.166
<i>townsendi</i> Curran, 1939b, <i>Chrysotachina</i>	104.82
<i>townsendi</i> Curran, 1925c, <i>Pcleteria</i>	104.45
<i>townsendi</i> Curran, 1929, <i>Signosomopsis</i>	104.72
<i>townsendi</i> Guimarães, nomen novum, <i>Actinotachina</i>	104.124
<i>townsendi</i> Guimarães, nomen novum, <i>Cylindromyia</i>	104.16



townsendi Guimarães, nomen novum, Euphorocera	104.158
townsendi Guimarães, nomen novum, Lindigepalpus	104.68
townsendi (Vimmer & Soukup), 1940 (Chaetoandinomyia), Andinomyia	104.75
townsendi (Williston), 1896 (Atrophopoda), Ceratomyiella	104.114
Trafoia Brauer & Bergenstamm, 1893	104.83
transcendens Townsend, 1940b, Trochilochaeta	104.88
translucens (Macquart), 1846 (Jurinia), Archytas	104.52
transiens (Walker), 1849 (Tachina), Cryptopalpus	104.62
transita Townsend, 1931a, Tachinophasia	104.17
transita Townsend, 1919b, Thelairophasia	104.200
transita Townsend, 1939c, Trichophasia	104.114
transitionalis Townsend, 1915m, Punamyia	104.80
transversus (Walker), 1852 (Tachina), Cryptopalpus	104.62
travassosi Guimarães, 1961a, Archytas	104.52
travassosi Tavares, 1964, Ormiophasia	104.22
Trepophryx Townsend, 1908	104.213
Triachora Townsend, 1908	104.184
triangularis (Aldrich), 1934, (Lispidea), Elfia	104.167
triangularis Thompson, 1963d, Phasiopsis	104.200
trianguli Walker, 1849 (Tachina)	104.68
triangulifer (Loew), 1863 (Hyalomyia), Hyalomyyodes	104.18
triangulifera (Bigot), 1889 (Homodexia), Ptilodegeeria	104.147
triangulifera (Jacobs), 1900 (Hystriicia), Lypha	104.86
triangulifera Thompson, 1963, Argrophylax	104.162
triangulifera Townsend, 1927a, Paraprosena	104.38
Trichinochaeta Townsend, 1917a	104.151
<i>Trichiopoda</i> , emend	104.9
Trichoceronia Cortés, 1945e	104.26
Trichodejeania Townsend, 1913e	104.45
Trichodischia Bigot, 1885c	104.100
Trichodura Macquart, 1843	104.103
Trichodurini	104.103
<i>Trichoduropsis</i> Townsend, 1919b	104.103
trichoneura Wulp, 1890 (Masicera)	104.216
Trichopelopus Townsend, 1914	104.72
Trichopareilini	104.171
Trichophasia Townsend, 1939c	104.14
Trichophora Macquart, 1847	104.72
Trichophoropsis Townsend, 1914	104.80
Trichopoda Berthold, 1827	104.9
trichopoda (Wulp), 1891 (Morinia), Ebenia	104.109
Trichopodini	104.5
<i>Trichopodopsis</i> Townsend, 1913	104.10
<i>Trichoprosopa</i> Cortés, 1963, error	104.25
Trichoprosopini	104.25
Trichoprosopus Macquart, 1843	104.26
trichops Townsend, 1935d, Trinitodexia	104.97
trichopsis Bigot, 1887, Leschenaultia	104.186
Trichopyrrhosia Townsend, 1927a	104.119
Trichosaundersia Townsend, 1914	104.72
trichosoma (Wulp), 1890 (Myothyria), Ceraclia	104.153
Trichopteryx Townsend, 1919a	104.170
Trichschizotachina Townsend, 1935d	104.171




tricincta (Fabricius), 1805 (Musca), Comatacta	104.120
tricincta Reinhard, 1968, Erynnia	104.173
tricolor Coquillett, 1899, Metadexia	104.101
tricolor (Wulp), 1890 (Exorista), Winthemia	104.197
tridens Curran, 1947, Adejcania	104.47
tridens Townsend, 1927a, Triodontopyga	104.213
trifaria (Wiedemann), 1824 (Stomoxys), Genea	104.117
trifasciata Walker, 1837 (Tachina)	104.153
trifasciata (Wulp), 1890 (Prospheysa), Paralispes	104.145
trifida (Wulp), 1890 (Brachycoma), Drino	104.189
trifilata Wulp, 1890, (Anisia)	104.215
triformis (Walker), 1852 (Tachina), Rhachoealpus	104.71
trifurca Wulp, 1890 (Telothyria)	104.153
trigonalis (Wulp), 1892 (Xanthomelana), Xanthomelanopsis ...	104.11
trilineata Reinhard, 1954, Prosenoides	104.32
trinidadensis Thompson, 1966, Drino	104.189
trinidadensis Thompson, 1968, Neominthoidea	104.15
trinidadensis Thompson, 1968, Pelecotheca	104.157
trinidadensis Townsend, 1935d, Siphocrocota	104.170
trinitas Thompson, 1963a, Neossarromyia	104.79
trinitas Townsend, 1935d, Trichschizotachina	104.171
trinitatis Thompson, 1963a, Archytas	104.52
trinitatis Thompson, 1964, Calocarcelia	104.202
trinitatis Thompson, 1963b, Cubaemyiopsis	104.188
trinitatis Thompson, 1963a, Deopalpus	104.77
trinitatis Thompson, 1963a, Panacemyia	104.144
trinitatis Thompson, 1963d, Paraphasiopsis	104.200
trinitatis Thompson, 1963a, Peleteria	104.45
trinitatis Thompson, 1964, Pseudochaeta	104.163
trinitatis Thompson, 1968, Stomatomyia	104.160
trinitatis Thompson, 1963b, Sturmiellina	104.192
trinitatis Thompson, 1963a, Tamanamyia	104.173
trinitatis Thompson, 1963a, Telothyria	104.121
trinitatis (Thompson), 1963a (Neotrafoia), Trafoia	104.83
trinitatis Thompson, 1963c, Triodontopyga	104.213
trinitatis (Thompson), 1963a (Uromacquantia), Uramya	104.104
trinitatis Thompson, 1963a, Winthemia	104.198
Trinitodexia Townsend, 1935d	104.97
Trinitodexia Townsend, 1935d	104.213
Trinitodoria Townsend, 1935d	104.213
Triodontopyga Townsend, 1927a	104.213
triquetra (Olivier), 1812 (Ocyptera), Ervia	104.101
trisetosa Curran, 1928a, Argyrophylax	104.162
trisetosa Wulp, 1888, (Trichophora), Copecrypta	104.76
triste Reinhard, 1961, Lispideosoma	104.168
tristis Aldrich, 1934, Lygaeomyia	104.168
trivittata (Curran), 1925, (Myiomima) Sarcocallirrhoe	104.34
trivittata (Wiedemann), 1890 (Exorista), Winthemia	104.195
trivittatus Reinhard, 1955a, Xanthomelanodes	104.10
Trixodes Coquillett, 1902	104.22
Trixodini	104.22
Trixodopsis Townsend, 1933	104.25
trixoides Townsend, 1917a (Euthresioptis), Tyreomma	104.38
Trochilochaeta Townsend, 1940b	104.88



<i>Trochilodexia</i> Townsend, 1915b	104.30
<i>Trochiloglossa</i> Townsend, 1919a	104.119
<i>Trochiloleskia</i> Townsend, 1917b	104.119
<i>Trochiloglossa</i> Townsend, 1927a	104.119
<i>Tromodesiana</i> Townsend, 1931a	104.102
<i>Tromodesiopsis</i> Townsend, 1927a	104.102
<i>trophocyon</i> Aldrich, 1928b, <i>Echinopyrrhosia</i>	104.62
<i>tropica</i> Townsend, 1915, <i>Phantasiosiphona</i>	104.169
<i>tropica</i> Townsend, 1934a, <i>Tapajohoughia</i>	104.213
<i>tropica</i> Townsend, 1919a, <i>Trichotopteryx</i>	104.170
<i>tropica</i> Townsend, 1919a, <i>Trochiloglossa</i>	104.119
<i>tropica</i> Townsend, 1927a, <i>Tropidopsiomorpha</i>	104.36
<i>tropicalis</i> Townsend, 1897c, <i>Trichopoda</i> (<i>Trichopoda</i>)	104.9
<i>Tropidodexia</i> Townsend, 1915e	104.35
<i>Tropidopsis</i> Brauer & Bergenstamm, 1889	104.42
<i>Tropidopsiomorpha</i> Townsend, 1927a	104.36
<i>truncata</i> Aldrich, 1934, <i>Lypha</i>	104.86
<i>truncaticornis</i> (Wulp), 1888, <i>Austeniops</i>	104.60
<i>Trypherini</i> , in part	104.123
<i>Trypheromyia</i> Reinhard, 1945	104.151
<i>Trypetidomima</i> Townsend, 1935b	104.112
<i>Tuberculocera</i> Townsend, 1927a	104.53
<i>tuchucheensi</i> Thompson, 1968, <i>Chaetona</i>	104.105
<i>tucumana</i> (Blanchard), 1963 (<i>Patelloapsis</i>)	104.211
<i>tucumanensis</i> (Sellers), 1943 (<i>Zenillia</i>), <i>Eusisyropa</i>	104.203
<i>tucumanus</i> Blanchard, 1941, <i>Rhachoepalpus</i>	104.71
<i>Tunapunia</i> Townsend, 1927a	104.193
<i>turbinata</i> Wulp, 1890 (<i>Hypostena</i>)	104.153
<i>turicai</i> Blanchard, 1958, <i>Allosturmia</i>	104.187
<i>typicus</i> Curran, 1947, <i>Eulasiopalpus</i>	104.54
<i>Tyreomma</i> Brauer & Bergenstamm, 1891	104.38
<i>ucayali</i> (Townsend), 1929 (<i>Chaetophlepsis</i>), <i>Homohypochaeta</i> ...	104.89
<i>Ucayalimyia</i> Townsend, 1927a	104.213
<i>umbra</i> Walker, 1849, <i>Trichopoda</i> (<i>Trichopoda</i>)	104.9
<i>umbratilis</i> Reinhard, 1935a, <i>Pseudeuantha</i>	104.104
<i>umbrifera</i> (Walker), 1852 (<i>Tachina</i>), <i>Ptilodegeeria</i>	104.147
<i>umbrifera</i> Wulp, 1892, <i>Hyalomya</i>	104.13
<i>umbrina</i> Wulp, 1890, <i>Anisia</i>	104.125
<i>umbripennis</i> (Wulp), 1890 (<i>Hypostena</i>), <i>Ptilodegeeria</i>	104.147
<i>umbrosa</i> Wulp, 1892, <i>Hyalomya</i>	104.13
<i>uncana</i> (Fabricius), 1805 (<i>Dictya</i>), <i>Oestrophasiana</i>	104.20
<i>unguis</i> (Townsend), 1915m, (<i>Makasinocera</i>), <i>Archytas</i>	104.52
<i>unica</i> (Townsend), 1915m (<i>Sphalloglandulus</i>), <i>Incamyia</i>	104.136
<i>unicolor</i> (Aldrich), 1934 (<i>Actia</i>), <i>Ceromya</i>	104.166
<i>unicolor</i> (Aldrich), 1934, (<i>Xanthobasis</i>), <i>Paraxanthobasis</i>	104.110
<i>unicolor</i> Macquart, 1846, (<i>Miltogramma</i>)	104.216
<i>unicolor</i> (Wulp), 1888, (<i>Saundersia</i>), <i>Epalpus</i>	104.65
<i>unicolor</i> (Wulp), 1890 (<i>Exorista</i>), <i>Myiopharus</i>	104.142
<i>unifasciata</i> (Robineau-Desvoidy), 1830 (<i>Latreillia</i>), <i>Triachora</i> ..	104.185
<i>uniformis</i> Aldrich, 1926e, <i>Cylindromyia</i>	104.16
<i>uniformis</i> Curran, 1927, <i>Adejeania</i>	104.47
<i>unispinosa</i> Townsend, 1927a, <i>Ametadoria</i>	104.205




Uramya Robineau-Desvoidy, 1830	104.104
<i>Uramya</i> , emend.	104.104
Uramyiini	104.103
Uraporia Townsend, 1919b	104.105
urbanae Guimarães, 1966am Oromomasiphyia	104.199
ureophila Townsend, 1928b, Ushpayacua	104.102
urichi (Aldrich), 1932 (Exoristoides), Helioplagia	104.87
urna Townsend, 1934, Urucurymyia	104.10
Urochaetona Townsend, 1919a	104.109
Urodexiini	104.105
Urodexodes Townsend, 1919a	104.151
<i>Urophyllophyla</i> Townsend, 1927a	104.107
Urohypomyia Townsend, 1939	104.109
<i>Uruactia</i> Townsend, 1927a	104.164
Urucurymyia Townsend, 1939	104.10
urucurytuba Townsend, 1934, Tapajosia	104.9
Uruleskia Townsend, 1934	104.120
Urummyobia Townsend, 1934	104.120
Uruhuasia Townsend, 1914	104.73
Uruhuasiopsis Townsend, 1915m	104.73
uruhuasi (Townsend), 1927a (<i>Uruactia</i>), <i>Actia</i>	104.165
uruhuasi Townsend, 1914, <i>Chromoepalpus</i>	104.61
uruhuasi Townsend, 1927a, <i>Euclausicella</i>	104.134
uruhuasi Townsend, 1927a, <i>Exodexia</i>	104.29
uruhuasi Townsend, 1927a, <i>Exoernestia</i>	104.84
uruhuasi Townsend, 1927a, <i>Piximactia</i>	104.145
uruhuasi Townsend, 1927a, <i>Trichopyrrhosia</i>	104.119
uruhuasi Townsend, 1919a, <i>Xeoprosopa</i>	104.171
Ushpayacua Townsend, 1928a	104.102
ushpayacua Townsend, 1937b <i>Xiphophasia</i>	104.14
usta Giglio-Tos, 1893, (<i>Masicera</i>)	104.216
usta (Wiedemann), 1830 (<i>Tachina</i>), <i>Xylocamptomima</i>	104.84
vaciva (Wulp), 1890 (<i>Telothyria</i>), <i>Spathimeigenia</i>	104.149
vaga Curran, 1947, <i>Jurinella</i>	104.58
vagabunda Wulp, 1890 (<i>Didyma</i>)	104.215
valera Curran, 1934b, <i>Calodexia</i>	104.155
valida (Townsend), 1892e, (<i>Rhinophora</i>), <i>Microchaetina</i>	104.28
validinervis (Wulp), 1890, (<i>Didyma</i>), <i>Paradidyma</i>	104.115
validum Brauer & Bergenstamm, 1889, <i>Mochlosoma</i>	104.31
valparadisi Cortés, 1968a, <i>Baculocaptus</i>	104.123
<i>Valpogonia</i> Townsend, 1928c	104.174
Vanderwulpella Townsend, 1919b	104.18
vanderwulpi Townsend, 1892d, <i>Anisia</i>	104.125
vanderwulpi Williston, 1886, <i>Belvosia</i>	104.183
Vanderwulpia Townsend, 1891b	104.115
vargas Curran, 1942, <i>Hystricia</i>	104.41
vargas Curran, 1947, <i>Jurinella</i>	104.58
varia Curran, 1934b, <i>Calodexia</i>	104.155
varia (Walker), 1852 (<i>Tachina</i>), <i>Echinopyrrhosia</i>	104.62
varia (Wulp), 1890, <i>Distichona</i>	104.174
variabilis Aldrich, 1934, <i>Notodytes</i>	104.99
varians Curran, 1947, <i>Jurinella</i>	104.58




<i>variata</i> Curran, 1926b, Tachinomyia	104.160
<i>varicolor</i> Curran, 1947, Lindigia	104.58
<i>varicornis</i> Curran, 1928b, Archytas	104.52
<i>variegata</i> (Fabricius), 1805 (Stomoxys), Comatacta	104.120
<i>variegata</i> (Wulp), 1892 (Cistogaster), Gymnoclytia	104.5
<i>variegata</i> (Wulp), 1890 (Nemoraeca), Winthemia	104.198
<i>varipennis</i> (Curran), 1934 (Sturmia), Drino	104.189
<i>varipennis</i> Wulp, 1891, Scotiptera	104.31
<i>varipes</i> (Wulp), 1891, (Rhynchodexia), Ptilodexia	104.34
<i>vegetum</i> Reinhard, 1956b, Leucostoma	104.18
<i>velada</i> Reinhard, 1964b, Heliprosopa	104.78
<i>Velocia</i> Robineau-Desvoidy, 1863	104.151
<i>velutinus</i> Aldrich, 1934, Mellachnus	104.88
<i>velutina</i> (Wulp), 1888, (Hystricia), Chlorohystricia	104.40
<i>venatoria</i> (Fabricius), 1805, (Musca), Scotiptera	104.35
<i>venosa</i> Wulp, 1891, Megaparia	104.23
<i>veneris</i> Curran, 1934b, Calodexia	104.155
<i>ventripyga</i> Blanchard, 1966b, Protonotodytes	104.28
<i>venusa</i> Curran, 1928 (Phorocera)	104.161
<i>venusta</i> Cortés, 1967b, Schlingermiya	104.100
<i>venusta</i> Wulp, 1890, (Macquartia), Pseudeuantha	104.104
<i>verdans</i> Curran, 1947, Vertepalpus	104.73
<i>verna</i> (Curran), 1934c (Myobia), Myiobiopsis	104.118
<i>vernalis</i> Curran, 1928b, Archytas	104.52
<i>verrugana</i> (Townsend), 1914 (Trichodejeania), Adejeania	104.47
<i>Verrugomyia</i> Townsend, 1927a	104.214
<i>Verrugophryno</i> Townsend, 1927a	104.214
<i>versicolor</i> Robineau-Desvoidy, 1863, Jurinia	104.68
<i>versicolor</i> (Wulp), 1890, (Macquartia), Xanthophyto	104.84
<i>vertebrata</i> (Say), 1829 (Dexia), Zelia	104.103
<i>Vertepalpus</i> Curran, 1947	104.73
<i>verticale</i> (Curran), 1934c, (Lydella), Blondelia	104.127
<i>verticalis</i> (Reinhard), 1935a, Exoristopsis	104.87
<i>vestitus</i> (Townsend), 1916g (Xanthozonopsis), Euepalpus	104.66
<i>vetula</i> Reinhard, 1964, Acroglossa	104.173
<i>vexans</i> (Curran), 1934c (Pelixia), Avibrissosturmia	104.194
<i>vexans</i> Macquart not Wiedemann, (Stomoxys), Pachymyia	104.37
<i>vexans</i> (Wiedemann), 1830 (Stomoxys), Chaetogyne	104.36
<i>vexatrix</i> (Osten-Sacken), 1877 (Dejeania), Adejeania	104.47
<i>vexor</i> Curran, 1928b, Archytas	104.52
<i>vibrissata</i> (Aldrich), 1934 (Lipidea), Elfia	104.167
<i>vibrissata</i> (Wulp), 1891, (Dexiosoma), Microphthalma	104.24
<i>Vibrissoepalpus</i> Townsend, 1915n	104.73
<i>Vibrissomyia</i> Townsend, 1912a	104.80
<i>Vibrissovoria</i> Townsend, 1919a	104.178
<i>vicina</i> Wulp, 1890 (Telothyria)	104.153
<i>vidua</i> Schiner, 1868, Trichodura	104.103
<i>vierecki</i> Curra, 1947, Lindigia	104.58
<i>vilis</i> (Wulp), 1890 (Prospheysa), Chaetogaedia	104.178
<i>villaricana</i> Reinhard, 1951, Belvosia	104.183
<i>villosa</i> Wulp, 1892, Hyalomya	104.13
<i>violacea</i> Wulp, 1890, (Mystacella), Bolomyia	104.187




<i>violaceiventris</i> (Brauer), 1898 (<i>Hyalomya</i>), <i>Alophorella</i>	104.12
<i>violascens</i> Townsend, 1897, <i>Hyalomya</i>	104.13
<i>virens</i> Aldrich, 1926a, <i>Microphthalma</i>	104.24
<i>virens</i> (Aldrich), 1928b (<i>Selenomyia</i>), <i>Psecacera</i>	104.26
<i>virescens</i> Macquart, 1843, <i>Gonia</i>	104.176
<i>viridis</i> Townsend, 1915m, <i>Chrysoexorista</i>	104.207
<i>viridis</i> (Townsend), 1912a, (<i>Oeestropsis</i>), <i>Euestropsis</i>	104.25
<i>virilis</i> (Aldrich & Webber), 1924, (<i>Zenillia</i>), <i>Eusisyropa</i>	104.203
<i>Visayalydina</i> Townsend, 1926n	104.88
<i>vittata</i> Aldrich, 1951, <i>Belvosia</i>	104.183
<i>vittata</i> Curran, 1937a, <i>Trichodura</i>	104.103
<i>vittata</i> (Guérin-Ménéville), in Duperrey 1831, (<i>Sophia</i>), <i>Scotip- tera</i>	104.35
<i>vittata</i> Thompson, 1963b, <i>Atactosturmia</i>	104.180
<i>vittata</i> Townsend, 1915m, <i>Minthopsis</i>	104.141
<i>vittata</i> Walker, 1852, (<i>Tachina</i>)	104.215
<i>vittatus</i> Curran, 1947, <i>Eulasioipalpus</i>	104.54
<i>vittatus</i> (Walker), 1852 (<i>Tachina</i>), <i>Epalpodes</i>	104.63
<i>vittiger</i> Curran, 1934c, <i>Ptilodexia</i>	104.34
<i>vittigerus</i> (Bigot), 1889 (<i>Homodexia</i>), <i>Thelairoides</i>	104.95
<i>Vivianoidea</i> Thompson, 1968	104.157
<i>vivida</i> (Wiedemann), 1830 (<i>Tachina</i>), <i>Euphorocera</i>	104.158
<i>volucris</i> Wulp, 1890, (<i>Didyma</i>)	104.215
<i>vorax</i> (Wiedemann), 1839 (<i>Tachina</i>), <i>Triodontopyga</i>	104.213
<i>Voria</i> Robineau-Desvoidy, 1830	104.93
<i>Vorialia</i> Curran, 1934d	104.92
<i>Voriini</i>	104.89
<i>vulgaris</i> Curran, 1928, <i>Archytas</i>	104.48
<i>vulgata</i> Walker, 1852, (<i>Tachina</i>)	104.153
<i>vultur</i> Curran, 1942, <i>Hystricia</i>	104.41
<i>Wagneria</i> , authors	104.92
<i>wagneri</i> (Blanchard), 1941, (<i>Archytodejeania</i>), <i>Archytas</i>	104.52
<i>weyenberghiana</i> Wulp, 1883, <i>Belvosia</i>	104.83
<i>wheeleri</i> Reinhard, 1942a, <i>Cowania</i>	104.90
<i>wiedemanni</i> Aldrich, 1928a, <i>Belvosia</i>	104.183
<i>williamsi</i> Aldrich, 1928a, <i>Belvosia</i>	104.183
<i>williamsi</i> Arnaud, 1966, <i>Polistiopsis</i>	104.17
<i>willinki</i> (Blanchard), 1954 (<i>Eubelvosiopsis</i>), <i>Belvosia</i>	104.183
<i>willistoni</i> Curran, 1925e, <i>Archytas</i>	104.52
<i>willistoni</i> Curran, 1939b, <i>Chrysotachina</i>	104.82
<i>willistoni</i> Townsend, 1894 (<i>Ocypterosipha</i>), <i>Beskia</i>	104.15
<i>Williston</i> Brauer & Bergenstamm, 1889	104.181
<i>Winthemia</i> Robineau-Desvoidy, 1830	104.195
<i>Winthemiini</i>	104.194
<i>Winthemiopsis</i> Blanchard, 1963	104.196
<i>wolcott</i> Sabrosky, 1953, <i>Ormia</i>	104.22
<i>Wulpia</i> Brauer & Bergenstamm, 1893	104.114
<i>wulp</i> Townsend, 1913a, <i>Bathydextia</i>	104.36
<i>wulp</i> Townsend, 1927a, <i>Eggonia</i>	104.216
<i>wygodzinskyi</i> Guimarães, 1966b, <i>Adejeania</i>	104.47
<i>wygodzinskyi</i> Tavares, 1953, <i>Ormia</i>	104.22



<i>xanthipes</i> Blanchard, in [Marchionatto], (1940), <i>Hypoproxylops</i>	104.136
<i>Xanthobasis</i> Aldrich, 1934	104.110
<i>xanthocera</i> (Wiedemann), 1830 (<i>Tachina</i>), <i>Winthemia</i>	104.198
<i>Xanthochaetona</i> Townsend, 1934a	104.109
<i>Xanthodexia</i> Wulp, 1891	104.96
<i>Xanthoepalpodes</i> Townsend, 1914	104.73
<i>Xanthoepalpus</i> Townsend, 1913a	104.73
<i>xanthogastra</i> Wulp, 1891, <i>Camarona</i>	104.36
<i>Xanthohystricia</i> Townsend, 1913a	104.42
<i>Xanthomelana</i> Wulp, 1892	104.10
<i>Xanthomelanodes</i> Townsend, 1893	104.10
<i>xanthomelanoides</i> Townsend, 1934, <i>Xenophasia</i>	104.11
<i>Xanthomelanopsis</i> Townsend, 1917c	104.11
<i>Xanthopelta</i> Aldrich, 1934	104.88
<i>Xanthophyllophylla</i> Townsend, 1927a	104.109
<i>Xanthophyto</i> Townsend, 1916e	104.84
<i>xanthopilosa</i> Guimarães, 1966b, <i>Adejeania</i>	104.47
<i>Xanthotheresia</i> Townsend, 1931a	104.38
<i>Xanthotrichius</i> Townsend, 1934a	104.14
<i>Xanthozona</i> Townsend, 1908	104.74
<i>Xanthozonella</i> Townsend, 1931b	104.62
<i>Xanthozonopsis</i> Townsend, 1916g	104.66
<i>xanthura</i> Wulp, 1890 (<i>Phorocera</i>)	104.194
<i>Xenophasia</i> Townsend, 1934	104.11
<i>Xenophysis</i> , error	104.86
<i>Xenoplagia</i> Townsend, 1914	104.93
<i>Xenopyxis</i> Townsend, 1940	104.86
<i>xenos</i> Townsend, 1934a, <i>Xanthotrichius</i>	104.14
<i>Xeoprosopa</i> Townsend, 1919a	104.171
<i>Xiphomyia</i> Townsend, 1917c	104.151
<i>Xiphophasia</i> Townsend, 1937b	104.14
<i>ychus</i> (Walker), 1849 (<i>Tachina</i>), <i>Lespesia</i>	104.209
<i>Xylocamptomima</i> Townsend, 1927a	104.84
<i>Xystotrix</i> Townsend, 1927a	104.27
<i>yahuarmayana</i> Townsend, 1927a, <i>Bolodoria</i>	104.127
<i>yahuarmayana</i> Townsend, 1927a, (<i>Epiphyllaphila</i>), <i>Microchaetona</i>	104.107
<i>yahuarmayensis</i> Townsend, 1927a, <i>Didymops</i>	104.132
<i>Yahuarmayola</i> Townsend, 1927a	104.102
<i>Yahuarphryno</i> Townsend, 1927a	104.211
<i>yahuarphrynoides</i> (Townsend), 1927a, (<i>Yahuartachina</i>), <i>Palpe-xorista</i>	104.159
<i>Yahuartachina</i> Townsend, 1927a	104.159
<i>yanesi</i> Marnef, 1968, <i>Lafuentemyia</i>	104.98
<i>Ypophaemyia</i> Townsend, 1916d	104.208
<i>Ypophaemyiopsis</i> Townsend, 1935d	104.214
<i>ypsiliformis</i> Blanchard, 1966b, <i>Ptilodexia</i>	104.34
<i>ypsilon</i> Curran, 1947, <i>Adejeania</i>	104.47
<i>Zelia</i> Robineau-Desvoidy, 1830	104.102
<i>Zeliini</i>	104.101
<i>Zenillia</i> Robineau-Desvoidy, 1830	104.214




zeteki Curran, 1947, <i>Jurinella</i>	104.58
zikani Guimarães, 1961b, <i>Archytas</i>	104.52
<i>Ziziphomyia</i> Townsend, 1916f	104.193
<i>Zonalia</i> Curran, 1934d	104.96
zonata (Bigot), 1889 (<i>Ceromasia</i>), <i>Spathimeigenia</i>	104.149
<i>Zonoepalpus</i> Townsend, 1927a	104.74
zonula Reinhard, 1951a, <i>Hemyda</i>	104.16
<i>Zuanalia</i> Curran, 1934d	104.25
zuercheri Townsend, 1928b, <i>Sarcocalirrhoe</i>	104.34
<i>Zygofrontina</i> Townsend, 1915m	104.208
<i>Zygosturmia</i> Townsend, 1911	104.188
<i>Zygozenillia</i> Townsend, 1927a	104.193






SciELO



SciELO₉



SciELO

DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

105

FAMILY CUTEREBRIDAE

J. H. GUIMARÃES

*Departamento de Zoologia, Secretaria da Agricultura
São Paulo, Brasil*

These large and robust flies, known as rodent warble-flies, are myiasis-producing Diptera. Their larvae develop mainly in the subcutaneous tissues of rodents and lagomorphs. These flies are restricted to the New World and are well represented in the Neotropical Region. A strange habit is found in *Dermatobia hominis* (Linnaeus Jr.), the human warble-fly, for the transmission of its eggs to the host — the adult female glues her eggs on other insects, especially mosquitoes, which thus carry them to the host.

Satisfactory systematic concepts for the family have not yet been established and the group is still in need of a critical review. Among the authors who made important contributions to this family in the Neotropical Region, especial mention is to be made to Lutz (1917), Bau (1906, 1931), Guimarães & Carrera (1941), Austen (1933). The biology of some species in Brazil have been dealt with by Fonseca (1938) and Neiva & Gomes (1917). An annotated bibliography of *Dermatobia hominis* has been prepared by Guimarães & Papavero (1963).

The suprageneric arrangement adopted here follows in its main lines that proposed by Townsend (1935).

I wish to express my sincere gratitude to Dr. Curtis W. Sabrosky, of the U.S. Department of Agriculture at Washington, for his generous cooperation in the preparation of the manuscript.

Tribe CUTEREBRINI

Genus ALOUATTAMYIA Townsend

ALOUATTAMYIA Townsend, 1931:319. Type-species, *Cuterebra baeri* Shannon & Greene (orig. des.).

BAERI (Shannon & Greene), 1926:286, pls. 152-153 (*Cuterebra*). Type-locality: British Guiana, Bartica District, Kartabo. Distr. — Costa Rica, Panama, British Guiana. Ref. — Zeledón, Jimenez & Brennes (1957; larva).

Genus CUTEREBRA Clark

CUTEREBRA Clark, 1815:64, 70. Type-species, *Musca americana* Fabricius (Desmarest, 1859:274).

105.1

DEPARTAMENTO DE ZOOLOGIA
BIBLIOTECA
Secretaria da Agricultura

Cutiterebra, emend.

Trypoderma Wiedemann, 1830:256. Type-species, *Musca americana* Fabricius (Coquillett, 1910:618).

Bogeria Austen, 1895:391. Type-species, *princeps* Austen (orig. des.).

Atrypoderma Townsend, 1919:592. Type-species, *Musca americana* Fabricius (orig. des.).

Orthocuterebra Bau, 1929b:543. Type-species, *Cuterebra lepusculi* Townsend (Aldrich, 1931:117) = *princeps* (Austen).

Paracuterebra Bau, 1929b:543. Type-species, *Musca americana* Fabricius (Aldrich, 1931:117).

Metacuterebra Bau, 1929b:544. Type-species, *Cuterebra apicalis* Guérin-Méneville (Bau, 1931:208). N. SYN.

Pseudobogeria Bau, 1931:206. Type-species, *Musca buccata* Fabricius (orig. des.).

Oricuterebra Bau, 1931:210. Type-species, *Cuterebra sarcophagoides* Lutz (orig. des.). N. SYN.

Baucuterebra Guimarães & Carrera, 1941:2. Type-species, *Cuterebra schroederi* Enderlein (Townsend, 1942: Addenda et corrigenda, p. 333). N. SYN.

Refs. — Bau, 1931 (key to Central and South America);
Lutz, 1917 (Brazilian species).

ALMEIDAI (Guimarães & Carrera), 1941:2, figs. 1, 3 (*Baucuterebra*).
Type-locality: Brazil, Amazonas, Cachoeira Taurino. Distr. — Brazil (Amazonas). N. COMB.

APICALIS Guérin-Méneville, 1835: pl. 101, fig. 1; 1844:547. Type-locality: "America". Distr. — Mexico to Patagonia. Refs. — Bau, 1931:208; Fonseca, 1938; d'Andretta Jr. & Jardim, 1954 (biology); Henriksen, 1942 (hosts, larva); Forattini & Lenko, 1959 (pupal stage).

analis Macquart, 1843:22, pl. 2, fig. 5. Type-locality: "Brazil".

ATROX Clark, 1848: fig. 5. Type-locality: "Mexico". Distr. — Mexico. Ref. — Brauer, 1863b:241.

BAUI (Fonseca), 1940:667, figs. 1-3 (*Oricuterebra*). Type-locality: Brazil, São Paulo, Capital, Pinheiros. Distr. — Brazil (São Paulo).

CAYENNENSIS Macquart, 1843:23. Type-locality: French Guiana, Cayenne. Distr. — French Guiana (Cayenne). Ref. — Brauer, 1863:240.


COMETES Shannon & Del Ponte, 1926:42. Type-locality: Argentina, Tucumán, Quebrada de Lulcs. Distr. — Argentina (Tucumán).

CONFLANS CONFLANS Bau, 1929a:303, figs. 1-2. Type-locality: Peru, Chanchamayo. Distr. — Peru, Brazil (Pará). Ref. — Bau, 1931:235.

ssp. OBIDOSIANA Bau, 1929a:305. Type-locality: Brazil, Pará, Óbidos. Ref. — Bau, 1931:233.

DETRUDATOR Clark, 1848: fig. 4. Type-locality: "South America". Distr. — "South America". Ref. — Bau, 1931:233.

- DORSALIS Bau, 1929a:305, fig. 3 (1930:83). Type-locality: "South America".
Distr. — "South America". Ref. — Bau, 1929a:305, fig. 3 (head).
- EPIHIPTIUM Latreille, 1818:271. Type-locality: French Guiana, Cayenne.
Distr. — French Guiana (Cayenne). Ref. — Bau, 1931:283.
- FLAVIVENTRIS (Bau), 1931:219, pl. 1, fig. 10 (*Metacuterebra*). Type-locality:
"Mexico". Distr. — Mexico.
- FUNEBRIS (Austen), 1895:378, pl. 13, figs. 1, 1b (*Cutiterebra*). Type-locality:
"Trinidad". Distr. — Trinidad. Refs. — Bau, 1931:230;
Austen, 1933: 704.
- HISTRIO Coquillett, 1902:103. Type-locality: Mexico, Guanajuato. Distr.
— Mexico (Guanajuato).
- INFULATA Lutz, 1917:99, pl. 27, fig. 4, pl. 29, fig. 4. Type-locality: Brazil,
Rio de Janeiro, Petrópolis. Distr. — Brazil (Rio de Janeiro).
- LEPIDA Austen, 1933: 707, pl. 1, fig. 8. Type-locality: "Nicaragua". Distr.
— Nicaragua.
- LUTZI Bau, 1930:81. Type-locality: "Peru". Distr. — Peru, Brazil.
- MACULOSA Knab, 1914:187. Type-locality: Panama, Canal Zone, Ancon
and Cristobal. Distr. — Panama (Canal Zone).
- MEGASTOMA Brauer, 1863b:247, pl. 4, fig. 5, pl. 6, fig. 12. Type-locality:
"South America". Distr. — "South America".
- NIGRICANS Lutz, 1917:99, pl. 27, fig. 2, pl. 29, fig. 2. Type-locality: Brazil,
São Paulo, Pôrto Martins. Distr. — Brazil (São Paulo).
- NIGRICINCTA (Austen), 1895:388, pl. 13, figs. 4, 4a (*Cutiterebra*). Type-
locality: Brazil, Pará. Distr. — Brazil (Pará). Refs. — Bau,
1931:233; Austen, 1933:710.
- ORNATA Bau, 1928a:50. Type-locality: "Costa Rica". Distr. — Costa Rica.
- PATAGONA Guérin-Ménéville, 1844:548. Type-locality: Argentina, Patagonia.
Distr. — Argentina.
- PESSOAI Guimarães & Carrera, 1941:4, figs. 4, 6. Type-locality: Brazil,
Amazonas, Rio Negro. Distr. — Brazil (Amazonas).
- PRAEGRANDIS Austen, 1933:706, pl. 1, figs. 6, 7. Type-locality: "Peru". Distr.
— Peru.
- PRINCEPS (Austen), 1895:393, pl. 13, figs. 5, 5b (*Bogeria*). Type-locality:
Mexico, Sonora. Distr. — U.S.A. (Washington to South Dakota,
south to Arizona, Texas), Mexico (Sonora).
- lepuseuli* Townsend, 1897:8. Type-locality: U.S.A., New Mexico.
- albifrons* Swenk, 1905:183. Type-locality: U.S.A., Wyoming.
- PULCHRA Bau, 1930:86, pl. 1, fig. 5. Type-locality: "South America".
Distr. — "South America".
- PUSILLA Bau, 1931:212, pl. 1, fig. 7, pl. 2, fig. 16. Type-locality: Bolivia,
Mapiri, Sarampiuni. Distr. — Bolivia (Mapiri, Sarampiuni).
- PYGMAEA Bau, 1931:215, pl. 5, figs. 2, 5. Type-locality: Brazil, Pará,



- Ilha de Marajó, Caldeirão. Distr. — Brazil (Pará, Ilha de Marajó).
- RIBEIROI (Artigas & d'Andretta Jr.), 1951:60, figs. 1-3 (*Alouattamyia*). Type-locality: Brazil, Goiás, Aruanã, Rio Vermelho). Distr. — Brazil (Goiás). N. comb.
- RUBIGINOSA Bau, 1931:214, pl. 1, fig. 9, pl. 2, fig. 14. Type-locality: Brazil, Rio Grande do Sul, Santa Cruz. Distr. — Brazil (Rio Grande do Sul).
- RUFIVENTRIS RUFIVENTRIS Macquart, 1843:21, pl. 2, figs. 4, 4a. Type-locality: Brazil, Pará. Distr. — Ecuador, Peru (Chanchamayo), Brazil (Pará). Ref. — Auten, 1933:705.
- ssp. ELEGANS Bau, 1931:227. Type-locality: Peru, Chanchamayo.
- SARCOPHAGOIDES Lutz, 1917:99, pl. 27, fig. 5, pl. 29, fig. 5. Type-locality: Brazil, São Paulo, Jacutinga. Distr. — Brazil (São Paulo).
- SCHMALZI Lutz, 1917: 100, pl. 27, fig. 3, pl. 29, fig. 3. Type-locality: Brazil, Santa Catarina, Joinville. Distr. — Brazil (Santa Catarina), Argentina (Misiones).
- SCHROEDERI (Enderlein), 1909:202 (*Cutiterebra*). Type-locality: Peru, Chanchamayo. Distr. — Peru. Ref. — Bau, 1929a:307, fig. 5.
- SEMILUTEA Bau, 1929c:1, figs. 1-2. Type-locality: "Brazil". Distr. — Brazil".
- SIMULANS Austen, 1933:709. Type-locality: "Peru". Distr. — Peru.
- TERRISONA Walker, 1849:683. Type-locality: "Guatemala". Distr. — Guatemala. Ref. — Austen, 1895:382, pl. 13, figs. 2, 2a (redesc. of Walker's type).
- TOWNSENDI (Fonseca), 1941:483, fig. 1 (*Orieuterebra*). Type-locality: Brazil, São Paulo, Capital. Distr. — Brazil (São Paulo).
- VERSICOLORA VERSICOLORA Bau, 1930:85, fig. 4. Type-locality: "Surinam". Distr. — Peru, Surinam.
- ssp. BICINCTA Bau, 1931:221, pl. 1, fig. 6, pl. 2, fig. 17. Type-locality: "Peru".
- WORONTZOWI Guimarães & Carrera, 1941:6, figs. 7-9. Type-locality: Brazil, Amazonas, Rio Paruarí. Distr. — Brazil (Amazonas).

Genus ROGENHOFERA Brauer

- ROGENHOFERA Brauer, 1863a:325 (1863b:215, 217). Type-species, *trigonophora* Brauer (mon.).
- Guerinioestrus* Townsend, 1931:318. Type-species, *Cephenemyia grandis* Guérin-Ménéville (orig. des.). N. SYN.
- Ref. — Bau, 1932 (key).
- BONAERENSIS (Dcl Pontc), 1939:529 (*Guerinioestrus*). Type-locality: Argentina, Buenos Aires. Distr. — Argentina (Buenos Aires).

- DAGUERREI (Del Ponte), 1939:530 (*Guerinioestrus*). Type-locality: "Argentina". Distr. — "Argentina".
- DASYPODA Brauer, 1896:272. Type-locality: Brazil, Espírito Santo. Distr. — Brazil (Espírito Santo, Rio de Janeiro). Refs. — Lutz, 1917:100, pl. 28, fig. 8; Bau, 1928:300.
- GILVOPILOSA Bau, 1932:11. Type-locality: Peru, Arequipa, Puno. Distr. — Peru (Arequipa).
- GRANDIS (Guérin-Méneville), 1844:548 (*Cephenemyia*). Type-locality: Argentina, Patagonia. Distr. — Southern Brazil to Patagonia. Refs. — Berg, 1876:272 (hosts). Del Ponte, 1939:527. N. comb.
- TRIGONOPHORA Brauer, 1863a:326. Type-locality: Brazil, Bahia. Distr. — Brazil (Bahia). Ref. — Lutz, 1917:101.
trigonocephala Lutz, 1917:96 (error).

Tribe DERMATOBINI

Genus DERMATOBIA Brauer

- DERMATOBIA Brauer, 1861:783. Type-species, *Cuterebra cyaniventris* Macquart (Guimarães & Papavero, 1966:228) = *hominis* (Linnaeus Jr., in Pallas). Refs. — Neiva & Gomes, 1917 (biology); Guimarães & Papavero, 1966 (annotated bibliography).
- HOMINIS (Linnaeus Jr.) in Pallas, 1781:158 (*Oestrus*). Type-locality: "Peru" Distr. — From Mexico to Argentina, and Chile.
humanus Howship, 1833:181 (*Oestrus*). Type-locality: Colombia, Mariquita.
- cyaniventris* Macquart, 1843:23 (*Cuterebra*). Type-locality: "Brazil".
noxialis Goudot, 1845:229, pl. 4 (*Cuterebra*). Type-locality: Colombia, Marmato.

Tribe PSEUDOGAMETINI

Genus PSEUDOGAMETES Bischof

- PSEUDOGAMETES Bischof, 1900a:131 (1900b:490). Type-species, *hermanni* (orig. des.).
- Xylocopodes* Townsend, 1927:207. Type-species, *Musca semiatra* Wiedemann (orig. des.). N. SYN.
- HERMANNI Bischof, 1900a:131 (1900b:490, figs. 1-3). Type-locality: Brazil, Minas Gerais. Distr. — Brazil (Minas Gerais).
- SEMIATRA (Wiedemann), 1830:421 (*Musca*). Type-locality: "Brazil". Distr. — Brazil (Minas Gerais and Espírito Santo, south to Santa Catarina). Ref. — Lutz, 1917:102. N. comb.

Genus MONTEMYIA Dalmat

- MONTEMYIA Dalmat, 1942:25. Type-species, *bureni* Dalmat (orig. des.).
- BURENI Dalmat, 1942:25, pl. 2. Type-locality: Costa Rica, Monte Redondo. Distr. — Costa Rica (Monte Redondo).



LITERATURE CITED

ALDRICH, J. M.

1931. Notes on Diptera, n° 5. *Proc. Ent. Soc. Washington* 33(5): 116-121.

ARTIGAS, P. T. & C. D'ANDRETTA JR.

1951. *Alouattamyia ribeiroi*, nova espécie de Cuterebridae (Diptera) do Brasil. *Arq. Mus. Nac.*, Rio de Janeiro 42:59-62, 3 pls.

AUSTEN, E. E.

1895. On the specimens of the genus *Cutiterebra* and its allies (Family Oestridae) in the collection of the British Museum. with descriptions of a new genus and three new species. *Ann. & Mag. Nat. Hist.* (6) 15:377-396, 1 pl.
1933. New and little-known species of *Cuterebra* Clark and *Bogeria* Austen (Diptera: Family Oestridae). *Proc. Zool. Soc. London* 3:699-713, 1 pl.

BAU, A.

1906. Diptera. Fam. Muscaridae. Subfam. Oestrinae, in P. Wytzman: *Genera Insectorum* 43:31 pp., 2 pls., Bruxelles.
- 1928a. *Cuterebra ornata*, spec. nov. (Diptera, Cuterebrinae). *Konowia* 7:50-51.
- 1928b. Ueber *Aulacocephala* und *Rogenhoferia*. *Konowia* 7:298-302.
- 1929a. *Cuterebra conflans* und *subbuccata*, spec. novae, sowie Bemerkung über *C. schroederi* Enderlein (Diptera, olim Oestridae). *Stettin. Ent. Ztg.* 90:303-307, 1 pl.
- 1929b. Versuch einer Teilung der Gattung *Cuterebra* (Diptera, olim Oestridae) in vier Untergattungen. *Centbl. f. Bakt. Parasitenk. u. Infektionskrank.* (2) 77:542-544, 1 fig.
- 1929c. *Cuterebra semilutea*, sp. n. (Ins. Dipt.), sowie Schlüssel zur Bestimmung der *Cuterebra*-Arten. *Senckenbergiana* 11:1-7, 2 figs.
1930. Vier neue "*Cuterebra* Arten" aus Südamerika (Diptera, olim Oestridae). *Konowia* 9:81-89, 1 pl.
1931. Ueber das Genus *Cuterebra* Clark (Diptera, Oestridae). Einteilung desselben in sechs Untergattungen. Beschreibung neuer Species und Aufstellung einer Bestimmungstabelle der mittel — und südamerikanischen Arten. *Konowia* 10:197-240, figs., 2 pls.
1932. *Rogenhoferia gilvopilosa* spec. nov. und *R. grandis* (Guérin) (Dipt. Cuterebr.). *Mitt. deutsch. ent. Ges.* 3:11-15.

BERG, C.

1876. Naturgeschichte der *Rogenhoferia grandis*, einer Fliege aus der Familie der Oestriden. *Stettin. Ent. Ztg.* 37:268-272.

BISCHOF, J.

- 1900a. Vorläufige Charakteristik einiger neuen Gattungen von Muscarien *Anz. k. Akad. Wiss. Wien* 37:131.
- 1900b. Einige neue Gattungen von Muscarien. *Sitzber. K. Akad. Wiss., Math. Nat. Cl.* 109:490-497, 5 figs.



BRAUER, F.

- [1861]. Ueber die Larven der Gattung *Cuterebra* Clk. *Verh. zool.-bot. Ges. Wien* 10 (1860): 777-786.
1863a. *Rogenhoferia*, eine neue Gattung aus der Familie der Oestriden. *Verh. zool.-bot. Ges. Wien* 13:325-326.
1863b. *Monographie der Oestriden*, 292 pp., 10 pls. Wien.
1896. Beiträge zur Kenntniss aussereuropäischer Oestriden. *Denk. K. Akad. Wiss., Mathem. Naturwiss. Cl.* 64:260-282, 1 pl.

CLARK, B.

1815. *An essay of the bots of horses and other animals*, 72 pp., 2 col. pls. London.
1848. *Idem*, Addenda, 1 pl., 5 figs.

COQUILLET, D. W.

1902. New Diptera from North America. *Proc. U.S. Nat. Mus.* 25 (1280): 83-126.
1910. The type-species of the North American genera of Diptera. *Proc. U.S. Nat. Mus.* 37 (1719): 499-647.

D'ANDRETTA JR., C. & J. L. JARDIM

1954. Contribuição ao conhecimento do ciclo biológico de *Metacuterebra apicalis* (Guérin, 1829/38) (Diptera, Cuterebridae). *An. Fac. Farm. Odont. Univ. S. Paulo* 12:67-73.

DALMAT, H. T.

1942. New genus and species of Cuterebridae from Costa Rica and other taxonomic notes (Diptera). *Trans. Amer. Ent. Soc.* 68:23-29, 1 pl.

DEL PONTE, E.

1939. Revisión de los "Oestridae" argentinos. *Physis*, Buenos Aires 17:525-534.

DESMAREST, E.

1859. Annelés, 312 pp., 279 figs., 40 pls., in J. C. Chenu: *Encyclopédie d'Histoire Naturelle*. Paris.

ENDERLEIN, G.

1909. *Cutiterebra schroederi*, eine neue parasitäre Fliege (Oestridae, olim). *Stettin. Ent. Ztg.* 70:202-203.

FONSECA, F.

1938. Observação de uma fase do ciclo evolutivo de *Cuterebra apicalis* Guérin (Diptera, Oestridae). *Bol. Biol., São Paulo (N.S.)* 3:166-167, fig. 1.
1940. Nova espécie de *Orienterebra* do Brasil (Dipt. Oestridae). *Rev. Entom., Rio de Janeiro* 11:662-671, 3 figs.
1941. Cuterebrineo novo parasito de Didelfídeo (Diptera, Oestridae). *Rev. Entom., Rio de Janeiro* 12:480-486, 1 fig., 2 pls.

FORATTINI, O. P. & K. LENKO

1959. Nota biológica sobre *Metacuterebra apicalis* (Guérin 1829/38) (Diptera, Cuterebridae). *Arq. Fac. Hig. Saúde Públ. Univ. S. Paulo* 13:155-158, 2 figs.



GOUDOT, J.

1845. Observations sur un diptère exotique dont la larve nuit aux boeufs (*Cuterebra noxialis*). *Ann. Sc. Nat.* 3:221-230.

GUÉRIN-MÉNEVILLE, F. E.

1835. Planches des animaux invertébrés, Insectes, 104 pls., in *Iconographie du règne animal de G. Cuvier*, vol. 2. Paris.

- (1844). Douzième ordre. Les Diptères: 531-559, in *Iconographie du règne animal de G. Cuvier*, vol. 3. Texte explicatif, Insectes, 576 pp., Paris.

GUIMARÃES, J. H. & N. PAPAVERO

1966. A tentative annotated bibliography of *Dermatobia hominis* (Linnaeus Jr., 1781) (Diptera, Cuterebridae). *Arq. Zool. Estado de São Paulo* 14:223-294.

GUIMARÃES, L. R. & M. CARRERA

1941. Contribuição ao conhecimento dos Cuterebrídeos do Brasil. *Arq. Zool. Estado de São Paulo* 3(1):1-12, 3 pls., 9 figs.

HENRIKSEN, K. L.

1942. A note upon two South American species of warbleflies (*Cuterebra apicalis* Guérin and *schmalzi* Lutz) parasites of rodents. *Ent. Meddel.* 22:284-289, 5 figs.

HOWSHIP, P.

1833. An account of two cases of inflammatory tumour produced by a deposit of the larva of a large fly (*Oestrus humanus*) beneath the cutis in the human subject; accompanied with drawings of the larva. *Proc. Royal Soc. London* 12:181.

KNAB, F.

1914. A new *Cuterebra* from Panama (Diptera). *Ins. Insc. Menstr.* 2(12):187-188.

LATREILLE, P. A.

1818. In *Nouveau dictionnaire d'histoire naturelle appliqué aux arts, principalement à l'agriculture et à l'économie rurale et domestique*. Nouvelle édition (2ème) 23:612 pp., 6 pls. Paris.

LUTZ, A.

1917. Contribuições ao conhecimento dos Oestrideos brasileiros. *Mem. Inst. Oswaldo Cruz* 9:94-113, pls. 27-29.

MACQUART, J.

1843. *Diptères exotiques nouveaux ou peu connus* 2(3):5-304, 36 pls. Paris.

NEIVA, A. & J. FLORENCIO GOMES

1917. Biologia da mosca do Berne (*Dermatobia hominis*) observada em todas as suas fases. *Ann. Paulistas Med. Cirur.* 8(9):197-209.

PALLAS, P. S.

1781. Mittheilungen über *Oestrus hominis*. *Neue nordische Beytr. Phys. Geogr. Erde y. Volk.* 1:157-158.

SHANNON, R. C. & C. T. GREENE

1926. A bot-fly parasitic in monkeys. *Zoopathologica*, New York 1:285-290, 2 pls., 2 figs.

SHANNON, R. C. & E. DEL PONTE

1926. Sinopsis parcial de los Muscoideos argentinos. *Rev. Inst. Bacteriol.* 5:1-44, 4 pls.

SWENK, M. H.

1905. The North American species of *Cuterebra*. *Jour. N. Y. Ent. Soc.* 13:181-185.

TOWNSEND, C. H. T.

1897. Description of the bot-fly of the cotton-tail rabbit in New Mexico, *Cuterebra lepusculi* n. sp. *Psyche* 8:8-9.

1919. New genera and species of muscoid flies. *Proc. U. S. Nat. Mus.* 56:541-592.

1927. Synopse dos generos museoideos da região humida tropical da America, com generos e especies novas. *Rev. Mus. Paulista* 15:205-385, 7 figs.

1931. New genera and species of American Oestromuseoid flies. *Rev. Entom., Rio de Janeiro* 1:313-354.

1935. *Manual of Myiology, in twelve parts* 2:1-289, pls. 1-9, Itaquaquecetuba.

1942. *Idem* 12:1-349, pls. 7-84; Addenda and corrigenda, 8 pp. Itaquaquecetuba.

WALKER, F.

1849. *List of the specimens et dipterous insects in the collection et the British Museum* 3:485-687. London.

WIEDEMANN, C. R. W.

1830. *Aussercuropäische zweiflügelige Insekten* 2: xii + 684 pp., 5 pls. Hamm.

ZELEDÓN, R., O. JIMÉNEZ Q. & R. R. BRENES M.

1957. *Cuterebra bacri* Shannon y Greene 1926 en el mono aullador de Costa Rica. *Rev. Biol. Trop.* 5:129-134, 11 figs.


INDEX


<i>albifrons</i> Swenk, 1905, <i>Cuterebra</i>	105.3
<i>almeidai</i> (Guimarães & Carrera), 1941 (<i>Baucuterebra</i>), <i>Cuterebra</i>	105.2
<i>Alouattamyia</i> Townsend, 1931	105.1
<i>analís</i> Macquart, 1843, <i>Cuterebra</i>	105.2
<i>apicalis</i> Guérin-Méneville, 1835, <i>Cuterebra</i>	105.2

<i>atrox</i> Clark, 1848, <i>Cuterebra</i>	105.2
<i>Atrypoderma</i> Townsend, 1919	105.2
<i>baeri</i> (Shannon & Greene), 1926 (<i>Cuterebra</i>), <i>Alouattamyia</i>	105.1
<i>Baucuterebra</i> Guimarães & Carrera, 1941	105.2
<i>baui</i> (Fonseca), 1940 (<i>Oriuterebra</i>), <i>Cuterebra</i>	105.2
<i>bicineta</i> Bau, 1931, <i>Cuterebra</i> , subsp. of <i>versicolora</i>	105.4
<i>Bogeria</i> Austen, 1895	105.2
<i>bonaerensis</i> (Del Ponte), 1939 (<i>Guerinioestrus</i>), <i>Rogenhoferia</i>	105.4
<i>bureni</i> Dalmat, 1942, <i>Montemyia</i>	105.5
<i>caayennensis</i> Macquart, 1843, <i>Cuterebra</i>	105.2
<i>eometes</i> Shannon & Del Ponte, 1926, <i>Cuterebra</i>	105.2
<i>conflans</i> Bau, 1929, <i>Cuterebra</i>	105.2
<i>Cuterebra</i> Clark, 1815	105.1
<i>Cuterebrini</i>	105.1
<i>Cutiterebra</i> , emend.	105.2
<i>cyaniventris</i> (Macquart), 1843 (<i>Cuterebra</i>), <i>Dermatobia</i>	105.5
<i>daguerrei</i> (Del Ponte), 1939 (<i>Guerinioestrus</i>), <i>Rogenhoferia</i>	105.5
<i>dasyopoda</i> Brauer, 1896, <i>Rogenhoferia</i>	105.5
<i>Dermatobia</i> Brauer, 1861	105.5
<i>Dermatobiini</i>	105.5
<i>detrudator</i> Clark, 1848, <i>Cuterebra</i>	105.2
<i>dorsalis</i> Bau, 1929, <i>Cuterebra</i>	105.3
<i>elegans</i> Bau, 1931, <i>Cuterebra</i> , subsp. of <i>rufiventris</i>	105.4
<i>ephippium</i> Latreille, 1818, <i>Cuterebra</i>	105.3
<i>flaviventris</i> (Bau), 1931 (<i>Metacuterebra</i>), <i>Cuterebra</i>	105.3
<i>funebis</i> (Austen, 1895 (<i>Cutiterebra</i>), <i>Cuterebra</i>	105.3
<i>gilvopilosa</i> Bau, 1932, <i>Rogenhoferia</i>	105.5
<i>grandis</i> (Guérin-Ménéville), 1844 (<i>Cephenemyia</i>), <i>Rogenhoferia</i> ..	105.5
<i>Guerinioestrus</i> Townsend, 1931	105.4
<i>hermanni</i> Bisehof, 1901, <i>Pseudogametes</i>	105.5
<i>hominis</i> (Linnaeus Jr.), 1781 (<i>Oestrus</i>), <i>Dermatobia</i>	105.5
<i>histrion</i> Coquillett, 1902, <i>Cuterebra</i>	105.3
<i>humanus</i> (Howship), 1833 (<i>Oestrus</i>), <i>Dermatobia</i>	105.5
<i>infulata</i> Lutz, 1917, <i>Cuterebra</i>	105.3
<i>lepida</i> Austen, 1933, <i>Cuterebra</i>	105.3
<i>lepusculi</i> Townsend, 1897, <i>Cuterebra</i>	105.3
<i>lutzi</i> Bau, 1930, <i>Cuterebra</i>	105.3
<i>maeulosa</i> Knab, 1914, <i>Cuterebra</i>	105.3
<i>megastoma</i> Brauer, 1863, <i>Cuterebra</i>	105.3
<i>Montemyia</i> Dalmat, 1942	105.5
<i>Metacuterebra</i> Bau, 1929	105.2
<i>noxiialis</i> (Goudot), 1845 (<i>Cuterebra</i>), <i>Dermatobia</i>	105.5



nigrieans Lutz, 1917, Cuterebra	105.3
nigricincta (Austen), 1895 (Cutiterebra), Cuterebra	105.3
obidosiana Bau, 1929, Cuterebra, subsp. of conflans	105.2
ornata Bau, 1928, Cuterebra	105.3
<i>Oricuterebra</i> Bau, 1931	105.2
<i>Orthocuterebra</i> Bau, 1929	105.2
<i>Paracuterebra</i> Bau, 1929	105.2
patagona Guérin-Méneville, 1844, Cuterebra	105.3
pessoai Guimarães & Carrera, 1941, Cuterebra	105.3
praegrandis Austen, 1933, Cuterebra	105.3
princeps (Austen), 1895 (Bogeria), Cuterebra	105.3
<i>Pseudobogeria</i> Bau, 1931	105.2
Pseudogametes Bischof, 1900	105.5
Pseudogametini	105.5
pulchra Bau, 1930, Cuterebra	105.3
pusilla Bau, 1931, Cuterebra	105.3
pygmaea Bau, 1931, Cuterebra	105.3
ribeiroi (Artigas & d'Andretta Jr.), 1951 (Alouattamyia), Cuterebra	105.4
Rogenhoferi Brauer, 1863	105.4
rubiginosa Bau, 1931, Cuterebra	105.4
rufiventris Macquart, 1843, Cuterebra	105.4
sarcophagoides Lutz, 1917, Cuterebra	105.4
schmalzi Lutz, 1917, Cuterebra	105.4
schroederi (Enderlein), 1909 (Cutiterebra), Cuterebra	105.4
semitra (Wiedemann), 1830 (Musca), Pseudogametes	105.5
semilutea Bau, 1929, Cuterebra	105.4
simulans Austen, 1933, Cuterebra	105.4
terrisona Walker, 1849, Cuterebra	105.4
townsendi (Fonseca), 1941 (Oricuterebra), Cuterebra	105.4
trigonocephala Lutz, 1917, Rogenhoferi (error)	105.5
trigonophora Brauer, 1863, Rogenhoferi	105.5
<i>Trypoderma</i> Wiedemann, 1830	105.2
versicolora Bau, 1930, Cuterebra	105.4
<i>Xylocopodes</i> Townsend, 1927	105.5
worontzowi Guimarães & Carrera, 1941, Cuterebra	105.4





SciELO

DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

106

FAMILY OESTRIDAE
(INCLUDING HYPODERMATIDAE)

J. H. GUIMARÃES

*Departamento de Zoologia, Secretaria da Agricultura
São Paulo, Brasil*

These are myiasis-producing flies, whose larvae are exclusively endoparasitic in mammals. For the classification of the group see Zumpt (1957) and for a general account see James (1948).

Subfamily OESTRINAE

The larvae live parasitically in the head and throat of various Artiodactyla and Perissodactyla. In the New World only one species occurs, whose larvae parasitize sheep and goats, and occasionally attacking man.

Genus OESTRUS Linnaeus

OESTRUS Linnaeus, 1758:584. Type-species, *ovis* Linnaeus (Curtis, 1826: pl. 106).

Cephalemyia Latreille, 1818:273. Type-species, *Oestrus ovis* Linnaeus (mon.).

ovis Linnaeus, 1758:585. Type-locality: "Sweden". Distr. — cosmopolitan.

Subfamily CEPHENEMYIINAE

Genus CEPHENEMYIA Latreille

CEPHENEMYIA Latreille, 1818:271. Type-species, *Oestrus trompe* Modeer (mon.).

Cephenomyia, emend.

Cephalemyia Curran, 1934:414 (error).

The larvae develop in the nasal and pharyngeal cavities of various Cervidae. For references to geographical distribution, redescriptions of adult and larvae, see Bennett & Sabrosky (1962).

PRATT Hunter, 1916:170. Type-locality: U.S.A., Texas. Distr. — U.S.A. (Oregon to Montana, south to Arizona), Mexico (north) and Texas.

106.1



Subfamily HYPODERMATINAE

Genus HYPODERMA Latreille

HYPODERMA Latreille, 1818:272. Type-species, *Oestrus bovis* Linnaeus (mon.).

Lithohypoderma Townsend, 1917:129. Type-species, *Musca ascarides* Scudder (orig. des.).

Subgenus HYPODERMA Latreille

The mature larvac develop in the skin of Artiodactyla.

BOVIS (Linnaeus), 1758:584 (*Oestrus*). Type-locality: "Europe". Distr. — Holarctic, northern Asia, north Africa. Neotropical region: Puerto Rico (from imported cattle), Jamaica (on imported cattle), Chile (on imported cattle; not established).

LINEATUM (Villers), 1789:349 (*Oestrus*). Type-locality: "Europe". Distr. — Holarctic, cosmopolitan. Neotropical Region: Dominican Republic, Puerto Rico, Chile (introduced, not established).

supplens Walker, 1849:685 (*Oestrus*). Type-locality: Nova Scotia.

bonassi Brauer, 1875:75. Type-locality: U.S.A. Colorado.

Subgenus OEDEMAGENA Latreille

OEDEMAGENA Latreille, 1818:272 (as genus). Type-species, *Oestrus tarandi* Linnaeus (mon.).

The mature larvac develop in the skin of reindeers.

TARANDI (Linnaeus), 1758:584 (*Oestrus*). Type-locality: "Sweden". Distr. — Arctic and subarctic regions ou Eurasia and America. In the Neotropical region it was recorded from Argentina, Tierra del Fuego, as parasite of imported reindeer (Del Ponte, 1939:533); not established.

terranoae (Knab), 1913:155 (*Oedemagena*). Type-locality: Canada, Newfoundland.

LITERATURE CITED

BENNETT, G. F. & C. W. SABROSKY

1962. The Nearctic species of the genus *Cephenemya* (Diptera, Oestridae). *Canad. Jour. Zool.* 40:431-448, figs. 1-7.

BRAUER, F.

1875. Beschreibung neuer und ungenügend bekannter Phryganiden und Oestriden. *Verh. zool.-bot. Ges. Wien* 25:69-78, 1 pl.



CURRAN, C. II.

1934. *The families and genera of North American Diptera*, 512 pp., 235 figs., 2 pls. New York.

CURTIS, J.

1826. *British Entomology: being illustrations and descriptions of the genera of insects found in Great Britain and Ireland* 3: pls. 99-146, London.

DEL PONTE, E.

1939. Revisión de los "Ocstridae" argentinos. *Physis* 17:525-534.

HUNTER, W. D.

1916. A new species of *Cephenemyia* from the United States (Diptera, Ocstridae). *Proc. Ent. Soc. Washington* 17:169-173, 1 pl.

JAMES, M. T.

- (1948). The flies that cause myiasis in man. *U. S. Dept. Agric. Misc. Publ.* (1947)631:175 pp., 98 figs.

KNAB, F.

1913. A new bot-fly from reindeer (Diptera: Museoidae). *Proc. Biol. Soc. Washington* 26:155-156.

LATREILLE, P. A.

1818. In: *Nouveau Dictionnaire d'Histoire Naturelle, appliqué aux arts, principalement à l'agriculture et à l'économie rurale et domestique* 4:575 pp., pls. B1-B12, B14-B15. Société de Naturalistes et d'Agriculteurs, Paris.

LINNAEUS, C.

1758. *Systema naturae per regna tria naturae*. Ed. 10. Vol. 1:824 pp., Holmiae.

TOWNSEND, C. H. T.

- (1917). *Lithohypoderma*, a new fossil genus of oestrids. *Ins. Insc. Menstr.* (1916) 4:128-130.

VILLERS, C. DE

1789. *Caroli Linnaci entomologia, faunae succicae descriptionibus aucta*. Vol. 3, 657 pp. 4 pls., Lugduni.

WALKER, F.

1849. *List of the specimens of dipterous insects in the collection of the British Museum* 3:485-687, London.

ZUMPT, F.

1957. Some remarks on the classification of the Ocstridae s. lat. (Diptera). *Jour. Ent. Soc. South Africa* 20:154-161, 1 pl.

INDEX


<i>bonassi</i> Brauer, 1875, Hypoderma	106.2
<i>bovis</i> (Linnaeus), 1758 (Oestrus), Hypoderma	106.2
<i>Cephaemyia</i> Curran, 1934 (error)	106.1


106.3



<i>Cephalemyia</i> Latreille, 1818	106.1
<i>Cephenemyia</i> Latreille, 1818	106.1
<i>Cephenemyiinae</i>	106.1
<i>Cephenomyia</i> , emend.	106.1
<i>Hypoderma</i> Latreille, 1818	106.2
<i>Hypodermatinae</i>	106.2
<i>lineatum</i> (Villers), 1789 (<i>Oestrus</i>), <i>Hypoderma</i>	106.2
<i>Lithohypoderma</i> Townsend, 1917	106.2
<i>Oedemagena</i> Latreille, 1818	106.2
<i>Oestrinae</i>	106.1
<i>Oestrus</i> Linnaeus, 1758	106.1
<i>ovis</i> Linnaeus, 1758, <i>Oestrus</i>	106.1
<i>pratti</i> Hunter, 1916, <i>Cephenemyia</i>	106.1
<i>supplens</i> (Walker), 1849 (<i>Oestrus</i>), <i>Hypoderma</i>	106.2
<i>tarandi</i> (Linnaeus), 1758 (<i>Oestrus</i>), <i>Oedemagena</i>	106.2
<i>terranae</i> (Knab), 1913 (<i>Oedemagena</i>), <i>Hypoderma</i>	106.2







SciELO

